



**HAL**  
open science

**Persistence exponents via perturbation theory:  
MA(1)-processes**

Frank Aurzada, Dieter Bothe, Pierre-Étienne Druet, Marvin Kettner,  
Christophe Profeta

► **To cite this version:**

Frank Aurzada, Dieter Bothe, Pierre-Étienne Druet, Marvin Kettner, Christophe Profeta. Persistence exponents via perturbation theory: MA(1)-processes. 2024. hal-04644436

**HAL Id: hal-04644436**

**<https://hal.science/hal-04644436v1>**

Preprint submitted on 11 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Persistence exponents via perturbation theory: MA(1)-processes

Frank Aurzada<sup>1</sup>    Dieter Bothe<sup>1</sup>    Pierre-Étienne Druet<sup>1</sup>  
Marvin Kettner<sup>1</sup>    Christophe Profeta<sup>2</sup>

July 11, 2024

## Abstract

For the moving average process  $X_n = \rho\xi_{n-1} + \xi_n$ ,  $n \in \mathbb{N}$ , where  $\rho \in \mathbb{R}$  and  $(\xi_i)_{i \geq -1}$  is an i.i.d. sequence of normally distributed random variables, we study the persistence probabilities  $\mathbb{P}(X_0 \geq 0, \dots, X_N \geq 0)$ , for  $N \rightarrow \infty$ . We exploit that the exponential decay rate  $\lambda_\rho$  of that quantity, called the persistence exponent, is given by the leading eigenvalue of a concrete integral operator. This makes it possible to study the problem with purely functional analytic methods. In particular, using methods from perturbation theory, we show that the persistence exponent  $\lambda_\rho$  can be expressed as a power series in  $\rho$ . Finally, we consider the persistence problem for the Slepian process, transform it into the moving average setup, and show that our perturbation results are applicable.

*Keywords:* moving average process; eigenvalue problem; integral equation; persistence; perturbation theory; Slepian process

*2020 Mathematics Subject Classification:* Primary 45C05; Secondary 47A55, 33C45, 60J05, 60G15

---

<sup>1</sup>Technical University of Darmstadt, Schlossgartenstraße 7, 64287 Darmstadt, Germany. E-mail: aurzada@mathematik.tu-darmstadt.de, bothe@mma.tu-darmstadt.de, druet@mma.tu-darmstadt.de, marvinkettner@web.de

<sup>2</sup>Université Paris-Saclay, CNRS, Univ Evry, Laboratoire de Mathématiques et Modélisation d'Evry, 91037, Evry-Courcouronnes, France. E-mail: christophe.profeta@univ-evry.fr

# 1 Introduction

## 1.1 Persistence probabilities for moving average processes

Let  $(\xi_i)_{i \geq -1}$  be a sequence of i.i.d. random variables and  $\rho \in \mathbb{R}$ . A moving average process of order one (MA(1)-process) is given by

$$X_n := \rho \xi_{n-1} + \xi_n, \quad \text{for } n \in \mathbb{N}.$$

Throughout the paper, we consider standard normally distributed random variables  $(\xi_i)_i$ , i.e.  $\xi_0$  has the density  $\phi(x) := (2\pi)^{-1/2} \exp(-x^2/2)$ ,  $x \in \mathbb{R}$ . We are interested in the persistence probabilities

$$\mathbb{P}(X_0 \geq 0, \dots, X_N \geq 0), \quad \text{for } N \rightarrow \infty, \quad (1)$$

and in particular in the exponential rate of decay of this quantity, which is called persistence exponent. Non-exit probabilities are a classic and fundamental topic in probability with numerous applications in finance, insurance, queueing, and other subjects. The question is also studied intensively in theoretical physics; there, the rationale is that the persistence exponent is a measure of how fast the underlying physical system returns to equilibrium. We refer to the survey [8] and the monograph [22] for an overview on the relevance of the question to physical systems and to [6] for a survey of the mathematical literature.

The persistence problem for moving average processes was studied before in [19, 17, 5]. For MA-processes, the persistence question can be rewritten as a non-exit problem for a two-dimensional Markov chain (see e.g. [5, Section 2.1]). It is well-known that non-exit probabilities of Markov chains have close relations to eigenvalues of operators (see [3, 5, 13], also see [11, 9, 21, 27, 28] for the quasi-stationary approach). The purpose of this paper is establish and to use the connection between the persistence exponent  $\lambda_\rho$ , i.e. the exponential decay rate of (1), and the leading eigenvalue of a suitable operator. Then, powerful tools from functional analysis can be applied. In particular, methods of perturbation theory in the spirit of [14] can be used to obtain a series representation of the eigenvalue in the parameter  $\rho$ . This ansatz was previously used in [4] for autoregressive processes.

## 1.2 The eigenvalue problem

In the recent work [5, Section 2.1] it is shown that for  $\rho \neq -1$  the persistence probability (1) of the MA(1)-process decays exponentially fast and that the

exponential decay rate, i.e. the persistence exponent, is the leading eigenvalue of the following explicit integral operator:

$$S_\rho: \mathcal{B}(\mathbb{R}) \rightarrow \mathcal{B}(\mathbb{R}), \quad S_\rho f(x) := \int_{-\rho x}^{\infty} f(y)\phi(y) dy, \quad (2)$$

where  $\mathcal{B}(\mathbb{R})$  is the space of bounded measurable real-valued functions on  $\mathbb{R}$ . Bearing this connection in mind, the purpose of this paper is to study the largest solution  $\lambda = \lambda_\rho$  of the eigenvalue equation

$$\lambda f(x) = \int_{-\rho x}^{\infty} f(y)\phi(y) dy, \quad x \in \mathbb{R}, f \in \mathcal{B}(\mathbb{R}). \quad (3)$$

The approach taken in this paper is to show that a modification of the integral operator  $S_\rho$  can be represented as a power series in  $\rho$  if we consider the operator on a suitable space of functions. The definition of this function space (see Section 2) is motivated by the following observation:

Assume that  $f$  is analytic, that is  $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$ , for  $x \in \mathbb{R}$ . Further, if we assume that  $\lim_{x \rightarrow \infty} (f\phi)^{(n-1)}(x) = 0$  for all  $n \geq 1$ , we can write  $(f\phi)^{(n-1)}(0) = (-1) \int_0^{\infty} (f\phi)^{(n)}(y) dy$ . Then,

$$\begin{aligned} S_\rho f(x) &= \int_{-\rho x}^{\infty} f(y)\phi(y) dy \\ &= \int_0^{\infty} f(y)\phi(y) dy + (-1) \int_0^{-\rho x} f(y)\phi(y) dy \\ &= \int_0^{\infty} f(y)\phi(y) dy + \sum_{n=1}^{\infty} \rho^n (-1)^{n+1} x^n \frac{(f\phi)^{(n-1)}(0)}{n!} \\ &= \int_0^{\infty} f(y)\phi(y) dy + \sum_{n=1}^{\infty} \rho^n (-1)^n x^n \frac{1}{n!} \int_0^{\infty} (f\phi)^{(n)}(y) dy \\ &= \sum_{n=0}^{\infty} \rho^n (-1)^n x^n \frac{1}{n!} \int_0^{\infty} (f\phi)^{(n)}(y) dy. \end{aligned} \quad (4)$$

Hence, under the above conditions on  $f$ , the expression  $S_\rho f(x)$  can be written as a power series in  $\rho$ .

With this observation in mind, we will consider a specific space of analytic functions such that we obtain a well-defined holomorphic operator (cf. Theorem 1). From this, by using perturbation techniques in the spirit of [14], we can conclude that the leading eigenvalue and the corresponding eigenfunction are holomorphic in  $\rho$ , too. This is the contents of our main result,

Theorem 2. In other words, the persistence exponent and the eigenfunction admit a power series representation in  $\rho$ , respectively. Additionally, we have iterative formulas for the coefficients of the power series representation of the persistence exponent and the eigenfunction, respectively; and we compute the first coefficients, cf. Theorem 3.

### 1.3 Application to the Slepian process

As a further application, we look at the persistence problem for the so-called Slepian process. Let  $(B_t)_{t \geq 0}$  be a standard Brownian motion and define the Slepian process by

$$S_t := B_{t+1} - B_t, \quad t \geq 0. \quad (5)$$

In his seminal paper [26], D. Slepian computed the distribution of the supremum of  $S$  on  $[0, 1]$  and found:

$$\mathbb{P} \left( \sup_{u \in [0,1]} S_u \leq a \right) = \Phi(a)^2 - \phi(a) (a\Phi(a) + \phi(a)), \quad (6)$$

where again  $\phi$  is the standard normal density and  $\Phi$  is the corresponding cumulative distribution function. The general formula for the distribution of  $\sup_{u \in [0,t]} S_u$  for any  $t \geq 0$  was then obtained by Shepp in [25]. Shepp leaves it as an open question to study of the asymptotics

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P} \left( \sup_{u \in [0,N]} S_u \leq a \right), \quad (7)$$

because his formulas, which involve iterated integrals, are not well-suited for such computations. The existence of this limit was then obtained by Li & Shao in [18], and numerical computations have been proposed recently by Noonan & Zhigljavsky [23]. The contribution of this paper is the observation that one can rewrite the persistence problem for the Slepian process as a persistence problem for an MA(1)-process. Further, we can show that our perturbation results are applicable.

The outline of this paper is as follows. In Section 2, our main results are stated. In particular, we show that the leading eigenvalue  $\lambda_\rho$  of (3) can be expanded into a power series in  $\rho$  and we discuss the coefficients of this power series. In Section 3, we consider the persistence problem for the Slepian process, transform it into the MA setup, and show that our main results are applicable. Section 4 is devoted to the proofs of the main theorems. Finally, in Section 5 we deal with the radius of convergence of the series for  $\lambda_\rho$ .

## 2 Main results

Let  $\gamma$  be the standard Gaussian measure on  $\mathbb{R}$  and let  $h_n$  denote the  $n$ -th Hermite polynomial given by

$$h_n(x) := (-1)^n e^{\frac{x^2}{2}} \frac{d^n}{dx^n} e^{-\frac{x^2}{2}}.$$

For the facts on Hermite polynomials that we use in this paper, we refer the reader to [2, Chapter 6]. Further we set  $\widehat{h}_n(x) := (n!)^{-1/2} h_n(x)$ . Here, the normalization is chosen such that  $\|\widehat{h}_n\|_{L^2(\mathbb{R}, \gamma)} = 1$ .

Fix  $0 < q < 1$  and let  $(a_n)_{n \in \mathbb{N}} \subseteq \mathbb{C}$  be a sequence such that  $\sum_{n=0}^{\infty} |a_n|^2 q^{-n} < \infty$ . By [1, Theorem 2], it holds that  $\sum_{n=0}^{\infty} |a_n \widehat{h}_n(x)|$  converges uniformly on compact subsets of  $\mathbb{R}$ . Hence, we can define an analytic function  $f: \mathbb{R} \rightarrow \mathbb{C}$  via  $f(x) := \sum_{n=0}^{\infty} a_n \widehat{h}_n(x)$ . In particular,  $\Re(f)$  and  $\Im(f)$  are analytic, where  $\Re(f)$  and  $\Im(f)$  are the real and the imaginary part of  $f$ , respectively. Let

$$\mathcal{H}_q := \left\{ f: \mathbb{R} \rightarrow \mathbb{C}, f(x) = \sum_{n=0}^{\infty} a_n \widehat{h}_n(x) \text{ with } \sum_{n=0}^{\infty} |a_n|^2 q^{-n} < \infty \right\}$$

and set

$$\langle f, g \rangle_{\mathcal{H}_q} := \sum_{n=0}^{\infty} a_n \overline{b_n} q^{-n} \text{ for } f = \sum_{n=0}^{\infty} a_n \widehat{h}_n, g = \sum_{n=0}^{\infty} b_n \widehat{h}_n.$$

We note that  $(\mathcal{H}_q, \langle \cdot, \cdot \rangle_{\mathcal{H}_q})$  is a Hilbert space of functions [1, Proposition 1]. In fact, we will see that it is a reproducing kernel Hilbert space and we will exploit this structure for the proofs. Note that we consider a complex Hilbert space instead of a real one, since a complex space is necessary for applying the powerful methods of perturbation theory in the spirit of [14].

We set

$$T_\rho: \mathcal{H}_q \rightarrow \mathcal{H}_q, \quad T_\rho f(x) := \int_{-\rho x}^{\infty} f(y) \phi(y) dy.$$

Note that this is the version of the operator  $S_\rho$  on the space  $\mathcal{H}_q$ , cf. (2), in the sense that  $S_\rho$  acts on bounded real valued functions, while  $T_\rho$  acts on complex valued functions in  $\mathcal{H}_q$ . Here, we set, for a complex valued function  $f: \int_{-\rho x}^{\infty} f(y) \phi(y) dy = \int_{-\rho x}^{\infty} \Re(f)(y) \phi(y) dy + i \int_{-\rho x}^{\infty} \Im(f)(y) \phi(y) dy$ . Our first main result states that  $T_\rho$  is a holomorphic operator.

**Theorem 1.** Fix  $0 < q < 1$ . Let  $-\sqrt{\frac{1-q}{1+q^{-1}}} < \rho < \sqrt{\frac{1-q}{1+q^{-1}}}$  and define for  $n \in \mathbb{N}$  the integral operator

$$T^{(n)}: \mathcal{H}_q \rightarrow \mathcal{H}_q, \quad T^{(n)} f(x) := (-1)^n x^n \frac{1}{n!} \int_0^{\infty} (f\phi)^{(n)}(y) dy.$$

The operator  $T_\rho$  is well-defined, bounded, compact and admits the representation

$$T_\rho = \sum_{n=0}^{\infty} \rho^n T^{(n)}. \quad (8)$$

**Remark.** To optimize the radius of convergence of the power series for  $T_\rho$ , the best choice of  $0 < q < 1$  for the Hilbert space  $\mathcal{H}_q$  is  $q^* := \sqrt{2} - 1$ . Then,  $T_\rho$  can be represented as a power series for  $-(\sqrt{2} - 1) < \rho < \sqrt{2} - 1$ .

Let

$$r(T_\rho) := \sup\{|\lambda| : \lambda \in \Sigma(T_\rho)\}$$

be the spectral radius of  $T_\rho$ , where  $\Sigma(T_\rho)$  denotes the spectrum of  $T_\rho$ . Our second main result deals with the leading eigenvalue  $\lambda_\rho$  of (3) on  $\mathcal{H}_q$ , i.e. the persistence exponent for moving average processes, and states that  $\lambda_\rho$  can be expanded into a power series.

**Theorem 2.** For  $-\sqrt{\frac{1-q}{1+q^{-1}}} < \rho < \sqrt{\frac{1-q}{1+q^{-1}}}$  we have

$$\mathbb{P}(X_0 \geq 0, \dots, X_N \geq 0) = \lambda_\rho^{N+o(N)},$$

where  $\lambda_\rho := r(T_\rho) \in (0, 1)$  is the largest eigenvalue of  $T_\rho$ . The corresponding eigenfunction  $f_\rho$  is non-negative, i.e.  $f_\rho(x) \geq 0$  for all  $x \in \mathbb{R}$ .

There are numbers  $K_n \in \mathbb{R}$  such that the quantity  $\lambda_\rho$  admits the representation

$$\lambda_\rho = \sum_{n=0}^{\infty} \rho^n K_n,$$

for all  $|\rho| < r_0$ , where  $r_0 > 0.332$ .

As an application, we are going to see in Section 3 that one can transform the persistence problem for the Slepian process into the setup of persistence of MA(1)-processes and that this case can be covered by Theorem 2, cf. Proposition 4.

As the third and last main result, we determine the coefficients  $K_n$ ,  $n \in \mathbb{N}$ , of the power series of the persistence exponent  $\lambda_\rho$ .

By Theorem 1 the operator  $T_\rho$  and by Theorem 2 the eigenvalue  $\lambda_\rho$  can be expressed as a power series in  $\rho$ , respectively. Additionally, the corresponding eigenfunction  $f_\rho$  can be expressed as a power series in  $\rho$  (see [4, Theorem 4]). Let us write

$$T_\rho = \sum_{k=0}^{\infty} \rho^k T^{(k)}, \quad \lambda_\rho = \sum_{k=0}^{\infty} \rho^k K_k, \quad f_\rho = \sum_{m=0}^{\infty} \rho^m g_m.$$

Note that for  $\rho = 0$  the MA(1)-process is a sequence of i.i.d. random variables, so that

$$K_0 = \lambda_0 = \mathbb{P}(\xi_0 \geq 0) = \frac{1}{2}.$$

**Theorem 3.** *For all  $m \in \mathbb{N}$  the function  $g_m$  is a polynomial of degree at most  $m$ . A full iterative description of the  $g_m$ ,  $m \in \mathbb{N}$ , is given by the equations  $g_0 = \mathbb{1}$ ,  $K_0 = \frac{1}{2}$ ,  $g_m(0) = 0$  for  $m \geq 1$ , and*

$$g_m = \frac{1}{K_0} \left( \sum_{j=1}^m T^{(j)} g_{m-j} - \sum_{j=1}^{m-1} T^{(0)} g_j \cdot g_{m-j} \right), \quad m \geq 1. \quad (9)$$

Further, the  $K_n$ ,  $n \geq 1$ , can be computed using

$$K_n = T^{(0)} g_n. \quad (10)$$

The first coefficients are given by

$$\begin{aligned} K_0 &= \frac{1}{2}, \\ K_1 &= \frac{1}{\pi}, \\ K_2 &= -\frac{2}{\pi^2}, \\ K_3 &= -\frac{5}{6\pi} + \frac{8}{\pi^3}, \\ K_4 &= \frac{13}{3\pi^2} - \frac{40}{\pi^4}, \\ K_5 &= \frac{23}{40\pi} - \frac{28}{\pi^3} + \frac{224}{\pi^5}, \\ K_6 &= -\frac{1069}{180\pi^2} + \frac{580}{3\pi^4} - \frac{1344}{\pi^6}, \\ K_7 &= -\frac{37}{112\pi} + \frac{842}{15\pi^3} - \frac{4144}{3\pi^5} + \frac{8448}{\pi^7}, \\ K_8 &= \frac{943}{168\pi^2} - \frac{1535}{3\pi^4} + \frac{10080}{\pi^6} - \frac{54912}{\pi^8}. \end{aligned}$$

It would be very interesting to obtain a closed-form expression for the coefficients ( $K_n$ ). We suspect that the last term of each  $K_n$ , respectively, is given by  $\tau_n := (-1)^{n-1} 2^{n-1} \binom{2(n-1)}{n-1} \frac{1}{n} \pi^{-n}$ . The second to last term of the  $K_n$ , respectively, seems to be of the form  $-\tau_{n-2} \frac{8(n-3)+5}{6}$ . Obtaining more terms seems complicated.



### 3 Slepian process

In this section, we consider the persistence problem for the Slepian process, show how this can be transformed into a persistence question for an MA(1)-process, and prove that the above main results can be applied.

Recall that the Slepian process  $(S_t)$  was defined in (5). Let us denote by

$$F_N(a) := \mathbb{P} \left( \sup_{u \in [0, N]} S_u \leq a \right)$$

the persistence probability for the Slepian process, where  $a \in \mathbb{R}$  and  $N \in \mathbb{N}$ .

**Proposition 4.** *The persistence probability  $F_N(a)$  may be written as*

$$F_N(a) = \mathbb{P}(X_0 \leq b, \dots, X_{N-1} \leq b)$$

where  $(X_n)_{n \geq 1}$  is a MA(1)-process with standard normally distributed random variables and with parameter

$$\hat{\rho} = \frac{\sqrt{1 - 4 \cos^2(2\pi F_2(\hat{a}))} - 1}{2 \cos(2\pi F_2(\hat{a}))} \simeq 0.3186,$$

where  $\hat{a} = F_1^{-1}(1/2)$  and  $b = \Phi^{-1}(F_1(a))\sqrt{1 + \hat{\rho}^2}$ .

When  $b = 0$ , i.e.  $a = \hat{a}$ , Theorem 2 shows that the desired exponential decay rate of the persistence probability of the Slepian process can be expressed as a power series since  $\hat{\rho} < 0.332$ . For arbitrary  $a \in \mathbb{R}$  a shifted version of  $T_{\hat{\rho}}$  needs to be considered. Presumably, as in Theorem 1 and Theorem 2, a power series representation can be obtained.

**Remark.** *The value of  $F_1$  was originally computed by [26] and is given in (6). Similarly, following the computation of [23], the value of  $F_2(\hat{a})$  is given by*

$$\begin{aligned} F_2(\hat{a}) &= \Phi(\hat{a}) - \Phi^3(\hat{a}) - \frac{1}{\sqrt{2}} \int_0^\infty \Phi(\hat{a} - y) \varphi(\sqrt{2}y) \left( \Phi(\sqrt{2}y) - \frac{1}{2} \right) dy \\ &\quad + \frac{\varphi^2(\hat{a})}{2} (\Phi(\hat{a})(\hat{a}^2 + 1) + \hat{a}\varphi(\hat{a})) + \int_0^\infty \Phi^2(\hat{a} - y) \varphi(y + \hat{a}) dy. \end{aligned}$$

*Proof of Proposition 4.* We start by writing the decomposition:

$$F_N(a) = \mathbb{P}(M_0 \leq a, \dots, M_{N-1} \leq a),$$

where the random variables  $(M_i)_{i \geq 0}$  are defined by

$$M_i := \sup_{u \in [0,1]} S_{u+i}.$$

Using the stationarity and the independence of the increments of Brownian motion, it is clear that the random variables  $(M_i)$  are identically distributed with common distribution  $F_1$ , and are such that for any  $i$  and  $j$  with  $|i-j| \geq 2$ ,  $M_i$  and  $M_j$  are independent. Define next the centered Gaussian random variables

$$Z_i := \Phi^{-1}(F_1(M_i)), \quad i \geq 0,$$

and observe that the sequence  $(Z_i)$  is stationary, with covariance matrix  $V$  given by

$$V = \begin{pmatrix} 1 & s & 0 & 0 & \dots \\ s & 1 & s & 0 & \dots \\ 0 & s & 1 & s & \dots \\ 0 & 0 & s & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \quad \text{where } s = \mathbb{E}[Z_0 Z_1].$$

Setting  $b_0 := \Phi^{-1}(F_1(a))$ , we are thus led to compute

$$F_N(a) = \mathbb{P}(Z_0 \leq b_0, Z_1 \leq b_0, \dots, Z_{N-1} \leq b_0), \quad (11)$$

where the density of the Gaussian vector  $(Z_0, \dots, Z_{N-1})$  depends only on  $s$ . To compute the value of  $s$ , observe that taking  $N = 2$  and  $b_0 = 0$ , we have

$$\begin{aligned} F_2(\hat{a}) &= \mathbb{P}(Z_0 \leq 0, Z_1 \leq 0) \\ &= \frac{\sqrt{1-s^2}}{2\pi} \int_{-\infty}^0 \int_{-\infty}^0 \exp\left(-\frac{1}{2}(x^2 + y^2 - 2sxy)\right) dx dy \\ &= \frac{1}{4} + \frac{1}{2\pi} \arctan\left(\frac{s}{\sqrt{1-s^2}}\right). \end{aligned}$$

Inverting this relation yields the value of  $s$  (note that  $2\pi F_2(\hat{a}) \in [\frac{\pi}{2}, \pi]$  so that we can use the last formula):

$$s = \frac{\tan\left(2\pi F_2(\hat{a}) - \frac{\pi}{2}\right)}{\sqrt{1 + \tan^2\left(2\pi F_2(\hat{a}) - \frac{\pi}{2}\right)}} = -\cos\left(2\pi F_2(\hat{a})\right).$$

Finally, let  $(\xi_i)_{i \geq -1}$  be an i.i.d. sequence of random variables with distribution  $\xi_0 \sim \mathcal{N}(0, 1)$  and set

$$\hat{\rho} := \frac{1 - \sqrt{1 - 4s^2}}{2s} \quad \text{so that} \quad s = \frac{\hat{\rho}}{1 + \hat{\rho}^2}.$$

Therefore, the sequence  $(Z_i)$  has the same distribution as the sequence

$$\frac{\widehat{\rho}\xi_{i-1} + \xi_i}{\sqrt{1 + \widehat{\rho}^2}}, \quad i \geq 0;$$

and going back to (11), we obtain

$$F_N(a) = \mathbb{P} \left( \widehat{\rho}\xi_{-1} + \xi_0 \leq b_0\sqrt{1 + \widehat{\rho}^2}, \dots, \widehat{\rho}\xi_{N-2} + \xi_{N-1} \leq b_0\sqrt{1 + \widehat{\rho}^2} \right).$$

This is exactly the statement of Proposition 4 after setting  $X_i := \widehat{\rho}\xi_{i-1} + \xi_i$  for  $i \in \mathbb{N}$ .  $\square$

## 4 Proofs of the main theorems

The following lemma makes it legitimate to use the computation (4) and provides a helpful representation of the inner product and the norm on  $\mathcal{H}_q$ .

**Lemma 5.** *Let  $f, g \in \mathcal{H}_q$ . It holds that*

- (a)  $\|f^{(k)}\|_{L^2(\mathbb{R}, \gamma)} < \infty$ , for all  $k \in \mathbb{N}$ ,
- (b)  $\lim_{x \rightarrow \infty} (f\phi)^{(n-1)}(x) = 0$ , for all  $n \geq 1$ ,
- (c)  $\langle f, g \rangle_{\mathcal{H}_q} = \sum_{k=0}^{\infty} \frac{(q^{-1}-1)^k}{k!} \langle f^{(k)}, g^{(k)} \rangle_{L^2(\mathbb{R}, \gamma)}$ ,
- (d)  $\|f\|_{\mathcal{H}_q}^2 = \sum_{k=0}^{\infty} \frac{(q^{-1}-1)^k}{k!} \|f^{(k)}\|_{L^2(\mathbb{R}, \gamma)}^2$ .

*Proof.* (a) Let  $f \in \mathcal{H}_q$ , i.e. we suppose that  $f(x) = \sum_{n=0}^{\infty} a_n \widehat{h}_n(x)$  for  $x \in \mathbb{R}$ , with  $\sum_{n=0}^{\infty} |a_n|^2 q^{-n} < \infty$ . Note that  $\widehat{h}_n^{(k)}(x) = \sqrt{\frac{n!}{(n-k)!}} \widehat{h}_{n-k}(x)$  for  $k \leq n$  (see e.g. [2, Section 6.1]). Thus, the derivatives of  $f$  are given by

$$f^{(k)}(x) = \sum_{n=0}^{\infty} a_n \widehat{h}_n^{(k)}(x) = \sum_{n=0}^{\infty} a_{n+k} \sqrt{\frac{(n+k)!}{n!}} \widehat{h}_n(x).$$

Recall that  $(\widehat{h}_n)_n$  is an orthonormal basis for  $L^2(\mathbb{R}, \gamma)$ . By Parseval's identity we get

$$\|f^{(k)}\|_{L^2(\mathbb{R}, \gamma)}^2 = \sum_{n=0}^{\infty} \left( |a_{n+k}| \sqrt{\frac{(n+k)!}{n!}} \right)^2 \leq \sum_{n=0}^{\infty} |a_{n+k}|^2 (n+k)^k < \infty.$$

(b) Note that  $(f\phi)^{(n-1)} = (\Re(f)\phi)^{(n-1)} + i(\Im(f)\phi)^{(n-1)}$ . We have

$$\begin{aligned}
\int_0^\infty |(\Re(f)\phi)^{(n-1)}(x)| dx &= \int_0^\infty \left| \sum_{k=0}^{n-1} \binom{n-1}{k} \Re(f)^{(k)}(x) \phi^{(n-1-k)}(x) \right| dx \\
&= \int_0^\infty \left| \sum_{k=0}^{n-1} \binom{n-1}{k} \Re(f)^{(k)}(x) h_{n-1-k}(x) \right| d\gamma(x) \\
&\leq \sum_{k=0}^{n-1} \binom{n-1}{k} \int_{\mathbb{R}} |\Re(f)^{(k)}(x) h_{n-1-k}(x)| d\gamma(x) \\
&\leq \sum_{k=0}^{n-1} \binom{n-1}{k} \|\Re(f)^{(k)}\|_{L^2(\mathbb{R}, \gamma)} \|h_{n-1-k}\|_{L^2(\mathbb{R}, \gamma)} \\
&< \infty,
\end{aligned}$$

by using Hölder's inequality in the last but one step and statement (a) in the last step. Therefore, if  $\lim_{x \rightarrow \infty} (\Re(f)\phi)^{(n-1)}(x)$  exists, then the limit must be zero. The limit exists since

$$\begin{aligned}
(\Re(f)\phi)^{(n-1)}(x) &= \int_0^x (\Re(f)\phi)^{(n)}(y) dy + (\Re(f)\phi)^{(n-1)}(0) \\
&\xrightarrow{x \rightarrow \infty} \int_0^\infty (\Re(f)\phi)^{(n)}(y) dy + (\Re(f)\phi)^{(n-1)}(0).
\end{aligned}$$

We conclude similarly that  $\lim_{x \rightarrow \infty} (\Im(f)\phi)^{(n-1)}(x) = 0$ . Hence, the assertion follows.

(c) Let  $f(x) = \sum_{n=0}^\infty a_n \widehat{h}_n(x)$ ,  $g(x) = \sum_{n=0}^\infty b_n \widehat{h}_n(x)$  for  $x \in \mathbb{R}$ . As in the proof of statement (a) we have  $f^{(k)}(x) = \sum_{n=k}^\infty a_n \sqrt{\frac{n!}{(n-k)!}} \widehat{h}_{n-k}(x)$  and  $g^{(k)}(x) = \sum_{n=k}^\infty b_n \sqrt{\frac{n!}{(n-k)!}} \widehat{h}_{n-k}(x)$  for  $x \in \mathbb{R}$  and  $k \in \mathbb{N}$ . Using that  $(\widehat{h}_n)_n$  is an orthonormal basis for  $L^2(\mathbb{R}, \gamma)$ , we compute

$$\begin{aligned}
\sum_{k=0}^\infty \frac{(q^{-1} - 1)^k}{k!} \langle f^{(k)}, g^{(k)} \rangle_{L^2(\mathbb{R}, \gamma)} &= \sum_{k=0}^\infty \frac{(q^{-1} - 1)^k}{k!} \sum_{n=k}^\infty \frac{n!}{(n-k)!} a_n \overline{b_n} \\
&= \sum_{n=0}^\infty a_n \overline{b_n} \sum_{k=0}^n \binom{n}{k} (q^{-1} - 1)^k \\
&= \sum_{n=0}^\infty a_n \overline{b_n} (q^{-1} - 1 + 1)^n \\
&= \langle f, g \rangle_{\mathcal{H}_q}.
\end{aligned}$$

(d) This statement follows directly from (c).  $\square$

Combining Lemma 5(b) with the fact that functions of  $\mathcal{H}_q$  are analytic, it holds that, as in (4),

$$T_\rho f(x) = \sum_{n=0}^{\infty} \rho^n (-1)^n x^n \frac{1}{n!} \int_0^{\infty} (f\phi)^{(n)}(y) dy$$

for all  $f \in \mathcal{H}_q$  and  $x \in \mathbb{R}$ , i.e. (8) holds.

*Proof of Theorem 1.* We want to compute an upper bound for the operator norm of  $T^{(n)}$  for  $n \in \mathbb{N}$ , which simultaneously shows that these operators are well-defined. Let  $m_n(x) := x^n$ , for  $x \in \mathbb{R}$ . Note that

$$\begin{aligned} \|T^{(n)} f\|_{\mathcal{H}_q} &= \|(-1)^n \frac{1}{n!} \int_0^{\infty} (f\phi)^{(n)}(y) dy \cdot m_n\|_{\mathcal{H}_q} \\ &= \left| \frac{1}{n!} \int_0^{\infty} (f\phi)^{(n)}(y) dy \right| \|m_n\|_{\mathcal{H}_q}. \end{aligned}$$

We recall the inverse explicit formula for the Hermite polynomials (see e.g. [24, Section 2]):

$$m_n = n! \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \frac{h_{n-2j}}{2^j j! (n-2j)!} = \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n!}{2^j j! \sqrt{(n-2j)!}} \widehat{h}_{n-2j}.$$

On one hand, by using the inequality  $(2j)! \leq 2^{2j} j!^2$  for  $j \in \mathbb{N}$ , we have

$$\begin{aligned} \|m_n\|_{\mathcal{H}_q}^2 &= \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \left( \frac{n!}{2^j j! \sqrt{(n-2j)!}} \right)^2 q^{-(n-2j)} \\ &= n! \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n!}{2^{2j} j!^2 (n-2j)!} q^{-(n-2j)} \\ &\leq n! \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n!}{(2j)! (n-2j)!} q^{-(n-2j)} \\ &= n! \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2j} q^{-(n-2j)} \end{aligned}$$

$$\begin{aligned}
&\leq n! \sum_{j=0}^n \binom{n}{j} q^{-(n-j)} \\
&= n!(1 + q^{-1})^n.
\end{aligned}$$

Hence,

$$\|m_n\|_{\mathcal{H}_q} \leq \sqrt{n!} \cdot (1 + q^{-1})^{n/2}.$$

On the other hand, we get, by using Hölder's inequality and Lemma 5(d),

$$\begin{aligned}
&\left| \frac{1}{n!} \int_0^\infty (f\phi)^{(n)}(y) \, dy \right| \\
&\leq \frac{1}{n!} \int_0^\infty \sum_{k=0}^n \binom{n}{k} |f^{(k)}(y)\phi^{(n-k)}(y)| \, dy \\
&= \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} \int_0^\infty |f^{(k)}(y)h_{n-k}(y)\phi(y)| \, dy \\
&\leq \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} \int_{\mathbb{R}} |f^{(k)}(y)h_{n-k}(y)| \, d\gamma(y) \\
&\leq \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} \|f^{(k)}\|_{L^2(\mathbb{R},\gamma)} \cdot \|h_{n-k}\|_{L^2(\mathbb{R},\gamma)} \\
&= \sum_{k=0}^n \frac{1}{k!(n-k)!} \|f^{(k)}\|_{L^2(\mathbb{R},\gamma)} \sqrt{(n-k)!} \\
&= \sum_{k=0}^n \frac{(q^{-1}-1)^{k/2}}{\sqrt{k!}} \|f^{(k)}\|_{L^2(\mathbb{R},\gamma)} \cdot \frac{1}{\sqrt{k!(n-k)!(q^{-1}-1)^{k/2}}} \\
&\leq \left( \sum_{k=0}^n \frac{(q^{-1}-1)^k}{k!} \|f^{(k)}\|_{L^2(\mathbb{R},\gamma)}^2 \right)^{1/2} \cdot \left( \sum_{k=0}^n \frac{1}{k!(n-k)!(q^{-1}-1)^k} \right)^{1/2} \\
&\leq \|f\|_{\mathcal{H}_q} \left( \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} \frac{1}{(q^{-1}-1)^k} \right)^{1/2} \\
&= \|f\|_{\mathcal{H}_q} \frac{1}{\sqrt{n!}} \left( \frac{1}{1-q} \right)^{n/2}.
\end{aligned}$$

Taking these computations together, we obtain

$$\|T^{(n)}\| \leq \left( \frac{1}{1-q} \right)^{n/2} \cdot (1 + q^{-1})^{n/2} = \left( \frac{1 + q^{-1}}{1 - q} \right)^{n/2}. \quad (12)$$

Therefore, we get that for  $|\rho| < \sqrt{\frac{1-q}{1+q^{-1}}}$

$$\sum_{n=0}^{\infty} \|\rho^n T^{(n)}\| < \infty.$$

The set of all linear and bounded operators on  $\mathcal{H}_q$  is a Banach space and thus,  $T_\rho = \sum_{n=0}^{\infty} \rho^n T^{(n)}$  is a linear and bounded operator on  $\mathcal{H}_q$ . For the compactness of  $T_\rho$ , note that  $T^{(n)}$  is a finite-rank operator for all  $n \in \mathbb{N}$ , namely of rank 1, i.e. the range of  $T^{(n)}$  is one-dimensional. As a finite-rank operator,  $T^{(n)}$  is a compact operator. The subset of all compact operators in the Banach space of the linear and bounded operators on  $\mathcal{H}_q$  is itself a Banach space (see e.g. [14, III. Theorem 4.7]). Hence,  $T_\rho = \sum_{n=0}^{\infty} \rho^n T^{(n)}$  is compact.  $\square$

*Proof of Theorem 2.* Let  $-\sqrt{\frac{1-q}{1+q^{-1}}} < \rho < \sqrt{\frac{1-q}{1+q^{-1}}}$ . We begin by relating the eigenvalue problem of  $T_\rho$  to the persistence problem of the MA(1)-process. First, note that  $S_\rho^N(\mathbf{1}) = T_\rho^N(\mathbf{1})$  for all  $N \in \mathbb{N}$ . By [5, Section 2.1] we can rewrite the persistence probability as follows:

$$\mathbb{P}(X_0 \geq 0, \dots, X_N \geq 0) = \int_S T_\rho^N(\mathbf{1})(x_2) d(\gamma \otimes \gamma)(x_1, x_2),$$

with  $S := \{(x_1, x_2) \in \mathbb{R}^2: \rho x_1 + x_2 \geq 0\}$ . Let  $r(T_\rho)$  be the spectral radius of  $T_\rho$ . We need to show that

$$\int_S T_\rho^N(\mathbf{1})(x_2) d(\gamma \otimes \gamma)(x_1, x_2) = r(T_\rho)^{N+o(N)}.$$

A priori we cannot exclude that  $r(T_\rho) = 0$ . For the upper bound note that

$$\|f\|_{L^1(\mathbb{R}, \gamma)} \leq \|f\|_{L^2(\mathbb{R}, \gamma)} \leq \|f\|_{\mathcal{H}_q}$$

for all  $f \in \mathcal{H}_q$  due to Lemma 5(d). Using this, we obtain

$$\begin{aligned} \int_S T_\rho^N(\mathbf{1})(x_2) d(\gamma \otimes \gamma)(x_1, x_2) &\leq \|T_\rho^N(\mathbf{1})\|_{L^1(\mathbb{R}, \gamma)} \\ &\leq \|T_\rho^N(\mathbf{1})\|_{\mathcal{H}_q} \\ &\leq \|T_\rho^N\| \cdot \|\mathbf{1}\|_{\mathcal{H}_q} \\ &= r(T_\rho)^{N+o(N)}, \end{aligned}$$

where the last step is due to Gelfand's formula, i.e.  $r(T_\rho) = \lim_{N \rightarrow \infty} \|T_\rho^N\|^{\frac{1}{N}}$ . Now, we turn to the lower bound. We need to consider two cases. If  $r(T_\rho) = 0$ , then clearly

$$\int_S T_\rho^N(\mathbb{1})(x_2) d(\gamma \otimes \gamma)(x_1, x_2) \geq r(T_\rho)^N.$$

If  $r(T_\rho) > 0$ , we show the lower bound by using the Krein-Rutman theorem (see [16], [12, Theorem 19.2]). For this purpose, let us define the cone  $C := \{f \in \mathcal{H}_q : f(x) \geq 0 \text{ for all } x \in \mathbb{R}\}$ . From [1, Proposition 1] it follows that  $\mathcal{H}_q$  is a reproducing kernel Hilbert space with reproducing kernel

$$K_q(x, y) := \sum_{n=0}^{\infty} q^n \widehat{h}_n(x) \widehat{h}_n(y) = \frac{1}{\sqrt{1-q^2}} e^{-\frac{q^2 x^2 + q^2 y^2 - 2qxy}{2(1-q^2)}},$$

where the last equality is due to Mehler's formula (see [20]). It holds that  $K_q^y(\cdot) := K_q(\cdot, y) \in C$  for all  $y \in \mathbb{R}$ . Since  $\text{span}\{K_q^y : y \in \mathbb{R}\}$  is dense in  $\mathcal{H}_q$  (see e.g. [7]), the closure of  $C + (-C)$  is equal to  $\mathcal{H}_q$ . Further, we have  $T_\rho(C) \subseteq C$ . Therefore, the Krein-Rutman theorem can be applied and yields the existence of an eigenfunction  $g \in C$  with eigenvalue  $r(T_\rho)$ . Note that any eigenfunction of  $T_\rho$  is bounded since

$$|T_\rho f(x)| \leq \int_{-px}^{\infty} |f(y) \phi(y)| dy \leq \|f\|_{L^1(\mathbb{R}, \gamma)} \leq \|f\|_{\mathcal{H}_q},$$

for all  $f \in \mathcal{H}_q$  and  $x \in \mathbb{R}$ . Hence,  $\|g\|_\infty < \infty$ . We obtain

$$\begin{aligned} \int_S T_\rho^N(\mathbb{1})(x_2) d(\gamma \otimes \gamma)(x_1, x_2) &\geq \int_S T_\rho^N \left( \frac{g}{\|g\|_\infty} \right) (x_2) d(\gamma \otimes \gamma)(x_1, x_2) \\ &= r(T_\rho)^N \int_S \frac{g(x_2)}{\|g\|_\infty} d(\gamma \otimes \gamma)(x_1, x_2) \\ &= r(T_\rho)^{N+o(N)}. \end{aligned}$$

Thus,

$$\mathbb{P}(X_0 \geq 0, \dots, X_N \geq 0) = r(T_\rho)^{N+o(N)}.$$

Now, note that [5, Proposition 2.3] implies that  $r(T_\rho) > 0$ . Hence,  $\lambda_\rho := r(T_\rho) > 0$  is the largest eigenvalue of  $T_\rho$  by the Krein-Rutman theorem. Further, we have  $\mathbb{P}(X_0 \geq 0, \dots, X_N \geq 0) \leq \mathbb{P}(\min_{0 \leq n \leq \lfloor \frac{N}{2} \rfloor} X_{2n} \geq 0)$ . Note that the random variables  $\{X_{2n} : 0 \leq n \leq \lfloor \frac{N}{2} \rfloor\}$  are independent. Hence,

$$\mathbb{P}(X_0 \geq 0, \dots, X_N \geq 0) \leq \mathbb{P}(X_0 \geq 0)^{\lfloor \frac{N}{2} \rfloor + 1},$$



which implies  $\lambda_\rho < 1$ .

It remains to show that  $\lambda_\rho$  admits a representation as a power series. We will see that this follows by [4, Theorem 4], which is based on the classical work of [14]. From the eigenvalue equation

$$\lambda f(x) = T_0 f(x) = \mathbb{1} \int_0^\infty f(y) \phi(y) dy \quad \text{for all } x \in \mathbb{R}, \quad (13)$$

it follows that the largest eigenvalue of  $T_0$  is given by  $\lambda_0 = \frac{1}{2}$ . To obtain the analyticity of the eigenvalue at 0 in  $\rho$  by methods of perturbation theory, it is necessary to show that the algebraic multiplicity of  $\lambda_0$  is equal to one.

For this purpose, let  $P_{\lambda_0}$  be the spectral projection of  $\lambda_0$ . The algebraic multiplicity is defined by the dimension of  $P_{\lambda_0} L^2(\mathbb{R}, \gamma)$ . Due to the compactness of  $T_0$ , we get  $P_{\lambda_0} L^2(\mathbb{R}, \gamma) = \ker(\lambda_0 - T_0)^v$ , where  $v \in \mathbb{N}$  is the smallest natural number such that  $\ker(\lambda_0 - T_0)^v = \ker(\lambda_0 - T_0)^{v+1}$  (see e.g. [10]).

From the eigenvalue equation (13), we see that  $\ker(\lambda_0 - T_0)^1$  is equal to the constant functions and therefore one-dimensional.

If we prove that  $\ker(\lambda_0 - T_0)^1 = \ker(\lambda_0 - T_0)^2$ , it follows that the algebraic multiplicity of  $\lambda_0$  is one. Let  $g \in \ker(\lambda_0 - T_0)^2$ . Then,

$$\begin{aligned} 0 &= (\lambda_0 - T_0)^2(g) \\ &= (\lambda_0 - T_0)(\lambda_0 g - T_0(g)) \\ &= \lambda_0^2 g - 2\lambda_0 T_0(g) + T_0(T_0(g)). \end{aligned}$$

Since  $T_0(g)$  is constant, the above equation yields that  $g$  is constant, i.e.  $g \in \ker(\lambda_0 - T_0)^1$ . By [4, Theorem 4], it follows that  $\lambda_\rho$  can be represented as a power series for  $|\rho| < r_0$  for some  $r_0 > 0$ . The question of the radius of convergence is tricky and will be tackled in Section 5. In particular, the only thing that is still to be proved is the bound for the radius of convergence, which can be found in Corollary 11 combined with Lemma 12.  $\square$

**Remark.** *We remark that the operator  $T_\rho$  is not normal. If it were, further results from perturbation theory would be applicable. In particular, a concrete bound for the radius of convergence would follow from Corollary 6 in [4] together with the bound (12).*

*Proof of Theorem 3.* The eigenvalue equation  $T_\rho(f_\rho) = \lambda_\rho f_\rho$  reads:

$$\sum_{k=0}^{\infty} \rho^k T^{(k)} \left( \sum_{m=0}^{\infty} \rho^m g_m \right) = \sum_{k=0}^{\infty} \rho^k K_k \cdot \sum_{m=0}^{\infty} \rho^m g_m.$$

Sorting this in powers of  $\rho$  (which is allowed within the radius of convergence) gives

$$\sum_{n=0}^{\infty} \rho^n \sum_{k=0}^n T^{(k)} g_{n-k} = \sum_{n=0}^{\infty} \rho^n \sum_{k=0}^n K_k g_{n-k}.$$

Since this holds for any  $\rho \in (-\rho_0, \rho_0)$  with  $\rho_0 > 0$ , we must have

$$\sum_{k=0}^n T^{(k)} g_{n-k} = \sum_{k=0}^n K_k g_{n-k}, \quad \text{for all } n \in \mathbb{N}. \quad (14)$$

For  $n = 0$ , this is

$$T^{(0)} g_0 = K_0 g_0.$$

Since the left-hand side is a constant, by the definition of  $T^{(0)} = T_0$ , we know that  $g_0$  must be constant, so w.l.o.g. (multiplication of the eigenfunction) take

$$g_0 = \mathbb{1}.$$

Fix  $n \geq 1$ . We now analyse the iterative structure of (14):

$$\sum_{k=1}^n T^{(k)} g_{n-k} + T^{(0)} g_n = K_n g_0 + \sum_{k=1}^{n-1} K_k g_{n-k} + K_0 g_n. \quad (15)$$

Observe that, by the definition of the operators  $T^{(k)}$ , the first term on the left is a polynomial. Further, the second term on the left and the first term on the right are constants. The second term on the right involves only the  $g_\ell$ ,  $\ell < n$ . Therefore, inductively we know that  $g_n$  is a polynomial of degree at most  $n$ .

Let us denote the coefficients of the polynomials  $g_\ell$  by  $g_\ell^i$ , i.e.  $g_\ell(x) =: \sum_{i=0}^{\ell} g_\ell^i x^i$ . Then comparing the coefficients of  $x^0$  in (15) gives

$$0 + T^{(0)} g_n = K_n g_0 + \sum_{k=1}^{n-1} K_k g_{n-k}^0 + K_0 g_n^0.$$

Further, by the definition of  $T^{(0)}$  and the representation of  $g_n$ , we have

$$T^{(0)} g_n = \sum_{i=0}^n g_n^i \int_0^\infty y^i \phi(y) dy.$$

Combining the last two equations and using that  $\int_0^\infty y^0 \phi(y) dy = K_0$ , we see that the terms involving  $g_n^0$  cancel; giving

$$\sum_{i=1}^n g_n^i \int_0^\infty y^i \phi(y) dy = K_n g_0 + \sum_{k=1}^{n-1} K_k g_{n-k}^0. \quad (16)$$

None of the other polynomial terms in (15) uses  $g_n^0$  either. Therefore,  $g_n^0$  is in fact arbitrary and can be chosen to be zero. This however simplifies (15) in the sense that

$$g_n = \frac{1}{K_0} \left( \sum_{k=1}^n T^{(k)} g_{n-k} - \sum_{k=1}^{n-1} K_k g_{n-k} \right), \quad (17)$$

because these are the parts of (15) involving the coefficients of  $x^i$ ,  $i > 0$ , only. Note that (17) computes  $g_n$  with the help of  $g_\ell$  and  $K_\ell$  for  $\ell < n$  only. Similarly, now (16) simplifies (using  $g_\ell^0 = 0$  and  $g_0 = \mathbb{1}$ ) to (10). Putting this back into (17) shows (9).  $\square$

## 5 Radius of convergence

This section is devoted to the study of the radius of convergence of the series in Theorem 2. In particular, we would like to finish the proof of that theorem by showing that the radius of convergence is at least 0.332.

For this purpose, we first give an alternative description of the leading eigenvalue of the eigenvalue equation (3), which might be interesting in its own right, cf. (28). We also show that the two descriptions must have the same radius of convergence. Using the alternative description, we can prove a lower bound for the radius of convergence, cf. Corollary 11.

We start with a re-formulation of the eigenvalue equation (3) that gets rid of the eigenfunction.

**Lemma 6.** *Let  $\rho \in [0, 1]$ . The leading eigenvalue  $\lambda = \lambda_\rho$  of the eigenvalue equation (3) is the largest positive root of the equation*

$$\lambda = \sum_{k=0}^{\infty} \frac{\kappa_k(\rho)}{\lambda^k}, \quad (18)$$

where  $\kappa_0 \equiv \frac{1}{2}$  and for  $k \geq 1$ :

$$\begin{aligned} \kappa_k(\rho) &:= \frac{1}{(2\pi)^{\frac{k+1}{2}}} \int_0^\infty \int_{-\rho s_0}^0 \cdots \int_{-\rho s_{k-2}}^0 \int_{-\rho s_{k-1}}^0 \exp\left(-\frac{1}{2} \sum_{i=0}^k s_i^2\right) ds_k \dots ds_0 \quad (19) \\ &= \frac{\rho^{\frac{k(k+1)}{2}}}{(2\pi)^{\frac{k+1}{2}}} \int_0^\infty \int_{-s_0}^0 \cdots \int_{-s_{k-2}}^0 \int_{-s_{k-1}}^0 \exp\left(-\frac{1}{2} \sum_{i=1}^k (\rho^i s_i)^2\right) e^{-\frac{s_0^2}{2}} ds_k \dots ds_0. \end{aligned}$$

Before we give the proof of Lemma 6, we note a useful technical bound for the coefficients  $(\kappa_k)$ .

**Lemma 7.** For the  $(\kappa_k)$  defined in (19), we have for all  $k \geq 0$ :

$$|\kappa_k(\rho)| = \frac{\rho^{\frac{k(k+1)}{2}}}{(2\pi)^{\frac{k+1}{2}}} \int_0^\infty \int_0^{s_0} \cdots \int_0^{s_{k-2}} \int_0^{s_{k-1}} \exp\left(-\frac{1}{2} \sum_{i=1}^k (\rho^i s_i)^2\right) e^{-\frac{s_0^2}{2}} ds_k \cdots ds_0$$

$$\leq \frac{\rho^{\frac{k(k+1)}{2}}}{2^{k+1} \pi^{\frac{k}{2}} \Gamma\left(\frac{k}{2} + 1\right)}.$$
(20)

*Proof of Lemma 7.* Bounding all the exponentials in (20) by one, except the last one, we obtain:

$$|\kappa_k(\rho)| \leq \frac{\rho^{\frac{k(k+1)}{2}}}{(2\pi)^{\frac{k+1}{2}} k!} \int_0^\infty s^k e^{-\frac{s^2}{2}} ds = \frac{\rho^{\frac{k(k+1)}{2}}}{2\pi^{\frac{k+1}{2}} k!} \Gamma\left(\frac{k+1}{2}\right) = \frac{\rho^{\frac{k(k+1)}{2}}}{2^{k+1} \pi^{\frac{k}{2}} \Gamma\left(\frac{k}{2} + 1\right)},$$

where we used the Legendre duplication formula  $\Gamma(2t) = 2^{2t-1} \Gamma(t) \Gamma(t + 1/2) / \sqrt{\pi}$  for  $t = (k+1)/2$  in the last step.  $\square$

**Remark.** A result similar to Lemma 7 (and subsequently a result similar to Lemma 6) can be proved in the same way as long as the density  $\phi$  has a superexponential decay.

**Remark.** We cannot expect explicit values for the coefficients  $\kappa_k(\rho)$ . In fact, they are persistence probabilities themselves:

$$\kappa_k(\rho) = \mathbb{P}(0 \leq \xi_0 \leq \rho \xi_1 \leq \dots \leq \rho^k \xi_k),$$

where the  $(\xi_i)$  are independent random variables with density  $\phi$ .

*Proof of Lemma 6.* The only purpose of the proof is to transform the eigenvalue equation:

$$\lambda f(x) = \int_{-\rho x}^\infty f(s) \phi(s) ds = \int_0^\infty f(s) \phi(s) ds + \int_{-\rho x}^0 f(s) \phi(s) ds, \quad x \in \mathbb{R}.$$
(21)

Further, we set w.l.o.g.  $f(0) := 1$ . Inserting this into (21) gives

$$\lambda = \int_0^\infty f(s) \phi(s) ds.$$
(22)

Replacing the corresponding term in (21) gives

$$f(x) = 1 + \lambda^{-1} \int_{-\rho x}^0 f(s) \phi(s) ds.$$
(23)

Inserting (23) into (22) we obtain

$$\begin{aligned}
\lambda &= \int_0^\infty \left[ 1 + \lambda^{-1} \int_{-\rho s_0}^0 f(s_1) \phi(s_1) ds_1 \right] \phi(s_0) ds_0 \\
&= \int_0^\infty \phi(s_0) ds_0 + \lambda^{-1} \int_0^\infty \int_{-\rho s_0}^0 f(s_1) \phi(s_1) ds_1 \phi(s_0) ds_0 \\
&= \frac{1}{2} + \lambda^{-1} \int_0^\infty \int_{-\rho s_0}^0 f(s_1) \phi(s_1) ds_1 \phi(s_0) ds_0.
\end{aligned}$$

Iterating this procedure gives

$$\lambda = \sum_{k=0}^{N-1} \lambda^{-k} \kappa_k(\rho) + \lambda^{-N} \int_0^\infty \int_{-\rho s_0}^0 \dots \int_{-\rho s_{N-1}}^0 f(s_N) \phi(s_N) ds_N \dots \phi(s_1) ds_1 \phi(s_0) ds_0.$$

We now let  $N \rightarrow \infty$  and show that the remainder term vanishes. For this purpose, we use that the eigenfunction  $f$  is bounded and we employ the bound from Lemma 7:

$$\begin{aligned}
&\left| \lambda^{-N} \int_0^\infty \int_{-\rho s_0}^0 \dots \int_{-\rho s_{N-1}}^0 f(s_N) \phi(s_N) ds_N \dots \phi(s_1) ds_1 \phi(s_0) ds_0 \right| \\
&\leq \|f\|_\infty \cdot \frac{\text{const}^N}{\Gamma(N/2 + 1)} \rightarrow 0. \quad \square
\end{aligned}$$

We are going to re-write the eigenvalue equation in the following way. Define

$$\Psi_\rho(\xi) := \sum_{k=1}^{\infty} 2^{k+1} \kappa_k(\rho) (1 + \xi)^{-k}.$$

Note that finding solutions of (18) for  $\lambda \geq \frac{1}{2}$  is equivalent to finding solutions of the following equation for  $\xi \geq 0$ :

$$\xi = \Psi_\rho(\xi), \quad \lambda = \frac{1}{2}(1 + \xi). \quad (24)$$

That means, we are looking for the largest root  $\xi \geq 0$  of equation (24). Next, we define

$$\Gamma_\rho(\xi) := \frac{\xi}{\Psi_\rho(\xi)}, \quad \xi \geq 0.$$

Let us state some important properties of the function  $\Gamma_\rho$  that will allow us to invert the function.

**Lemma 8.** *We have  $\Gamma_\rho(0) = 0$ ,  $\lim_{\xi \rightarrow \infty} \Gamma_\rho(\xi) = \infty$ . Further, for  $\rho \in [0, 1]$ , the function  $\xi \mapsto \Gamma_\rho(\xi)$  is strictly increasing.*

*Proof.* The first two properties follow immediately from Lemma 7. To see the last property, note that the derivative of  $\Gamma_\rho$  has the same sign as:

$$\begin{aligned}
\Psi_\rho(\xi) - \xi\Psi'_\rho(\xi) &= \sum_{k=1}^{\infty} 2^{k+1} \kappa_k(\rho) (1+\xi)^{-k} + \sum_{k=1}^{\infty} 2^{k+1} \kappa_k(\rho) k\xi (1+\xi)^{-k-1} \\
&= 4\kappa_1(\rho) \frac{1+2\xi}{(1+\xi)^2} + \sum_{k=2}^{\infty} 2^{k+1} \kappa_k(\rho) \frac{1+(k+1)\xi}{(1+\xi)^{k+1}} \\
&\geq 4\kappa_1(\rho) \frac{1+2\xi}{(1+\xi)^2} - \sum_{k=2}^{\infty} \frac{\rho^{\frac{k(k+1)}{2}}}{\pi^{\frac{k}{2}}} \frac{1}{\Gamma(\frac{k}{2}+1)} \frac{1+(k+1)\xi}{(1+\xi)^{k+1}} \\
&= 4 \frac{1+2\xi}{(1+\xi)^2} \left[ \frac{1}{2\pi} \arctan(\rho) - \sum_{k=2}^{\infty} \frac{\rho^{\frac{k(k+1)}{2}}}{\pi^{\frac{k}{2}}} \frac{1}{\Gamma(\frac{k}{2}+1)} \frac{1+(k+1)\xi}{(1+\xi)^{k-1}(1+2\xi)} \right], \tag{25}
\end{aligned}$$

where differentiating under the sum in the first step is allowed due to Lemma 7 and we note that

$$\kappa_1(\rho) = \frac{1}{2\pi} \int_0^\infty \int_{-\rho z}^0 e^{-x^2/2} e^{-z^2/2} dx dz = \frac{1}{2\pi} \arctan(\rho) > 0. \tag{26}$$

To see the last equality, differentiate with respect to  $\rho$ . Now note that, for  $k \geq 2$ , the function

$$\xi \mapsto \frac{1+(k+1)\xi}{(1+\xi)^{k-1}(1+2\xi)}$$

is strictly decreasing, which can be seen by differentiation. Therefore, we can replace it in (25) by its value at  $\xi = 0$  and thus get the estimate:

$$\begin{aligned}
\Psi_\rho(\xi) - \xi\Psi'_\rho(\xi) &\geq 4 \frac{1+2\xi}{(1+\xi)^2} \left[ \frac{1}{2\pi} \arctan(\rho) - \sum_{k=2}^{\infty} \frac{\rho^{\frac{k(k+1)}{2}}}{\pi^{\frac{k}{2}}} \frac{1}{\Gamma(\frac{k}{2}+1)} \right] \\
&\geq 4 \frac{1+2\xi}{(1+\xi)^2} \left[ \frac{1}{2\pi} \arctan(\rho) - \sum_{k=2}^{\infty} \frac{\rho^{k+1}}{\pi^{\frac{k}{2}}} \right] \\
&= 4 \frac{1+2\xi}{(1+\xi)^2} \left[ \frac{1}{2\pi} \arctan(\rho) - \frac{\rho^3}{\sqrt{\pi}(\sqrt{\pi}-\rho)} \right],
\end{aligned}$$

which is strictly positive for all  $\rho \in (0, 1]$  and all  $\xi \geq 0$ .  $\square$

**Corollary 9.** *The largest solution of (18) admits the following representation*

$$\lambda = \frac{1}{2} + \frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n!} \left( \frac{\partial^{n-1}}{\partial \xi^{n-1}} [\Psi_{\rho}(\xi)^n] \right) \Big|_{\xi=0}. \quad (27)$$

*Proof.* We use the Lagrange-Bürmann formula, a variant of the Lagrange inversion formula. Knowing that  $\Gamma_{\rho}(0) = 0$ ,  $\Gamma_{\rho}(\infty) = \infty$ , and that it is strictly increasing, we can resolve the equation  $\Gamma_{\rho}(\xi) = 1$ , i.e. compute the inverse of  $\Gamma_{\rho}$  at 1. The result from the Lagrange-Bürmann formula is precisely the statement of the corollary. Note that (18) admits several solutions in  $\lambda$ , while here we show that there is only one solution in  $\xi \geq 0$ , i.e. one solution in  $\lambda$  with  $\lambda \geq \frac{1}{2}$ .  $\square$

**Corollary 10.** *The largest solution of (18) admits the following representation*

$$\lambda = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{(-1)^{n-1} 2^{n-1}}{n!} \sum_{k_1=1}^{\infty} \dots \sum_{k_n=1}^{\infty} \left( \prod_{j=1}^n 2^{k_j} \kappa_{k_j}(\rho) \right) \frac{\Gamma(k_1 + \dots + k_n + n - 1)}{\Gamma(k_1 + \dots + k_n)}. \quad (28)$$

*Proof.* To compute the successive derivatives in (27), we write  $\Psi_{\rho}$  as a Laplace transform

$$\Psi_{\rho}(\xi) = \sum_{k=1}^{\infty} 2^{k+1} \kappa_k(\rho) \frac{1}{\Gamma(k)} \int_0^{\infty} e^{-\xi x} e^{-x} x^{k-1} dx = \int_0^{\infty} e^{-\xi x} \psi_{\rho}(x) dx,$$

where we can exchange the sum and integral, due to Lemma 7, and we set

$$\psi_{\rho}(x) := e^{-x} \sum_{k=1}^{\infty} \frac{2^{k+1}}{\Gamma(k)} \kappa_k(\rho) x^{k-1}.$$

As a consequence,

$$\begin{aligned} \left( \frac{\partial^{n-1}}{\partial \xi^{n-1}} (\Psi_{\rho}(\xi))^n \right) \Big|_{\xi=0} &= \left( \int_0^{\infty} (-1)^{n-1} x^{n-1} e^{-\xi x} \psi_{\rho}^{*(n)}(x) dx \right) \Big|_{\xi=0} \\ &= (-1)^{n-1} \int_0^{\infty} x^{n-1} \psi_{\rho}^{*(n)}(x) dx, \end{aligned}$$

where  $*$  denotes the usual convolution product. For the first term, we have

$$\psi_{\rho}^{*(1)}(x) = \psi_{\rho}(x) = e^{-x} \sum_{k_1=1}^{\infty} \frac{2^{k_1+1}}{\Gamma(k_1)} \kappa_{k_1}(\rho) x^{k_1-1}.$$

For the second term, we get

$$\begin{aligned}
\psi_\rho^{*(2)}(x) &= e^{-x} \int_0^x \sum_{k_1=1}^{\infty} \frac{2^{k_1+1}}{\Gamma(k_1)} \kappa_{k_1}(\rho) (x-y)^{k_1-1} \sum_{k_2=1}^{\infty} \frac{2^{k_2+1}}{\Gamma(k_2)} \kappa_{k_2}(\rho) y^{k_2-1} dy \\
&= e^{-x} \sum_{k_1=1}^{\infty} \frac{2^{k_1+1}}{\Gamma(k_1)} \kappa_{k_1}(\rho) \sum_{k_2=1}^{\infty} \frac{2^{k_2+1}}{\Gamma(k_2)} \kappa_{k_2}(\rho) x^{k_1+k_2-1} \int_0^1 (1-y)^{k_1-1} y^{k_2-1} dy \\
&= e^{-x} \sum_{k_1=1}^{\infty} \sum_{k_2=1}^{\infty} 2^{k_1+k_2+2} \kappa_{k_1}(\rho) \kappa_{k_2}(\rho) \frac{x^{k_1+k_2-1}}{\Gamma(k_1+k_2)},
\end{aligned}$$

where exchanging sums and integral is permitted by Lemma 7. By induction, we obtain

$$\psi_\rho^{*(n)}(x) = e^{-x} \sum_{k_1=1}^{\infty} \dots \sum_{k_n=1}^{\infty} 2^{k_1+\dots+k_n+n} \left( \prod_{j=1}^n \kappa_{k_j}(\rho) \right) \cdot \frac{x^{k_1+\dots+k_n-1}}{\Gamma(k_1+\dots+k_n)}$$

and so

$$\begin{aligned}
&\int_0^\infty x^{n-1} \psi_\rho^{*(n)}(x) dx \\
&= \sum_{k_1=1}^{\infty} \dots \sum_{k_n=1}^{\infty} \left( \prod_{j=1}^n 2^{k_j+1} \kappa_{k_j}(\rho) \right) \cdot \frac{\Gamma(k_1+\dots+k_n+n-1)}{\Gamma(k_1+\dots+k_n)}. \quad \square
\end{aligned}$$

Together with the bound in Lemma 7, it is now possible to obtain a bound for the radius of convergence – despite the fact that we cannot determine the  $\kappa_n(\rho)$  more explicitly.

**Corollary 11.** *The representation in (28) converges absolutely for  $|\rho| < 0.332$ .*

*Proof.* Observe that the coefficient  $\kappa_k(\rho)$  admits a series expansion of the form

$$\kappa_k(\rho) =: \frac{\rho^{\frac{k(k+1)}{2}}}{(2\pi)^{\frac{k+1}{2}}} \sum_{n=0}^{\infty} \psi_n^{(k)} \rho^n.$$

By putting absolute values everywhere, we thus obtain the following bound for the absolute value of the term in (28):

$$\frac{1}{2} + \frac{1}{2} \sum_{n=1}^{\infty} \frac{2^n}{n!} \sum_{k_1=1}^{\infty} \dots \sum_{k_n=1}^{\infty} \left( \prod_{j=1}^n \frac{\rho^{\frac{k_j(k_j+1)}{2}} 2^{k_j}}{(2\pi)^{\frac{k_j+1}{2}}} \sum_{p=0}^{\infty} |\psi_p^{(k_j)}| |\rho|^p \right) \frac{\Gamma(k_1+\dots+k_n+n-1)}{\Gamma(k_1+\dots+k_n)}. \quad (29)$$



For the remainder of this proof, we consider  $\rho > 0$  only to avoid the absolute value signs. To compute the multiple sum, we first bound the ratio of Gamma functions. Note that, by the binomial theorem, for  $x \in \mathbb{N}$

$$\begin{aligned} \frac{\Gamma(x+n-1)}{\Gamma(x)} &= \frac{(x+n-2)!}{(x-1)!} = (n-1)! \frac{(x+n-2)!}{(n-1)!(x-1)!} \\ &= (n-1)! \binom{x+n-2}{n-1} \leq (n-1)! 2^{x+n-2}. \end{aligned}$$

As a consequence, also using the Fubini-Tonelli theorem, we get the upper bound for the term in (29):

$$\begin{aligned} &\frac{1}{2} + \frac{1}{2} \sum_{n=1}^{\infty} \frac{2^n}{n!} \sum_{k_1=1}^{\infty} \cdots \sum_{k_n=1}^{\infty} \left( \prod_{j=1}^n \frac{\rho^{\frac{k_j(k_j+1)}{2}} 4^{k_j}}{(2\pi)^{\frac{k_j+1}{2}}} \sum_{p=0}^{\infty} |\psi_p^{(k_j)}| \rho^p \right) (n-1)! 2^{n-2} \\ &= \frac{1}{2} + \frac{1}{8} \sum_{n=1}^{\infty} \frac{4^n}{n} \left( \sum_{k=1}^{\infty} \frac{\rho^{\frac{k(k+1)}{2}} 4^k}{(2\pi)^{\frac{k+1}{2}}} \sum_{p=0}^{\infty} |\psi_p^{(k)}| \rho^p \right)^n. \end{aligned} \quad (30)$$

We thus arrive at the following sufficient condition for the sum in (30) to converge:

$$\sum_{k=1}^{\infty} \frac{\rho^{\frac{k(k+1)}{2}} 4^k}{(2\pi)^{\frac{k+1}{2}}} \sum_{p=0}^{\infty} |\psi_p^{(k)}| \rho^p < \frac{1}{4}. \quad (31)$$

We now study  $\sum_{p=0}^{\infty} |\psi_p^{(k)}| \rho^p$ . For  $k=1$ , we have due to (26):

$$\kappa_1(\rho) = \frac{\arctan(\rho)}{2\pi} = \frac{1}{2\pi} \sum_{n=0}^{\infty} (-1)^n \frac{\rho^{2n+1}}{2n+1};$$

hence, the definition  $\kappa_1(\rho) = \frac{\rho}{2\pi} \sum_{p=0}^{\infty} \psi_p^{(1)} \rho^p$  yields

$$\sum_{n=0}^{\infty} |\psi_n^{(1)}| \rho^n = \sum_{n=0}^{\infty} \frac{\rho^{2n}}{2n+1} = \frac{1}{2\rho} \log \left( \frac{1+\rho}{1-\rho} \right).$$

Turning to the terms for  $k \geq 2$  in (31), similarly to the proof of Lemma 7, we have

$$\sum_{n=0}^{\infty} |\psi_n^{(k)}| \rho^n \leq \int_0^{\infty} \int_0^{s_0} \cdots \int_0^{s_{k-1}} \int_0^{s_k} \exp \left( \frac{1}{2} \sum_{i=1}^k (\rho^i s_i)^2 \right) e^{-\frac{s_0^2}{2}} ds_k \cdots ds_0$$

$$\begin{aligned}
&\leq \int_0^\infty \frac{s^k}{k!} \exp\left(\frac{1}{2} \sum_{i=1}^k (\rho^i s)^2\right) e^{-\frac{s^2}{2}} ds \\
&\leq \int_0^\infty \frac{s^k}{k!} \exp\left(-\frac{s^2}{2} \frac{1-2\rho^2}{1-\rho^2}\right) ds \\
&= \frac{1}{2 \cdot k!} \Gamma\left(\frac{k+1}{2}\right) 2^{\frac{k+1}{2}} \left(\frac{1-\rho^2}{1-2\rho^2}\right)^{\frac{k+1}{2}}. \tag{32}
\end{aligned}$$

Therefore, (31) is implied by

$$\frac{1}{\pi} \log\left(\frac{1+\rho}{1-\rho}\right) + \sum_{k=2}^\infty \frac{\rho^{\frac{k(k+1)}{2}} 4^k}{\pi^{\frac{k+1}{2}}} \frac{1}{2 \cdot k!} \Gamma\left(\frac{k+1}{2}\right) \left(\frac{1-\rho^2}{1-2\rho^2}\right)^{\frac{k+1}{2}} < \frac{1}{4}.$$

One can check numerically that this is true at least for all  $\rho < 0.332$ .  $\square$

Note that the bound in the last corollary can be improved by evaluating more terms in the sum (31) explicitly instead of using the estimate (32).

**Lemma 12.** *The representation in (28) holds if and only if the representation  $\lambda_\rho = \sum_{i=0}^\infty K_i \rho^i$  holds, where the  $(K_i)$  are as in Theorem 2.*

*Proof.* First observe that the  $\kappa_k(\rho)$  can be written as a series in  $\rho$ . Now, within the radius of convergence, one may rearrange all sums in (28). Therefore, the representation in (28) may be re-written as a series in  $\rho$ . Since the representation  $\lambda_\rho = \sum_{i=0}^\infty K_i \rho^i$  is analytic in a vicinity of 0, the coefficients of the two series have to agree. Therefore, the radius of convergence has to be identical.  $\square$

We mention that a large portion of the results of this manuscript are part of the PhD thesis [15].

## References

- [1] S. T. Ali and M. Engliš. Hermite polynomials and quasi-classical asymptotics. *Journal of Mathematical Physics*, 55(4):042102, 2014.
- [2] G. E. Andrews, R. Askey, and R. Roy. *Special functions*, volume 71 of *Encyclopedia of mathematics and its applications*. Cambridge University Press, 1999.
- [3] F. Aurzada and Ch. Baumgarten. Survival probabilities of weighted random walks. *ALEA, Latin American Journal of Probability and Mathematical Statistics*, 8:235–258, 2011.

- [4] F. Aurzada and M. Kettner. Persistence exponents via perturbation theory: AR(1)-processes. *Journal of Statistical Physics*, 177(4):651–665, 2019.
- [5] F. Aurzada, S. Mukherjee, and O. Zeitouni. Persistence exponents in Markov chains. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 57:1411–1441, 2021.
- [6] F. Aurzada and T. Simon. Persistence probabilities and exponents. In *Lévy matters. V*, volume 2149 of *Lecture Notes in Math.*, pages 183–224. Springer, Cham, 2015.
- [7] N. Aronszajn. Theory of reproducing kernels. *Transactions of the American Mathematical Society*, 68(3):337–404, 1950.
- [8] A. J Bray, S. N. Majumdar, and G. Schehr. Persistence and first-passage properties in nonequilibrium systems. *Advances in Physics*, 62(3):225–361, 2013.
- [9] P. Collet, S. Martínez, and J. San Martín. *Quasi-stationary distributions: Markov chains, diffusions and dynamical systems*. Springer Science & Business Media, 2012.
- [10] J. B. Conway. *A course in functional analysis*, volume 96 of *Graduate texts in mathematics*. Springer Science & Business Media, 2007.
- [11] N. Champagnat and D. Villemonais. General criteria for the study of quasi-stationarity. *Electron. J. Probab.* 28, Paper No. 22, 2023.
- [12] K. Deimling. *Nonlinear functional analysis*. Springer, 1985.
- [13] G. Hinrichs, M. Kolb, and V. Wachtel. Persistence of one-dimensional AR(1)-sequences. *Journal of Theoretical Probability*, 33(1):65–102, 2020.
- [14] T. Kato. *Perturbation theory for linear operators*, volume 132 of *Classics in mathematics*. Springer Science & Business Media, 1966.
- [15] M. Kettner. Persistence exponents via perturbation theory: autoregressive and moving average processes. PhD Dissertation. Technische Universität Darmstadt. 2021. DOI: 10.26083/tuprints-00017566
- [16] M. G. Krein and M. A. Rutman. Linear operators leaving invariant a cone in a Banach space. *Uspekhi Matematicheskikh Nauk*, 3(1):3–95, 1948.

- [17] M. Krishna and M. Krishnapur. Persistence probabilities in centered, stationary, Gaussian processes in discrete time. *Indian J. Pure Appl. Math.*, 47(2), 183–194, 2016.
- [18] W. V. Li and Q.-M. Shao. Lower tail probabilities for Gaussian processes. *The Annals of Probability*, 32(1A):216–242, 2004.
- [19] S. N. Majumdar, A. J. Bray, and G.C.M.A. Ehrhardt. Persistence of a continuous stochastic process with discrete-time sampling. *Physical Review E*, 64, 015101(R), 2001.
- [20] F. G. Mehler. Ueber die Entwicklung einer Function von beliebig vielen Variablen nach Laplaceschen Functionen höherer Ordnung. *Journal für die reine und angewandte Mathematik*, 66:161–176, 1866.
- [21] S. Méléard and D. Villemonais. Quasi-stationary distributions and population processes. *Probability Surveys*, 9:340–410, 2012.
- [22] R. Metzler, G. Oshanin and S. Redner. *First-passage phenomena and their applications*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2014.
- [23] J. Noonan and A. Zhigljavsky. Approximating Shepp’s constants for the Slepian process. *Statistics & Probability Letters*, 153:21–31, 2019.
- [24] K. Y. Patarroyo. A digression on Hermite polynomials. *arXiv preprint arXiv:1901.01648*, 2019.
- [25] L. A. Shepp. First passage time for a particular Gaussian process. *The Annals of Mathematical Statistics*, pages 946–951, 1971.
- [26] D. Slepian. First passage time for a particular Gaussian process. *The Annals of Mathematical Statistics*, 32(2):610–612, 1961.
- [27] R. L. Tweedie. Quasi-stationary distributions for Markov chains on a general state space. *Journal of Applied Probability*, 11(4):726–741, 1974.
- [28] R. L. Tweedie. R-theory for Markov chains on a general state space I: solidarity properties and R-recurrent chains. *The Annals of Probability*, 2(5):840–864, 1974.