



**HAL**  
open science

# Sharp functional calculus for the Taibleson operator on non-archimedean local fields

Cédric Arhancet, Christoph Kriegler

► **To cite this version:**

Cédric Arhancet, Christoph Kriegler. Sharp functional calculus for the Taibleson operator on non-archimedean local fields. 2024. hal-04644652

**HAL Id: hal-04644652**

**<https://hal.science/hal-04644652v1>**

Preprint submitted on 12 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Sharp functional calculus for the Taibleson operator on non-archimedean local fields

Cédric Arhancet and Christoph Kriegler

## Abstract

For any non-archimedean local field  $\mathbb{K}$  and any integer  $n \geq 1$ , we show that the Taibleson operator admits a bounded  $H^\infty(\Sigma_\theta)$  functional calculus for any angle  $\theta > 0$  on the Banach space  $L^p(\mathbb{K}^n)$ , where  $\Sigma_\theta = \{z \in \mathbb{C}^* : |\arg z| < \theta\}$  and  $1 < p < \infty$ , and even a bounded Hörmander functional calculus of order  $\frac{3}{2}$  (with striking contrast to the Euclidean Laplacian on  $\mathbb{R}^n$ ). In our study, we explore harmonic analysis on locally compact Spector-Vilenkin groups establishing the  $R$ -boundedness of a family of convolution operators. Our results enhance the understanding of functional calculi of operators acting on  $L^p$ -spaces associated to totally disconnected spaces and have implications for the maximal regularity of the fundamental evolution equations associated to the Taibleson operator, relevant in various physical models.

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Preliminaries</b>	<b>5</b>
<b>3</b>	<b>Locally compact Spector-Vilenkin groups and radial functions</b>	<b>10</b>
<b>4</b>	<b>Maximal inequalities</b>	<b>15</b>
<b>5</b>	<b><math>R</math>-boundedness of some family of convolution operators</b>	<b>17</b>
<b>6</b>	<b>Application to functional calculus</b>	<b>20</b>
	<b>Bibliography</b>	<b>27</b>

## 1 Introduction

Over the past four decades, the realm of  $q$ -adic analysis has garnered increasing interest due to its interdisciplinary applications spanning fields such as medicine, biology, and physics. For an in-depth exploration of advancements in  $q$ -adic analysis and its applications over this period, we refer to the books [AKS10], [Koc01], [KKZ18], [Tai75], [VVZ94], [Zun16] and to the surveys [BrF93] and [DKKVZ17]. Similar to how phenomena in nature are traditionally modeled using complex numbers and differential equations like the heat equation, certain processes including fluid dynamics in porous media [KOJ16a], [KOJ16b], macromolecules [ABO04], models of spin glasses [ABK99] and protein dynamics [BiZ23] can be effectively described using non-archimedean local fields like the  $q$ -adic numbers field. Mathematically, the majority of these models describe the time evolution of a complex system using a «master equation», a type of parabolic equation. This equation governs the evolution of a transition function for a Markov

---

2020 *Mathematics subject classification*: 46S10, 47S10, 11S80.

*Key words*: non-archimedean local fields, Spector-Vilenkin locally compact groups, Hörmander functional calculus,  $H^\infty(\Sigma_\theta)$  functional calculus,  $R$ -boundedness.

process in an ultrametric space. In recent years, numerous topics have been explored from the perspective of ultrametric analysis, including the Navier-Stokes equations [KhK20], quantum physics [ViV89], [Zun22], [Zun24b], quantum field theory [FGZ22] and neural networks [ZaZ23], [Zun24a].

A cornerstone of ultrametric analysis, the Taibleson operator  $D^\alpha$  is an ultrametric counterpart to the fractional Laplace operator  $(-\Delta)^\alpha$  on  $\mathbb{R}^n$ . We refer to [Sti19] and references therein for information on this last operator. Recall that one method to define this linear operator is to use the formula

$$(1.1) \quad (-\Delta)^\alpha f(x) = \frac{4^\alpha \Gamma(\frac{n}{2} + \alpha)}{\pi^{\frac{n}{2}} \Gamma(-\alpha)} \text{p. v.} \int_{\mathbb{R}^n} \frac{f(y) - f(x)}{|y - x|^{2\alpha+n}} dy, \quad 0 < \alpha < 1, f \in \mathcal{S}(\mathbb{R}^n), x \in \mathbb{R}^n.$$

If  $\mathbb{K}$  is a non-archimedean local field (as the field of  $q$ -adic numbers), i.e. a non-discrete totally disconnected locally compact topological field, the Taibleson operator is defined by the formula

$$(1.2) \quad (D^\alpha f)(x) \stackrel{\text{def}}{=} \frac{1 - q^\alpha}{1 - q^{-\alpha-n}} \int_{\mathbb{K}^n} \frac{f(y) - f(x)}{\|y - x\|_{\mathbb{K}^n}^{\alpha+n}} dy, \quad \alpha > 0, x \in \mathbb{K}^n,$$

for any suitable function  $f: \mathbb{K}^n \rightarrow \mathbb{C}$ , where  $q$  is some prime number related to the field  $\mathbb{K}$ . See [Koc01, (2.8)] for the case  $n = 1$  and [RoZ08, (2.3)] for the case of  $\mathbb{Q}_q^n$ . Here

$$(1.3) \quad \|x\|_{\mathbb{K}^n} \stackrel{\text{def}}{=} \max_{1 \leq i \leq n} |x_i|_{\mathbb{K}}, \quad x = (x_1, \dots, x_n) \in \mathbb{K}^n.$$

This operator can be seen as a Fourier multiplier with symbol  $\|\cdot\|_{\mathbb{K}^n}^\alpha$ , i.e.

$$(D^\alpha f)(x) = \mathcal{F}^{-1}(\|\cdot\|_{\mathbb{K}^n}^\alpha \hat{f}), \quad \alpha > 0, x \in \mathbb{K}^n,$$

where we use the Fourier transform  $\mathcal{F}$ . We also refer to [DeV22, Definition 1.12] for a generalization for a graded  $\mathbb{K}$ -Lie group. Thus, paralleling the critical role of the Laplacian in physics and geometry, the Taibleson operator stands as a central figure in the investigation of non-archimedean analysis and in modeling natural phenomena with ultrametric spaces.

Our main result is the following theorem where  $\Sigma_\theta \stackrel{\text{def}}{=} \{z \in \mathbb{C}^* : |\arg(z)| < \theta\}$  (see Figure 1).

**Theorem 1.1** *Let  $\mathbb{K}$  be a non-archimedean local field and  $n \geq 1$  be an integer. Suppose that  $1 < p < \infty$ . The Taibleson operator  $D^\alpha$  admits a bounded  $H^\infty(\Sigma_\theta)$  functional calculus for any angle  $\theta > 0$  on the Banach space  $L^p(\mathbb{K}^n)$ , as well as a Hörmander  $\text{Hör}_2^s(\mathbb{R}_+^*)$  functional calculus for any  $s > \frac{3}{2}$ .*

The first part of our result says that

$$(1.4) \quad \|f(D^\alpha)\|_{L^p(\mathbb{K}^n) \rightarrow L^p(\mathbb{K}^n)} \lesssim \|f\|_{H^\infty(\Sigma_\theta)}$$

for any suitable function  $f$  of the algebra  $H_0^\infty(\Sigma_\theta)$  of all bounded holomorphic functions  $f: \Sigma_\theta \rightarrow \mathbb{C}$  which are «regularly decaying» at 0 and at  $\infty$ , i.e. for which there exist two positive real numbers  $s, K > 0$  such that

$$(1.5) \quad |f(z)| \leq K \max\{|z|^s, |z|^{-s}\}, \quad z \in \Sigma_\theta.$$

The operator  $f(D^\alpha)$  is defined by integrating on the boundary of a «sector», see (2.10). It is worth noting that the Laplacian  $-\Delta$  also admits an  $H^\infty(\Sigma_\theta)$  functional calculus for any  $\theta > 0$

on the Banach space  $L^p(\mathbb{R}^n)$  or even on any Bochner space  $L^p(\mathbb{R}^n, X)$  for any UMD Banach space  $X$  by [HvNVW23, Theorem 10.2.25 p. 391]. Moreover, it admits a Hörmander Hör $_2^s(\mathbb{R}_+^*)$  functional calculus of order  $s > \frac{n}{2}$  on the scalar  $L^p(\mathbb{R}^n)$  space (which is a sharp condition for a parameter  $s$  valid for all  $1 < p < \infty$ ).

Actually, we prove a vector-valued generalization, replacing the scalar  $L^p$ -space  $L^p(\mathbb{R}^n)$  by any Bochner space  $L^p(\mathbb{R}^n, X)$  where  $X$  is a UMD Banach function space. The result on the  $H^\infty(\Sigma_\theta)$  functional calculus is sharp. We refer also to our recent paper [AHKP24], where a similar result is proved for the harmonic oscillator on the Moyal-Groenewold plane.

**Application to maximal regularity of evolution equations** Our study establishes the maximal regularity of the Taibleson operator. We will outline the broad context of this concept and direct readers to [Den21], [KuW04], [HvNVW23] (see also [Are04]) and references therein for a comprehensive discussion of this classical topic that has received considerable attention.

If  $A$  is a closed densely defined linear operator on a Banach space  $X$ , we can consider the following inhomogeneous autonomous Cauchy problem

$$(1.6) \quad \begin{cases} y'(t) + A(y(t)) = f(t), & t \in ]0, T[ \\ y(0) = x_0 \end{cases}$$

where  $f: ]0, T[ \rightarrow X$  is a function whose class belongs to the Banach space  $L^1(]0, T[, X)$ , with  $0 < T < \infty$  and  $x_0 \in X$ . If  $-A$  is the generator of a strongly continuous semigroup  $(T_t)_{t \geq 0}$  on a Banach space  $X$  then by [HvNVW23, Theorem G.3.2 p. 531] or [ABHN11, Proposition 3.1.16 p. 118] the Cauchy problem (1.6) admits a unique *mild* solution given by

$$(1.7) \quad y(t) = T_t(x_0) + \int_0^t T_{t-s}(f(s)) ds, \quad 0 \leq t < T.$$

Suppose that  $1 < p < \infty$ . If the right-hand side  $f$  of (1.6) belongs to the Banach space  $L^p(]0, T[, X)$ , «optimal regularity» would imply that both  $y'$  and  $Ay$  should also belong to the space  $L^p(]0, T[, X)$ . We are led to say, following [HvNVW23, Definition 17.2.4 p. 577] and [HvNVW23, Proposition 17.2.11 p. 581], that the operator  $A$  has maximal  $L^p$ -regularity on  $]0, T[$  if for any vector-valued function  $f \in L^p(]0, T[, X)$  the mild solution of the Cauchy problem (1.6) is a «strong  $L^p$ -solution», i.e. satisfies the two following conditions:

1.  $y$  takes values in  $\text{dom } A$  almost everywhere,
2.  $Ay \in L^p(]0, T[, X)$ .

A classical result of Dore [HvNVW23, Theorem 17.2.15 p. 586] says that this property implies that the operator  $-A$  generates an analytic semigroup on  $X$ . Moreover, this property is independent of  $1 < p < \infty$  by [HvNVW23, Theorem 17.2.31 p. 604].

A consequence of maximal  $L^p$ -regularity is the «a priori» estimate [HvNVW23, p. 577]

$$(1.8) \quad \|y'\|_{L^p(]0, T[, X)} + \|Ay\|_{L^p(]0, T[, X)} \leq C \|f\|_{L^p(]0, T[, X)},$$

where  $C \geq 0$  denotes a constant which is independent of  $f$ . Using linearization techniques, this estimate allows an effective approach to quasilinear problems via the contraction mapping principle. For a detailed description of this technique, we refer to [HvNVW23, Chapter 18].

It is known [HvNVW23, Corollary 17.3.6 p. 631] (combined with [HvNVW23, Lemma 17.2.16 p. 590]) that an operator admitting a bounded  $H^\infty(\Sigma_\theta)$  functional calculus of angle  $\theta < \frac{\pi}{2}$  on a UMD Banach space  $X$  has maximal  $L^p$ -regularity on  $]0, T[$ . Consequently, our main result

(Theorem 1.1) implies that the Taibleson operator  $D^\alpha$  defined in (1.2) has maximal  $L^p$ -regularity on  $]0, T[$  on the Banach space  $L^q(\mathbb{K}^n)$  for any  $1 < p, q < \infty$ . It is interesting to note that the equation  $y'(t) + D^\alpha(y(t)) = f(t)$  is the «master equation» of some concrete models, see [KKZ18, p. 117] and [ABO03].

**Approach of the proof of Theorem 1.1** Recall that the opposite of the Taibleson operator generates a Markovian semigroup  $(T_t)_{t \geq 0}$  on  $L^\infty(\mathbb{K}^n)$ . A fairly general result (Theorem 2.4) concerning generators of strongly continuous semigroups of positive contractions acting on  $L^p$ -spaces shows that the Taibleson operator admits a bounded  $H^\infty(\Sigma_\theta)$  functional calculus on the Banach space  $L^p(\mathbb{K}^n)$  for any angle  $\theta > \frac{\pi}{2}$ .

According to a classical result (Theorem 2.6), it is sufficient to demonstrate that the semigroup is  $R$ -analytic at angle  $\frac{\pi}{2}$  as this reduces the angle required for the functional calculus. This probabilistic condition is stronger than the property of bounded analyticity and, roughly speaking, means that the family  $(T_t)_{t > 0}$  can be extended to a bounded holomorphic family  $(T_z)_{\operatorname{Re} z > 0}$  which is « $R$ -bounded» on each sector  $\Sigma_\theta$  for any  $\theta < \frac{\pi}{2}$ . We refer to (2.7) for the definition of  $R$ -boundedness.

We then establish in Theorem 5.4 an  $R$ -boundedness result for a family of convolution operators on Vilenkin groups. This result is relevant since the additive group  $(\mathbb{K}^n, +)$  of  $\mathbb{K}^n$  is an abelian locally compact Vilenkin group, and since each operator  $T_z$  is a convolution operator, relies on a maximal inequality and is of independent interest.

To apply this result, it is necessary to estimate the  $L^1$ -norm of the kernel  $K_z$  of the convolution operator  $T_z$ , which leads to the estimate

$$(1.9) \quad \|K_z\|_{L^1(\mathbb{K}^n)} \lesssim_q \frac{|z|}{\operatorname{Re} z}.$$

with sharp contrast with the well-known analogue estimate [Are04, p. 153] (or [HvNVW18, p. 543]) saying that

$$\|h_z\|_{L^1(\mathbb{R}^n)} \leq \left( \frac{|z|}{\operatorname{Re} z} \right)^{\frac{n}{2}}, \quad \operatorname{Re} z > 0$$

of the kernel  $h_z$  of the heat semigroup on  $\mathbb{R}^n$ . This result of  $R$ -analyticity, enables us to obtain the boundedness of the functional calculus for any angle  $\theta > 0$ . For the Hörmander functional calculus, one only needs to check the conditions of the abstract result [KrW18, Theorem 7.1 (1) p. 424] (see Proposition 2.7).

**Locally compact Vilenkin groups** Initially, the groups investigated by Vilenkin in [Vil63] were the zero-dimensional compact *abelian* groups satisfying the second axiom of countability. These groups can be described as being the topological groups such that there exists a strictly decreasing sequence  $(G_n)_{n \geq 0}$  of compact open subgroups with  $\cup_{n \geq 0} G_n = G$  and  $\cap_{n \geq 0} G_n = \{0\}$ . These groups are precisely the infinite metrizable totally disconnected compact abelian groups. Examples of such groups include the Cantor group and countable products of finite abelian groups, see Example 3.6.

Nowadays, the class of locally compact abelian groups such that there exists a strictly decreasing sequence  $(G_n)_{n \in \mathbb{Z}}$  of compact open subgroups such that  $\cup_{n \in \mathbb{Z}} G_n = G$  and  $\cap_{n \in \mathbb{Z}} G_n = \{0\}$  is the class of locally compact Vilenkin groups, terminology introduced by Onneweer, see [SBSW15a, Chap. 7]. Nevertheless this class of groups was already considered by some other authors before. Indeed, we note that in the classical book [EdG77, p. 59] the locally compact Vilenkin groups are called «groups having a suitable family of compact open subgroups» and

in the book [AVDR81] the locally compact Vilenkin groups are called «locally compact 0-dimensional topologically periodic abelian groups with second axiom of countability».

Our harmonic analysis results, which we establish to prove our main result, build upon previous research efforts aimed at advancing analysis on these locally compact groups. For instance, the dyadic analysis on the Cantor group (and similar groups) is well-developed, see [SWS90], [SBSW15a] and [SBSW15b].

In [Wei07], the almost everywhere convergence of Banach space-valued Vilenkin-Fourier series is investigated. In [KaV22], Hardy inequalities were explored. Multiplier theory was investigated in [OnQ89]. Most recently, in [DHW24], the convergence of Fejér means on some noncommutative Vilenkin groups was examined. Actually, our results are valid for a slightly broader class of groups, which we refer to as Spector-Vilenkin groups because they were studied by Spector in [Spe70].

It is interesting to note that compact abelian Vilenkin groups frequently emerge when attempting to prove a property for all locally compact groups by reducing the problem to specific types of groups through case-by-case reasoning. See, for example [Arh12] and [ArK23, Section 7.2] for illustrations.

**Structure of the paper** Section 2 provides background information on local fields, operator theory, functional calculus, and harmonic analysis. Section 3 explores locally compact Spector-Vilenkin groups and their radial functions. In particular, we introduce the notion of least radially decreasing majorant. In Section 4, we establish a maximal inequality related to averaging operators on locally compact Spector-Vilenkin groups. This result plays a crucial role in Section 5, where Proposition 5.2 sets forth a sufficient condition for the  $R$ -boundedness of a family of operators acting on the  $L^p$ -space  $L^p(G)$  of a Vilenkin group, dominated by the maximal operator. In Theorem 5.3, we deduce an  $R$ -boundedness result on some specific family of convolution operators. Section 6 details our proof of the estimate (1.9) on the kernel. Finally, we present our principal result in Theorem 6.8.

## 2 Preliminaries

We begin by providing information and background on local fields.

**Local fields** Let  $\mathbb{K}$  be a commutative locally compact field. For any  $x \in \mathbb{K}^*$ , we denote by  $|x|$  the module of the automorphism  $\mathbb{K} \rightarrow \mathbb{K}$ ,  $y \mapsto xy$  of the additive group of  $\mathbb{K}$ . In addition, we let  $|0| = 0$ . By [Bou04, Proposition 12 VII.21], the function  $\mathbb{K} \rightarrow \mathbb{R}^+$ ,  $x \mapsto |x|$  is continuous and for any  $x, y \in \mathbb{K}$  we have  $|xy| = |x||y|$ .

Recall that a local field is a non-discrete locally compact field. We warn the reader that the terminology «local field» does not always have the same meaning in the literature. Several definitions coexist. In this case, by [Bou04, Proposition 13 VII.22] the subsets  $\{x \in \mathbb{K} : |x| \leq K\}$  for  $K > 0$  define a fundamental system of compact neighbourhoods of 0 in  $\mathbb{K}$ . Moreover, by [Bou04, Corollary VII.22], a local field is a second-countable topological space.

If for any  $x, y \in \mathbb{K}$  we have  $|x + y| \leq \max\{|x|, |y|\}$ , we say [RaV99, p. 137] that  $\mathbb{K}$  is non-archimedean and that  $|\cdot|$  is an ultrametric. In this case, by [RaV99, Lemma 4.14 p. 145] the set

$$(2.1) \quad \mathcal{O} \stackrel{\text{def}}{=} \{x \in \mathbb{K} : |x| \leq 1\}$$

is the unique maximal compact subring of  $\mathbb{K}$  and is called ring of integers [Tai75, p. 6]. By [RaV99, Lemma 4.15 p. 145], the set

$$(2.2) \quad \mathfrak{p} \stackrel{\text{def}}{=} \{x \in \mathbb{K} : |x| < 1\}$$

is the unique maximal ideal of  $\mathcal{O}$  and we have  $\mathfrak{p} = \pi\mathcal{O}$  for some element  $\pi$  of  $\mathfrak{p}$ . The same reference states that the quotient  $\mathcal{O}/\mathfrak{p}$  is isomorphic to a finite field with cardinal  $q = p^m$  for some prime  $p$  and some positive integer  $m \geq 1$  and is called the residue field. For any  $x \in \mathbb{K}^* = \mathbb{K} - \{0\}$ , by [RaV99, p. 146] we have  $|x| = q^k$  for some  $k \in \mathbb{Z}$ . That means that  $|\mathbb{K}^*|$  is a discrete subgroup of  $\mathbb{R}_*^+$ .

**Example 2.1 (fields of  $q$ -adic numbers)** Here  $q$  will denote a prime number. Recall that the field  $\mathbb{Q}_q$  of  $q$ -adic numbers is defined as the completion of the field  $\mathbb{Q}$  of rational numbers with respect to the  $q$ -adic norm  $|\cdot|_q$  which is defined by

$$(2.3) \quad |x|_q \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } x = 0 \\ \frac{1}{q^k} & \text{if } x = q^k \frac{a}{b} \end{cases}$$

where  $a$  and  $b$  are integers coprime with  $q$ . This field is of characteristic 0. We refer to the books [Gou20] and [Rob00] for more information. A particular case of [RaV99, Lemma 4.17 p. 146] (or [AKS10, p. 17], [Fol16, Proposition 2.8 p. 39]) says that any non-zero  $q$ -adic number  $x$  has a unique expansion of the form

$$(2.4) \quad x = q^\gamma \sum_{j=0}^{\infty} x_j q^j$$

where  $\gamma \in \mathbb{Z}$  and  $x_j \in \{0, 1, \dots, q-1\}$ . Following [Rob00, p. 40], the fractional part  $\{x\}_q$  of an element  $x$  of  $\mathbb{Q}_q$  is the element of  $\mathbb{Q}$  defined by

$$\{x\}_q \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } x = 0 \text{ or } \gamma \geq 0 \\ q^\gamma \sum_{j=0}^{|\gamma|-1} x_j q^j & \text{if } \gamma < 0 \end{cases}.$$

The ring of integers  $\mathcal{O}$  is equal to the closure  $\mathbb{Z}_q$  of the subspace  $\mathbb{Z}$  in the space  $\mathbb{Q}_q$  and the unique maximal ideal  $\mathfrak{p}$  of  $\mathcal{O}$  is equal to  $q\mathbb{Z}_q$ . The residue field is  $\mathcal{O}/\mathfrak{p} = \mathbb{Z}_q/q\mathbb{Z}_q = \mathbb{Z}/q\mathbb{Z}$ . Finally, recall that each  $x \in \mathbb{Z}_q$  can be uniquely written as  $x = \sum_{j=0}^{\infty} x_j q^j$  with  $x_j \in \{0, 1, \dots, q-1\}$ .

**Example 2.2 (fields of formal Laurent series over finite fields)** Let  $p$  be a prime number. If  $n \geq 1$  is an integer, we denote by  $\mathbb{F}_{p^n}$  the finite field with cardinal  $p^n$ . The field

$$\mathbb{F}_{p^n}((X)) \stackrel{\text{def}}{=} \left\{ \sum_{j \geq m} c_j X^j : m \in \mathbb{Z}, c_j \in \mathbb{F}_{p^n} \right\},$$

of formal Laurent series over the finite field  $\mathbb{F}_{p^n}$  is a non-archimedean local field of characteristic  $p$ . Recall that the product of  $\sum_{j \geq n} b_j X^j$  with  $\sum_{j \geq m} c_j X^j$  is given by the formal Laurent series  $\sum_j (\sum_k b_k c_{j-k}) X^j$ .

If  $x = \sum_{j \geq m} c_j X^j$  is a non-zero element of the field  $\mathbb{F}_{p^n}((X))$  and if  $m$  is the smallest integer such that  $c_m \neq 0$ , we have  $|x| = p^{-nm}$ .

Its ring of integers  $\mathcal{O}$  is the ring  $\mathbb{F}_{p^n}[[X]] = \left\{ \sum_{j \geq 0} c_j X^j : c_j \in \mathbb{F}_{p^n} \right\}$  of formal power series over the field  $\mathbb{F}_{p^n}$ . The unique maximal ideal of the ring  $\mathcal{O}$  is  $\mathfrak{p} = X\mathbb{F}_{p^n}[[X]]$  and the residue field  $\mathcal{O}/\mathfrak{p}$  is isomorphic to  $\mathbb{F}_{p^n}$ .

Indeed, it is known, e.g. [RaV99, Theorem 4.12 p. 140], that non-archimedean local fields are precisely the fields  $\mathbb{F}_{p^n}((X))$  of formal Laurent series over the finite fields of cardinal  $p^n$  and the finite extensions of the fields  $\mathbb{Q}_q$  of  $q$ -adic numbers where  $q$  is a prime number.

**Pontryagin duality** Let  $\mathbb{K}$  be a local field. Let  $\chi: \mathbb{K} \rightarrow \mathbb{C}$  be a non-constant character of the additive group of  $\mathbb{K}$ . Let  $n \geq 1$  be an integer. For any  $x, y \in \mathbb{K}^n$ , we let  $x \cdot y \stackrel{\text{def}}{=} x_1 y_1 + \cdots + x_n y_n$ . We write  $\chi_x(y) \stackrel{\text{def}}{=} \chi(x \cdot y) = \chi(x_1 y_1 + \cdots + x_n y_n)$ . For each  $x \in \mathbb{K}^n$ , the map  $\chi_x: \mathbb{K}^n \rightarrow \mathbb{C}$  is a character of  $\mathbb{K}^n$ . Then by [Bou67] the map  $\mathbb{K}^n \rightarrow \widehat{\mathbb{K}^n}$ ,  $x \mapsto \chi_x$  is a topological isomorphism of the additive group of  $\mathbb{K}^n$  onto its Pontryagin dual  $\widehat{\mathbb{K}^n}$ .

**Example 2.3** Here, we use the notations of Example 2.1. Consider the map  $\chi_q: \mathbb{Q}_q \rightarrow \mathbb{C}$  defined by

$$(2.5) \quad \chi_q(x) \stackrel{\text{def}}{=} e^{2\pi i \{x\}_q}, \quad x \in \mathbb{Q}_q.$$

By [Fol16, pp. 99-100], the map  $\chi_q$  is a non-zero character of the additive group of  $\mathbb{Q}_q$  with kernel  $\mathbb{Z}_q$ . In particular, we have the equality  $\chi_q(x+y) = \chi_q(x)\chi_q(y)$  for any  $x, y \in \mathbb{Q}_q$ .

**Fourier transform** In the sequel, we will always consider the Haar measure  $\mu$  on the additive group of  $\mathbb{K}^n$  normalized by the condition  $\mu(\mathcal{O}^n) = 1$ . Let again  $\chi: \mathbb{K} \rightarrow \mathbb{C}$  be a fixed non-constant character. The Fourier transform of a complex function  $f \in L^1(\mathbb{K}^n)$  is defined by

$$(2.6) \quad \hat{f}(x) \stackrel{\text{def}}{=} \int_{\mathbb{K}^n} \chi(-x \cdot y) f(y) d\mu(y).$$

In the sequel, we will write  $dy$  for  $d\mu(y)$ .

We continue with background on operator theory.

**$R$ -boundedness** Suppose that  $1 < p < \infty$ . Following [HvNVW18, Definition 8.1.1, Remark 8.1.2 p. 165], we say that a set  $\mathcal{F}$  of bounded operators on a Banach space  $X$  is  $R$ -bounded provided that there exists a constant  $C \geq 0$  such that for any finite families  $T_1, \dots, T_n$  in  $\mathcal{F}$  and  $x_1, \dots, x_n$  in  $X$ , we have

$$(2.7) \quad \left\| \sum_{k=1}^n \varepsilon_k \otimes T_k(x_k) \right\|_{L^p(\Omega, X)} \leq C \left\| \sum_{k=1}^n \varepsilon_k \otimes x_k \right\|_{L^p(\Omega, X)}$$

where  $(\varepsilon_i)_{i \geq 1}$  is a sequence of independent Rademacher variables on some probability space  $\Omega$ . This property is independent of  $p$ . The best constant in (2.7) is denoted by  $\mathcal{R}_p(\mathcal{F})$  and is referred to as the  $R$ -bound of the family  $\mathcal{F}$ . If  $X$  is a complex Banach lattice with finite cotype, a classical theorem of Maurey asserts that we have a uniform equivalence

$$(2.8) \quad \left\| \sum_{k=1}^n \varepsilon_k \otimes x_k \right\|_{L^p(\Omega, X)} \approx \left\| \left( \sum_{k=1}^n |x_k|^2 \right)^{\frac{1}{2}} \right\|_X$$

for any vector  $x_1, \dots, x_n$  of  $X$ , see e.g. [DJT95, Theorem 16.18 p. 338] (or [HvNVW18, Definition 8.1.1 p. 165] for the real case).



**$H^\infty$  functional calculus** For detailed information we refer the reader to the books [HvNVW18], [HvNVW23], [Haa06] to the survey papers [Are04], [KuW04] and to references therein. Recall the main definitions. We will use the notation  $\Sigma_\theta \stackrel{\text{def}}{=} \{z \in \mathbb{C}^* : |\arg(z)| < \theta\}$  for any  $\theta > 0$ , see Figure 1. Let  $A : \text{dom } A \subset X \rightarrow X$  be a closed linear operator on a Banach space  $X$  with dense domain. We say that  $A$  is a sectorial operator of type  $\mu \in (0, \pi)$  if its spectrum  $\sigma(A)$  is a subset of the closed sector  $\overline{\Sigma_\mu}$  and if for any angle  $\nu \in (\mu, \pi)$ , the set

$$(2.9) \quad \left\{ zR(z, A) : z \in \mathbb{C} \setminus \overline{\Sigma_\nu} \right\}$$

is bounded in  $B(X)$ , where  $R(z, A) \stackrel{\text{def}}{=} (z\text{Id} - A)^{-1}$  is the resolvent operator. We let

$$\omega(A) \stackrel{\text{def}}{=} \inf\{\omega \in (0, \pi) : A \text{ is sectorial of type } \mu\}.$$




Figure 1: sector  $\Sigma_\theta$  of angle  $\theta$

For any angle  $\theta \in (0, \pi)$ , we consider the algebra  $H_0^\infty(\Sigma_\theta)$  of all bounded holomorphic functions  $f : \Sigma_\theta \rightarrow \mathbb{C}$  for which there exist two positive real numbers  $s, K > 0$  such that (1.5) holds. Let  $0 < \mu < \theta < \pi$  and consider a function  $f \in H_0^\infty(\Sigma_\theta)$ . Then for any  $\nu \in (\mu, \theta)$ , we introduce the operator

$$(2.10) \quad f(A) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\partial\Sigma_\nu} f(z)R(z, A) dz,$$

where the boundary  $\partial\Sigma_\nu$  is oriented counterclockwise. The sectoriality condition ensures that this integral is absolutely convergent and defines a bounded operator on  $X$ . Using Cauchy's theorem, it is possible to show that this definition does not depend on the choice of the chosen angle  $\nu$ . The resulting map  $H_0^\infty(\Sigma_\theta) \rightarrow B(X)$ ,  $f \mapsto f(A)$  is an algebra homomorphism which is consistent with the usual functional calculus for rational functions.

We say that  $A$  admits a bounded  $H^\infty(\Sigma_\theta)$  functional calculus if the latter homomorphism is bounded, i.e. if there exists a constant  $K \geq 0$  such that

$$(2.11) \quad \|f(A)\|_{X \rightarrow X} \leq K \|f\|_{H^\infty(\Sigma_\theta)}, \quad f \in H_0^\infty(\Sigma_\theta).$$

If  $A$  has dense range and admits a bounded  $H^\infty(\Sigma_\theta)$  functional calculus, then the previous homomorphism naturally extends to a bounded homomorphism  $f \mapsto f(A)$  from the algebra  $H^\infty(\Sigma_\theta)$  into the algebra  $B(X)$ . We let

$$\omega_{H^\infty}(A) \stackrel{\text{def}}{=} \inf\{\omega \in (\omega(A), \pi) : A \text{ admits a bounded } H^\infty(\Sigma_\omega) \text{ functional calculus}\}.$$

The following result is [HvNVW18, Theorem 10.7.12 p. 462].

**Theorem 2.4** *Let  $\Omega$  be a measure space and let  $X$  be a UMD Banach space. Suppose that  $1 < p < \infty$ . Let  $-A$  be the infinitesimal generator of a strongly continuous semigroup  $(T_t)_{t \geq 0}$  of positive contractions on  $L^p(\Omega)$ . Then (the closure of)  $A \otimes \text{Id}_X$  admits a bounded  $H^\infty(\Sigma_\theta)$  functional calculus on the Bochner space  $L^p(\Omega, X)$  for any angle  $\theta > \frac{\pi}{2}$ .*

The next result is a particular case of [HvNVW18, Proposition 10.3.3 p. 400] and connects  $R$ -sectoriality and  $R$ -analyticity. Recall that an operator  $A$  is called  $R$ -sectorial of type  $\sigma$  if it is sectorial of type  $\sigma$  and if for any angle  $\theta \in (\sigma, \pi)$  the set

$$(2.12) \quad \{zR(z, A) : z \notin \overline{\Sigma_\theta}\}$$

is  $R$ -bounded. We let  $\omega_R(A) \stackrel{\text{def}}{=} \inf\{\theta \in (\omega(A), \pi) : \text{the subset of (2.12) is } R\text{-bounded}\}$ .

**Theorem 2.5** *Let  $(T_t)_{t \geq 0}$  be a strongly continuous bounded analytic semigroup with infinitesimal generator  $-A$ . If  $(T_z)_{z \in \Sigma_\omega}$  is  $R$ -bounded for some angle  $\omega \in (0, \frac{\pi}{2})$ , then  $A$  is  $R$ -sectorial and*

$$\omega_R(A) = \frac{\pi}{2} - \sup\{\omega \in (0, \frac{\pi}{2}) : (T_z)_{z \in \Sigma_\omega} \text{ is } R\text{-bounded}\}.$$

We will use the following result [HvNVW18, Corollary 10.4.10 p. 422], [KaW01, Theorem 5.3 p. 335]. Every UMD Banach space has the triangular contraction  $(\Delta)$  property by [HvNVW18, Corollary 7.5.9 p. 137].

**Theorem 2.6** *Let  $X$  be a Banach space with property  $(\Delta)$ . Let  $A$  be a sectorial operator on  $X$  with a bounded  $H^\infty$  functional calculus. Then  $A$  is  $R$ -sectorial and one has  $\omega_R(A) = \omega_{H^\infty}(A)$ .*

**Hörmander functional calculus** If  $s > 0$ , using the usual Sobolev norm

$$(2.13) \quad \|f\|_{W^{s,2}(\mathbb{R})} \stackrel{\text{def}}{=} \left\| \left[ \hat{f}(t)(1+t^2)^{\frac{s}{2}} \right] \right\|_{L^2(\mathbb{R})}$$

and the inhomogeneous Sobolev space  $W^{s,2}(\mathbb{R}) \stackrel{\text{def}}{=} \{f \in \mathcal{S}'(\mathbb{R}^d) : \|f\|_{W^{s,2}(\mathbb{R})} < \infty\}$ , we introduce the norm

$$(2.14) \quad \|f\|_{\text{Hör}_2^s(\mathbb{R}_+^*)} \stackrel{\text{def}}{=} \sup_{t>0} \|x \mapsto \eta(x) \cdot f(tx)\|_{W^{s,2}(\mathbb{R})} < \infty,$$

where  $\eta: ]0, \infty[ \rightarrow \mathbb{R}$  is any non-zero function of class  $C^\infty$  with compact support. The following is a slight variant of [AHKP24, Proposition 10.7] and is a direct consequence of [KrW18, Theorem 7.1 (1) p. 424]. Note hereby that it is well-known that on a reflexive Banach space  $X$ , a sectorial operator  $A$  can be turned into one with dense range by using the splitting  $X = \text{Ker } A \oplus \overline{\text{Ran } A}$  [HvNVW18, p. 361], and that a complex Banach lattice with finite cotype has Pisier's property  $(\alpha)$  by [HvNVW18, Theorem 7.5.20 p. 145].

**Proposition 2.7** *Suppose  $s > 0$ . Let  $-A$  be the generator of a bounded holomorphic semigroup  $(T_z)_{\text{Re } z > 0}$  acting on a reflexive complex Banach lattice  $X$  with finite cotype. Suppose that  $A$  has a bounded  $H^\infty(\Sigma_\omega)$  functional calculus for some angle  $\omega \in (0, \pi)$ . Assume in addition that the family*

$$(2.15) \quad \{(\cos \arg z)^\beta T_z : \text{Re } z > 0\}$$

*of operators is  $R$ -bounded over the space  $X$ . Then the operator  $A$  admits a bounded  $\text{Hör}_2^s(\mathbb{R}_+^*)$  Hörmander calculus of any order  $s > \beta + \frac{1}{2}$ , i.e.*

$$(2.16) \quad \|f(A)\|_{X \rightarrow X} \lesssim \|f\|_{\text{Hör}_2^s(\mathbb{R}_+^*)}, \quad f \in \text{Hör}_2^s(\mathbb{R}_+^*).$$

**Conditional expectations** Suppose that  $(\Omega, \mathcal{A}, \mu)$  is a measure space and that  $(\Omega_j)_{j \in J}$  is an at most countable and measurable partition of  $\Omega$ , i.e. a partition of  $\Omega$  such that  $\Omega_j \in \mathcal{A}$  for each  $j \in J$ . We suppose that each  $\Omega_j$  is of finite measure  $\mu(\Omega_j) > 0$  for any  $j \in J$ , then a slight generalization [Ste70, p. 89] of [AbA02, Exercise 4 p. 227] says that for any  $f \in L^1(\Omega, \mathcal{A}, \mu) \cap L^\infty(\Omega, \mathcal{A}, \mu)$  we have

$$(2.17) \quad \mathbb{E}(f|\mathcal{A}) = \sum_{j \in J} \left[ \frac{1}{\mu(\Omega_j)} \int_{\Omega_j} f \, d\mu \right] 1_{\Omega_j}.$$

### 3 Locally compact Spector-Vilenkin groups and radial functions

We begin by recalling some general topology concepts. Following [En89, p. 360], a topological space  $Y$  is said to be hereditarily disconnected if its connected components are reduced to singletons. Unfortunately in the literature, for instance [HeR79, p. 11], hereditarily disconnected spaces are sometimes called totally disconnected spaces. Following [En89, p. 360], a topological space  $Y$  is called zero-dimensional if  $Y$  is a non-empty  $T_1$ -space and has a basis consisting of open-and-closed subsets. Finally, a space is said to be totally disconnected if the quasi-component of any point  $x \in Y$  consists of the point  $x$  alone. By [En89, p. 369], every zero-dimensional space is totally disconnected and every totally disconnected space is hereditarily disconnected. Conversely, by [En89, Theorem 6.2.9 p. 362], every hereditarily disconnected locally compact space is zero-dimensional. So these notions are equivalent for the class of locally compact spaces and we will use the terminology «totally disconnected».

We will use the following elementary result [En89, Corollary 3.1.5 p. 124].

**Proposition 3.1** *Let  $U$  be an open subset of a compact topological space  $X$ . If a family  $(F_k)_{k \in I}$  of closed subsets of  $X$  satisfies  $\bigcap_{k \in I} F_k \subset U$ , then there exists a finite subset  $\{k_1, \dots, k_l\}$  of  $I$  such that  $F_1 \cap \dots \cap F_{k_l} \subset U$ .*

Now, we define the class of groups on which we will focus in this paper inspired by [Vil63, p. 3] and [Spe70, p. 61].

**Definition 3.2** *A Hausdorff topological group  $G$  is said to be a locally compact Spector-Vilenkin group if there exists a strictly decreasing sequence  $(G_k)_{k \in I}$  with  $I = \mathbb{Z}$ ,  $I = \mathbb{N}$  or  $I = -\mathbb{N}$  of compact open subgroups of  $G$  such that  $\bigcup_{k \in I} G_k = G$  and satisfying*

$$(3.1) \quad \bigcap_{k \in I} G_k = \{e\}.$$

Note the following result.

**Lemma 3.3** *Let  $G$  be a Hausdorff topological group admitting a strictly decreasing sequence  $(G_k)_{k \in I}$  with  $I = \mathbb{Z}$ ,  $I = \mathbb{N}$  or  $I = -\mathbb{N}$  of compact open subgroups such that  $\bigcap_{k \in I} G_k = \{e\}$ . Then the family  $(G_k)_{k \in I}$  forms a basis of neighborhoods at  $e$ .*

*Proof* : Each  $G_k$  contains  $e$  and is open, so each  $G_k$  is a neighborhood at  $e$ . It remains to show that any neighborhood of  $e$  contains one of the  $G_k$ . Let  $U$  be such a neighborhood. Clearly,  $U$  can be supposed to be open, and if replaced by the smaller neighborhood  $U \cap G_0$  if necessary, we can also suppose that  $U \subset G_0$ .

If  $I = -\mathbb{N}$ , then there is nothing to prove since  $G_0 = \{e\}$  and consequently  $U = G_0$ .

If  $I = \mathbb{N}$  or  $I = \mathbb{Z}$  we have  $\bigcap_{k \in I} G_k = \{e\} \subset U$ . As  $G_0$  is compact, we can use Proposition 3.1 to deduce that there exists a finite subset  $\{k_1, \dots, k_l\}$  of  $I$  such that  $G_{k_1} \cap \dots \cap G_{k_l} \subset U$ . This means that  $G_{\max\{k_1, \dots, k_l\}} \subset U$ . ■

Let  $G$  be a locally compact Spector-Vilenkin group with associated sequence  $(G_k)_{k \in I}$ . By Lemma 3.3, the family  $(G_k)_{k \in I}$  forms a basis of neighborhoods at  $e$ . Consequently, the topology on  $G$  is the unique «subgroup topology» defined by the family  $(G_k)_{k \in I}$  as defined in [Bou98, Example p. 223] or [Hal50].

Suppose  $I = \mathbb{Z}$  or  $I = \mathbb{N}$ . Since  $G_{k+1}$  is open in  $G_k$  for each  $k \in I$ , the homogeneous space  $G_k/G_{k+1}$  is discrete by [Bou98, Proposition 14 p. 231]. Furthermore  $G_k$  is compact, so the homogeneous space  $G_k/G_{k+1}$  is also compact by [Bou98, Proposition 11 p. 257]. Consequently, each  $G_k/G_{k+1}$  is discrete and compact, hence finite. For any integer  $k \in I$ , let  $d_k$  be the order of the space  $G_k/G_{k+1}$ . Then  $d_k \geq 2$ . If we also have

$$\sup_{k \in I} |G_k/G_{k+1}| < \infty,$$

we say that  $G$  is a bounded-order locally compact Spector-Vilenkin group. If  $|G_k/G_{k+1}|$  is constant, we say that  $G$  is a constant-order locally compact Spector-Vilenkin group. If  $I = -\mathbb{N}$ , we leave it to the reader to adapt the previous discussion by replacing  $G_{k+1}$  by  $G_{k-1}$ .

It is well-known that a locally compact group  $G$  is totally disconnected if and only if the compact open subgroups form a basis of neighbourhoods of the identity  $e$ , see e.g. [HoM13, Theorem 1.34 p. 22] and [HeR79, Theorem 7.7 p. 62]. In other words, this means that  $G$  has arbitrarily small compact open subgroups. If  $G$  is compact, we can suppose that the subgroups are normal, again by the cited results. Consequently, a locally compact Spector-Vilenkin group is totally disconnected. Moreover, such a group is infinite since the sequence of subgroups is strictly decreasing and metrizable by the Birkhoff-Kakutani theorem [En89, p. 437] which says that it is equivalent to saying that the group  $G$  is first countable.

Finally, if each subgroup  $G_k$  is *normal*, the locally compact Spector-Vilenkin group  $G$  is pro-discrete by [ArK23, Proposition 6.36 p. 119], and unimodular by [Bou04, Proposition 3 p. 20] which says that the existence of a compact neighborhood of  $e$  invariant by the inner automorphisms implies the unimodularity of the group.

**Example 3.4** If  $I = \mathbb{N}$ , we have a countable chain of strictly decreasing inclusions

$$\{e\} \subset \dots \subset G_k \subset \dots \subset G_1 \subset G_0 = G.$$

Since  $G = G_0$ , such a group is compact and we drop the word «locally». By [HoM13, Theorem 1.34 p. 22] and [HeR79, Theorem 7.7 p. 62], we can suppose that the groups  $G_k$  are normal. So, we recover the notion of compact Vilenkin group of [DeV22, Definition 2.1]. The assumption «profinite» is useless in [DeV22, Definition 2.1] since a topological group is compact and totally disconnected if and only if it is profinite by [Wil98, Corollary 1.2.4 p. 19].

**Example 3.5** The subclass of locally compact *abelian* Spector-Vilenkin groups with  $I = \mathbb{N}$  is the class of groups considered in the paper [Vil63]. These groups are precisely the infinite metrizable totally disconnected compact abelian groups (also stated in [Spe70, p. 61]). Indeed, Vilenkin assumes in [Vil63] the second axiom of countability instead of our weaker assumption of metrizability, but it is equivalent for compact groups by [DH16, Theorem 2.A.10 p. 15]. Note that the dual  $\hat{G}$  of such a group is discrete by [HeR79, Theorem 23.17 p. 362]. Moreover,  $\hat{G}$  is a torsion group (i.e. periodic) by [HeR79, Corollary 24.20 p. 383] and is countable by [HeR79, Theorem 24.15 p. 382].

**Example 3.6** For any integer  $m \geq 2$ , we denote by  $\mathbb{Z}/m\mathbb{Z}$  the cyclic group of order  $m$ . Let  $p = (p_0, p_1, \dots)$  be a sequence of natural numbers  $p_k$  with  $p_k \geq 2$ . By [Wil98, Corollary 1.2.4 p. 19], the group  $G = \prod_{k=0}^{\infty} \mathbb{Z}/p_k\mathbb{Z}$  equipped with the product topology is a totally disconnected compact abelian group which is of course infinite and metrizable since the countable product of metric spaces is metrizable by [En89, Theorem 4.2.2 p. 259]. So it is an example of a compact *abelian* Spector-Vilenkin group with  $I = \mathbb{N}$ . It is easy to define a suitable sequence  $(G_k)_{k \geq 0}$ .

**Example 3.7** Let  $q$  be a prime number. Then the group  $\mathbb{Z}_q$  defined in Example 2.1 is another compact abelian Spector-Vilenkin group under the topology induced by the one of  $\mathbb{Q}_q$ . Indeed, according to [DeE14, Proposition 13.1.8 p. 253], for each integer  $k \geq 0$

$$G_k \stackrel{\text{def}}{=} q^k \mathbb{Z}_q = \{x \in \mathbb{Q}_q : |x|_q \leq \frac{1}{q^k}\}$$

is a compact open subgroup of the additive group  $\mathbb{Q}_q$  and the sequence  $(G_k)_{k \geq 0}$  is a basis of neighborhoods at 0 in the additive group  $\mathbb{Q}_q$  and consequently also in the open subset  $\mathbb{Z}_q$ . Of course,  $\bigcup_{k \geq 0} G_k = G_0 = \mathbb{Z}_q$  and  $\bigcap_{k \geq 0} G_k = \{0\}$ . Moreover, the sequence  $(G_k)_{k \geq 0}$  is clearly decreasing. It is even strictly decreasing. Indeed, since  $|q^k|_q \stackrel{(2.3)}{=} \frac{1}{q^k}$  the element  $q^k$  belongs to  $G_k$  and does not belong to  $G_{k+1}$ .

**Example 3.8** If  $I = \mathbb{Z}$ , we «essentially» recover the notion of «locally compact Vilenkin group» of [DeV22, Definition 1.1]. We have a countable chain of strictly decreasing inclusions

$$\{e\} \subset \dots \subset G_{k+1} \subset G_k \subset \dots \subset G.$$

Indeed, the authors add the additional assumptions «unimodular» and «type I».

**Example 3.9** In the classical book [EdG77, p. 59], the locally compact *abelian* Spector-Vilenkin groups with  $I = \mathbb{Z}$  are called «groups having a suitable family of compact open subgroups». We also refer to the book [AVDR81], which appears to contain interesting information, but is, unfortunately, not available in English.

**Example 3.10** Our *bounded order* locally compact *abelian* Spector-Vilenkin groups with  $I = \mathbb{Z}$  coincide with the «locally compact Vilenkin groups» of [SBSW15a, Chap. 7, Definition 7.1 p. 248].

**Example 3.11** If  $n \geq 1$  and if  $\mathbb{K}$  is a non-archimedean local field then the additive group  $(\mathbb{K}^n, +)$  is a locally compact Vilenkin group. Recall that the quotient  $\mathcal{O}/\mathfrak{p}$  is isomorphic to a finite field with cardinal  $q$ . Then we can consider the family  $(G_k)_{k \in \mathbb{Z}}$  of subsets defined by

$$(3.2) \quad G_k \stackrel{\text{def}}{=} \{x \in \mathbb{K}^n : \|x\|_{\mathbb{K}^n} \leq \frac{1}{q^k}\},$$

where  $\|\cdot\|_{\mathbb{K}^n}$  is defined in (1.3). By [Tai75, p. 8], each  $G_k$  is a compact open subgroup of  $(\mathbb{K}^n, +)$ . In particular, if  $q$  is a prime number then the additive group  $(\mathbb{Q}_q, +)$  of  $q$ -adic numbers is a locally compact Vilenkin group.

**Remark 3.12** In [DHW24, p. 1], it is stated that any totally disconnected compact group is isomorphic to the complete direct product of some finite groups. It is false<sup>1</sup> since a topological group is compact and totally disconnected if and only if it is profinite by [Wil98, Corollary 1.2.4 p. 19].

---

1. G. Hong has been notified by email.

**Remark 3.13** It is stated without proof in [Spe70, p. 61], that the locally compact *abelian* Spector-Vilenkin groups coincide with the  $\sigma$ -compact infinite totally disconnected metrizable locally compact abelian groups having a compact open subgroup such that the quotient is with torsion.

**Example 3.14** If  $I = -\mathbb{N}$ , we have a countable chain of strictly decreasing inclusions

$$G \supset \dots G_{-k} \supset \dots \supset G_{-2} \supset G_{-1} \supset G_0 = \{e\}.$$

The following result is written in [Spe70, p. 61] without proof. We add a proof here for the convenience of the reader. Recall that a group  $G$  is said to be a torsion group if every element of  $G$  has finite order.

**Proposition 3.15** *The subclass of locally compact abelian Spector-Vilenkin groups with  $I = -\mathbb{N}$  is the class of torsion countable abelian discrete groups.*

*Proof* : Suppose first that  $G$  is a locally compact abelian Spector-Vilenkin group with  $I = -\mathbb{N}$ . As  $G_0 = \{e\}$  is open, the group  $G$  is discrete by [Bou98, Proposition 5 p. 227]. Moreover, for any  $k \in -\mathbb{N}$ ,  $G_k \cong G_k/G_0$  is a subgroup of finite order. Then as  $G = \bigcup_{k \in -\mathbb{N}} G_k$  is the countable union of finite sets,  $G$  is countable. Now let  $g \in G$  be arbitrary. Then there exists some  $k \in -\mathbb{N}$  such that  $g \in G_k$ . As  $G_k$  is finite,  $g$  is of finite order, hence  $G$  is torsion.

Conversely, suppose now that  $G$  is a torsion countable abelian discrete group. Let  $(g_k)_{k \in -\mathbb{N}}$  be any enumeration of the elements in  $G$ . Then let  $G_0 \stackrel{\text{def}}{=} \{e\}$ . For any  $k \in -\mathbb{N}$ , define inductively  $G_k \stackrel{\text{def}}{=} \langle g_1, \dots, g_{-n} \rangle$  as the subgroup generated by the first  $n$  elements of the group, where  $n$  is the minimal value such that  $G_k \neq G_{k+1}$ . Note that each  $G_k$  is finite (what allows us to define the sequence by recursion since  $G$  is infinite). Indeed, we have  $G_k = \{g_1^{m_1} \cdots g_{-n}^{m_n} : 0 \leq m_j \leq (\text{order of } g_j) - 1\}$ . Then  $|G_k| \leq \prod_{j=1}^n (\text{order of } g_j) < \infty$  since  $G$  is a torsion group. So we obtained a strictly increasing sequence  $\{e\} = G_0 \subset G_{-1} \subset G_{-2} \subset \dots \subset G_k \subset \dots \subset G$ . Moreover, for any element  $g \in G$ , there exists  $k \in -\mathbb{N}$  such that  $g = g_k$ . Hence  $G = \bigcup_{k \in -\mathbb{N}} G_k$ . Each  $G_k$  is finite, hence compact and, by discreteness of  $G$ , open. ■

**Example 3.16** Let  $q \neq 2$  be a prime number. By [DeV22, Example 1.4], the  $2n+1$ -dimensional Heisenberg group over  $\mathbb{Q}_q$  is a constant-order locally compact Vilenkin group.

**Example 3.17** Let  $q \neq 2$  be a prime number. By [DeV22, Example 1.6], the 4-dimensional  $q$ -adic Lie group  $\mathbb{E}_4(\mathbb{Q}_q)$  over  $\mathbb{Q}_q$  is a constant-order locally compact Vilenkin group.

Finally, we refer to [DeV22, Example 1.5] for the case of graded groups.

**Crowns and radial functions** Following [Spe70, Definition V.1.1] (see also [Omm88, p. 193] and [Que87, p. 114]), we introduce the following definition. We leave it to the reader to adapt for the case of  $I = -\mathbb{N}$ .

**Definition 3.18** *Let  $G$  be a locally compact Spector-Vilenkin group with respect to a strictly decreasing sequence  $(G_k)_{k \in I}$ . A crown of  $G$  is a part of  $G$  of the form  $G_k - G_{k+1}$  for some  $k \in I$ . We say that a function  $f: G \rightarrow \mathbb{C}$  is radial if it is constant on each crown.*

Recall that if  $\phi \in L^1(\mathbb{R}^n)$  the least radially decreasing majorant of  $\phi$  is a function  $\mathcal{R}_\phi$  defined in [HvNVW16, Theorem 2.3.8 p. 103] by  $\mathcal{R}_\phi(x) \stackrel{\text{def}}{=} \text{esssup}_{|x| \leq |y|} |\phi(y)|$  for almost all  $x \in \mathbb{R}^n$ . This

positive function belongs to the space  $L^1(\mathbb{R}^n)$ . Such a function satisfies  $\mathcal{R}_\phi(x) = \eta(|x|)$  for some decreasing function  $\eta: [0, \infty) \rightarrow [0, \infty)$  and  $|\phi(x)| \leq \mathcal{R}_\phi(x)$  for almost all  $x \in \mathbb{R}^n$ . Now, we introduce the following definition whose second part is an analogue of this notion for locally compact Spector-Vilenkin groups, greatly inspired by [Omn88, p. 193].

**Definition 3.19** *Let  $G$  be a locally compact Spector-Vilenkin group and  $(G_k)_{k \in I}$  be the associated sequence. We consider a left Haar measure  $\mu_G$  on  $G$  normalized by the condition  $\mu_G(G_0) = 1$ .*

1. *Let  $s \in G$  and  $k \in I$ . If  $s \in G_k - G_{k+1}$ , we let*

$$(3.3) \quad |s| = \mu_G(G_k).$$

*Moreover, we let  $|e| = 0$ .*

2. *The radially decreasing majorant of a complex measurable function  $\phi: G \rightarrow \mathbb{C}$  is the function  $\mathcal{R}_\phi: G \rightarrow \mathbb{R}_+$  defined by*

$$(3.4) \quad \mathcal{R}_\phi(s) = \operatorname{esssup}_{|s| \leq |r|} |\phi(r)|.$$

We have

$$(3.5) \quad |\phi(s)| \leq \mathcal{R}_\phi(s).$$

Note that  $\mathcal{R}_\phi(s)$  depends only on  $|s|$  and is a decreasing function of  $|s|$ . For any element  $s \in G \setminus \{e\}$  and any integer  $k \in I$ , we have

$$(3.6) \quad |s| \leq \mu_G(G_k) \iff s \in G_k.$$

**Example 3.20** Consider the additive group  $G = (\mathbb{K}^n, +)$  of  $\mathbb{K}^n$ . Equipped with the family  $(G_k)_{k \in \mathbb{Z}}$  of subgroups defined in (3.2), it is a locally compact Spector-Vilenkin group by Example 3.11. According to [RaV99, p. 146], the function  $\|\cdot\|_{\mathbb{K}^n}$  takes only values in the set  $q^{\mathbb{Z}} \cup \{0\}$ . We also have  $G_k - G_{k+1} \stackrel{(3.2)}{=} \{x \in \mathbb{K}^n : \|x\|_{\mathbb{K}^n} = \frac{1}{q^k}\}$ . By [Tai75, Lemma 4.1 p. 128] reproduced in (6.10), we have

$$(3.7) \quad \mu_G(G_k - G_{k+1}) = \int_{\|x\|_{\mathbb{K}^n} = q^{-k}} d\mu_G(x) = q^{-kn}(1 - q^{-n}).$$

Using the disjoint union  $G_k = (G_k - G_{k+1}) \cup (G_{k+1} - G_{k+2}) \cup \dots$ , we obtain

$$\mu_G(G_k) = \sum_{l=k}^{\infty} \mu(G_l - G_{l+1}) \stackrel{(3.7)}{=} (1 - q^{-n}) \sum_{l=k}^{\infty} q^{-ln} = q^{-kn}.$$

For any  $x \in G_k - G_{k+1}$ , we infer that

$$(3.8) \quad |x| \stackrel{(3.3)}{=} \mu_G(G_k) = \frac{1}{q^{kn}} = \|x\|_{\mathbb{K}^n}^n.$$

In the particular case  $n = 1$  and  $\mathbb{K} = \mathbb{Q}_q$ , we obtain that  $|\cdot| = |\cdot|_q$  and  $\mu_{\mathbb{Q}_q}(G_k) = \frac{1}{q^k}$  which is also stated in [VVZ94, (2.3) p. 40].



## 4 Maximal inequalities

Let  $I$  be a nonempty countable totally ordered set. Let  $(\mathcal{A}_i)_{i \in I}$  be an increasing sequence of  $\sigma$ -algebras on a measure space  $(\Omega, \mathcal{A}, \mu)$  such that  $\mu$  is  $\sigma$ -finite on each  $\mathcal{A}_i$ . If  $1 < p < \infty$ , the classical Doob inequality says that for any  $f \in L^p(\Omega)$  we have

$$\left\| \sup_{i \in I} |\mathbb{E}(f | \mathcal{A}_i)| \right\|_{L^p(\Omega)} \lesssim_p \|f\|_{L^p(\Omega)},$$

where  $\mathbb{E}(\cdot | \mathcal{A}_i) : L^p(\Omega) \rightarrow L^p(\Omega)$  is the conditional expectation. See e.g. [Pis16, Remark 1.39 p. 27] and [Ste70, Theorem 6 p. 91]. Actually, if  $n \geq 1$  is an integer there exists an  $\ell_n^2$ -extension of this inequality providing the estimate (with constant independent of  $n$ )

$$(4.1) \quad \left\| \left( \sup_{i \in I} |\mathbb{E}(f_k | \mathcal{A}_i)| \right)_{1 \leq k \leq n} \right\|_{L^p(\Omega, \ell_n^2)} \lesssim \|(f_k)_{1 \leq k \leq n}\|_{L^p(\Omega, \ell_n^2)},$$

for any element  $f = (f_k)_{1 \leq k \leq n}$  of the Bochner space  $L^p(\Omega, \ell_n^2)$ . We refer to [Pis16, Remark 1.38 p. 27]. See also [HvNVW16, Theorem 3.2.7 p. 178]. We need a consequence of this inequality for a unimodular locally compact Spector-Vilenkin group  $G$  with strictly decreasing sequence  $(G_k)_{k \in I}$ , where  $I = \mathbb{Z}$ ,  $I = \mathbb{N}$  or  $I = -\mathbb{N}$ , equipped with a Haar measure. Let  $(A_i)_{i \in I}$  be the family of averaging operators defined by

$$(4.2) \quad (A_i f)(s) \stackrel{\text{def}}{=} \frac{1}{\mu_G(G_i)} \int_{G_i} f(t^{-1}s) d\mu_G(t), \quad f \in L^p(G), \quad s \in G, \quad i \in I.$$

If  $f \in L^p(G)$  is a scalar function, we also let

$$(4.3) \quad (Mf)(s) \stackrel{\text{def}}{=} \sup_{i \in I} (A_i(|f|))(s), \quad s \in G.$$

Now, we can prove the following result.

**Proposition 4.1** *Let  $G$  be a unimodular locally compact Vilenkin group equipped with a Haar measure. Suppose that  $1 < p < \infty$ . Let  $f = (f_k)_{k \geq 0}$  be an element of the Bochner space  $L^p(G, \ell^2)$ . We have*

$$(4.4) \quad \|(Mf_k)_{1 \leq k \leq n}\|_{L^p(G, \ell_n^2)} \lesssim \|(f_k)_{1 \leq k \leq n}\|_{L^p(G, \ell_n^2)}.$$

*Proof* : Let  $\mathcal{A}$  be the  $\sigma$ -algebra of Borel sets on  $G$ . For each integer  $i \in I$ , we define  $\mathcal{A}_i$  to be the  $\sigma$ -subalgebra of  $\mathcal{A}$  generated by the right cosets

$$\{G_i s : s \in G\}$$

of the group  $G_i$ . Let  $i \in I$ . Since  $G_i/G_{i+1}$  is finite, we can write  $G_i = G_{i+1}t_1 \cup \dots \cup G_{i+1}t_N$  for some elements  $t_1, \dots, t_N \in G_i$ . Then for all  $s \in G$ , we have  $G_i s = G_{i+1}t_1 s \cup \dots \cup G_{i+1}t_N s$ . Hence  $G_i s \in \mathcal{A}_{i+1}$ . Thus,  $\mathcal{A}_i \subset \mathcal{A}_{i+1}$  if the integer  $i$  belongs to  $I$ .

It is easy to see that there are only countably many cosets of  $G_i$  in  $G$ . Consider a family  $(G_i s_{i,j})_{j \in J_i}$  of right coset representatives. The conditional expectation  $\mathbb{E}(\cdot | \mathcal{A}_i)$  with respect to the  $\sigma$ -algebra  $\mathcal{A}_i$  is given by

$$(4.5) \quad \mathbb{E}(f | \mathcal{A}_i) \stackrel{(2.17)}{=} \sum_{j \in J_i} \left[ \frac{1}{\mu(G_i s_{i,j})} \int_{G_i s_{i,j}} f d\mu \right] 1_{G_i s_{i,j}}.$$



If  $s \in G$ , recall that  $s \in G_i s_{i,j}$  if and only if we have  $G_i s_{i,j} = G_i s$ . In this case, we obtain that

$$\begin{aligned}\mathbb{E}(f|\mathcal{A}_i)(s) &= \sum_{j \in J_i} \left[ \frac{1}{\mu(G_i s_{i,j})} \int_{G_i s_{i,j}} f \, d\mu \right] 1_{G_i s_{i,j}}(s) \\ &= \frac{1}{\mu(G_i)} \int_{G_i s} f \, d\mu = \frac{1}{\mu(G_i)} \int_{G_i} f(t^{-1}s) \, d\mu(t) \stackrel{(4.2)}{=} (A_i f)(s).\end{aligned}$$

Appealing to the generalization (4.1) of Doob's inequality, for any element  $(f_k)_{1 \leq k \leq n}$  of the Bochner space  $L^p(G, \ell_n^2)$ , we deduce that

$$\begin{aligned}\|(Mf_k)_{1 \leq k \leq n}\|_{L^p(G, \ell_n^2)} &\stackrel{(4.3)}{=} \left\| \left( \sup_{i \in I} A_i(|f_k|) \right)_{1 \leq k \leq n} \right\|_{L^p(G, \ell_n^2)} \\ &\stackrel{(4.2)}{=} \left\| \left( \sup_{i \in I} |A_i(|f_k|)| \right)_{1 \leq k \leq n} \right\|_{L^p(G, \ell_n^2)} \\ &\stackrel{(4.1)}{\lesssim} \|(f_k)_{1 \leq k \leq n}\|_{L^p(G, \ell_n^2)}.\end{aligned}$$

■

In a similar manner to Proposition 4.1, we obtain the following variant where the space  $\ell^2$  is replaced by any UMD Banach function space.

Given a (complete)  $\sigma$ -finite measure space  $(\Omega', \mathcal{A}, \mu)$ , a Banach function space  $Y$  over  $(\Omega', \mathcal{A}, \mu)$  [BeS88, Chapter I] (see also [LiT79, Definition 1.b.17. p. 28] for a variant) is a subspace consisting of equivalence classes, modulo equality almost everywhere, of locally integrable functions on  $\Omega'$ , equipped with a norm  $\|\cdot\|_Y$ , such that the following three properties hold.

1. If  $f \in Y$  and  $g \in L^0(\Omega)$  with  $|g| \leq |f|$  a.e., then  $g \in Y$  with  $\|g\|_Y \leq \|f\|_Y$ .
2. For any increasing sequence  $(f_n)_{n \geq 1}$  of positive functions of  $Y$  with  $\sup_{n \geq 1} \|f_n\|_Y < \infty$  such that  $f_n \rightarrow f$  almost everywhere, we have  $f \in Y$  and  $\|f\|_Y = \sup_{n \geq 1} \|f_n\|_Y$ .
3. For every  $A \in \mathcal{A}$  with  $\mu(A) < \infty$  the characteristic function  $1_A$  of  $A$  belongs to  $Y$ .

Every Banach function space is a complex Banach lattice in the obvious order, i.e.  $f \geq 0$  if  $f(\omega') \geq 0$  almost everywhere.

Recall that a complex Banach lattice  $Y$  is UMD if and only if  $Y$  and  $Y^*$  have the Hardy-Littlewood property. Let  $Y = Y(\Omega')$  be a Banach function space. Further, let  $(\mathcal{A}_i)_{i \in I}$  be a filtration on a measure space  $(\Omega, \mathcal{A}, \mu)$  such that  $\mu$  is  $\sigma$ -finite on each  $\mathcal{A}_i$  and denote by  $\mathbb{E}_i$  the corresponding conditional expectation operators.

If  $Y$  has the Hardy-Littlewood property, then according to [DKK21, Lemma 3.4] the lattice maximal function

$$(4.6) \quad M(f)(x, \omega') \stackrel{\text{def}}{=} \sup_{i \in I} |\mathbb{E}_i f(x, \omega')|, \quad x \in \Omega, \quad \omega' \in \Omega',$$

defines a bounded operator on the Bochner space  $L^p(\Omega, Y)$  for any  $p \in (1, \infty)$ . For a locally compact Spector-Vilenkin group  $G = \bigcup_{i \in I} G_i$ , we let

$$(4.7) \quad M(f)(s, \omega) \stackrel{\text{def}}{=} \sup_{i \in I} (A_i(|f(\cdot, \omega')|))(s), \quad f \in L^p(G, Y(\Omega')), \quad s \in G, \omega' \in \Omega'.$$

**Proposition 4.2** *Let  $G$  be a locally compact Spector-Vilenkin group. Suppose that  $1 < p < \infty$ . Let  $Y = Y(\Omega')$  be a UMD Banach function space. For any element  $f$  of the Bochner space  $L^p(G, Y)$ , we have*

$$\|M(f)\|_{L^p(G, Y)} \lesssim \|f\|_{L^p(G, Y)}.$$

*Proof* : We consider the  $\sigma$ -algebras  $\mathcal{A}$  and  $\mathcal{A}_i$  as in the proof of Proposition 4.1. Note that  $Y$  has the Hardy-Littlewood property. Using the previously described result [DKK21, Lemma 3.4], we obtain

$$\left\| \sup_i (A_i(|f(\cdot, \omega')|))(s) \right\|_{L^p(G, Y)} \lesssim \|f\|_{L^p(G, Y)}.$$

■

## 5 $R$ -boundedness of some family of convolution operators

If  $G$  is a unimodular locally compact group equipped with a Haar measure  $\mu_G$ , recall that the convolution product of two functions  $f$  and  $g$  is given, when it exists, by

$$(5.1) \quad (g * f)(s) \stackrel{\text{def}}{=} \int_G g(t)f(t^{-1}s) d\mu_G(t) = \int_G g(st^{-1})f(t) d\mu_G(t).$$

The following is an analogue of [HvNVW18, Proposition 2.3.9 p. 104] and [Gra14, p. 82 and p. 84]. See also [Lor19] for related results. Recall that the radially decreasing majorant  $\mathcal{R}_\phi$  is defined in (3.4).

**Proposition 5.1** *Let  $G$  be a unimodular locally compact Vilenkin group. Let  $f \in L^1_{\text{loc}}(G)$  and  $\phi \in L^1(G)$ . For almost all  $s \in G$ , we have*

$$(5.2) \quad |(\phi * f)(s)| \leq \|\mathcal{R}_\phi\|_{L^1(G)} (Mf)(s).$$

*Proof* : Since  $\mathcal{R}_\phi(s)$  depends only on  $|s|$  and is a decreasing function of  $|s|$ , there exists some sequence  $(u_k)_{k \in \mathbb{Z}}$  of positive real numbers such that

$$(5.3) \quad \mathcal{R}_\phi(s) = \sum_{k \in \mathbb{Z}, |s| \leq \mu_G(G_k)} u_k, \quad s \in G.$$

Using Fubini's theorem in the second equality, we see that

$$(5.4) \quad \begin{aligned} \sum_{k \in \mathbb{Z}} \mu_G(G_k) u_k &= \sum_{k \in \mathbb{Z}} \left( \int_G 1_{G_k} d\mu_G \right) u_k \\ &= \int_G \sum_{k \in \mathbb{Z}} 1_{G_k}(s) u_k d\mu_G(s) \\ &\stackrel{(3.6)}{=} \int_G \sum_{|s| \leq \mu_G(G_k)} u_k d\mu_G(s) \\ &\stackrel{(5.3)}{=} \int_G \mathcal{R}_\phi(s) d\mu_G(s) \\ &= \|\mathcal{R}_\phi\|_{L^1(G)}. \end{aligned}$$

For almost all  $s \in G$ , using again Fubini's theorem in the fourth equality, we obtain

$$\begin{aligned}
|(\phi * f)(s)| &\stackrel{(5.1)}{=} \left| \int_G \phi(t) f(t^{-1}s) \, d\mu_G(t) \right| \leq \int_G |\phi(t)| |f(t^{-1}s)| \, d\mu_G(t) \\
&\stackrel{(3.5)}{\leq} \int_G \mathcal{R}_\phi(t) |f(t^{-1}s)| \, d\mu_G(t) \stackrel{(5.3)}{=} \int_G \left( \sum_{|t| \leq \mu_G(G_k)} u_k \right) |f(t^{-1}s)| \, d\mu_G(t) \\
&\stackrel{(3.6)}{=} \int_G \sum_{k \in \mathbb{Z}} 1_{G_k}(t) u_k |f(t^{-1}s)| \, d\mu_G(t) = \sum_{k \in \mathbb{Z}} \left( \int_{G_k} |f(t^{-1}s)| \, d\mu_G(t) \right) u_k \\
&= \sum_{k \in \mathbb{Z}} \mu_G(G_k) \left( \frac{1}{\mu_G(G_k)} \int_{G_k} |f(t^{-1}s)| \, d\mu_G(t) \right) u_k \\
&\stackrel{(5.4)}{\leq} \|\mathcal{R}_\phi\|_{L^1(G)} \sup_{k \in \mathbb{Z}} \frac{1}{\mu_G(G_k)} \int_{G_k} |f(t^{-1}s)| \, d\mu_G(t) \\
&\stackrel{(4.2)}{=} \|\mathcal{R}_\phi\|_{L^1(G)} \sup_{k \in \mathbb{Z}} (A_k(|f|))(s) \stackrel{(4.3)}{=} \|\mathcal{R}_\phi\|_{L^1(G)} (Mf)(s).
\end{aligned}$$

■

The following is an analogue of [HvNVW18, Proposition 8.2.1 p. 184] which gives the same result when  $G$  is replaced by  $\mathbb{R}^n$ . Recall that  $M$  is defined in (4.3).

**Proposition 5.2** *Let  $G$  be a locally compact Spector-Vilenkin group. Suppose that  $1 < p < \infty$ . The family  $\mathcal{T} \stackrel{\text{def}}{=} \{T: L^p(G) \rightarrow L^p(G) : |T(f)| \leq M(f)\}$  is  $R$ -bounded with*

$$(5.5) \quad \mathcal{R}_p(\mathcal{T}) \leq c_{G,p}.$$

Moreover, if  $Y$  is a UMD Banach function space, then the family  $\mathcal{T}_Y \stackrel{\text{def}}{=} \{T: L^p(G, Y) \rightarrow L^p(G, Y) : |T(f)(s, \omega)| \leq M|f|(s, \omega)\}$  is  $R$ -bounded over the Bochner space  $L^p(G, Y)$ .

*Proof* : Using Proposition 4.1 in the inequality, we obtain for any integer  $n \geq 1$  and any functions  $f_1, \dots, f_n \in L^p(G)$

$$\begin{aligned}
&\left\| \sum_{k=1}^n \varepsilon_k \otimes T_k(f_k) \right\|_{L^p(\Omega, L^p(G))} \stackrel{(2.8)}{\approx} \left\| \left( \sum_{k=1}^n |T_k(f_k)|^2 \right)^{\frac{1}{2}} \right\|_{L^p(G)} \\
&\leq \left\| \left( \sum_{k=1}^n (M(f_k))^2 \right)^{\frac{1}{2}} \right\|_{L^p(G)} \\
&= \left\| (M(f_k))_{1 \leq k \leq n} \right\|_{L^p(G, \ell_n^2)} \stackrel{(4.4)}{\lesssim_{G,p}} \left\| (f_k)_{1 \leq k \leq n} \right\|_{L^p(G, \ell_n^2)} \\
&= \left\| \left( \sum_{k=1}^n |f_k|^2 \right)^{\frac{1}{2}} \right\|_{L^p(G)} \stackrel{(2.8)}{\approx} \left\| \sum_{k=1}^n \varepsilon_k \otimes f_k \right\|_{L^p(\Omega, L^p(G))}.
\end{aligned}$$

We turn to the last sentence of the statement. Since a UMD Banach space has finite cotype by [HvNVW18, Proposition 7.3.15 p. 99], the equivalence (2.8) holds also in the space  $L^p(G, Y)$ . Note that when  $Y = Y(\Omega)$  is a UMD Banach function space, then also the vector-valued space

$$Y(\Omega, \ell^2) \stackrel{\text{def}}{=} \{f: \Omega \rightarrow \ell^2 : f \text{ is strongly measurable and } \omega \mapsto \|f(\omega)\|_{\ell^2} \text{ belongs to } Y\},$$

with norm  $\|f\|_{Y(\ell^2)} = \|\|F(\cdot)\|_{\ell^2}\|_Y$ , is a UMD Banach function space (see e.g. [DK23, Lemma 2.9]). Then argue as before, this time using Proposition 4.2 with the UMD Banach function space  $Y(\Omega, \ell^2)$  instead of using Proposition 4.1. ■

If  $1 < p < \infty$ , note that a sufficient condition for the  $R$ -boundedness of a family  $(C_K)_{K \in \mathcal{K}}$  of integral operators  $C_K: L^p(\mathbb{R}^n, X) \rightarrow L^p(\mathbb{R}^n, X)$ ,  $f \mapsto K * f$  on the Bochner space  $L^p(\mathbb{R}^n, X)$  is well-known if  $X$  is UMD *Banach function space*. Indeed, if  $M$  is the Hardy-Littlewood maximal operator and if

$$\mathcal{K} \stackrel{\text{def}}{=} \{K \in L^1(\mathbb{R}^n) : |K| * |f| \leq Mf \text{ a.e. for all simple } f: \mathbb{R}^n \rightarrow \mathbb{C}\}$$

then the family  $(C_K)_{K \in \mathcal{K}}$  of operators is  $R$ -bounded on the Bochner space  $L^p(\mathbb{R}^n, X)$ , see e.g. [Lor19, Proposition 3.1]. Note that this class of kernels contains any radially decreasing  $K \in L^1(\mathbb{R}^n)$  with  $\|K\|_{L^1(\mathbb{R}^n)} \leq 1$ . The following is an analogue of [HvNVW18, Proposition 8.2.3 p. 184].

**Theorem 5.3** *Let  $G$  be a unimodular locally compact Spector-Vilenkin group. Let  $\mathcal{K}$  the subset of all  $K \in L^1(G)$  such that*

$$\int_G \operatorname{esssup}_{|s| \leq |r|} |K(r)| \, d\mu_G(s) \leq 1.$$

*Suppose that  $1 < p < \infty$ . Then the family  $\mathcal{T}_{\mathcal{K}} \stackrel{\text{def}}{=} \{f \mapsto K * f : K \in \mathcal{K}\}$  of convolution operators acting on the space  $L^p(G)$  is  $R$ -bounded and*

$$(5.6) \quad \mathcal{R}(\mathcal{T}_{\mathcal{K}}) \leq c_{G,p}.$$

*Proof* : Let  $f \in L^p(G)$ . If  $K \in \mathcal{K}$ , by Proposition 5.1, we have for almost all  $s \in G$

$$|(K * f)(s)| \stackrel{(5.2)}{\leq} \|\mathcal{R}_K\|_{L^1(G)} (Mf)(s) \leq (Mf)(s).$$

since  $\mathcal{R}_K(s) \stackrel{(3.4)}{=} \operatorname{esssup}_{|s| \leq |r|} |K(r)|$ . Now, it suffices to use Proposition 5.2. ■

In a similar manner to the proof of Theorem 5.3, by applying Proposition 4.2 instead of Proposition 4.1, we obtain the following «UMD Banach function space-valued» variant.

**Theorem 5.4** *Let  $G$  be a unimodular locally compact Spector-Vilenkin group. Let  $Y$  be a UMD Banach function space. Suppose that  $1 < p < \infty$ . Let  $\mathcal{K}$  be the same subset of  $L^1(G)$  as in the statement of Theorem 5.3 and  $\mathcal{T}_{\mathcal{K}}$  the associated family of convolution operators acting on  $L^p(G, Y)$ . Then this family is  $R$ -bounded and*

$$(5.7) \quad \mathcal{R}(\mathcal{T}_{\mathcal{K}}) \leq c_{G,p,Y}.$$

*Proof* : We have again for  $f \in L^p(G, Y)$ , according to Proposition 5.1 for almost all  $s \in G$

$$|(K * f)(s)| \leq (Mf)(s).$$

Now, it suffices to use the second part of Proposition 5.2. ■

## 6 Application to functional calculus

**Gamma function** Let  $n \geq 1$  be an integer. We will use an  $n$ -dimensional generalization  $\Gamma_q^{(n)}$  of the Gelfand-Graev Gamma function  $\Gamma_q$ . This function is defined by

$$(6.1) \quad \Gamma_q^{(n)}(z) \stackrel{\text{def}}{=} \frac{1 - q^{z-n}}{1 - q^{-z}}, \quad z \in \mathbb{C} - \frac{2\pi i}{\ln(q)}\mathbb{Z}.$$

The function  $\Gamma_q^{(n)}$  is meromorphic, with its set of zeros<sup>2</sup> (all simple) being  $n + \frac{2\pi i}{\ln(q)}\mathbb{Z}$ , and its set of poles (all simple) being  $z \in \frac{2\pi i}{\ln(q)}\mathbb{Z}$ . If  $z$  is a complex number such that  $\text{Re } z > 0$  and if  $u \in \mathbb{K}^n$  satisfies  $|u| = 1$ , it is shown in [Tai75, Theorem 4.2 p. 128] that<sup>3</sup>

$$(6.2) \quad \Gamma_q^{(n)}(z) = \int_{\mathbb{K}^n} \|x\|_{\mathbb{K}^n}^{z-n} \bar{\chi}(u \cdot x) \, dx.$$

Recall that by [Tai75, Remark p. 128] we have

$$(6.3) \quad \Gamma_q^{(n)}(z)\Gamma_q^{(n)}(n-z) = 1.$$

If  $n = 1$ , we recover the Gelfand-Graev Gamma function  $\Gamma_q: \mathbb{C}^* \rightarrow \mathbb{C}$  which admits the simple formula

$$\Gamma_q(z) = \frac{1 - q^{z-1}}{1 - q^{-z}}, \quad z \in \mathbb{C} - \frac{2\pi i}{\ln(q)}\mathbb{Z}.$$

If  $z$  is a complex number such that  $\text{Re } z > 0$ , formula (6.2) gives in particular

$$\Gamma_q(z) \stackrel{\text{def}}{=} \int_{\mathbb{K}} |x|^{z-1} \bar{\chi}(x) \, dx.$$

which is stated in [Tai75, p. 49]. We also refer to [BrF93, p. 16] for more information on this function.

**Lévy-Khinchin formula** Let  $n \geq 1$  and  $0 < \alpha < 2$ . Recall the well-known classical formula [BeF75, p. 184]:

$$(6.4) \quad \|x\|_{\mathbb{R}^n}^\alpha = \frac{\alpha 2^{\alpha-1} \Gamma(\frac{\alpha+n}{2})}{\pi^{\frac{n}{2}} \Gamma(1 - \frac{\alpha}{2})} \int_{\mathbb{R}^n - \{0\}} \frac{1 - \cos\langle x, y \rangle}{\|y\|_{\mathbb{R}^n}^{\alpha+n}} \, dy, \quad x \in \mathbb{R}^n$$

where  $\|x\|_{\mathbb{R}^n} \stackrel{\text{def}}{=} (\sum_{i=1}^n x_i^2)^{\frac{1}{2}}$ . This is a Lévy-Khinchin representation of the conditionally negative definite function  $\|\cdot\|_{\mathbb{R}^n}^\alpha$  as a continuous sum of elementary conditionally negative definite functions<sup>4</sup>, which is a particular case of [BeF75, Corollary 18.20 p. 184]. Replacing  $\mathbb{R}$  by a non-archimedean local field  $\mathbb{K}$ , we will establish in Proposition 6.1 the folklore analogue of this formula for  $0 < \alpha < \infty$  instead of  $0 < \alpha < 2$  with sharp contrast, for the sake of completeness.

The function  $\|\cdot\|_{\mathbb{K}^n}^{\alpha-n}$  determines a distribution<sup>5</sup> on  $\mathbb{K}^n$  acting as

$$\langle \|\cdot\|_{\mathbb{K}^n}^{\alpha-n}, f \rangle = \int_{\mathbb{K}^n} \|x\|_{\mathbb{K}^n}^{\alpha-n} f(x) \, dx, \quad f \in \mathcal{D}(\mathbb{K}^n).$$

2. It is stated in [Tai75, p. 128] that the function  $\Gamma_q^{(n)}$  has a unique zero and a unique pole, which is false.

3. Here the integral is understood to be the limit  $\lim_{k \rightarrow \infty} \int_{q^{-k} \leq \|x\|_{\mathbb{K}^n} \leq q^k} \cdots \, dx$ .

4. If  $G$  is a locally compact abelian group and if  $\chi$  is a character of  $G$ , recall that a function  $G \rightarrow \mathbb{R}$ ,  $s \mapsto 1 - \text{Re } \chi(s)$  is conditionally negative definite by [BeF75, Corollary 7.7 p. 41] and [BeF75, Proposition 7.4 (ii) p. 40].

5. For the definition of the space of test functions  $\mathcal{D}(\mathbb{K}^n)$  and distributions  $\mathcal{D}'(\mathbb{K}^n)$ , we refer to [Tai75, Section II.3 and III.3].

Actually, the map  $\alpha \mapsto \|\cdot\|_{\mathbb{K}^n}^{\alpha-n}$  extends to a meromorphic distribution-valued function on the subset  $\mathbb{C} - \{n + \frac{2\pi i}{\ln q} \mathbb{Z}\}$  given by

$$\langle \|\cdot\|_{\mathbb{K}^n}^{\alpha-n}, f \rangle \stackrel{\text{def}}{=} \frac{1 - q^{-n}}{1 - q^{-\alpha}} f(0) + \int_{\|x\|_{\mathbb{K}^n} > 1} \|x\|^{\alpha-n} f(x) dx + \int_{\|x\|_{\mathbb{K}^n} \leq 1} \|x\|^{\alpha-n} [f(x) - f(0)] dx.$$

If  $\text{Re } \alpha > 0$  and if  $f \in \mathcal{D}(\mathbb{K}^n)$ , it is known that

$$(6.5) \quad \langle \|\cdot\|_{\mathbb{K}^n}^{-\alpha-n}, f \rangle = \int_{\mathbb{K}^n} [f(x) - f(0)] \frac{dx}{\|x\|_{\mathbb{K}^n}^{\alpha+n}}.$$

The Fourier transform<sup>6</sup> of the distribution  $\|\cdot\|_{\mathbb{K}^n}^{\alpha-n}$  is known. Actually, by [Tai68, Theorem 2 p. 194], [Tai75, Theorem 4.5 p. 130], [VVZ94, (4.3) p. 136], we have in  $\mathcal{D}'(\mathbb{K}^n)$

$$(6.6) \quad \mathcal{F}(\|\cdot\|_{\mathbb{K}^n}^{\alpha-n}) = \Gamma_q^{(n)}(\alpha) \|\cdot\|_{\mathbb{K}^n}^{-\alpha}, \quad \alpha \in \mathbb{C} - \{n + \frac{2\pi i}{\ln q} \mathbb{Z}\}.$$

Now, we can prove the following result.

**Proposition 6.1** *Let  $\mathbb{K}$  be a non-archimedean local field and  $n \geq 1$ . For any  $0 < \alpha < \infty$ , we have*

$$(6.7) \quad \|x\|_{\mathbb{K}^n}^{\alpha} = \frac{-1}{\Gamma_q^{(n)}(-\alpha)} \int_{\mathbb{K}^n - \{0\}} \frac{1 - \chi(x \cdot y)}{\|y\|_{\mathbb{K}^n}^{\alpha+n}} dy, \quad x \in \mathbb{K}^n.$$

In particular, the function  $\mathbb{K}^n \rightarrow \mathbb{R}$ ,  $x \mapsto \|x\|_{\mathbb{K}^n}^{\alpha}$  is a conditionally negative definite function on the additive group of  $\mathbb{K}^n$ .

*Proof* : For any function  $f \in \mathcal{D}(\mathbb{K}^n)$ , using (6.6) with  $-\alpha$  instead of  $\alpha$ , we have

$$\begin{aligned} \langle \|\cdot\|_{\mathbb{K}^n}^{\alpha}, f \rangle &\stackrel{(6.6)}{=} \Gamma_q^{(n)}(\alpha+n) \langle \mathcal{F}^{-1}(\|\cdot\|_{\mathbb{K}^n}^{-\alpha-n}), f \rangle = \Gamma_q^{(n)}(\alpha+n) \langle \|\cdot\|_{\mathbb{K}^n}^{-\alpha-n}, \check{f} \rangle \\ &\stackrel{(6.3)(6.5)}{=} \frac{1}{\Gamma_q^{(n)}(-\alpha)} \int_{\mathbb{K}^n} [\check{f}(x) - \check{f}(0)] \frac{dx}{\|x\|_{\mathbb{K}^n}^{\alpha+n}} \\ &\stackrel{(2.6)}{=} \frac{1}{\Gamma_q^{(n)}(-\alpha)} \int_{\mathbb{K}^n} \left( \int_{\mathbb{K}^n} (\chi(x \cdot y) - 1) f(y) dy \right) \frac{dx}{\|x\|_{\mathbb{K}^n}^{\alpha+n}}. \end{aligned}$$

Using Fubini's theorem, we get

$$\langle \|\cdot\|_{\mathbb{K}^n}^{\alpha}, f \rangle = \frac{1}{\Gamma_q^{(n)}(-\alpha)} \int_{\mathbb{K}^n} f(y) \left( \int_{\mathbb{K}^n} \frac{\chi(x \cdot y) - 1}{\|x\|_{\mathbb{K}^n}^{\alpha+n}} dx \right) dy.$$

■

**Remark 6.2** Note that (6.7) remains true if we replace  $\chi(x \cdot y)$  by  $\text{Re } \chi(x \cdot y)$ . Indeed, the integral  $\int_{\mathbb{K}^n - \{0\}} \frac{\text{Im } \chi(x \cdot y)}{\|y\|_{\mathbb{K}^n}^{\alpha+n}} dy$  vanishes which can be seen in the following way. Note that in the first equality, we use that  $\mathbb{K}^n$  is unimodular as a commutative (additive) group, and in the second

6. For any distribution  $T \in \mathcal{D}'(\mathbb{K}^n)$  its Fourier transform, denoted by  $\mathcal{F}(T)$ , is the distribution defined as

$$\langle \mathcal{F}(T), f \rangle = \langle T, \mathcal{F}(f) \rangle, \quad f \in \mathcal{D}(\mathbb{K}^n).$$

that  $\operatorname{Im} \chi(-x \cdot y) = -\operatorname{Im} \chi(x \cdot y)$ . Indeed, since  $\chi$  is a character, we have  $\chi(-x \cdot y) = \overline{\chi(x \cdot y)}$ . We infer that  $\operatorname{Im} \chi(-x \cdot y) = \operatorname{Im} \overline{\chi(x \cdot y)} = -\operatorname{Im} \chi(x \cdot y)$ . We deduce that

$$\begin{aligned} \int_{\mathbb{K}^n - \{0\}} \frac{\operatorname{Im} \chi(x \cdot y)}{\|y\|_{\mathbb{K}^n}^{\alpha+n}} dy &= \int_{\mathbb{K}^n - \{0\}} \frac{\operatorname{Im} \chi(-x \cdot y)}{\| -y \|_{\mathbb{K}^n}^{\alpha+n}} dy \\ &= \int_{\mathbb{K}^n - \{0\}} \frac{-\operatorname{Im} \chi(x \cdot y)}{\|y\|_{\mathbb{K}^n}^{\alpha+n}} dy. \end{aligned}$$

Thus the previous integral equals its negative and is therefore equal to 0.

**Remark 6.3** If  $n = 1$  and  $\mathbb{K} = \mathbb{Q}_q$ , we recover the formula

$$|x|_q^\alpha = \frac{q^\alpha - 1}{1 - q^{-1-\alpha}} \int_{\mathbb{Q}_q - \{0\}} \frac{1 - \operatorname{Re} \chi(x \cdot y)}{|y|_q^{\alpha+1}} dy, \quad x \in \mathbb{Q}_q.$$

of the papers [Har90, p. 699] and [Hara93, p. 908]. By (2.5), we have  $\operatorname{Re} \chi(x \cdot y) = \cos 2\pi \{x \cdot y\}_q$ .

Consequently, by essentially [BeF75, Theorem 8.3], we have a weak\* semigroup  $(T_t)_{t \geq 0}$  of selfadjoint contractive positive Fourier multipliers on the algebra  $L^\infty(\mathbb{K}^n)$ . Note that the opposite of the infinitesimal generator, i.e. the Fourier multiplier by the symbol  $\|\cdot\|_{\mathbb{K}^n}^\alpha$ , is called Taibleson operator and sometimes Vladimirov operator. Each  $T_t$  is a convolution map  $K_t * \cdot$  by the kernel

$$K_t(x) \stackrel{\text{def}}{=} \int_{\mathbb{K}^n} e^{-t\|\xi\|_{\mathbb{K}^n}^\alpha} \overline{\chi(x \cdot \xi)} d\xi, \quad t \geq 0, x \in \mathbb{K}^n.$$

More generally, we need the kernel for complex times

$$(6.8) \quad K_z(x) \stackrel{\text{def}}{=} \int_{\mathbb{K}^n} e^{-z\|\xi\|_{\mathbb{K}^n}^\alpha} \overline{\chi(x \cdot \xi)} d\xi, \quad \operatorname{Re} z > 0, x \in \mathbb{K}^n.$$

The following is a slight extension of [RoZ08, Lemma 2] (see also [Koc01, (4.5) p. 147]). Again, we refer to [Ber52], [Fel71, Lemma 1 p. 583] and [Jan22, Theorem 4.1] for the analogous result in the literature for  $\mathbb{R}^n$ . For the proof, for any integer  $k \in \mathbb{Z}$  and any  $x \in \mathbb{K}^n$  we need the value of the classical integral [Tai75, Lemma 4.1 p. 128]

$$(6.9) \quad \int_{\|y\|_{\mathbb{K}^n} = q^{-k}} \chi(x \cdot y) dy = \begin{cases} q^{-kn}(1 - q^{-n}) & \text{if } \|x\|_{\mathbb{K}^n} \leq q^k \\ -q^{-n(k+1)} & \text{if } \|x\|_{\mathbb{K}^n} = q^{k+1} \\ 0 & \text{if } \|x\|_{\mathbb{K}^n} \geq q^{k+2} \end{cases}.$$

In particular, we have

$$(6.10) \quad \int_{\|y\|_{\mathbb{K}^n} = q^{-k}} dy = q^{-kn}(1 - q^{-n}) = q^{-kn} - q^{-(k+1)n},$$

see also [Vla99, (15.4) p. 64].

**Proposition 6.4** *Let  $\alpha > 0$ . For any complex number  $z \in \mathbb{C}$  with  $\operatorname{Re} z > 0$  and any  $x \in \mathbb{K}^n - \{0\}$ , we have*

$$(6.11) \quad K_z(x) = \sum_{k=1}^{\infty} \frac{(-z)^k \Gamma_q^{(n)}(k\alpha + n)}{k! \|x\|_{\mathbb{K}^n}^{k\alpha+n}}.$$

*Proof* : Let  $x \in \mathbb{K}^n - \{0\}$ . We have  $\|x\|_{\mathbb{K}^n} = q^{n_0}$  for some  $n_0 \in \mathbb{Z}$ . We expand as usual the integral over  $\mathbb{K}^n$  into crowns and we get

$$\begin{aligned}
K_z(x) &\stackrel{(6.8)}{=} \int_{\mathbb{K}^n} e^{-z\|\xi\|_{\mathbb{K}^n}^\alpha} \overline{\chi(x \cdot \xi)} d\xi = \sum_{k \in \mathbb{Z}} \int_{\|\xi\|_{\mathbb{K}^n} = q^{-k}} e^{-zq^{-k\alpha}} \overline{\chi(x \cdot \xi)} d\xi \\
&= \sum_{k \in \mathbb{Z}} e^{-zq^{-k\alpha}} \int_{\|\xi\|_{\mathbb{K}^n} = q^{-k}} \overline{\chi(x \cdot \xi)} d\xi \\
&\stackrel{(6.9)}{=} (1 - q^{-n}) \left( \sum_{\|x\|_{\mathbb{K}^n} \leq q^k} e^{-zq^{-k\alpha}} q^{-kn} \right) - e^{-z\frac{q^\alpha}{\|x\|_{\mathbb{K}^n}^\alpha}} \|x\|_{\mathbb{K}^n}^{-n} \\
&= (1 - q^{-n}) \left( \sum_{k \geq n_0} e^{-zq^{-k\alpha}} q^{-kn} \right) - e^{-z\frac{q^\alpha}{\|x\|_{\mathbb{K}^n}^\alpha}} \|x\|_{\mathbb{K}^n}^{-n} \\
&= (1 - q^{-n}) \left( \sum_{k \geq 0} e^{-zq^{-k\alpha}} \|x\|_{\mathbb{K}^n}^{-\alpha} q^{-kn} \|x\|_{\mathbb{K}^n}^{-n} \right) - e^{-z\frac{q^\alpha}{\|x\|_{\mathbb{K}^n}^\alpha}} \|x\|_{\mathbb{K}^n}^{-n}.
\end{aligned}$$

We expand in the series over  $k$  the exponential function into a series over  $l$  and get

$$K_z(x) = \frac{1 - q^{-n}}{\|x\|_{\mathbb{K}^n}^n} \left( \sum_{k \geq 0} \sum_{l \geq 0} \frac{(-z)^l}{l!} \left( \frac{q^{-k\alpha}}{\|x\|_{\mathbb{K}^n}^\alpha} \right)^l q^{-kn} \right) - e^{-z\frac{q^\alpha}{\|x\|_{\mathbb{K}^n}^\alpha}} \|x\|_{\mathbb{K}^n}^{-n}.$$

This double series converges absolutely. Indeed, we can bound  $\sum_{k \geq 0} \sum_{l \geq 0} |\dots|$  by

$$\sum_{k \geq 0} \exp\left(|z| \frac{q^{-k\alpha}}{\|x\|_{\mathbb{K}^n}^\alpha}\right) q^{-kn} < \infty.$$

Thus we can interchange the order of summation. Expanding the second exponential and summing a geometric series, we deduce that

$$\begin{aligned}
K_z(x) &= (1 - q^{-n}) \sum_{l \geq 0} \frac{(-z)^l}{l!} \frac{1}{\|x\|_{\mathbb{K}^n}^{l\alpha+n}} \sum_{k \geq 0} q^{-l\alpha k - kn} - \sum_{l \geq 0} \frac{(-z)^l}{l!} \frac{q^{l\alpha}}{\|x\|_{\mathbb{K}^n}^{l\alpha+n}} \\
&= \sum_{l \geq 0} \frac{(-z)^l}{l!} \frac{1}{\|x\|_{\mathbb{K}^n}^{l\alpha+n}} \left( (1 - q^{-n}) \left( \sum_{k \geq 0} q^{-k(l\alpha+n)} \right) - q^{l\alpha} \right) \\
&= \sum_{l \geq 0} \frac{(-z)^l}{l!} \frac{1}{\|x\|_{\mathbb{K}^n}^{l\alpha+n}} \left( \frac{1 - q^{-n}}{1 - q^{-(l\alpha+n)}} - q^{l\alpha} \right) \\
&= \sum_{l \geq 0} \frac{(-z)^l}{l!} \frac{1}{\|x\|_{\mathbb{K}^n}^{l\alpha+n}} \left( \frac{1 - q^{l\alpha}}{1 - q^{-(l\alpha+n)}} \right) \stackrel{(6.1)}{=} \sum_{l \geq 0} \frac{(-z)^l}{l!} \frac{1}{\|x\|_{\mathbb{K}^n}^{l\alpha+n}} \Gamma_q^{(n)}(l\alpha + n).
\end{aligned}$$

Finally, recall that  $\Gamma_q^{(n)}(n) = 0$ . So the first term is zero.  $\blacksquare$

The following is a generalization of [Koc01, Lemma 4.1 p. 147] which gives for  $n = 1$  the estimate  $|K_t(x)| \lesssim \frac{t}{(t^\frac{1}{\alpha} + \|x\|_{\mathbb{K}})^{\alpha+1}}$  for any  $t > 0$ . See also [RoZ08, Lemma 3] for the  $n$ -dimensional generalization. We also refer to the papers [BGPV14], [BGHH21] and to the survey [Gri23] for heat kernel estimates in a larger ultrametric context.

**Lemma 6.5** *Let  $\alpha > 0$ . For any  $x \in \mathbb{K}^n$  and any complex number such that  $\operatorname{Re} z > 0$ , we have*

$$(6.12) \quad |K_z(x)| \lesssim_{q,n} \frac{|z|}{[(\operatorname{Re} z)^\frac{1}{\alpha} + \|x\|_{\mathbb{K}^n}]^{\alpha+n}}.$$



*Proof* : Let  $k \in \mathbb{Z}$  be an integer such that

$$(6.13) \quad q^{k-1} \leq (\operatorname{Re} z)^{\frac{1}{\alpha}} \leq q^k.$$

Let  $p$  be a prime element of  $\mathbb{K}$  that has modulus  $\|p\|_{\mathbb{K}} = q^{-1}$ . Then using the change of variable  $\eta = p^{1-k}\xi$  in the third equality, we obtain

$$(6.14) \quad \begin{aligned} |K_z(x)| &\stackrel{(6.8)}{=} \left| \int_{\mathbb{K}^n} e^{-z\|\xi\|_{\mathbb{K}^n}^{\alpha}} \overline{\chi(x \cdot \xi)} d\xi \right| \leq \int_{\mathbb{K}^n} \left| e^{-z\|\xi\|_{\mathbb{K}^n}^{\alpha}} \overline{\chi(x \cdot \xi)} \right| d\xi \\ &\leq \int_{\mathbb{K}^n} e^{-\operatorname{Re}(z)\|\xi\|_{\mathbb{K}^n}^{\alpha}} d\xi \stackrel{(6.13)}{\leq} \int_{\mathbb{K}^n} e^{-q^{\alpha(k-1)}\|\xi\|_{\mathbb{K}^n}^{\alpha}} d\xi = \int_{\mathbb{K}^n} e^{-\|p^{1-k}\xi\|_{\mathbb{K}^n}^{\alpha}} d\xi \\ &= \frac{1}{q^{(k-1)n}} \int_{\mathbb{K}^n} e^{-\|\eta\|_{\mathbb{K}^n}^{\alpha}} d\eta = \frac{1}{q^{kn}} \left( q^n \int_{\mathbb{K}^n} e^{-\|\eta\|_{\mathbb{K}^n}^{\alpha}} d\eta \right) \stackrel{(6.13)}{\lesssim_{q,n}} \frac{1}{(\operatorname{Re} z)^{\frac{n}{\alpha}}}. \end{aligned}$$

On the other hand, according to (6.11), we have for any complex number  $z$  with  $\operatorname{Re} z > 0$ ,

$$\begin{aligned} K_z(x) &\stackrel{(6.11)}{=} \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} \frac{\Gamma_q^{(n)}(k\alpha + n)}{\|x\|_{\mathbb{K}^n}^{k\alpha+n}} \stackrel{(6.1)}{=} \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} \frac{1 - q^{k\alpha}}{1 - q^{-k\alpha-n}} \frac{1}{\|x\|_{\mathbb{K}^n}^{k\alpha+n}} \\ &= \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} \frac{1}{1 - q^{-k\alpha-n}} \frac{1}{\|x\|_{\mathbb{K}^n}^{k\alpha+n}} - \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} \frac{q^{k\alpha}}{1 - q^{-k\alpha-n}} \frac{1}{\|x\|_{\mathbb{K}^n}^{k\alpha+n}} \\ &= \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} \sum_{l=0}^{\infty} q^{-lk\alpha - ln} \frac{1}{\|x\|_{\mathbb{K}^n}^{k\alpha+n}} - \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} \sum_{l=0}^{\infty} q^{k\alpha} q^{-lk\alpha - ln} \frac{1}{\|x\|_{\mathbb{K}^n}^{k\alpha+n}} \\ &= \sum_{l=0}^{\infty} q^{-ln} \|x\|_{\mathbb{K}^n}^{-n} \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} \frac{q^{-lk\alpha}}{\|x\|_{\mathbb{K}^n}^{k\alpha}} - \sum_{l=0}^{\infty} q^{-ln} \|x\|_{\mathbb{K}^n}^{-n} \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} q^{k\alpha} q^{-lk\alpha} \frac{1}{\|x\|_{\mathbb{K}^n}^{k\alpha}} \\ &= \sum_{l=0}^{\infty} q^{-ln} \|x\|_{\mathbb{K}^n}^{-n} \left( \exp(-q^{-l\alpha} \|x\|_{\mathbb{K}^n}^{-\alpha} z) - \exp(-q^{-l\alpha+\alpha} \|x\|_{\mathbb{K}^n}^{-\alpha} z) \right). \end{aligned}$$

Note that the elements  $q^{-l\alpha} \|x\|_{\mathbb{K}^n}^{-\alpha} z$  and  $q^{-l\alpha+\alpha} \|x\|_{\mathbb{K}^n}^{-\alpha} z$  are complex numbers with real part  $> 0$ . We infer by the mean value theorem<sup>7</sup>

$$(6.15) \quad |K_z(x)| \leq \sum_{l=0}^{\infty} q^{-ln} \|x\|_{\mathbb{K}^n}^{-n} \|x\|_{\mathbb{K}^n}^{-\alpha} |z| |q^{-l\alpha}(1 - q^{\alpha})| \lesssim \frac{|z|}{\|x\|_{\mathbb{K}^n}^{\alpha+n}}.$$

We will examine two cases.

*First case.* If  $\|x\|_{\mathbb{K}^n} \leq (\operatorname{Re} z)^{\frac{1}{\alpha}}$ , we have

$$|K_z(x)| \stackrel{(6.14)}{\lesssim} \frac{\operatorname{Re} z}{[(\operatorname{Re} z)^{\frac{1}{\alpha}}]^{\alpha+n}} \leq \frac{\operatorname{Re} z}{[\frac{1}{2}(\operatorname{Re} z)^{\frac{1}{\alpha}} + \frac{1}{2}\|x\|_{\mathbb{K}^n}]^{\alpha+n}} \lesssim \frac{\operatorname{Re} z}{[(\operatorname{Re} z)^{\frac{1}{\alpha}} + \|x\|_{\mathbb{K}^n}]^{\alpha+n}}.$$

*Second case.* Now, if  $\|x\|_{\mathbb{K}^n} \geq (\operatorname{Re} z)^{\frac{1}{\alpha}}$  then

$$|K_z(x)| \stackrel{(6.15)}{\lesssim} \frac{|z|}{\|x\|_{\mathbb{K}^n}^{\alpha+n}} \leq \frac{|z|}{[\frac{1}{2}(\operatorname{Re} z)^{\frac{1}{\alpha}} + \frac{1}{2}\|x\|_{\mathbb{K}^n}]^{\alpha+n}} \lesssim \frac{|z|}{[(\operatorname{Re} z)^{\frac{1}{\alpha}} + \|x\|_{\mathbb{K}^n}]^{\alpha+n}}.$$

7. Note that  $e^{-\operatorname{Re} \zeta} \leq 1$  if  $\operatorname{Re} \zeta > 0$ .

■

We will use the following estimate. It is stated in [Koc01, Lemma 4.3 p. 151] that for any  $\alpha > 0$  and any  $b > 0$  we have

$$(6.16) \quad \int_{\mathbb{K}} \frac{1}{[b + \|x\|_{\mathbb{K}}]^{\alpha+1}} dx \lesssim \frac{1}{b^\alpha}.$$

For the convenience of the reader, we include a short proof, also for the generalization [RoZ08, Proposition 2 p. 335] with  $\mathbb{K}$  replaced by  $\mathbb{K}^n$ .

**Lemma 6.6** *Let  $n \in \mathbb{N}$ ,  $\alpha > 0$  and  $b > 0$ . We have*

$$(6.17) \quad \int_{\mathbb{K}^n} \frac{1}{[b + \|x\|_{\mathbb{K}^n}]^{\alpha+n}} dx \lesssim \frac{1}{b^\alpha}.$$

*Proof* : First, suppose that  $b = q^{k_0}$  for some integer  $k_0 \in \mathbb{Z}$ . Then

$$(6.18) \quad \begin{aligned} \int_{\mathbb{K}^n} \frac{1}{[q^{k_0} + \|x\|_{\mathbb{K}^n}]^{\alpha+n}} dx &= \sum_{k \in \mathbb{Z}} \int_{\|x\|_{\mathbb{K}^n} = q^k} \frac{1}{[q^{k_0} + q^k]^{\alpha+n}} dx \\ &\leq \sum_{k=-\infty}^{k_0} \int_{\|x\|_{\mathbb{K}^n} = q^k} \frac{1}{q^{k_0(\alpha+n)}} dx + \sum_{k=k_0+1}^{\infty} \int_{\|x\|_{\mathbb{K}^n} = q^k} \frac{1}{q^{k(\alpha+n)}} dx \\ &= \frac{1}{q^{k_0(\alpha+n)}} \sum_{k=-\infty}^{k_0} \int_{\|x\|_{\mathbb{K}^n} = q^k} dx + \sum_{k=k_0+1}^{\infty} \frac{1}{q^{k(\alpha+n)}} \int_{\|x\|_{\mathbb{K}^n} = q^k} dx \\ &\stackrel{(6.10)}{=} \frac{1}{q^{k_0(\alpha+n)}} \sum_{k=-\infty}^{k_0} (q^{kn} - q^{(k-1)n}) + \sum_{k=k_0+1}^{\infty} \frac{1}{q^{k(\alpha+n)}} q^{kn} (1 - q^{-n}) \\ &= \frac{1}{q^{k_0(\alpha+n)}} q^{k_0 n} + \frac{1}{q^{(k_0+1)\alpha}} \frac{1}{1 - q^{-\alpha}} (1 - q^{-n}) \cong \frac{1}{q^{k_0 \alpha}}. \end{aligned}$$

Then for an arbitrary real number  $b > 0$ , consider the integer  $k_0 \in \mathbb{Z}$  such that  $q^{k_0} \leq b < q^{k_0+1}$ . We deduce from the previous inequality that

$$\int_{\mathbb{K}^n} \frac{1}{[b + \|x\|_{\mathbb{K}^n}]^{\alpha+n}} dx \leq \int_{\mathbb{K}^n} \frac{1}{[q^{k_0} + \|x\|_{\mathbb{K}^n}]^{\alpha+n}} dx \stackrel{(6.18)}{\lesssim} \frac{1}{q^{k_0 \alpha}} \lesssim \frac{1}{b^\alpha}.$$

■

The following result is an analogue of the classical estimate  $\|K_z\|_{L^1(\mathbb{R}^n)} \leq \left(\frac{|z|}{\operatorname{Re} z}\right)^{\frac{n}{2}}$  [ABHN11, Remark 3.7.10 p. 153] (or [HvNVW18, p. 543]) on the kernel  $K_z$  of the heat semigroup on  $\mathbb{R}^n$  for any complex number  $z$  with  $\operatorname{Re} z > 0$ .

**Corollary 6.7** *Let  $\alpha > 0$  and  $n \in \mathbb{N}$ . For any complex number  $z$  such that  $\operatorname{Re} z > 0$ , we have*

$$(6.19) \quad \|K_z\|_{L^1(\mathbb{K}^n)} \lesssim_q \frac{|z|}{\operatorname{Re} z}.$$

*The same estimate also holds for  $K_z$  replaced by its radially decreasing majorant  $\mathcal{R}_{K_z}(x)$   $\stackrel{(3.4)}{=} \operatorname{esssup}_{|x| \leq |r|} |K_z(r)|$ .*

*Proof* : For any complex number  $z$  such that  $\operatorname{Re} z > 0$ , we have

$$\|K_z\|_{L^1(\mathbb{K}^n)} = \int_{\mathbb{K}^n} |K_z(x)| \, dx \stackrel{(6.12)}{\lesssim_q} \int_{\mathbb{K}^n} \frac{|z|}{[(\operatorname{Re} z)^{\frac{1}{\alpha}} + \|x\|_{\mathbb{K}^n}]^{\alpha+n}} \, dx \stackrel{(6.17)}{\lesssim} \frac{|z|}{\operatorname{Re} z}.$$

We turn to the last sentence in the statement. Note that according to Example 3.20, we have  $|x| \stackrel{(3.8)}{=} \|x\|_{\mathbb{K}^n}^n$  for any  $x \in \mathbb{K}^n$ . Thus, the estimate (6.12) bounds  $K_z(x)$  by a radial and radially decreasing term in  $|x|$  (as well as in  $\|x\|_{\mathbb{K}^n}$ ). We infer that (6.12) holds with the left-hand side  $K_z$  replaced by  $\mathcal{R}_{K_z}$ , and the calculation hereabove finishes the proof.  $\blacksquare$

**Theorem 6.8** *Let  $\mathbb{K}$  be a non-archimedean local field and  $n \geq 1$ . Consider a UMD Banach function space  $Y$ . Suppose that  $1 < p < \infty$ . The operator  $A_p$  of the infinitesimal generator  $-A_p$  of the strongly continuous semigroup  $(T_t \otimes \operatorname{Id}_Y)_{t \geq 0}$  of operators acting on the Banach space  $L^p(\mathbb{K}^n, Y)$  admits a bounded  $H^\infty(\Sigma_\omega)$  functional calculus at any angle  $\omega > 0$ , and a bounded Hörmander  $\operatorname{Hör}_2^s(\mathbb{R}_+^*)$  calculus for any  $s > \frac{3}{2}$ .*

*Proof* : By Theorem 2.4, the operator  $A_p$  admits a bounded  $H^\infty(\Sigma_\sigma)$  functional calculus for any angle  $\sigma > \frac{\pi}{2}$ . Fix an angle  $\theta \in (0, \frac{\pi}{2})$  and recall that  $(T_z \otimes \operatorname{Id}_Y)_{z \in \Sigma_\theta}$  denotes the complex time semigroup with generator  $-D^\alpha \otimes \operatorname{Id}_Y$ . Note that each operator  $T_z$  is given by convolution with the integral kernel  $K_z$  from (6.8). Let  $\mathcal{R}(x) \stackrel{(3.4)}{=} \operatorname{esssup}_{|x| \leq |r|} |K_z(r)|$  denote its radially decreasing majorant. Using Corollary 6.7 together with Theorem 5.4 (or Theorem 5.3 in the scalar case), we obtain that the family  $(T_z \otimes \operatorname{Id}_Y)_{|\arg z| \leq \theta}$  is  $R$ -bounded over the Bochner space  $L^p(\mathbb{K}^n, Y)$ . Indeed, for  $|\arg z| \leq \theta$ , we have that

$$\frac{|z|}{\operatorname{Re} z} = \frac{1}{\cos \arg z}$$

is bounded above by  $\frac{1}{\cos \theta} < \infty$ . Then Theorem 2.6 and Theorem 2.6 imply that the operator  $A_p$  has a bounded  $H^\infty(\Sigma_\omega)$  functional calculus for any angle  $\omega > \frac{\pi}{2} - \theta$ . Note that  $\theta$  can be chosen arbitrarily close to  $\frac{\pi}{2}$ , so  $\omega$  arbitrarily close to 0.

Actually, note that

$$\frac{|z|}{\operatorname{Re} z} = \frac{1}{\cos \arg z} \cong \left(\frac{\pi}{2} - |\arg z|\right)^{-1}.$$

Now, with Proposition 2.7, we conclude that the operator  $A_p$  has a bounded Hörmander  $\operatorname{Hör}_2^s(\mathbb{R}_+^*)$  functional calculus for any  $s > 1 + \frac{1}{2} = \frac{3}{2}$ .  $\blacksquare$

**Declaration of interest** None.

**Competing interests** The author declares that he has no competing interests.

**Data availability** No data sets were generated during this study.

**Acknowledgements and Grants** None.

## Bibliography

- [AbA02] Y. A. Abramovich and C. D. Aliprantis. An invitation to operator theory. Graduate Studies in Mathematics, 50. American Mathematical Society, Providence, RI, 2002. [10](#)
- [AVDR81] G. N. Agaev, N. Y. Vilenkin, G. M. Dzhaferli, A. I. Rubinshtein. Multiplicative systems of functions and harmonic analysis on zero-dimensional groups (Russian). Elm, Baku, 1981. [5](#), [12](#)
- [AKS10] S. Albeverio, A. Y. Khrennikov and V. M. Shelkovich. Theory of  $p$ -adic distributions: linear and nonlinear models. London Math. Soc. Lecture Note Ser., 370. Cambridge University Press, Cambridge, 2010. [1](#), [6](#)
- [Are04] W. Arendt. Semigroups and evolution equations: functional calculus, regularity and kernel estimates. Evolutionary equations. Vol. I, 1–85, Handb. Differ. Equ., North-Holland, Amsterdam, 2004. [3](#), [4](#), [8](#)
- [ABHN11] W. Arendt, C. J. K. Batty, M. Hieber and F. Neubrander. Vector-valued Laplace transforms and Cauchy problems. Second edition. Monographs in Mathematics, 96. Birkhäuser/Springer Basel AG, Basel, 2011. [3](#), [25](#)
- [Arh12] C. Arhancet. Unconditionality, Fourier multipliers and Schur multipliers. Colloq. Math. 127 (2012), no. 1, 17–37. [5](#)
- [ArK23] C. Arhancet and C. Kriegler. Projections, multipliers and decomposable maps on noncommutative  $L^p$ -spaces. Mémoires de la Société Mathématique de France (N.S.) (2023), no. 177. [5](#), [11](#)
- [AHKP24] C. Arhancet, L. Hagedorn, C. Kriegler and P. Portal. The harmonic oscillator on the Moyal-Groenewold plane: an approach via Lie groups and twisted Weyl tuples. Preprint, 67 pages. [arXiv:2312.06143](#) [3](#), [9](#)
- [ABK99] V. A. Avetisov, A. H. Bikulov and S. V. Kozyrev. Application of  $p$ -adic analysis to models of breaking of replica symmetry. J. Phys. A 32 (1999), no. 50, 8785–8791. [1](#)
- [ABO03] V. A. Avetisov, A. H. Bikulov and V. A. Osipov.  $p$ -adic description of characteristic relaxation in complex systems. J. Phys. A 36 (2003), no. 15, 4239–4246. [4](#)
- [ABO04] V. A. Avetisov, A. H. Bikulov and V. A. Osipov.  $p$ -adic models of ultrametric diffusion in the conformational dynamics of macromolecules. Tr. Mat. Inst. Steklova 245 (2004), 55–64. Proc. Steklov Inst. Math. (2004), no. 2, 48–57. [1](#)
- [BGPV14] A. D. Bendikov, A. Grigor’yan, K. Pittet and V. Voess. Isotropic Markov semigroups on ultra-metric space. Uspekhi Mat. Nauk 69 (2014), no. 4 (418), 3–102; translation in Russian Math. Surveys 69 (2014), no. 4, 589–680. [23](#)
- [BGHH21] A. Bendikov, A. Grigor’yan, E. Hu and J. Hu. Heat kernels and non-local Dirichlet forms on ultrametric spaces. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 22 (2021), no. 1, 399–461. [23](#)
- [BeS88] C. Bennett and R. Sharpley. Interpolation of operators. Pure Appl. Math., 129. Academic Press, Inc., Boston, MA, 1988. [16](#)
- [BeF75] C. Berg and G. Forst. Potential theory on locally compact abelian groups. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 87. Springer-Verlag, New York-Heidelberg, 1975. [20](#), [22](#)
- [Ber52] H. Bergström. On some expansions of stable distribution functions. Ark. Mat. 2 (1952), 375–378. [22](#)
- [BiZ23] A. K. Bikulov and A. P. Zubarev. Oscillations in  $p$ -adic diffusion processes and simulation of the conformational dynamics of protein.  $p$ -Adic Numbers Ultrametric Anal. Appl. 15 (2023), no. 3, 169–186. [1](#)
- [Bou67] N. Bourbaki. Éléments de mathématique. Fasc. XXXII. Théories spectrales. Chapitre I: Algèbres normées. Chapitre II: Groupes localement compacts commutatifs. (French) Actualités Scientifiques et Industrielles, No. 1332 Hermann, Paris 1967. [7](#)
- [Bou98] N. Bourbaki. General topology. Chapters 1–4. Translated from the French. Reprint of the 1989 English translation. Elements of Mathematics. (Berlin). Springer-Verlag, Berlin, 1998. [11](#), [13](#)
- [Bou04] N. Bourbaki. Integration. II. Chapters 7–9. Translated from the 1963 and 1969 French originals by Sterling K. Berberian. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004. [5](#), [11](#)
- [BrF93] L. Brekke and P. Freund.  $p$ -adic numbers in physics. Phys. Rep. 233 (1993), no. 1, 1–66. [1](#), [20](#)

- [DH16] Y. de Cornulier and P. de la Harpe. Metric geometry of locally compact groups. EMS Tracts in Mathematics 25, 2016. [11](#)
- [DeE14] A. Deitmar and S. Echterhoff. Principles of harmonic analysis. Second edition. Universitext Springer, Cham, 2014. [12](#)
- [DKK21] L. Deleaval, M. Kemppainen and C. Kriegler. Hörmander functional calculus on UMD lattice valued  $L^p$  spaces under generalized Gaussian estimates. J. Anal. Math. 145 (2021), no. 1, 177–234. [16](#), [17](#)
- [DK23] L. Deleaval and C. Kriegler. Maximal Hörmander functional calculus on  $L^p$  spaces and UMD lattices. Int. Math. Res. Not. IMRN (2023), no. 6, 4643–4694. [19](#)
- [DeV22] J. Delgado and J. P. Velasquez-Rodriguez. Fundamental solution of the Vladimirov-Taibleson operator on noncommutative Vilenkin groups. Preprint, arXiv:2204.07133. [2](#), [11](#), [12](#), [13](#)
- [Den21] R. Denk. An introduction to maximal regularity for parabolic evolution equations. Springer, Proc. Math. Stat., 346, Singapore, 2021, 1–70. [3](#)
- [DJT95] J. Diestel, H. Jarchow and A. Tonge. Absolutely summing operators. Cambridge Studies in Advanced Mathematics, 43. Cambridge University Press (1995). [7](#)
- [DHW24] F. Ding, G. Hong and X. Wang. A noncommutative maximal inequality for Fejér means on totally disconnected non-abelian groups. Preprint, arXiv:2403.09263. [5](#), [12](#)
- [DKKVZ17] B. Dragovich, A. Y. Khrennikov, S. V. Kozyrev, I. V. Volovich and E. I. Zelenov.  $p$ -adic mathematical physics: the first 30 years.  $p$ -Adic Numbers Ultrametric Anal. Appl. 9 (2017), no. 2, 87–121. [1](#)
- [EdG77] R. E. Edwards and G. I. Gaudry. Littlewood-Paley and multiplier theory. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 90. Springer-Verlag, Berlin-New York, 1977. [4](#), [12](#)
- [En89] R. Engelking. General topology. Translated from the Polish by the author. Second edition. Sigma Series in Pure Mathematics, 6. Heldermann Verlag, Berlin, 1989. [10](#), [11](#), [12](#)
- [Fel71] W. Feller. An introduction to probability theory and its applications. Vol. II. John Wiley & Sons, Inc., New York-London-Sydney, 1971. [22](#)
- [Fol16] G. Folland. A course in abstract harmonic analysis. Second edition. Textbooks in Mathematics. CRC Press, Boca Raton, FL, 2016. [6](#), [7](#)
- [FGZ22] A. R. Fuquen-Tibatá, H. García-Compeán, W. A. Zúñiga-Galindo. Euclidean quantum field formulation of  $p$ -adic open string amplitudes. Nuclear Phys. B 975 (2022), Paper No. 115684, 27 pp. [2](#)
- [Gra14] L. Grafakos. Classical Fourier analysis. Third edition. Graduate Texts in Mathematics, 249. Springer, New York, 2014. [17](#)
- [Gou20] F. Q. Gouvêa.  $p$ -adic numbers. Universitext, Springer, Cham, 2020. [6](#)
- [Gri23] A. Grigor’yan. Analysis on ultra-metric spaces via heat kernels.  $p$ -Adic Numbers Ultrametric Anal. Appl. 15 (2023), no. 3, 204–242. [23](#)
- [Haa06] M. Haase. The functional calculus for sectorial operators. Operator Theory: Advances and Applications, 169. Birkhäuser Verlag (2006). [8](#)
- [Hal50] M. Jr. Hall. A topology for free groups and related groups. Ann. of Math. (2) 52 (1950), 127–139. [11](#)
- [Har90] S. Haran. Riesz potentials and explicit sums in arithmetic. Invent. Math. 101 (1990), no. 3, 697–703. [22](#)
- [Hara93] S. Haran. Analytic potential theory over the  $p$ -adics. Ann. Inst. Fourier (Grenoble) 43 (1993), no. 4, 905–944. [22](#)
- [HeR79] E. Hewitt and K. A. Ross. Abstract harmonic analysis. Vol. I. Structure of topological groups, integration theory, group representations. Second edition. Grundlehren der Mathematischen Wissenschaften, 115. Springer-Verlag, Berlin-New York, 1979. [10](#), [11](#)
- [HoM13] K. H. Hofmann and S. A. Morris. The structure of compact groups. 3rd Edition, Revised and Augmented. De Gruyter Studies in Mathematics 25, 2013. [11](#)
- [HvNVW16] T. Hytönen, J. van Neerven, M. Veraar and L. Weis. Analysis in Banach spaces, Volume I: Martingales and Littlewood-Paley theory. Springer, 2016. [13](#), [15](#)
- [HvNVW18] T. Hytönen, J. van Neerven, M. Veraar and L. Weis. Analysis in Banach spaces, Volume II: Probabilistic Methods and Operator Theory. Springer, 2018. [4](#), [7](#), [8](#), [9](#), [17](#), [18](#), [19](#), [25](#)

- [HvNVW23] T. Hytönen, J. van Neerven, M. Veraar and L. Weis. Analysis in Banach spaces, Volume III: Harmonic Analysis and Spectral Theory. Springer, 2023. [3](#), [8](#)
- [Jac89] N. Jacobson. Basic algebra. II. Second edition. W. H. Freeman and Company. New York, 1989.
- [Jan22] S. Janson. Stable distributions. Preprint, arXiv:1112.0220. [22](#)
- [KaW01] N. J. Kalton and L. Weis. The  $H^\infty$ -calculus and sums of closed operators. Math. Ann. 321 (2001), no. 2, 319–345. [9](#)
- [KaV22] A. Kassymov and J.P. Velasquez-Rodriguez. Hardy inequalities on constant-order noncommutative Vilenkin groups. Preprint, arXiv:2208.03495. [5](#)
- [KKZ18] A. Y. Khrennikov, S. V. Kozyrev and W. A. Zuniga-Galindo. Ultrametric pseudodifferential equations and applications. Encyclopedia of Mathematics and its Applications, 168. Cambridge University Press, Cambridge, 2018. [1](#), [4](#)
- [KOJ16a] A. Y. Khrennikov, K. Oleschko, M. de Jesús Correa López. Application of  $p$ -adic wavelets to model reaction-diffusion dynamics in random porous media. J. Fourier Anal. Appl. 22 (2016), no. 4, 809–822. [1](#)
- [KOJ16b] A. Khrennikov, K. Oleschko and M. de Jesús Correa López. Modeling fluid’s dynamics with master equations in ultrametric spaces representing the treelike structure of capillary networks. Entropy 18 (2016), no. 7, Paper No. 249, 28 pp. [1](#)
- [KhK20] A. Y. Khrennikov and A. N. Kochubei. On the  $p$ -adic Navier-Stokes equation. Appl. Anal. 99 (2020), no. 8, 1425–1435. [2](#)
- [Koc01] A. N. Kochubei. Pseudo-differential equations and stochastics over non-Archimedean fields. Monographs and Textbooks in Pure and Applied Mathematics, vol. 244, Marcel Dekker Inc., New York, 2001. [1](#), [2](#), [22](#), [23](#), [25](#)
- [KrW18] C. Kriegler and L. Weis. Spectral multiplier theorems via  $H^\infty$  calculus and  $R$ -bounds. Math. Z. 289 (2018), no. 1-2, 405–444. [4](#), [9](#)
- [KuW04] P. C. Kunstmann and L. Weis. Maximal  $L_p$ -regularity for parabolic equations, Fourier multiplier theorems and  $H^\infty$ -functional calculus. Functional analytic methods for evolution equations, 65–311, Lecture Notes in Math., 1855, Springer, Berlin, 2004. [3](#), [8](#)
- [LiT79] J. Lindenstrauss and L. Tzafriri. Classical Banach spaces. II. Function spaces. Ergeb. Math. Grenzgeb., 97 [Results in Mathematics and Related Areas] Springer-Verlag, Berlin-New York, 1979. [16](#)
- [Lor19] E. Lorist. The  $\ell^s$ -boundedness of a family of integral operators on UMD Banach function spaces. Positivity and noncommutative analysis, 365–379, Trends Math., Birkhäuser/Springer, Cham, 2019. [17](#), [19](#)
- [OnQ89] C. W. Onneweer and T. S. Quek. Multipliers on weighted  $L^p$ -spaces over locally compact Vilenkin groups. Proc. Amer. Math. Soc. 105 (1989), no. 3, 622–631. [5](#)
- [Onn88] C. W. Onneweer. Weak  $L^p$ -spaces and weighted norm inequalities for the Fourier transform on locally compact Vilenkin groups. Proceedings of the analysis conference, Singapore 1986, 191–201, North-Holland Math. Stud., 150, North-Holland, Amsterdam, 1988. [13](#), [14](#)
- [Pis16] G. Pisier. Martingales in Banach spaces. Cambridge Studies in Advanced Mathematics, 155. Cambridge University Press, Cambridge, 2016. [15](#)
- [Que87] T. S. Quek. Weighted norm inequalities for the Fourier transform on certain totally disconnected groups. Proc. Amer. Math. Soc. 101 (1987), no. 1, 113–121. [13](#)
- [RaV99] D. Ramakrishnan and R. J. Valenza. Fourier Analysis on Number Fields. Graduate Texts in Mathematics 186, Springer-Verlag, New York Inc. (1999). [5](#), [6](#), [7](#), [14](#)
- [Rob00] A. Robert. A course in  $p$ -adic analysis. Grad. Texts in Math., 198. Springer-Verlag, New York, 2000. [6](#)
- [RoZ08] J. J. Rodriguez-Vega and W. A. Zuniga-Galindo. Taibleson operators,  $p$ -adic parabolic equations and ultrametric diffusion. Pacific J. Math. 237 (2008), no. 2, 327–347. [2](#), [22](#), [23](#), [25](#)
- [SWS90] F. Schipp, W. R. Wade and P. Simon. Walsh series. An introduction to dyadic harmonic analysis. Adam Hilger, Ltd., Bristol, 1990. [5](#)
- [Spe70] R. Spector. Sur la structure locale des groupes abéliens localement compacts. (French). Bull. Soc. Math. France Suppl. Mém. 24 (1970), 94 pp. [5](#), [10](#), [11](#), [13](#)

- [SBSW15a] R. S. Stankovic, P. L. Butzer, F. Schipp and W. R. Wade. Dyadic Walsh analysis from 1924 onwards—Walsh-Gibbs-Butzer dyadic differentiation in science. Vol. 1. Foundations. A monograph based on articles of the founding authors, reproduced in full. In collaboration with the co-authors: Weiyi Su, Yasushi Endow, Sandor Fridli, Boris I. Golubov, Franz Pichler and Kees Onneweer. Atlantis Studies in Mathematics for Engineering and Science, 12. Atlantis Press, Paris, 2015. [4](#), [5](#), [12](#)
- [SBSW15b] R. S. Stankovic, P. L. Butzer, F. Schipp and W. R. Wade. Dyadic Walsh analysis from 1924 onwards—Walsh-Gibbs-Butzer dyadic differentiation in science. Vol. 2. Extensions and generalizations. A monograph based on articles of the founding authors, reproduced in full. In collaboration with the co-authors: Weiyi Su, Yasushi Endow, Sandor Fridli, Boris I. Golubov, Franz Pichler and Kees Onneweer. Atlantis Studies in Mathematics for Engineering and Science, 13. Atlantis Press, Paris, 2015. [5](#)
- [Ste70] E. M. Stein. Topics in harmonic analysis related to the Littlewood-Paley theory. Annals of Mathematics Studies, No. 63 Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1970. [10](#), [15](#)
- [Sti19] P. R. Stinga. User’s guide to the fractional Laplacian and the method of semigroups. Handbook of fractional calculus with applications. Vol. 2, 235–265. De Gruyter, Berlin, 2019. [2](#)
- [Tai68] M. Taibleson. Harmonic analysis on  $n$ -dimensional vector spaces over local fields. I. Basic results on fractional integration. Math. Ann. 176 (1968), 191–207. [21](#)
- [Tai75] M. Taibleson. Fourier analysis on local fields. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1975. [1](#), [6](#), [12](#), [14](#), [20](#), [21](#), [22](#)
- [Vel19] J. P. Velasquez-Rodriguez. Spectral properties of pseudo-differential operators over the compact group of  $p$ -adic integers and compact Vilenkin groups. Preprint, arXiv:1912.11407.
- [Vil63] N. J. Vilenkin. On a class of complete orthonormal systems. Amer. Math. Soc. Transl. (2) 28 (1963) 1–35. [4](#), [10](#), [11](#)
- [Vla99] V.S. Vladimirov. Tables of Integrals of Complex-valued Functions of  $p$ -Adic Arguments. Preprint, arXiv:math-ph/9911027. [22](#)
- [VIV89] V. S. Vladimirov and I. V. Volovich.  $p$ -adic quantum mechanics. Comm. Math. Phys. 123 (1989), no. 4, 659–676. [2](#)
- [VVZ94] V. S. Vladimirov, I. V. Volovich and E. I. Zelenov.  $p$ -adic analysis and mathematical physics. Ser. Soviet East European Math., 1, World Scientific Publishing Co., Inc., River Edge, NJ, 1994. [1](#), [14](#), [21](#)
- [Wei07] F. Weisz. Almost everywhere convergence of Banach space-valued Vilenkin-Fourier series. Acta Math. Hungar. 116 (2007), no. 1-2, 47–59. [5](#)
- [Wil98] J. S. Wilson. Profinite groups. London Mathematical Society Monographs. New Series, 19. The Clarendon Press, Oxford University Press, New York, 1998. [11](#), [12](#)
- [ZaZ23] B. A. Zambrano-Luna and W. A. Zuniga-Galindo.  $p$ -adic cellular neural networks: applications to image processing. Phys. D 446 (2023), Paper No. 133668, 11 pp. [2](#)
- [Zun16] W. A. Zuniga-Galindo. Pseudodifferential equations over non-Archimedean spaces. Lecture Notes in Mathematics, 2174. Springer, Cham, 2016. [1](#)
- [Zun22] W. A. Zuniga-Galindo. Non-Archimedean quantum mechanics via quantum groups. Nuclear Phys. B 985 (2022), Paper No. 116021, 21 pp. [2](#)
- [Zun24a] W. A. Zuniga-Galindo. Deep Neural Networks: A Formulation Via Non-Archimedean Analysis. Preprint, arXiv:2402.00094. [2](#)
- [Zun24b] W. A. Zuniga-Galindo.  $p$ -Adic Quantum Mechanics, the Dirac Equation, and the violation of Einstein causality. Preprint, arXiv:2312.02744. [2](#)

Cédric Arhancet  
 6 rue Didier Daurat  
 81000 ALBI  
 FRANCE  
 URL: <https://sites.google.com/site/cedricarhancet>  
 cedric.arhancet@protonmail.com  
 ORCID: 0000-0002-5179-6972

Christoph Kriegler  
Université Clermont Auvergne  
CNRS  
LMBP  
F-63000 CLERMONT-FERRAND  
FRANCE  
URL: <https://lmbp.uca.fr/~kriegler/indexenglish.html>  
christoph.kriegler@uca.fr  
ORCID: 0000-0001-8120-6251