



Radiative transfer: scattering processes

Benjamin Godard

► To cite this version:

| Benjamin Godard. Radiative transfer: scattering processes. Master. France. 2023. hal-04645954

HAL Id: hal-04645954

<https://hal.science/hal-04645954v1>

Submitted on 12 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Scattering processes

International Research Track

2023

Lecture description


Outline

- introduction
 - Thomson scattering
 - Rayleigh scattering
 - Mie scattering
 - Compton scattering
 - Monte-Carlo num method
 - other num method
 - conclusions
-
1. Introduction
 2. Thomson scattering
 3. Rayleigh scattering
 4. Mie scattering
 5. Compton scattering
 6. Monte-Carlo method
 7. Other methods
 8. Conclusions

Radiative transfer equation

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions



F. Levrier - radiative transfer lecture

$$\frac{dI_\nu}{dl} = - [\alpha_\nu + \sigma_\nu] I_\nu + j_\nu + \frac{1}{4\pi} \int_{\Omega'} \tilde{\sigma}_\nu(\mathbf{k}' \rightarrow \mathbf{k}) I_\nu(\mathbf{k}') d\Omega'$$

• absorption coefficient	$\alpha_\nu = \sum_a n_a s_{\nu,a}$	cm^{-1}
• diffusion coefficient	$\sigma_\nu = \sum_d n_d s_{\nu,d}$	cm^{-1}
• emissivity	j_ν	$\text{erg s}^{-1} \text{cm}^{-3} \text{Hz}^{-1} \text{sr}^{-1}$
• opacity	$d\tau_\nu = [\alpha_\nu + \sigma_\nu] dl$	
• source function	S_ν	$\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1} \text{sr}^{-1}$

Questions related to the scattering

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

what systems are involved in the scattering ?

- electrons, ions
- bound systems (atoms or molecules)
- macroscopic systems (dust, droplets, ...)

what are the properties of the microphysical processes ?

- elastic or inelastic scattering
- interaction cross section, angular dependence
- quantum and relativistic effects

how to solve problems with scattering

- analytical considerations
- numerical methods

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

Scattering processes

Field induced by a single charge in motion

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

consider a charge q in motion at a velocity $\mathbf{u}(t)$

→ Lienard-Wiechart retarded potentials $t' = t - \frac{|\mathbf{r} - \mathbf{r}'|}{c}$

$$\phi(\mathbf{r}, t) = \frac{q}{R(t') \left[1 - \frac{\mathbf{n}(t') \cdot \mathbf{u}(t')}{c} \right]}$$

$$\mathbf{A}(\mathbf{r}, t) = \frac{q\mathbf{u}}{cR(t') \left[1 - \frac{\mathbf{n}(t') \cdot \mathbf{u}(t')}{c} \right]}$$

→ for non-relativistic particles


$$\mathbf{E}_{\text{rad}} = \frac{q}{c^2 R(t')} \mathbf{n}(t') \times [\mathbf{n}(t') \times \dot{\mathbf{u}}]$$

$$\mathbf{B}_{\text{rad}} = \mathbf{n}(t') \times \mathbf{E}_{\text{rad}}$$

→ Larmor formula (total power emitted)

$$\frac{dP_{\text{rad}}}{d\Omega} = \frac{q^2 \dot{\mathbf{u}}^2}{4\pi c^3} \sin^2(\Theta)$$

$$P_{\text{rad}} = \frac{2q^2 \dot{\mathbf{u}}^2}{3c^3}$$



Dipole approximation

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

consider a system of charged particles

→ dipole moment

$$\mathbf{d}(t) = \sum_i q_i \mathbf{r}_i(t)$$

→ if all points emit in phase (i.e. $\lambda \gg L$)


$$\frac{dP_{\text{rad}}}{d\Omega} = \frac{\ddot{\mathbf{d}}^2}{4\pi c^3} \sin^2(\Theta)$$

$$P_{\text{rad}} = \frac{2\ddot{\mathbf{d}}^2}{3c^3}$$

→ emitted wavelength

$$\mathbf{d}(t) = \int_{-\infty}^{\infty} e^{-i\omega t} \hat{\mathbf{d}}(\omega) d\omega$$

$$\frac{dP_{\text{rad}}}{d\omega d\Omega} = \frac{\omega^4}{c^3} |\hat{\mathbf{d}}(\omega)|^2 \sin^2(\Theta)$$



Rybicki & Lightman

spectrum depends on oscillation frequencies of the dipole

Scattering by free electron - classical treatment

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

free electron at rest in interaction with radiation of pulsation ω

→ incident energy flux $\langle S \rangle = \frac{c}{8\pi} E_0^2$

→ classical treatment

dipole of amplitude

$$d_0 = \frac{e^2 E_0}{m_e \omega^2}$$



energy conservation

$$\frac{dP_{\text{rad}}}{d\Omega} = \langle S \rangle \frac{d\sigma}{d\Omega}$$


→ total cross section

$$\sigma_T = \frac{8\pi}{3} r_0^2 \sim 6.65 \times 10^{-25} \text{ cm}^2$$

$$r_0 = \frac{e^2}{mc^2} \sim 2.8 \times 10^{-13} \text{ cm}$$



Vertically polarized



Unpolarized

$$\frac{d\sigma}{d\Omega} = r_0^2 \sin^2(\Theta)$$

$$\frac{d\sigma}{d\Omega} = \frac{1}{2} r_0^2 [1 + \cos^2(\theta)]$$

Scattering by bound electrons

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

atom / molecule of polarizability η in interaction with radiation of pulsation ω

→ incident energy flux $\langle S \rangle = \frac{c}{8\pi} E_0^2$




→ classical treatment

dipole of amplitude $d_0 = \eta E_0$

energy conservation $\frac{dP_{\text{rad}}}{d\Omega} = \langle S \rangle \frac{d\sigma}{d\Omega}$

→ total cross section $\sigma = \frac{8\pi}{3} r_0^2$

$$r_0 = \frac{\omega^2 \eta}{c^2}$$



$$\frac{d\sigma}{d\Omega} = r_0^2 \sin^2(\Theta)$$


$$\frac{d\sigma}{d\Omega} = \frac{1}{2} r_0^2 [1 + \cos^2(\theta)]$$

From microscopic to macroscopic systems

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

diffusion by macroscopic systems (dust, water droplets, objects)



Malvern Instruments Ltd

- size parameter
- abs and scatt efficiencies
- index of refraction

$$\rightarrow x = \frac{2\pi a}{\lambda}$$

$$\rightarrow Q_{\text{abs}} = \frac{\sigma_{\text{abs}}}{\pi a^2}, Q_{\text{sca}} = \frac{\sigma_{\text{sca}}}{\pi a^2}$$

$$\rightarrow m(\lambda) = n(\lambda) + ik(\lambda)$$

scattering




absorption

From microscopic to macroscopic systems

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

spherical particles → Mie theory (1908)




From microscopic to macroscopic systems

Outline


- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

schematic view of scattering phase function




$$x = \frac{2\pi a}{\lambda}$$

example of Mie scatt phase function



simplified model: Henyey-Greenstein

$$p_g(\mu) = \frac{1}{2} \frac{1 - g^2}{(1 + g^2 - 2g\mu)^{3/2}}$$



Herbert Gross

Examples of classical scattering effects

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

Stellar Coronae




NASA / Aubrey Gemignani

Reflexion nebulae




NASA / Gary Stevens

Sky and cloud colors



circumstellar envelops & disks





model from RADMC3D - Dullemond

Scattering by free electrons - quantum treatment

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

free electron at rest in collision with a photon







→ quantum mechanics treatment

- recoil of the electron (inelastic)
- differential cross section (QED)
Klein-Nishina (unpolarized)

$$\lambda_1 - \lambda = \frac{h}{m_e c} (1 - \cos\theta)$$

$$\frac{d\sigma}{d\Omega} = \frac{1}{2} r_0^2 \left(\frac{\lambda}{\lambda_1} \right)^2 \left(\frac{\lambda_1}{\lambda} + \frac{\lambda}{\lambda_1} - \sin^2\theta \right)$$

→ simplifies to Thomson scattering for $\epsilon \ll m_e c^2$




Scattering by free electrons - quantum treatment

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

quantum effects on electron - photon scattering

- energy shift of the photon - loss of energy
- photon preferentially scattered forward
- reduction of the total cross section




Inverse Compton scattering

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

free electron in motion in collision with a photon



- step 1: from LF to ERF

$$\epsilon' = \epsilon \gamma (1 - \beta \cos \theta)$$

- step 2: scattering in ERF

$$\epsilon'_1 = \frac{\epsilon'}{1 + \frac{\epsilon'}{m_e c^2} (1 - \cos \theta')}$$

- step 3: from ERF to LF

$$\epsilon_1 = \epsilon'_1 \gamma (1 + \beta \cos \theta'_1)$$

if electron has a sufficient kinetic energy

→ transfer of energy to the photon : inverse Compton scattering

Inverse Compton scattering

Outline


- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

in the Thomson regime $\gamma\epsilon \ll m_e c^2$



→ energy gain $\frac{\epsilon_1}{\epsilon} = \gamma^2 (1 - \beta \cos\theta) (1 + \beta \cos\theta'_1)$

depends on angle but
 > 1 on average

strong gain for relativistic e⁻



→ schematic view of energy transfer for relativistic e⁻



Nick Keiser

Inverse Compton scattering

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

in the Thomson regime $\gamma\epsilon \ll m_e c^2$ and for isotropic radiation

→ single scattering radiated power

$$P_{\text{Comp}} = \frac{4}{3} \sigma_T c \gamma^2 \beta^2 U_{ph} \quad (\text{erg s}^{-1})$$

→ incident field at ν_0 only spectrum

→ thermal dist of non-relativistic e⁻

$$P_{\text{tot}} = \left(\frac{4kT}{m_e c^2} \right) \sigma_T c n_e U_{ph} \quad (\text{erg cm}^{-3} \text{s}^{-1})$$


energy transfer for repeated scattering in finite thermal medium

→ non-relativistic e⁻ : per scattering

$$\frac{\Delta\epsilon}{\epsilon} \sim \frac{(4kT - \epsilon)}{m_e c^2}$$

→ ultra-relativistic e⁻ : per scattering

$$\frac{\Delta\epsilon}{\epsilon} \sim \frac{4}{3} \gamma^2 = 16 \left(\frac{kT}{m_e c^2} \right)^2$$






Example of inverse Compton scattering

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

Sunyaev-Zeldovich effect



map of galaxy clusters
e- density & temperature
cluster size

Planck Collaboration (2015)

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

methods to solve transfer
with scattering


Complexity of radiative transfer

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

$$\frac{dI_\nu}{dl} = - [\alpha_\nu + \sigma_\nu] I_\nu + j_\nu + \frac{1}{4\pi} \int_{\Omega'} \tilde{\sigma}_\nu(\mathbf{k}' \rightarrow \mathbf{k}) I_\nu(\mathbf{k}') d\Omega'$$

- goal: find the source function or the mean intensity at each point
- complexity: scattering & thermochemistry couple all rays & cells
- simpler (yet complicated) question:
*how to solve the full radiative transfer
(including absorption & scattering)
knowing the thermal and chemical state
of the medium ?*



Dullemond (2012)

Scattering and random walk

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

the fate of a photon expressed in terms of probabilities

$$l_\nu = \frac{1}{\alpha_\nu + \sigma_\nu} \longrightarrow \text{mean free path}$$

$$\varepsilon_\nu = \frac{\alpha_\nu}{\alpha_\nu + \sigma_\nu} \longrightarrow \text{proba that a free path ends with abs}$$

$$\omega_\nu = 1 - \varepsilon_\nu = \frac{\sigma_\nu}{\alpha_\nu + \sigma_\nu} \longrightarrow \text{proba that a free path ends with scatt}$$

$$l_* = \sqrt{n} l_\nu \longrightarrow \text{rms net displacement for } n \text{ free paths}$$

for a medium of size L

$$\tau_\nu^a = \alpha_\nu L, \tau_\nu^s = \sigma_\nu L \longrightarrow \text{absorption and scattering optical depth}$$

$$N_\nu = \frac{1 - \varepsilon_\nu}{\varepsilon_\nu} \longrightarrow \langle \text{number} \rangle \text{ scatt before absorption}$$


$$N_\nu \approx \max(\tau_\nu^s, \tau_\nu^{s2}) \longrightarrow \langle \text{number} \rangle \text{ scatt before escape}$$

$$l_* = \min \left(L, \frac{1}{\sqrt{\alpha_\nu (\alpha_\nu + \sigma_\nu)}} \right) \longrightarrow \text{rms displacement in the cloud}$$

Monte Carlo basic algorithm

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions



Examples of probabilities and further readings

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

assign probability distributions to each step

- frequency of the photon packet $p(\nu) = \frac{L_\nu^*}{\int_0^\infty L_\nu^* d\nu}$
- emission direction (Lambertian source ? else ?)
- position of next scattering event $p(\tau) = \exp(-\tau)$
- scattering phase function
 - isotropic or more complex
 - $p(\mu), p(\phi) = \frac{1}{2}, \frac{1}{2\pi}$
 - $p_g(\mu), p(\phi) = \frac{1}{2} \frac{1 - g^2}{(1 + g^2 - 2g\mu)^{3/2}}, \frac{1}{2\pi}$

further readings on the subject


- amazing textbook by Dullemond (radiative transfer in astrophysics, 2012)
- Lucy (1999), Bjorkman & Wood (2001)
- Min et al. (2009), Robitaille et al. (2010)

Discrete ordinate method

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

class of method where all coordinates ($x, y, z, \nu, \theta, \phi$) are discretized



- examples of discrete ordinate methods
- lambda iteration (LI) or accelerated lambda iteration (ALI)
Cannon (1973), Olson et al. (1986)
 - decomposition of in Legendre polynomials + spectral method
Roberge (1983), Kopp (1996), Goicoechea & Le Bourlot (2007)

Moments method

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

reduce angular resolution by computing only the moments of I_ν

→ moments of I_ν transfer in 1D

$$J_\nu(x) = \frac{1}{4\pi} \int I_\nu(x, \mathbf{k}) d\Omega$$

$$\mu \frac{dI}{d\tau} = I - S$$

$$\mathbf{H}_\nu(x) = \frac{1}{4\pi} \int I_\nu(x, \mathbf{k}) \mathbf{k} d\Omega$$

$$\frac{dF}{d\tau} = 4\pi(J - S)$$

$$\mathbf{K}_\nu(x) = \frac{1}{4\pi} \int I_\nu(x, \mathbf{k}) \mathbf{k} \otimes \mathbf{k} d\Omega$$

$$c \frac{dp}{d\tau} = F$$

...

→ closure equation (e.g. Eddington approximation)

$$K_{i,j,\nu} = \frac{1}{3} \delta_{ij} J_\nu$$

$$p = \frac{4\pi}{3c} J$$

→ other closure equation: variable Eddington tensor

Outline

- introduction
- Thomson scattering
- Rayleigh scattering
- Mie scattering
- Compton scattering
- Monte-Carlo num method
- other num method
- conclusions

Scattering processes

- at the heart of radiative transfer (impact all astrophysics environments)
- microprocesses explained by multipolar decomposition
- bound / free charges, small / large λ , micro / macroscopic systems
- change the direction and the energy of the photon (Compton)
- makes radiative transfer complicated

Methods to solve the transfer with scattering require several hours lecture

- amazing textbook by Dullemond (radiative transfer in astrophysics)
- algorithm & optimization are active fields of current research
- intuition comes from practice