



**HAL**  
open science

## **Inhibition of anaerobic digestion of urban sewage sludge by sodium propionate: biogas production and microbial community shift in continuous reactors**

Joel Awinzure Agumah, Xiaojun Liu, Laura André, Sophie Thibault, Camille Auneau, Sabrina Guérin-Rechdaoui, Vincent Rocher, Carlyne Lacroix, Olivier Chapleur, Ariane Bize, et al.

### ► **To cite this version:**

Joel Awinzure Agumah, Xiaojun Liu, Laura André, Sophie Thibault, Camille Auneau, et al.. Inhibition of anaerobic digestion of urban sewage sludge by sodium propionate: biogas production and microbial community shift in continuous reactors. 18th IWA World Conference On Anaerobic Digestion, Jun 2024, Istanbul, Turkey. 18th IWA World Conference on Anaerobic Digestion CONFERENCE PROCEEDINGS, 2024. hal-04647882

**HAL Id: hal-04647882**

**<https://hal.science/hal-04647882>**

Submitted on 15 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Inhibition of anaerobic digestion of urban sewage sludge by sodium propionate: biogas production and microbial community shift in continuous reactors

J.A. Agumah<sup>1</sup>, X. Liu<sup>1</sup>, L. André<sup>2</sup>, C. Auneau<sup>4</sup>, S. Thibault<sup>4</sup>, S. Guérin<sup>3</sup>, V. Rocher<sup>3</sup>, C. Lacroix<sup>3</sup>, O. Chapleur<sup>4</sup>, A. Bize<sup>4\*</sup>, C. Roose-Amsaleg<sup>5</sup>, A. Pauss<sup>1</sup>, T. Ribeiro<sup>2</sup>

<sup>1</sup> Université de technologie de Compiègne, ESCOM, TIMR, Centre de recherche Royallieu CS 60319, 60203 Compiègne Cedex

<sup>2</sup> Institut Polytechnique UniLaSalle, Université d'Artois, ULR 7519, 60026 Beauvais, France

<sup>3</sup> Direction Innovation SIAAP – Service public pour l'assainissement francilien, 92700 Colombes, France

<sup>4</sup> Université Paris-Saclay, INRAE, PROSE, 92160, Antony, France

<sup>5</sup> CNRS, ECOBIO-UMR 6553, Université Rennes, 35000 Rennes, France

## CONTEXT


Anaerobic digestion (AD) is widely used to convert sewage sludge to biogas. Volatile fatty acid (VFA) accumulation, especially that of propionic acid, can inhibit AD and may lead to AD failure. Knowledge about the level of VFA concentration that inhibits the AD process as well as the microbial targets is limited. Further research is needed to understand this inhibition and to identify microbial biomarkers that predict possible reactor failure.

**OBJECTIVE:** Assess the impact of propionic acid (HPro) (its dissociated form, propionate (NaPro)) on biogas production and microbial community shift in order to identify potential microbial biomarkers

## MATERIALS AND METHODS


### 4 Continuous Stirred Anaerobic Reactors (R1, R2, R3 and R4)

- Substrate: Urban sewage sludge, with different amounts of sodium propionate
- HRT: 20 days
- Duration: 2HRT+Batch mode (no heating and feeding)
- Volume: 5 L




## RESULTS AND DISCUSSION

### EVOLUTION OF VFA AND BIOGAS PRODUCTION




- No VFA accumulation in R2 (1.95 g/L NaPro), negligible in R3 (7.78 g/L NaPro)
- Highest VFA accumulation observed in R4




- R4 produced less biogas at the end of 2<sup>nd</sup> HRT due to inhibition
- R4 showed a significant 40% reduction in CH<sub>4</sub> compared to expected value (data not shown)

### MICROBIAL COMMUNITY



- Identification of *peptococcaceae* family whose abundance increases at high propionate concentrations.



- The proportion of methanogenic archaea decreases according to the applied NaPro concentration

## CONCLUSION

- No inhibition at 1.95 g/L NaPro
- Slight inhibition at 7.78 g/L NaPro, inhibition at 13 g/L NaPro
- Drop in archaea abundance with addition of NaPro from 1.95 g/L NaPro
- Peptococcaceae* family could be a possible biomarker and could allow inhibition to be detected quickly

## ACKNOWLEDGMENTS

Thanks to MOCOPEE program for funding this work. Thanks to John Cockerill, Sources and Le Ministère de l'Enseignement Supérieur et de la Recherche (MESR) for funding this PhD grant

## REFERENCES

