



HAL
open science

Population genetic structure of *Schistosoma bovis* and *S. curassoni* collected from cattle in Mali

Assitan Diakité, Privat Agniwo, Abdoulaye Dabo, Bakary Sidibé, Boris A.E.S. Savassi, Ahristode Akplogan, Hassim Guindo, Laurent Dembélé, Moudachirou Ibikounlé, Safiatou Doumbo Niaré, et al.

► To cite this version:

Assitan Diakité, Privat Agniwo, Abdoulaye Dabo, Bakary Sidibé, Boris A.E.S. Savassi, et al.. Population genetic structure of *Schistosoma bovis* and *S. curassoni* collected from cattle in Mali. *Parasite*, 2024, 31, pp.36. 10.1051/parasite/2024035 . hal-04650137

HAL Id: hal-04650137

<https://hal.science/hal-04650137v1>

Submitted on 16 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Population genetic structure of *Schistosoma bovis* and *S. curassoni* collected from cattle in Mali

Assitan Diakit¹✉, Privat Agniwo^{1,2,3}✉, Abdoulaye Dabo¹, Bakary Sidibé¹, Boris A.E.S. Savassi^{2,3}, Ahristode Akplogan¹, Hassim Guindo¹, Laurent Dembélé¹, Moudachirou Ibikounlé³, Safiatou Doumbo Niaré², Saidou Tembely⁴, Jérôme Boissier²

1. Department of Epidemiology of Infectious Diseases, Faculty of Pharmacy, University of Sciences, Techniques and Technologies of Bamako
2. Interactions Hôtes-Pathogènes-Environnements (IHPE), Univ. Montpellier, CNRS, Ifremer, Univ. Perpignan Via Domitia, Perpignan, France.
3. Centre de Recherche pour la lutte contre les Maladies Infectieuses Tropicales (CReMIT/TIDRC), Université d'Abomey-Calavi
4. Académie des Sciences du Mali, Bamako, Mali

Correspondence: adabo@icermali.org

✉ These authors contributed equally to this work.

Abstract

Schistosomiasis is of medical and veterinary importance. Despite the critical situation of schistosomiasis in sub-Saharan Africa, few molecular epidemiological studies have been carried out to determine the role of animals in the transmission of schistosomiasis. In Mali, it has been over three decades since the last molecular study of animal schistosomes was carried out. It is now urgent to identify circulating strains of the parasite because of potential interactions with other schistosome species, which could complicate disease control. The aim of our work was to study the composition and genetic structure of schistosome populations collected from cattle. The prevalence of schistosome was 23.9%, with the prevalences of *Schistosoma bovis* (*Sb*) and *S. curassoni* (*Sc*) estimated at 12.6% and 9.8% respectively. No hybrid strains or *S. haematobium* were found. The parasites displayed distinct geographical distribution with *Sb* dominant in Bamako (78.8% and 98% in CBS and SS, respectively) and *Sc* dominant in Kayes (95.3%). Of the 476 parasites with a complete genetic profile, 60.4% were pure *Sc*, and were mainly from Kayes. We identified two clusters at the site level (*F*_{st} of 0.057 and 0.042 for *Sb* and *Sc*, respectively). Cluster 1 was predominantly composed of pure *Sb* parasites and cluster 2 was mainly composed of pure *Sc* parasites, from Bamako and Kayes, respectively.

Our study shows that cattle schistosomiasis remains endemic in Mali with *Sb* and *Sc*. A good genetic structure between the different schistosome populations was identified, which included two clusters based on the geographical distribution of the parasites.

Keys words: *Schistosoma bovis*, *S. curassoni*, slaughterhouse, cattle, genetic structure, Mali.

Introduction

Schistosomiasis continues to be a major public health threat in tropical and subtropical areas. This parasitosis belongs to a group of Neglected Tropical Diseases which collectively affect 1 billion people worldwide, and of which schistosomiasis accounts for infections in 250 million people in 78 countries [5]. It was estimated that worldwide, approximately 165 million animals are also infected with the parasite, which causes hemorrhagic

enteritis, anemia, cachexia and for most, death [10]. In Mali, compared to the numerous studies done on human schistosomiasis, livestock schistosomiasis has received little attention and the most recent studies were carried out more than 30 years ago, hence the need to update existing data [36]. Of the 23 species of *Schistosoma* known worldwide, only nine have been identified to be of significant veterinary importance especially for ruminants in Asia and Africa; these include: *Schistosoma mattheei*; *S. bovis* (*Sb*); *S. curassoni* (*Sc*); *S. spindale*; *S. indicum*; *S. nasale*; *S. incognitum*; *S. margrebowiei*; and *S. japonicum* [4, 22, 35]. Of these, two species, *Sb* and *Sc* are native to West Africa, six are widely distributed in Central and Eastern Africa [22, 26, 28, 31, 36, 41, 42] and one, *S. japonicum* is native to Asia [40]. The prevalence of *Sb* and *Sc* has been reported in some countries [26, 28, 36, 41, 42]. A study by Rollinson [36] in 1990 reported a high prevalence of schistosomiasis in cattle across three West African countries, namely Gambia (*S. bovis* : 14.3-27%), Senegal (*Sb* plus *Sc* and hybrids: 0 - 41.3%) and Mali (*Sb*, *Sc* and hybrids) with 62.8% in Bamako and 85.1% in Mopti. The two most recent molecular epidemiology studies of animal schistosomes were carried out in Senegal and in Cameroon [12, 21]. In Cameroon, a recent study showed a 19.5% prevalence of adult schistosome worms in cattle [12], and prevalences of 92% and 8% respectively were reported for *S. bovis* and *S. curassoni* in cattle in Senegal [21]. The control or elimination of cattle schistosomiasis will contribute to a substantial increase in the weight of the animals, thus improving animal productivity, which in turn will increase the incomes of livestock farmers and the availability of quality meat for consumption.

The schistosome parasite has a complex life cycle. They are dioecious rather than hermaphroditic as is the case with most other trematodes like liver flukes or tapeworms. Gonochorism allows interspecific interactions between males and females within their definitive hosts. Today, parasite hybridization is an emerging public health problem, as the cohabitation of human populations and domestic or wildlife animals. Increased mobility of populations creates opportunity for new agents to combine, and for parasites, this includes new interactions between parasites of different lineages or species within individual hosts [20, 29, 39]. The genetic variation of schistosome species and populations over time and space is based on the hybridization phenomenon and can only be described through molecular epidemiological studies. While many studies on population genetics of schistosomes in humans have been conducted [35], only few studies exist for animal populations. Parasite hybridization is a biological phenomenon that is gaining renewed interest as new molecular tools have enabled hybridization to be more easily detected, and because the repercussions in the evolution of infections are still largely understood [19]. Hybridization is known in some cases to increase the virulence of the parasite towards its vertebrate hosts and to allow the parasite to broaden its spectrum of intermediate and/or definitive hosts [19, 47]. The first cases of hybridization between schistosomes, notably between *Sb* and *Sc* in animals, were reported in Africa (Mali & Senegal) by Rollinson [36]. In Senegal, a recent study found a high occurrence of bidirectional hybridization between these animal schistosome species, and is the second piece of conclusive evidence of natural hybridization between *Sb* and *Sc* [44]. Also in the Niger River valley region, transmission of human and bovine schistosomiasis has been identified, but it seems that the transmission of the different species was limited [32].

The only population genetic study carried out on livestock schistosome was done in Cameroon [12]. The present study aims to understand the genetic composition and the population genetic structure of schistosomes collected from cattle in Mali. Specifically, we proposed (i) to determine the prevalence of schistosomes in cattle in Kayes and Bamako slaughterhouses; (ii) to identify the schistosome species and the eventual presence of *S.*

haematobium group hybrid parasites using genetic markers (ITS, 18S and Cox1); and (iii) to describe the genetic structure of schistosomes in cattle based on microsatellite data.

Material and methods

Study sites

Mali ranks first in the West African Economic and Monetary Union (UEMOA) and second in the [Economic Community of West African States \(ECOWAS\)](#) zone in terms of bovine breeding, with an estimated 12,474,462 heads of cattle in its national herd [34]. A cross-sectional study was performed for two collections, in September (end of the rainy season) and December (cold dry season) 2021 in two areas: at the Central Bamako Slaughterhouse (CBS) and Sabalibougou Slaughterhouses (SS) in District of Bamako (Koulikoro region) and at the Kayes Slaughterhouse (KS) in Kayes region (**Figure 1**). Unlike the two slaughterhouses located in the District of Bamako, where animals from all regions of the country converge, Kayes is an extensive livestock farming area. The two areas were specifically selected because they are known to be endemic for *S. haematobium* (*Sh*) [1], a human schistosome species. *Sb* and *Sc* are two schistosome species that infect animals [36]. In Mali, the Ministry of Livestock and Fisheries, through the National Veterinary Services Department, is the official body responsible for animal health. Anthrax, contagious bovine pneumonia, and plague of small ruminants are among the diseases managed by this service. No specific treatment is currently provided for schistosomiasis, even in small-scale dairy farms. Parasites were isolated and conditioned at the Department of Epidemiology and Parasitic Diseases (DEAP) of the Malaria Research and Training Center (MRTC) in Mali, and molecular manipulations were carried out at the Host-Pathogen-Environment Interactions (IHPE) laboratory in France.

Parasite collection

Collections of adult schistosomes were made between 0:00 am and 3:00 am from slaughtered cattle. Each collection session was followed by the identification of the animal from which the worms were collected, on a pre-established survey form. The site, the season, the number of worms recovered from each animal and the organ from which the worm was collected were recorded. In each slaughterhouse, intestines and caeca were randomly collected from slaughtered animals. The mesenteric veins surrounding the small intestines, and the veins which irrigate the caeca were carefully observed macroscopically by transparency for the presence of schistosome worms (whitish-colored worms through the vein). The worms were removed from the veins using forceps and placed in a 15 ml falcon tube containing physiological water, labelled with the bovine identifier and the removed organ. Once in the laboratory, the worms were isolated one by one using fine tweezers and stored individually in pure ethanol.

Extraction of genomic DNA (gDNA)

To extract the genomic DNA (gDNA), the ethanol was first removed using a pipette, and then the tube containing the parasite was kept open and dried using the Speed Vac set at 14000 rpm. The genomic DNA of 491 parasites was extracted using the DNeasy blood and tissue kit (*QIAGEN strosse1, 40724 Hilden, Germany*) following the manufacturer's protocol and the DNA was eluted in a total of 100 µl.

Rapid Diagnostic Polymerase Chain Reaction (RD-PCR) for Cox1 and Tetraprimer Amplification Refractory Mutations System - Polymerase Chain Reaction (T-ARMS-PCR) for ITS2 and 18S genotyping

Identification of Cox 1 profiles from 491 worms was done by Cox1 rapid diagnostic multiplex PCR (RD-PCR) with primers targeting two species groups: *Sh* and *Sb* or *Sc* [43]. In short, we used a universal reverse (Shmb.R) and two forward primers specific to three schistosome species: *ShF* (120 bp), *SbF* (260 bp – this band is the same for both *Sc* and *Sb*). Each PCR was carried out in a total volume of 10 µl containing 4 µL of ultra-pure water, 2 µL of buffer (Green GoTaq Flexi buffer, 5X; Promega; Madison, Wisconsin, USA), 0.6 µL of MgCl₂ (Promega) at 25 mM, 0.2 µL of mixture of dNTPs (Promega) at 10 mM; 1µL of 10X primer mix with 4 µl of each primer in 84µl of ultra-pure water), 0.2µL of GoTaq® Polymerase G2 Hot Start at 5U (Promega), and 2 µL of DNA extract. After 45 cycles of amplification, the PCR products (Cox1) were visualized on a 2% agarose gel at 135 V for 40 minutes using the 100 bp DNA size marker (Promega) to estimate the size of the bands. We used the T-ARMS-PCR method developed in a recent study [32] to detect the parasites' nuclear genotypes (ITS/18S). Each partial PCR amplification of nuclear DNA was performed in a total volume of 25 µl, containing 3 µl of DNA template, 5 µl of GoTaq® Flexy buffer (Promega; Madison, Wisconsin, USA), 1, 5 µl of 25mM MgCl₂, 0.5 µl of a mixture of 8 prediluted primers (12µM external primers and 15 µM for internal ones), 0.2 µl of dNTP solutions at 10 mM each, 0.5 µl of GoTaq® Taq polymerase G2 Hot Start (Promega) and 10.75 µl of milli-Q water. After 28 cycles of amplification, PCR products were examined on 1.8% agarose gels at 135 V for 40 min using the 100 bp DNA size marker (Promega) for size estimation. Four to six bands could be observed on the agarose gels which allowed for the differentiation between *Sc*, *Sb*, *Sh* and all hybrid combinations [4].

Sanger sequencing

Cox1: Because the RD-PCR does not differentiate between *Sc* and *Sb*, we sequenced Cox 1 partial region (~918 bp) for 188 parasites using a previously published protocol [23]. We considered the T-ARMS-PCR result as an indicator of sample selection for sequencing. We sequenced 107 and 81 parasites with either the *Sb* and *Sc* T-ARMS-PCR profile, respectively. The size fragment was checked on 5% agarose gel and PCR products were sent without purification to a subcontractor for sequencing (Genoscreen, Lille, France). Cox1 sequences were then compared to a reference sequence from GenBank (Accession number: AJ519521.1 and MT579424.1 for *Sb* and *Sc*, respectively).

18S: In order to double check several of the T-ARMS-PCR results, partial 18S regions (329 bp) were amplified and sequenced on 26 and 20, pure *Sb* and pure *Sc* profiles, as previously described [32]. The size fragment was checked on 5% agarose gel and PCR products were sent without purification to a subcontractor for sequencing (Genoscreen, Lille, France). The sequences obtained were compared with reference sequences (Accession number: AY157238.1 and AY157236.1 for *Sb* and *Sc*, respectively). At position 297, a C base identified *Sb* and a T base identified *Sc* [4].

Microsatellite genotyping

Microsatellite genotyping was performed on parasites collected from the three slaughterhouses. Only samples of parasites with a well-defined profile for each of the three genes (Cox1, ITS and 18S) were selected for microsatellite genotyping. The worms were individually genotyped using the PCR multiplex kit from Qiagen® and two panels of 8 loci : (*Sh9*, *Sh3*, C102, *Sh1*, *Sh14*, *Sh6*, C111, *Sh7*) & (*Sh13*, *Sh4*, *Sh11*, *Sh15*, *Sh2*, *Sh5*, *Sh10*,

Sh12) [45]. The PCR mixture consisted of 5µl of Qiagen MM 2X, 1µl of 10X microsatellite primer mix and 4µl of DNA extract for a final volume of 10µl. Thermal cycling was performed in a PerkinElmer 9600 Thermal Cycler (PerkinElmer, Waltham, Massachusetts, USA): pre-denaturation at 95°C for 15 minutes followed by 40 cycles. Microsatellite PCR products were sent to Genoscreen, Lille, France, for genotyping. After verification and correction using GeneMarker software, the data were then transferred to Excel to determine the position of the various loci.

Statistical analyses

Data were entered using Excel software. Calculations of prevalence of infection was performed using SPSS v 23.0 (IBM) software. Logistic regressions were used to infer the relationship between variables (slaughterhouse, season, and organ of sampling) and genetic profile. *p*-values less than 0.05 were considered as significant.

Genetic data analysis

Observed (H_o) and expected (H_e) heterozygosity, number of alleles (A), allelic richness (Ar) and inbreeding coefficient (F_{is}) at each microsatellite locus and for each population were calculated using FSTAT 2 software. 9.3.2 [16]. Population refers to the set of parasites collected from a given slaughterhouse (CBS, SS or KS). These parameters were compared according to the different populations or species using either Wilcoxon or Friedman pairwise rank tests followed by Nemenyi post-hoc tests.

First, population genetic structure was assessed using F_{st} indices [46]. Second, Genetix software [3] enabled us to visualize the potential genetic structure between individuals according to species using principal component analysis (PCA). Third, we evaluated the probability that the microsatellite database followed a defined number of clusters from $K=1$ to $K=4$ using Structure software [33]. We considered 10 runs for each trial, each of which consisted of a running period of 50,000 following by 250,000 MCMC iterations. The ΔK values were then calculated in R to determine the most likely number of clusters [13]. Finally, 10 additional runs were calculated for the best K using a running period of 50,000 and 500,000 iterations. The probability of each worm belonging to each cluster was averaged over the 10 runs and graphically represented using CLUMPP, version 1.1.2 [14] and DISTRUCT, version 1.1 [37].

Results

Schistosome prevalence in cattle

Animals slaughtered in slaughterhouses were mainly local Zebu breeds (*Bos taurus indicus*) and less commonly crossbreeds with N'Dama (*B. taurus taurus*). Post-mortem examination of veins surrounding the small intestine and the caecum of cattle from Bamako and Kayes slaughterhouses revealed the presence of adult schistosome worms. Out of 398 cattle examined at the three slaughterhouses (two in Bamako and one in Kayes), 23.9% (95/398) were found to be infected by schistosomes (**Table 1**). Of these 95 cattle, we obtained the species of parasite involved for 89 individuals, with 50 animals infected by *Sb* parasites and 39 animals infected by *Sc* parasites. We can therefore estimate prevalences of 12.6% (50/398) and 9.8% (39/398) for *Sb* and *Sc*, respectively. Note that 4/89 were co-infected by both species (1 from CBS and 3 from KS). The highest prevalence was recorded in KS (44.6%) compared to CBS (18.4%). More worms were recovered from the intestine (22.9%)

compared to caecum veins (5%). No statistical difference in the prevalence was observed between seasons ($p=0.12$). Sixteen bovines (6.5%) had parasites in both the small intestine and the caecum.

Schistosome species identification

Out of 616 adult worms collected, 491 were genetically analyzed. Correct genetic profiles for 491 and 476 worms were obtained using RD-PCR and T-ARMS-PCR, respectively. Results from these analyses indicated that the majority of the parasites originated from KS 58.4 % (278/476) (**Table 2**). The T-ARMS-PCR revealed either *Sc* or *Sb* homozygous profiles and no hybrids were identified (**Table 2**). All 18S sequences (26 and 20 with *Sb* and *Sc* T-ARMS-PCR profile, respectively) confirmed the T-ARMS-PCR profiles. The RD-PCR revealed only the *Sb/Sc* profile and not the *Sh* profile. All 107 parasites with a homozygous *Sb* T-ARMS-PCR profile showed a typical Cox1 *Sb* sequence and all 81 parasites with a homozygous *Sc* T-ARMS-PCR profile showed a typical Cox1 *Sc* sequence. Because we randomly selected 40% of the parasites for sequencing, we assumed that the T-ARMS-PCR (*SbxSb* or *ScxSc*) profile reflected the Cox1 profile (*Sb* or *Sc*). *Sb_SbxSb* profiles were predominantly present in Bamako with 78.8% in CBS and 98% in SS, respectively. All haplotypes are available on Genbank with access numbers for Cox1: *Sb* (PP654220 to PP654290); *Sc* (PP654291 to PP654330) and 18S: *Sb* (PP654446, PP654447, PP654448); *Sc* (PP654443, PP654444, PP654445).

Tropism of schistosome species

The vast majority of worms analyzed (81.9%) were localized in the veins surrounding the small intestine (Table 2). The small number of *Sc_ScxSc* parasites collected from Bamako were also principally localized in the veins surrounding the small intestine. In contrast, most parasites from KS exhibited an *Sc_ScxSc* profile (95.3%). The localization of these worms is more shared with 82% (390/476) localized in the veins surrounding the small intestine.

Microsatellite analysis

Genetic diversity

Among the 476 parasites with complete mitochondrial and nuclear profiles, 166 were genotyped using 16 microsatellite markers. Parasite genetic diversity indices (He, Ho, A, Ar and Fis) at each marker and for each slaughterhouse are presented in Table 3. The Friedman test showed a difference between populations for all calculated indices, He ($\chi^2 = 8$, $df = 2$, $P = 0.011$), Ar ($\chi^2 = 8.375$, $df = 2$, $p = 0.015$) and Fis ($\chi^2 = 12.5$, $df = 3$, $p = 0.0017$) (**Table 3**). The Nemenyi post-hoc tests revealed that the difference in Ar and He observed was significant between CBS and KS ($p=0.013$ and $p=0.009$, respectively). For Fis, significant difference was observed between SS and KS ($p=0.002$). Table 4 represents the overall gene diversity (He) and allelic richness (Ar) for *Sb_SbxSb* and *Sc_ScxSc*, Mean He and Ar are both higher for *Sb_SbxSb* than for *Sc_ScxSc* but the difference is only significant for He ($W=116$, $p=0.21$ for Ar, $W=116$, $p=0.01$ for He) (**Table 4**).

Population genetic structure

Whatever's the PCA (**Figure 2**) or Bayesian approach (**Figure 3**), two groups were clearly identified. The first group is mainly composed of parasites from Bamako (CBS or CS) and the second group is mainly composed of parasites from Kayes (KS). According to the parasite genotyping, this structure is more a result of

the differences between species than to the sampling site. The few parasites from Kayes group that clustered with the Bamako group are *Sb_SbxSb* and conversely the few parasites from Bamako group that clustered with the Kayes group are *Sc_ScxSc* parasites. The PCA first axes cluster the genotypes, and the second axes cluster the sites. The Bayesian analysis (**Figure 3**) also shows that the vast majority of parasites are either assigned to cluster 1 and possess a *Sb_SbxSb* profile or to cluster 2 and possess a *Sc_ScxSc* profile. A single parasite from Bamako shows a 50/50 assignment probability between cluster 1 and 2. This parasite has an *Sb_SbxSb* profile in both PCR and sequencing and thus has not been identified as a hybrid.

PCAs by species show that whatever the species, *Sb* (**Figure 4**) or *Sc* (**Figure 5**), the parasites are structured according to the collection site. Pairwise *Fst* value between *Sb_SbxSb* and *Sc_ScxSc* genotypes is 23%. Table 5 presents the *Fst* values between genotypes at different sites. Except for the *Sb_SbxSb* genotype between SS and CBS the *Fst* values range between 4.2% and 5.7%.

Discussion

Our study focused on the epidemiology, molecular characterization, and population genetics of *Schistosoma* spp. that infect cattle in Mali. The study was conducted in abattoirs in two regions, the CBS and the SS in the District of Bamako and the KS. The results showed that the livestock schistosomiasis was endemic in the two regions with a global prevalence of 23.9%. The highest rate was observed in KS (44.6%) compared to CBS and SS (18.4%). The prevalence of *Sc* in our study was estimated at 9.8%. This prevalence is similar to that reported in Cameroon (10.03%) in domestic ruminants, using morphological egg criteria [30] but lower than that observed in Bamako (Mali) for cattle (30.8%) using an iso-enzymatic method [36]. Using a molecular approach, *Sc* was also reported in Barkedji and Linguère in Senegal, with an estimated prevalence of 73% in sheep, 84% in goats and 8% in cattle [21]. The low prevalence of *Sc* in our study may be due to the final host examined (cattle). As previously observed in Senegal [21], *Sc* preferred goats and sheep as hosts. The prevalence of *Sb* in our study was estimated at 12.6%. The prevalence of *Sb* varies greatly from one study to another according to the host species concerned. In sheep, prevalence rates ranging from 4% to 20% have been observed in Senegal [21, 27]. In goats, a prevalence of 15% has been observed in Senegal [21]. Prevalence rates are usually higher in cattle, with 19.5% in Cameroon [12] and 92% in Senegal [21]. The presence of schistosomes is dependent on the presence of intermediate snail hosts. Unfortunately, too few studies exist on the distribution of snails in Africa, particularly those that transmit schistosomes that infect animals. In West African countries, the snail species that hosts *Sb* may be *Bulinus truncatus*, *B. forskalii* or *B. globosus* [25] whereas *B. umbilicatus* is described as the main host of *Sc* [11]. In Mali, no study has yet looked specifically at snails as intermediate hosts of *Sc* or *Sb*, even though various intermediate hosts of schistosomiasis have been documented (*B. truncatus*, *B. globosus*, *B. umbilicatus*, *B. forskalii* and *B. senegalensis*) [9, 24]. The snail *B. umbilicatus* has only been identified in the Mopti region but the authors of the study were unable to determine the species of schistosome it transmitted [24]. Despite the specific distribution of snails due to ecological factors, animals slaughtered in Kayes or Bamako can be moved over great distances (600 km separate the livestock supply areas of Mopti from Bamako and those of Kayes and Nara on the Mauritanian border), which can bring them into contact with different transmission sites. No significant variation in prevalence was found with respect to season. The seasonal movement of animals over large distances to find new pastures and water points (transhumance) in Mali promotes their exposure to a variety of infected water bodies.

During this study, we did not find any *SbxSc* hybrid species of bovine schistosome, nor any hybrid between *ShxSb* or *ShxSc*, as recently highlighted in Cameroon [12]. However, *SbxSc* hybrids have been reported in cattle in Senegal, Niger and Mali [6, 36, 44] and sheep and goats, in Niger [7, 8]. The absence of hybrid parasites could be explained by the fact that the transmission sites we sampled were characterized by a high prevalence of either *Sb* or *Sc* or by a low rate of co-infected cattle. In the latter case, interspecific encounters are infrequent and hybridization unlikely. In addition, we collected adult parasites without identifying whether we had inter-specific pairs. Genotyping the eggs might have enabled us to identify first-generation hybrid eggs, as observed in Benin [38].

Regardless of the method used (Fst, PCA or Structure), there is strong structuring between *Sb* and *Sc* species, and weak structuring between slaughterhouses within the same species. Concerning the genetic difference between species, most published studies are based on sequence comparisons, and in particular, involve sequencing the cytochrome oxidase gene [18]. In the latter study, the authors measured a genetic difference of 6.1% between *Sb* and *Sc*, in Senegal. Based on microsatellite markers, we measured a genetic differentiation of 4.2% between these two species. Concerning the difference between populations within the same species, we measured an Fst of 0.43% between the CBS and SS abattoirs for *Sb*. These two abattoirs are about 18 Km apart. This result is in agreement with the weak genetic structuring of *Sb* observed in Cameroon [12] and Cote d'Ivoire [15]. With only two sites it is difficult to provide more general results on the genetic structure of this parasite species in Mali. As Mali is the leading livestock-producing country in the West African Economic and Monetary Union, it's normal for livestock to move in search of pasture across the country and even across borders, which encourages the spread of parasites. During this transhumance, it is often possible for herders to sell along the way [2]. Moreover, cattle trade between countries is universally known in West Africa. It is possible that some of the cattle sampled in this study came from countries bordering Mali. As Bamako is the country's capital, it receives cattle from all areas, including Burkina-Faso, Niger and even Senegal. All of these movements may be at the root of the weak structuring within the various species. The strong presence of *Sc* in the Kayes slaughterhouse is easily explained by its proximity to Mauritania and Senegal, where this species has already been reported [17,44]. Being able to use the same panel of microsatellite markers on *Sh*, *Sb* and *Sc* is an opportunity to carry out more studies on animal parasites and to infer their possible role in schistosome transmission to humans. For instance, our study shows that the heterozygosity rate is higher for *Sb* than for *Sc*. Variations in heterozygosity have already been observed in previous studies between *Sb* and *Sh* in Cameroon [12] and in Cote d'Ivoire [15].

Our work has a number of limitations. Because the worms were grouped by animal and by organ, we were unable to isolate the pairs directly in the abattoir and we were unable to obtain the genotypes/species of each pair. It is also incredibly difficult to collect parasites in abattoirs. The time allowed per animal is always very limited so as not to disrupt the slaughter line. In addition, we limited our study to cattle. It would have been interesting to inspect sheep and goats.

Animal schistosomiasis is prevalent in cattle in the Kayes and Bamako abattoirs, with the highest prevalence found in Kayes. Our study focused on the genetic profile and diversity of schistosome adults collected from livestock in Mali. Two pure species were found (*Sb* and *Sc*). Parasites were well structured according to species, with *Sb* and *Sc* dominating in Bamako and Kayes, respectively. Further studies on the genome of animal schistosomes on a large scale and the potential impacts of hybrid strains on animals are required. Additional studies are also needed to expand our understanding in this field by increasing the sample size and by extending

the surveys to other slaughterhouses and to the farm livestock in Mali. In view of our findings, a strategic control program for cattle trematodes is warranted. An important component of such a control program is to water cattle in troughs rather than to let them water freely from rivers and ponds, in order to reduce fecal contamination.

Abbreviations

<i>Sb</i>	<i>Schistosoma bovis</i>
<i>Sc</i>	<i>Schistosoma curassoni</i>
ITS	Internal transcribed spacer
Cox1	Cytochrome oxidase subunit 1 gene
RD-PCR	Rapid diagnostic multiplex PCR
T-ARMS-PCR	Tetra-amplification refractory mutation system PCR
FTA	Flinders Technology Associates
DNA	Deoxyribonucleic acid
UEMOA	Union Economique et Monétaire Ouest Africaine
ECOWAS	Economic Community of West African States
<i>PRODEVIM</i>	Programme de Développement à l'Exportation de la Viande du Mali
PCA	Principal Component Analysis
CBS	Central Bamako Slaughterhouse
SS	Sabalibougou Slaughterhouses
KS	Kayes Slaughterhouse
He	Expected heterozygosity
Ho	Observed heterozygosity
A	Allele
Ar	Allelic richness
Fis	Inbreeding coefficient
<i>B</i>	<i>Bulinus</i>

Acknowledgments

We thank Dr Bonnie Webster for her constructive comments on the manuscript. We thank slaughterhouse managers and butchers for their frank collaboration.

The authors would particularly like to thank: Amadou Dabo dit Boua for technical support in the field; Mathias Dolo and Kadra Koné, the geographers; all slaughterhouse managers and butchers of all three sites; the staff of the MRTC (Malaria Research and Training Center); the staff of the Faculty of Medicine and Dentistry and the Faculty of Pharmacy. We would like to thank the government of Mali for funding this research, and the IHPE laboratory, Univ. Montpellier, CNRS, Ifremer, Univ. Perpignan Via Domitia, Perpignan, France for molecular analyses.

Author contributions

P.A., A.D., B.J., N.S.D., D.L, S.T., I.M. participated in the design of the study, updated the research methodology and contributed to the writing of the final document. P.A., A.D., A.D., J.B., N.S.D., L. D., S.B., A.A., H.G., coordinated the field trial. P.A., B.A.E.S, provided laboratory handling and sequence correction. P.A., J. B., and A.D. carried out the statistical analysis. B.A.E.S, L.D., M. I., A.D., S.T., J.B. reviewed the manuscript for submission.

Funding

This research was funded by the Competitive Fund for Research and Technological Innovation (FCRIT).

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Declarations

Ethics approval and consent to participate

The global protocol covering both human and animal schistosomiasis study was reviewed and approved by the Institutional Ethics Committee of the Faculty of Medicine, Pharmacy and Dentistry of the University of Sciences, Techniques and Technologies of Bamako under the number N° 2018/71/CE/FMPOS. The officials of the veterinary services and the managers of the slaughterhouses were informed before any sampling. To avoid the disagreements caused by our presence inside the slaughterhouses and the loss of time caused by the examination of the animals, the butchers were willing to cooperate with us over the course of this study.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Epidemiology of Infectious Diseases, Faculty of Pharmacy, Malaria Research and Training Center (MRTC), University of Sciences, Techniques and Technologies of Bamako, Environnement, Santé, Sociétés (USTTB/ UCAD/UGB/CNRST/CNRS), BP 1805, IRL 3189 Bamako, Mali. ²Interactions Hôtes-Pathogènes-Environnements (IHPE), Univ. Montpellier, CNRS, Ifremer, Univ. Perpignan Via Domitia, Perpignan, France. ³Centre de Recherche pour la lutte contre les Maladies Infectieuses Tropicales (CRMIT/TIDRC), Université d'Abomey-Calavi, Abomey-Calavi, Bénin. ⁴Academie des Sciences du Mali, Baco-Djicoroni ACI Ouest, Rue 619 Porte, 104, Bamako, Mali.

References

1. Agniwo P, Sidibé B, Diakité AT, Niaré SD, Guindo H, Akplogan A, Ibikounlé M, Boissier J, Dabo A. 2023. Ultrasound aspects and risk factors associated with urogenital schistosomiasis among primary school children in Mali. *Infectious Diseases of Poverty*, 12(1), 40-51.
2. Bassett T. 2009. Mobile pastoralism on the brink of land privatization in Northern Côte d'Ivoire. *Geoforum*, 40(5), 756–766.
3. Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme M. 2004. GENETIX 4.05, logiciel sous Windows TM pour la génétique des populations. Laboratoire Génome, Populations, Interactions, CNRS UMR 5000, Université de Montpellier II, Montpellier (France). Available at <http://www.univmontp2.fr/~genetix/genetix/genetix.htm> (accessed on 2022-07-20).
4. Blin M, Dametto S, Agniwo P, Webster BL, Angora E, Dabo A, Boissier J. 2023. A duplex tetra-primer ARMS-PCR assay to discriminate three species of the *Schistosoma haematobium* group: *Schistosoma curassoni*, *S. bovis*, *S. haematobium* and their hybrids. *Parasites and Vectors*, 16, 121-131.
5. Boissier J, Mouahid G, Moné H. 2019. *Schistosoma* spp, in *Water and Sanitation for the 21st Century: Health and Microbiological Aspects of Excreta and Wastewater Management (Global Water Pathogen Project)*, Rose JB, Jiménez-Cisneros B, Editors; in Part 3: Specific Excreted Pathogens: Environmental and Epidemiology Aspects - Section 4: Helminths, Robertson L, Editor, Michigan State University, E. Lansing, MI, UNESCO. <https://doi.org/10.14321/waterpathogens.45>.
6. Bremond P, Sellin B, Sellin E, Nameoua B, Labbo R, Theron A. 1993. Arguments for the modification of the genome (introgression) of the human parasite *Schistosoma haematobium* by genes from *S. bovis*, in Niger. *Comptes Rendus de l'Académie des Sciences. Serie III, Sciences de la vie*, 316(7), 667–670.
7. Brémond P. 1990. Application des techniques électrophorétiques à deux aspects de la biologie des populations de schistosomes africains : Caractérisation des parasites et de leurs hôtes intermédiaires ; détection des schistosomes hybrides, in *Les Schistosomes*, in *Conférence International sur la Situation Epidémiologique les Stratégies de Lutte contre les Schistosomiasés en Afrique l'Ouest*. IRD, Bobo Dioulasso : OCCGE, p. 182-189.
8. Brémond P, Mouchet F, Chevallier P, Sellin E, Vera C, Sellin B. 1990. Flux génique entre *Schistosoma bovis* et *S. curassoni* au Niger. *Bulletin de la Société Française de Parasitologie*, 8, 708.
9. Coulibaly G, Madsen H. 1990. Seasonal density fluctuations of intermediate hosts of schistosomes in two streams in Bamako, Mali. *Journal of African Zoology*, 104(3), 201-212.
10. De Bont J, Vercruyse J. 1997. The Epidemiology and Control of Cattle Schistosomiasis. *Parasitology Today*, 13(7), 255-262
11. Diaw O, Vassiliades G. 1987. Epidémiologie des schistosomoses du bétail au Sénégal. *Revue d'Élevage et de Médecine Vétérinaire des Pays Tropicaux*, 40 (3), 265-274.
12. Djuikwo-Teukeng FF, Kouam Simo A, Allienne JF, Rey O, Njyou Ngapagna A, Tchuem-Tchuente LA, Boissier J. 2019. Population genetic structure of *Schistosoma bovis* in Cameroon. *Parasites and Vectors*, 12, 56–67.
13. Evanno G, Regnaut S, Goudet J. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology*, 14(8), 2611–2620.
14. Francis RM. 2017. Pophelper: an R package and web app to analyse and visualize population structure.

Molecular Ecology Resources, 17(1), 27–32.

15. Giovanoli Evack J, Kouadio JN, Achi LY, Bonfoh B, N’Goran EK, Zinsstag J, Jürg Utzinger J, Balmer O. 2024. Genetic characterization of schistosome species from cattle in Côte d’Ivoire. *Parasites and Vectors*, 17, 122–138.
16. Goudet J, Perrin N, Waser P. 2002. Tests for sex-biased dispersal using bi-parentally inherited genetic markers. *Molecular Ecology*, 11(6), 1103–1114.
17. Grétilat S. 1962. Une nouvelle zoonose, la “Bilharziose Ouest Africaine” *Schistosoma curassoni* Brumpt, 1931 commune à l’homme et aux ruminants domestiques. *Comptes Rendus Hebdomadaires des Séances de l’Académie des Sciences*, 255(15), 1805.
18. Huyse T, Webster BL, Geldof S, Stothard JR, Diaw OT, Rollinson D. 2009. Bidirectional Introgressive Hybridization between a Cattle and Human Schistosome Species. *PLoS Pathogens*, 5(9), e1000571.
19. King KC, Stelkens RB, Webster JP, Smith DF, Brockhurst MA. 2015. Hybridization in Parasites: Consequences for Adaptive Evolution, Pathogenesis, and Public Health in a Changing World. *PLoS Pathogens*, 11(9), e1005098.
20. Lafferty KD. 2009. The ecology of climate change and infectious diseases. *Ecology*, 90(4), 888–900.
21. Léger E, Borlase A, Fall C, Diouf N, Diop S, Yasenev L, Catalano S, Thiam CT, Ndiaye A, Emery A, Morrell A, Rabone M, Ndao M, Faye B, Rollinson D, Rudge WJ, Sène M, Webster JP. 2020. Prevalence and distribution of schistosomiasis in human, livestock, and snail populations in northern Senegal: a One Health epidemiological study of a multi-host system. *Lancet Planetary Health*, 4(8), e330-e342.
22. Léger E, Webster JP. 2016. Hybridizations within the Genus *Schistosoma*: Implications for evolution, epidemiology and control. *Parasitology*, 144(1), 65–80.
23. Lockyer AE, Olson PD, Ostergaard P, Rollinson D, Johnston DA, Attwood SW, Southgate VR, Horak P, Snyder SD, Le TH, Agatsuma T, McManus DP, Carmichael AC, Naem S, Littlewood DTJ. 2003. The phylogeny of the Schistosomatidae based on three genes with emphasis on the interrelationships of *Schistosoma* Weinland, 1858. *Parasitology*, 126(3), 203–224.
24. Madsen H, Coulibaly G, Furu P. 1987. Distribution of freshwater snails in the river Niger basin in Mali with special reference to the intermediate hosts of schistosomes. *Hydrobiologia*, 146, 77–88.
25. Moné H, Mouahid G, Morand S. 1999. The Distribution of *Schistosoma bovis* Sonsino, 1876 in Relation to Intermediate Host Mollusc–Parasite Relationships. *Advances in Parasitology*, 44, 99–138.
26. Mouchet F, Bremond P, Théron, A. 1989. Preliminary observations on *Schistosoma curassoni* Brumpt, 1931 in Niger. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 83(6), 811.
27. Ndao M, Belot J, Zinsstag J, Pfister K. 1995. Épidémiologie des helminthoses gastro-intestinales des petits Ruminants dans la zone sylvo-pastorale au Sénégal. *Veterinary Research*, 26(2), 132–139.
28. Ndifon GT, Betterton C, Rollinson D. 1988. *Schistosoma curassoni* Brumpt, 1931 and *S. bovis* (Sonsino , 1876) in cattle in northern Nigeria. *Journal of Helminthology*, 62(1), 33-34.
29. Nichols GL, Andersson Y, Lindgren E, Devaux I, Semenza JC. 2014. European monitoring systems and data for assessing environmental and climate impacts on human infectious diseases. *International Journal of Environmental Research and Public Health*, 11(4) 3894–3936.
30. Ousseini S. 1990. Schistosomoses des ruminants domestiques au Cameroun septentrionale. Thèse de Médecine vétérinaire de l’Université Cheikh Anta Diop, Dakar, Sénégal. p. 112.

31. Panzner U, Boissier J. 2021. Natural Intra- and Interclade Human Hybrid Schistosomes in Africa with Considerations on Prevention through Vaccination. *Microorganisms*, 9(7), 1465.
32. Pennance T, Allan F, Emery A, Rabone M, Cable J, Garba AD, Hamidou AA, Webster JP, Rollinson D, Webster BL. 2020. Interactions between *Schistosoma haematobium* group species and their *Bulinus* spp. intermediate hosts along the Niger River Valley. *Parasites and Vectors*, 13, 268–282.
33. Pritchard J, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus genotype data. *Genetics*, 155(2), 945–959.
34. PRODEVIM. 2022. Mali Meat Export Development Program (PRODEVIM). Available at [https://bamada.net/La viande : Quatre usines de viande dans les starting blocs - Bamada.net](https://bamada.net/La%20viande%20-%20Quatre%20usines%20de%20viande%20dans%20les%20starting%20blocs%20-%20Bamada.net) (accessed 2023-06-12).
35. Rey O, Webster BL, Huyse T, Rollinson D, Van den Broeck F, Kincaid-Smith J, Onyekwere A, Boissier J. 2021. Population genetics of African *Schistosoma* species. *Infection, Genetics and Evolution*, 89, 104727.
36. Rollinson D, Southgate VR, Vercruysse J, Moore PJ. 1990. Observations on natural and experimental interactions between *Schistosoma bovis* and *S. curassoni* from West Africa. *Acta Tropica*, 47(2), 101–114.
37. Rosenberg NA. 2004. Distruct: A program for the graphical display of population structure. *Molecular Ecology Notes*, 4(1), 137–138.
38. Savassi BAES, Mouahid G, Lasica C, Mahaman SK, Garcia A, Courtin D, Allienne JF, Ibikounlé M, Moné H. 2020. Cattle as natural host for *Schistosoma haematobium* (Bilharz , 1852) Weinland , 1858 x *Schistosoma bovis* Sonsino, 1876 interactions , with new cercarial emergence and genetic patterns. *Parasitology Research*, 119, 2189–2205.
39. Shuman EK. 2010. Global climate change and infectious diseases. *New England Journal of Medicine*, 362(12), 1061–1063.
40. Tanaka H, Tsuji M. 1997. From discovery to eradication of Schistosomiasis in Japan: 1847-1996. *International Journal for Parasitology*, 27(12), 1465–1480.
41. Vera C, Mouchet F, Bremond P, Sidiki A, Sellin E, Sellin B. 1992. Natural infection of *Bulinus senegalensis* by *Schistosoma haematobium* in a temporary pool focus in niger: Characterization by cercarial emergence patterns. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 86(1), 62.
42. Vercruysse J, Southgate VR, Rollinson D. 1984. *Schistosoma curassoni* brumpt, 1931 in sheep and goats in senegal. *Journal of Natural History*, 18(6), 969–976.
43. Webster B, Rollinson D, Stothard J, Huyse T. 2010. Rapid diagnostic multiplex PCR (RD-PCR) to discriminate *Schistosoma haematobium* and *S. bovis*. *Journal of Helminthology*, 84(1), 107–114.
44. Webster B, Diaw O, Seye M, Webster J, Rollinson D. 2013. Introgressive Hybridization of *Schistosoma haematobium* Group Species in Senegal: Species Barrier Break Down between Ruminant and Human Schistosomes. *PLoS Neglected Tropical Diseases*, 7(4), e2110.
45. Webster BL, Rabone M, Pennance T, Webster JP. 2015. Development of novel multiplex microsatellite polymerase chain reactions to enable high-throughput population genetic studies of *Schistosoma haematobium*. *Parasites and Vectors*, 8, 432-436.
46. Weir B, Cockerham CC. 1984. Estimating F-Statistics for the Analysis of Population Structure. *Evolution*, 38, 1358–1370.
47. Wright CA, Southgate VR. 1976. Hybridization of schistosomes and some of its implications, in *Genetic aspects of host-parasite interactions*, Taylor A, Muller R, Editors. Blackwell Scientific, Oxford. p.55-86.

Table 1: Number (%) of worms recovered in cattle in three transmission sites in Mali according to the season of collection and their localization (veins surrounding the small intestine or the caecum).

		Cattle examined	Cattle infected	P-value	Worms collected in veins surrounding small intestine	p-value	Worms collected in veins surrounding the caecum	p-value
Area								
<i>Bamako</i>	CBS	140	31 (22.1)	<0.0001	28 (20.0)	<i>p</i> <0.0001	3 (2.1)	<i>p</i> <0.0001
	SS	175	27 (15.4)		27 (15.4)		0 (0)	
Total		315	58 (18.4)		55 (17.5)		3 (0.9)	
<i>Kayes</i>	KS	83	37 (44.6)		36 (43.4)		17 (20.5)	
	Total	83	37 (44.6)		36 (43.4)		17 (20.5)	
Season								
	Rainy	268	69 (25.7)	<i>p</i> =0.12	65 (24.3)	<i>p</i> =0.2	20 (7.5)	<i>p</i> <0.0001
	Dry	130	26 (20.0)		26 (20.0)		0 (0)	
Total		398	95 (23.9)		91 (22.9)		20 (5.0)	

Table 2: Multivariate analyses of adult worm species according to the site and the organ.

Regions			Total	<i>Sb_SbxSb</i>	<i>Sc_ScxSc</i>	<i>p</i>
Bamako	CBS	Small Intestine	95	75 (78.9)	20 (21.1)	<0.0001
		Caecum	4	3 (75)	1 (25)	
		<i>Total</i>	99	78 (78.8)	21 (21.2)	
	SS	Small Intestine	99	97 (98)	2 (2)	<0.5
		Caecum	0	0	0	
		<i>Total</i>	99	97 (98.0)	2 (2.0)	
Kayes	KS	Small Intestine	196	10 (5.1)	186 (94.9)	<0.0001
		Caecum	82	3 (3.7)	79 (96.3)	
		<i>Total</i>	278	13 (4.7)	265 (95.3)	
	Total		476	188 (39.5)	288 (60.5)	

Table 3: Genetic diversity indices of schistosome worms collected in cattle in three Malian slaughterhouses. Expected (He) and observed (Ho) heterozygosity, total number of alleles (A), allelic richness (Ar), inbreeding coefficient (Fis) for each microsatellite locus and for each sampled site in Mali.

Slaughterhouse	Sh9	Sh3	C102	Sh1	Sh14	Sh6	C111	Sh7	Sh13	Sh4	Sh11	Sh15	Sh2	Sh5	Sh10	Sh12	Moy	Std
CBS (n=55)																		
He exp	0.84	0.89	0.89	0.90	0.65	0.76	0.71	0.91	0.92	0.89	0.62	0.40	0.95	0.72	0.83	0.64	0.78	0.12
He Obs	0.40	0.71	0.24	0.67	0.67	0.33	0.44	0.66	0.90	0.87	0.48	0.32	0.96	0.29	0.71	0.58	0.57	0.22
A	13	16	13	13	8	8	8	20	17	18	10	7	22	7	12	10	12.63	3.88
Ar	10.83	13.94	12.42	11.84	6.51	7.06	6.99	16.11	15.64	14.87	8.37	5.74	19.55	6.83	10.55	7.69	10.93	3.48
Fis	0.51	0.20	0.73	0.25	-0.03	0.56	0.38	0.28	0.02	0.01	0.22	0.19	-0.02	0.60	0.14	0.09	0.26	0.19
SS (n=42)																		
He exp	0.75	0.87	0.88	0.90	0.67	0.76	0.67	0.88	0.93	0.89	0.52	0.23	0.93	0.68	0.76	0.58	0.74	0.14
He Obs	0.46	0.82	0.30	0.89	0.59	0.40	0.54	0.54	0.91	0.88	0.50	0.21	0.94	0.31	0.83	0.63	0.60	0.24
A	9	17	14	12	5	7	7	12	17	15	4	3	16	6	8	3	9.69	4.40
Ar	8.02	15.25	14.00	11.73	4.99	6.80	6.49	11.49	16.49	14.09	3.91	2.69	15.67	5.83	7.87	3.00	9.27	4.23
Fis	0.38	0.06	0.69	0.01	0.12	0.47	0.19	0.39	0.02	0.01	0.05	0.08	-0.02	0.54	-0.11	-0.09	0.17	0.20
KS (n=69)																		
He exp	0.56	0.83	0.69	0.16	0.36	0.27	0.22	0.21	0.94	0.91	0.81	0.19	0.82	0.28	0.68	0.83	0.55	0.27
He Obs	0.10	0.60	0.03	0.06	0.3	0.07	0.2	0.08	0.9	0.9	0.5	0.13	0.8	0.09	0.72	0.90	0.53	0.30
A	6	19	11	7	4	6	5	5	21	15	18	5	21	5	8	10	10.38	5.34
Ar	4.69	14.87	10.33	4.42	3.41	5.20	4.01	4.45	18.14	12.83	14.15	4.17	16.08	4.59	7.33	8.79	8.59	4.38
Fis	0.79	0.29	0.96	0.62	0.21	0.73	0.06	0.65	0.06	-0.02	0.33	0.27	0.01	0.69	-0.04	-0.09	0.34	0.30
Total (n=166)																		
A	15	25	19	14	8	10	10	22	24	20	18	7	26	8	13	11	15.63	5.58
Ar	8.66	15.29	14.02	11.00	5.26	7.05	5.96	13.08	17.39	13.70	10.79	4.36	19.61	7.02	10.50	7.95	10.73	3.63

Table 4: Allelic richness and genetic diversity for *Sc_ScxSc* and *Sb_SbxSb* genotypes.

	Allelic richness (Ar)		Gene diversity (He)	
	<i>Sc_ScxSc</i>	<i>Sb_SbxSb</i>	<i>Sc_ScxSc</i>	<i>Sb_SbxSb</i>
Sh9	2.86	12.06	50%	81%
Sh3	16.71	16.8	79%	88%
C102	10	16.74	62%	88%
Sh1	1	13.09	-	90%
Sh14	2.85	6.5	25%	70%
Sh6	5.18	7.91	27%	76%
C111	4.42	7.59	21%	69%
Sh13	20.92	15.71	94%	92%
Sh4	14.28	15.39	90%	87%
Sh11	16.5	3.99	82%	49%
Sh15	5.78	4	25%	29%
Sh2	17.91	17.38	81%	93%
Sh5	4.76	6.4	28%	74%
Sh12	9.64	4.75	81%	57%
Mean	9.49	10.59	57%	75%
SE	1.76	1.39	8%	5%

Table 5: Pairwise estimates of F_{st} values for microsatellite DNA genotype (below the diagonal) of *Sb* and *Sc* parasite populations as a function of slaughterhouse.

<i>Sb_SbxSb</i>	CBS	SS
SS	<i>0.0043</i>	-
KS	<i>0.044</i>	<i>0.057</i>
<i>Sc_ScxSc</i>		
KS	<i>0.042</i>	



Figure 1: Map of Mali showing selected slaughterhouses.

Figure 2: Principal component analysis (PCA) of schistosome worms parasite recovered from three slaughterhouses in Mali. Each point represents a parasite, and the color label corresponds to the originated population (CSB, SS or KS). The first and the second axis explained 86.09% and 13.91% of the genetic variation, respectively.

Figure 3: Bar plot showing the population genetic structure using Structure software of 491 schistosome worms collected in three slaughterhouses in Mali. Each column represents one worm. The colors show the proportion of contribution of each cluster to each genotype. The cluster structure is for $K = 2$. The geographic populations are CBS (1), SS (2) and KS (3).

Figure 4: Principal component analysis (PCA) of *Schistosoma bovis* worms recovered from three abattoirs in Mali. Each point represents a parasite, and the color corresponds to the population of origin (CSB, SS or KS). The first and second axes explain 53.67% and 46.33 % of genetic variation respectively.

Figure 5: Principal component analysis (PCA) of *Schistosoma curassoni* worms recovered from three abattoirs in Mali. Each point represents a parasite, and the color corresponds to the population of origin (CSB, SS or KS). The first and second axes explain 100.00 % and 0.00 % of genetic variation respectively.





* Slaughterhouse

Watercourse

District boundary

Source : MRTC GIS
Auteur : Dolo M








Fig. 2. Frequency of the three categories (CBS, SS, KS) in the three numbered positions (1, 2, 3).

