



**HAL**  
open science

## Synthesis, DNA binding properties and biological evaluation of novel oligo-meta-benzamides related to netropsin

Francoise Debart, Gilles Gosselin, Bernard Rayner, P Le Ber, C Auclair, J Balzarini, E de Clercq, C Paoletti, Jean-Louis Imbach

► **To cite this version:**

Francoise Debart, Gilles Gosselin, Bernard Rayner, P Le Ber, C Auclair, et al.. Synthesis, DNA binding properties and biological evaluation of novel oligo-meta-benzamides related to netropsin. *European Journal of Medicinal Chemistry*, 1991, 26 (3), pp.261-271. 10.1016/0223-5234(91)90058-U . hal-04651228

**HAL Id: hal-04651228**

**<https://hal.science/hal-04651228v1>**

Submitted on 17 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Synthesis, DNA binding properties and biological evaluation of novel oligo-*meta*-benzamides related to netropsin

F Debart<sup>1</sup>, G Gosselin<sup>1</sup>, B Rayner<sup>1</sup>, P Le Ber<sup>2</sup>, C Auclair<sup>2</sup>, J Balzarini<sup>3</sup>,  
E De Clercq<sup>3</sup>, C Paoletti<sup>2</sup>, JL Imbach<sup>1\*</sup>

<sup>1</sup>Laboratoire de Chimie Bio-Organique, case courrier 008, URA 488 et UMR 112 du CNRS,  
Université de Montpellier II, Sciences et Techniques du Languedoc, Place Eugène-Bataillon, 34095 Montpellier Cedex 5;

<sup>2</sup>Institut Gustave-Roussy, Laboratoire de Biochimie-Enzymologie, U 140 de l'INSERM, LA 147 du CNRS,  
rue Camille-Desmoulins, 94805 Villejuif Cedex, France

<sup>3</sup>Rega Institute for Medical Research, Katholieke Universiteit Leuven, Minderbroedersstraat 10, B-3000 Leuven, Belgium

**Summary** — A series of oligo-*meta*-benzamides structurally related to the natural agent netropsin have been synthesized. Their binding constants to double-stranded polynucleotides as well as their cytostatic activity against tumor cell lines and their *in vitro* activity against a wide variety of DNA and RNA viruses have been determined. Most of these analogues retain the DNA binding capacity of the parent compound but with a notable decrease of selectivity and affinity. Like netropsin, the evaluated analogues did not show significant cytostatic and antiviral activity.

*meta*-benzamide analogues / netropsin / DNA binding properties / biological activities

## Introduction

Rational design of sequence-specific DNA binding molecule is of great current importance not only from a fundamental biological point of view but also in a therapeutic approach as antiviral and anticancer agents [2]. In this regard, besides natural and modified synthetic antisense oligonucleotides [3], the structurally related antibiotics netropsin and distamycin (fig 1) have received much attention to date. These polypyrrolocarboxamides exert their biological activities by blocking the template function of DNA by strong non-intercalative binding to (A•T)<sub>n</sub> sequences in the minor groove of double-stranded B-DNA [4]. In spite of their reported antiviral and anticancer properties, these 2 molecules are too toxic for clinical use. In order to decrease the toxicity and increase the potency





Fig 1. Structure of netropsin and distamycin.


and selectivity of these agents, several attempts of structural modification have been reported. Examples of these modifications include variation of the number of pyrrole units and their replacement by other rings, changes in the mode of pyrrole linkages and substitution of the side chains (surveyed in [5, 6]) [7–15]. While a significant increase of potency compared with the parent natural compounds was achieved for only a few derivatives, it is noteworthy that the replacement of the pyrrole ring(s) by hydrogen bond accepting heterocycles such as imidazole resulted in an alteration of DNA base recognition from A•T to G•C [16–24].

As a part of our ongoing work in attempting to understand the structural requirements of oligopeptides for biological activities and DNA-binding properties we have previously described some analogues of netropsin showing substitution of the aminopropionamide side chain by either a polyamine or by mono-, di- or tri-L-alanyl residues [6]. In the

present study we report the synthesis, characterization and biological evaluation of new netropsin analogues which bear both 1,3-disubstituted benzene moieties in place of *N*-methylpyrrole and various side chains including aminopropionamide, aminoethylimidazole and spermidine functions. The rationale for these structural changes results from an attempt to assess the influence of both the backbone curvature of these molecules and the nature of the side chains on their DNA-binding selectivity and biological activities. Thus model building studies have shown that benzene rings in place of *N*-methylpyrroles increase the curvature of the molecule as compared to netropsin and distamycin. Such a modification, as well as the change in the natural side chains, might lead to different van der Waals and hydrogen-bond interactions, and consequently, altered DNA affinity and selectivity. There have been previous reports in the literature of synthesis of bis- and tris(*meta*-benzamido)netropsin or distamycin analogues [7, 8, 10, 25, 26]. However,



**Scheme 1.** Reaction conditions: (a) SOCl<sub>2</sub>, Δ; (b) H<sub>2</sub>N-(CH<sub>2</sub>)<sub>2</sub>-CN/pyridine; (c) HCl in CHCl<sub>3</sub>-C<sub>2</sub>H<sub>5</sub>OH, then dry NH<sub>3</sub> in C<sub>2</sub>H<sub>5</sub>OH; (d) H<sub>2</sub>, Pd/C in CH<sub>3</sub>OH-DMF; (e) 2/pyridine; (f) H<sub>2</sub>, Pd/C in C<sub>2</sub>H<sub>5</sub>OH; (g) HCl·H<sub>2</sub>NC(=NH)NHCH<sub>2</sub>CO<sub>2</sub>H, DCC in DMF.



**Scheme 2.** Reaction conditions: (a)  $\text{CH}_3\text{C}(=\text{O})\text{OC}(=\text{O})\text{H}$ ; (b)  $\text{H}_2\text{N}(\text{CH}_2)_2\text{NH}_2/\text{C}_2\text{H}_5\text{OH}$ ; (c)  $\text{HCl}\cdot\text{H}_2\text{NC}(=\text{NH})\text{NHCH}_2\text{CO}_2\text{H}$ , DCC in DMF.

so far no modification has been introduced in their side chains and only a few data on their physico-chemical and biological properties have been reported.

## Results and Discussion

### Chemistry

The preparation of the 1,3-disubstituted benzene netropsin homologue **9** was carried out from the commercial 3-nitrobenzene-1-carboxylic acid **1** essentially according to the strategy originally reported [25]; the main features are outlined in scheme 1.

The synthesis of derivatives **10** and **12** carrying a formyl or an ethylimidazoline function in place of the

natural guanidinoacetyl of propionamide side chains respectively was accomplished from the intermediate **8**, previously obtained during the preparation of **9**. On the one hand, the exocyclic amine of **8** was condensed with formic acetic anhydride [27] to afford the already known **10** [7] in good yield. On the other hand, the amidine function of **8** was first transformed with ethylenediamine [28] to an imidazoline function, after which the amine of the resulting **11** was reacted with guanidinoacetic acid hydrochloride [29] in the presence of dicyclohexylcarbodiimide (DCC) to yield the hitherto unknown analogue **12** (scheme 2).

Concerning the introduction of spermidine as a side chain, 2 procedures were applied for amide bond formation, namely either a direct reaction of an amine with an acid chloride in the presence of base or a one-step active ester procedure implementing benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP) reagent [30, 31].

Thus condensation of  $N^4,N^8$ -bis(*tert*-butoxycarbonyl)spermidine [32] with the acid chloride **2** in a pyridine-acetonitrile mixture gave **13**. The latter was reduced by catalytic hydrogenation to give the amine **14** which was acylated with **2** to afford **15**. This nitro compound **15** was reduced catalytically to give the amine **16** which was allowed to react with guanidinoacetic acid hydrochloride in the presence of DCC to give **17**. From the latter compound, removal of the *tert*-butoxycarbonyl protecting groups on treatment with trifluoroacetic acid proceeded satisfactorily to yield the target compound **18** (scheme 3).


Furthermore, the amine **16** was also transformed into its succinamido derivative **19**, which was subsequently condensed with  $N^4,N^8$ -bis(*tert*-butoxycarbonyl)spermidine using BOP as coupling reagent to yield the desired fully protected compound **20**. Finally, removal of the *tert*-butoxycarbonyl groups from **20** on treatment with trifluoroacetic acid yielded the last target analogue **21** (scheme 4).

It is noteworthy that the low yields observed during the synthesis of **9**, **12** and **17** were caused by a known side reaction [33] between the carbodiimide and aminobenzene moieties as exemplified by the isolation of  $N^1$ -([3-[3-( $N,N'$ -dicyclohexylguanilyl)-benzene-1-carboxamido]-benzen-1-yl]carbonyl)- $N^4,N^8$ -bis(*tert*-butoxycarbonyl)spermidine during the purification of the crude reaction mixture resulting from the coupling of the amine **16** with guanidinoacetic acid in the presence of DCC.


### Binding of compounds to polynucleotides

The binding constants of netropsin analogues **4**, **7–10**, **12**, **18** and **21** to poly(dA-dT)•poly(dA-dT), and poly(dG-dC)•poly(dG-dC) are presented in table I.

Compound **9** which differs from netropsin only by the substitution of the original *N*-methylpyrrole rings



**Scheme 3.** Reaction conditions: (a) H<sub>2</sub>N(CH<sub>2</sub>)<sub>3</sub>N(Boc)(CH<sub>2</sub>)<sub>4</sub>NHBoc/pyridine-CH<sub>3</sub>CN; (b) H<sub>2</sub>, Pd/C in CH<sub>3</sub>OH; (c) 2/pyridine-CH<sub>3</sub>CN; (d) HCl·H<sub>2</sub>N-C(=NH)-NHCH<sub>2</sub>CO<sub>2</sub>H, DCC in DMF; (e) CF<sub>3</sub>CO<sub>2</sub>H.



**Scheme 4.** Reaction conditions: (a) Succinic anhydride, DMAP,  $(\text{C}_2\text{H}_5)_3\text{N}/\text{CH}_2\text{Cl}_2$ ; (b)  $\text{H}_2\text{N}(\text{CH}_2)_3 \text{N}(\text{Boc})(\text{CH}_2)_4 \text{NH Boc}$ , BOP/ $\text{CH}_3\text{CN}$ ; (c)  $\text{CF}_3\text{CO}_2\text{H}$ .

by benzene moieties, displays a much lesser affinity for poly(dA-dT)•poly(dA-dT) than the parent molecule (decrease of  $K_{\text{app}}$ : more than 2 orders of magnitude). Since this compound is able to better bind to poly(dG-dC)•poly(dG-dC) than netropsin (increase of  $K_{\text{app}}$ :  $\approx 1$  order of magnitude), the result is a loss of sequence selectivity recognition for **9**. From this analogue **9**, change of the left guanidinoacetamido side chain by a nitro function to give **7** lowers the affinity constant for double-stranded polynucleotides, while its change by an amino or a formylamino func-

tion (compounds **8** and **10**) leads to the disappearance of significant binding capacity. On the other hand, cyclising the propionamidino right side chain of **9** into an ethylimidazoline function to give **12** does not appreciably modify the affinity for polynucleotides. Finally, it can be noted that the spermidine derivatives **18** and **21** keep the ability of the parent compound, netropsin, to bind to poly(dA-dT)•poly(dA-dT) albeit with reduced affinity. In addition, these spermidine derivatives also bind better to poly(dG-dC)•poly(dG-dC) than netropsin.

**Table I.** Association constants ( $K_{app}$ ) of compounds with polynucleotides.

Compd	$K_{app}, M^{-1}$		Ratio $K_{app}$ $poly(dA-dT)/$ $poly(dG-dC)$
	$poly(dA-dT)$	$poly(dG-dC)$	
4	$6.3 \times 10^3$	$1.2 \times 10^3$	5.25
7	$1.8 \times 10^4$	$3.4 \times 10^3$	5.29
8	$< 10^2$	$< 10^2$	
9	$3.5 \times 10^5$	$2.3 \times 10^5$	1.52
10	$< 10^2$	$< 10^2$	
12	$4.7 \times 10^5$	$2.1 \times 10^5$	2.24
18	$4.1 \times 10^6$	$6.7 \times 10^4$	61.19
21	$8.2 \times 10^6$	$5.4 \times 10^5$	15.19
Netropsin	$7.3 \times 10^8$	$3.0 \times 10^4$	23548
Distamycin	$8.5 \times 10^7$	$2.5 \times 10^4$	3400

**Table II.** Inhibitory effects of compounds on the proliferation of murine leukemia (L1210), human B-lymphoblast (Raji) and human T-lymphoblast (Molt/4F) cells.

Compd	$CD_{50}^a$ ( $\mu\text{g/ml}$ )		
	L1210	Raji	Molt/4F
4	$> 200$	$171 \pm 13$	$> 200$
7	$> 200$	$> 200$	$> 200$
8	$> 200$	$\geq 200$	$> 200$
9	$> 200$	$145 \pm 30$	$> 200$
10	$> 200$	$> 200$	$> 200$
12	$> 200$	$> 200$	$> 200$
18	$> 200$	$153 \pm 42$	$\geq 200$
21	$> 200$	$> 200$	$> 200$
Netropsin	$\geq 100$	$\geq 100$	$> 200$

<sup>a</sup>50% cytotoxic dose or dose required to inhibit cell proliferation by 50%.

## Biological evaluation

### Cytostatic activity

Among the 8 netropsin analogues tested for their inhibitory effects on the proliferation of exponentially growing murine leukemia L1210 and human B-lymphoblast Raji and T-lymphoblast Molt/4F cells, compounds **4**, **9** and **18** proved to be slightly cyto-

static to Raji tumor cells (table II). None of the other compounds showed any cytostatic activity at the highest concentration tested (200  $\mu\text{g/ml}$ ).

### Antiviral activity

The netropsin analogues were evaluated for their *in vitro* inhibitory effects on the replication of a number of DNA viruses (*ie* herpes simplex virus type 1 (KOS) and type 2 (G), vaccinia virus) and RNA viruses (vesicular stomatitis virus, Coxsackie virus B4, polio virus-1, parainfluenza virus-3, reovirus-1, Sindbis virus and Semliki forest virus) in primary rabbit kidney, HeLa and Vero B cell cultures (table III; for comparative purposes, the antiviral data of the well-known antiviral compounds tubercidin, (S)-DHPA, ribavirin, and carbocyclic 3-deazaadenosine are included). With a few exceptions, none of the analogues showed a marked antiviral effect at doses up to 200 or 400  $\mu\text{g/ml}$ . Compounds **7** and **18** were active at 100 and 300  $\mu\text{g/ml}$ , respectively, against vesicular stomatitis virus in HeLa cells; they were also effective at 150  $\mu\text{g/ml}$  against reovirus-1 in Vero B cells. None of the compounds showed any detectable alteration of host cell morphology at the highest concentration tested (400  $\mu\text{g/ml}$ ).

When evaluated in an anti-human immunodeficiency virus assay, none of the analogues proved effective against HIV-induced cytopathogenicity in human MT-4 cells at nontoxic concentrations (table IV).

## Conclusion

According to the present results, it is obvious that the change of the *N*-methylpyrrole rings by benzene moieties results in a loss of DNA-binding specificity, perhaps owing to the increase in the curvature of the molecules. However, the A•T specificity can be in part recovered when the original side chains are substituted by a spermidine function. While the evaluated analogues showed no significant biological activity, it is noteworthy that, in contrast to previously published data [34–40], netropsin itself was also inactive in our assays. These findings provide relevant information in the design of future sequence-specific agents structurally related to netropsin.

## Experimental protocols

### Chemical synthesis

General procedures and instrumentation used have been described in [6]. High-performance liquid chromatographic (HPLC) analyses were also carried out on a third Waters Associate Unit (apparatus C) equipped with 2 Model 510 EF

**Table III.** Antiviral activity of compounds against different viruses in different cell systems.

Compd	Minimum cytotoxic conc <sup>b</sup> µg/ml			Minimum inhibitory concentration, <sup>a</sup> µg/ml												
				Primary rabbit kidney (PRK) cells				HeLa cells			African green monkey kidney (Vero B) cells					
	PRK cells	HeLa cells	Vero B cells	Herpes simplex virus-1 (KOS)	Herpes simplex virus-2 (G)	Vaccinia virus	Vesicular stomatitis virus	Polio virus-1	Coxsackie virus B4	Sindbis virus	Parainfluenza virus-3	Reovirus	Semliki forest virus			
4	400	400	≥ 200	> 200	> 200	> 100	> 200	> 200	> 200	> 200	> 200	> 200	> 200	> 100	> 100	> 200
7	400	400	400	> 200	> 200	> 200	100	> 200	> 200	> 200	> 200	> 200	> 200	> 200	150	> 200
8	≥ 400	≥ 400	400	> 400	> 400	> 400	> 200	> 400	> 400	> 400	> 400	> 200	> 200	> 200	> 200	> 200
9	> 400	≥ 400	400	> 400	> 400	> 400	> 400	> 400	> 400	> 400	> 400	> 200	> 200	> 200	> 200	> 200
10	> 400	≥ 400	400	> 400	> 400	> 400	> 400	> 400	> 400	> 400	> 400	> 200	> 200	> 200	> 200	> 200
12	> 400	≥ 400	400	> 400	> 400	> 400	> 400	> 400	> 400	> 400	> 400	> 200	> 200	> 200	> 200	> 200
18	> 400	≥ 400	400	> 400	> 400	> 200	300	> 400	> 200	> 400	> 200	> 200	> 200	> 200	150	> 200
21	> 400	> 400	≥ 400	> 400	> 400	> 400	> 400	> 400	> 400	> 400	> 400	> 200	> 200	> 200	> 200	> 200
Netropsin	≥ 40	≥ 100	200	> 40	> 40	> 10	> 40	> 100	> 100	> 100	> 100	> 100	> 100	> 100	> 100	> 100
Tubercidin	≥ 0.4	≥ 1	0.4	> 0.1	> 0.1	> 0.1	0.2	0.04	0.2	0.07	> 0.1	> 0.1	> 0.1	> 0.1	> 0.1	> 0.1
(S)-DHPA	> 400	> 400	≥ 400	> 400	> 400	70	20	150	> 400	> 400	> 400	> 400	> 400	100	40	> 400
Ribavirin	> 400	≥ 200	> 400	> 400	> 400	20	300	20	150	70	> 400	70	150	200	70	
Carbocyclic 3-deazaadenosine	> 400	> 400	≥ 400	> 400	> 400	2	2	2	> 400	> 400	70	400	2	0.7	> 400	

<sup>a</sup>Required to reduce virus-induced cytopathogenicity by 50%. Viral cytopathogenicity was recorded as soon as it reached completion in the control virus-infected cell cultures. The multiplicity of infection (MOI) was invariably 100 x CCID<sub>50</sub>, ie 100 times the virus dose needed to infect 50% of the cells. <sup>b</sup>Required to cause a microscopically detectable alteration of normal cell morphology, when incubated with cells for the same duration as required to measure antiviral activity.

solvent delivery systems, a Model 680 automated gradient controller, a Model U6K sample injector, a Waters 990 photodiode array detector, and a NEC-APC IV microprocessor-controlled data system (standard program Waters 990+).

Compounds 2–9 were synthesized with only minor modifications following similar conditions as those used by Julia and Gombert during the original synthesis of 9 [25]. The strategy of their preparation is shown in scheme 1 and their physical properties are presented below.

### 3-(3-Nitrobenzene-1-carboxamido)propionitrile 3

Mp = 110–111°C (lit mp = 104–105°C [25]); <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 2.85 (t, 2H, CH<sub>2</sub>-CN), 3.6 (m, 2H, NH-CH<sub>2</sub>), 7.6–8.8 (m, 4H, C<sub>6</sub>H<sub>4</sub>), 9.25 (t, 1H, NH); MS (matrix, 3-nitrobenzyl alcohol) FAB > O: 220 (M+H)<sup>+</sup>.

### 3-(3-Nitrobenzene-1-carboxamido)-propionamide hydrochloride 4

Mp = 210–213°C (lit mp = 210–212°C [25]); <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 2.72 (t, 2H, NH-CH<sub>2</sub>-CN<sub>2</sub>), 3.6 (m, 2H, NH-

**Table IV.** Anti-HIV-1 and cytostatic activities of compounds in human T-lymphocyte (MT-4) cells.

Compd	CD <sub>50</sub> <sup>a</sup> (µg/ml)	ED <sub>50</sub> <sup>b</sup> (µg/ml)
4	97.6 ± 13.3	> 40
7	> 200	> 200
8	> 200	> 200
9	> 200	> 200
10	> 200	> 200
12	> 200	> 200
18	> 200	> 200
21	> 200	> 200
Netropsin	63.9 ± 17.9	> 40

<sup>a</sup>50% cytotoxic dose; <sup>b</sup>50% effective dose



CH<sub>2</sub>-CH<sub>2</sub>), 7.8–8.7 (m, 4H, C<sub>6</sub>H<sub>4</sub>), 8.5–9.1 [br s, 4H, (C=NH<sub>2</sub><sup>+</sup>)-NH<sub>2</sub>], 9.24 (t, 1H, NH); MS (matrix, glycerol) FAB > O: 237 [M+H-Cl]<sup>+</sup>.

### 3-[3-(3-Aminobenzene-1-carboxamido)propionitrile 5

Mp = 118–119°C (lit mp = 123–124°C [25]); <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 2.78 (t, 2H, CH<sub>2</sub>CN), 3.5 (m, 2H, NH-CH<sub>2</sub>), 5.3 (br s, 2H, NH<sub>2</sub>), 6.6–7.3 (m, 4H, C<sub>6</sub>H<sub>4</sub>), 8.6 (t, 1H, NH); MS (matrix, 3-nitrobenzyl alcohol) FAB > O: 190 [M+H]<sup>+</sup>.

### 3-[3-(3-Nitrobenzene-1-carboxamido)-benzene-1-carboxamido]propionitrile 6

Mp = 187–188°C (lit mp = 180°C [25]); <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 2.85 (t, 2H, CH<sub>2</sub>-CN), 3.6 (m, 2H, NH-CH<sub>2</sub>), 7.5–8.8 (m, 8H, 2 C<sub>6</sub>H<sub>4</sub>), 8.9 (t, 1H, NH-CH<sub>2</sub>), 10.8 (br s, 1H, NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix, 3-nitrobenzyl alcohol) FAB > O: 339 [M+H]<sup>+</sup>.

### 3-[3-(3-Nitrobenzene-1-carboxamido)-benzene-1-carboxamido]propionamide hydrochloride 7

Mp = 148–150°C (lit mp = 142–145°C [25]); <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 2.71 (t, 2H, CH<sub>2</sub>-CH<sub>2</sub>), 3.6 (m, 2H, NH-CH<sub>2</sub>), 7.4–8.5 and 8.83 (m and s, 7 and 1H respectively, 2 C<sub>6</sub>H<sub>4</sub>), 8.78 (t, 1H, NH-CH<sub>2</sub>), 8.7–9.2 [br s, 4H, (C=NH<sub>2</sub><sup>+</sup>)NH<sub>2</sub>], 10.8 (br s, 1H, NH, C<sub>6</sub>H<sub>4</sub>); MS (matrix, glycerol) FAB > O: 356 [M+H-Cl]<sup>+</sup>; HPLC retention time 14.9 min, spectrophotometric purity at 254 nm: > 97% [UV, λ<sub>max</sub> 224 and 256 (sh) nm] (Apparatus B; column C<sub>18</sub> 'Radial Pak' (100 x 8 mm id, 10 μm particle size) in a Waters Associates Radial compression module RCM 100, protected by a prefilter and a precolumn C<sub>18</sub> 'Guard Pak'; solvent A, 100% 0.1 M ammonium acetate buffer, pH 5.9; solvent B, 50% of acetonitrile in the same buffer; linear gradient 0–50% B in 10 min, then 50–100% B in 5 min; flow rate 3.0 ml/min).

### 3-[3-(3-Aminobenzene-1-carboxamido)-benzene-1-carboxamido]propionamide hydrochloride 8

<sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 2.71 (t, 2H, NH-CH<sub>2</sub>-CH<sub>2</sub>), 3.6 (m, 2H, NH-CH<sub>2</sub>), 5.3 (br s, 2H, NH<sub>2</sub>), 6.7–8.3 (m, 8H, 2 C<sub>6</sub>H<sub>4</sub>), 8.7 [m, 3H, NH-CH<sub>2</sub> and (C=NH<sub>2</sub><sup>+</sup>)NH<sub>2</sub>], 9.1 [br s, 2H, (C=NH<sub>2</sub><sup>+</sup>)NH<sub>2</sub>], 10.2 (s, 1H, NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix, 3-nitrobenzyl alcohol) FAB > O: 326 [M+H-Cl]<sup>+</sup>; HPLC retention time 11.4 min, spectrophotometric purity at 254 nm: > 96% (UV, λ<sub>max</sub> 227 nm) (same apparatus, column, solvents and conditions as described for the characterization of 7).

### 3-[3-[3-(Guanidinoacetamido)benzene-1-carboxamido]-benzene-1-carboxamido]propionamide bis(acetate) 9

This compound, first obtained as a sulfate salt [25] with a 42% yield, was then transformed in its acetate salt to improve its aqueous solubility. <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 1.65 (s, 6H, 2 CH<sub>3</sub>CO<sub>2</sub>), 2.62 (t, 2H, NH-CH<sub>2</sub>-CH<sub>2</sub>), 3.6 (m, 2H, NH-CH<sub>2</sub>-CH<sub>2</sub>), 4.08 (s, 2H, NH-CH<sub>2</sub>-CO), 7.4–8.3 (m, 8H, 2 C<sub>6</sub>H<sub>4</sub>), 8.4–10.5 [br s, 12H, 2(C=NH<sub>2</sub><sup>+</sup>)NH<sub>2</sub> and 4NH]; MS (matrix, glycerol) FAB > O: 425 [M+H-2CH<sub>3</sub>CO<sub>2</sub>H]<sup>+</sup>; HPLC retention time 11.3 min, spectrophotometric purity at 254 nm: > 95% (UV, λ<sub>max</sub> 248 nm) (Apparatus B; column, Ultrasphere XLODS cartridge Beckman, 70 x 4.6 mm id, 3 μm particle size, protected by prefilters and a XLODS precolumn (5 x 4.6 mm id, 3 μm); solvent A, 100% ammonium acetate buffered to pH 2.5 by formic acid; solvent B, 50% acetonitrile in the same buffer; linear gradient 0–50% B in 10 min, then 50–100% B in 5 min; flow rate 1 ml/min).

### 3-[3-[3-(Formylamino)benzene-1-carboxamido]-benzene-1-carboxamido]propionamide hemi(sulfate) 10

Acetic formic anhydride (0.203 ml) [27] was added to compound **8** (0.253 g, 0.7 mmol). The resulting solution was first stirred at room temperature for 1 h, then diluted with methanol (4 ml) and finally poured into diethyl ether. The obtained precipitate was collected, dissolved in water (1.5 ml) at 50°C and to the solution was added sodium sulfate (0.1 g, 0.7 mmol). By cooling to room temperature, the sulfate salt of **10** crystallized (0.228 g, 81% yield); mp: 187–189°C; <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 2.64 (t, 2H, NH-CH<sub>2</sub>-CH<sub>2</sub>), 3.6 (m, 2H, NH-CH<sub>2</sub>), 7.3–8.3 (m, 8H, 2 C<sub>6</sub>H<sub>4</sub>), 8.27 [s, 1H, H-C(=O)], 8.5–9.3 [m, 7H, (C=NH<sub>2</sub><sup>+</sup>)NH<sub>2</sub> and 3 NH]; MS (matrix, glycerol) FAB > O: 354 [M+H-1/2H<sub>2</sub>SO<sub>4</sub>]<sup>+</sup>; HPLC retention time 17.9 min, spectrophotometric purity at 254 nm: 100% (UV, λ<sub>max</sub> 229 nm) (same apparatus, column and solvents as described for the characterization of 7; linear gradient 0–50% B in 20 min, flow rate 3.0 ml/min).

### 2-[2-[3-(3-Aminobenzene-1-carboxamido)-benzene-1-carboxamido]ethyl]imidazoline bis(hydrochloride) 11

A solution of compound **8** (0.253 g, 0.7 mmol) and anhydrous ethylenediamine (0.051 ml, 0.77 mmol) in absolute ethanol (1 ml) was refluxed for 1.5 h [28] and then cooled to room temperature. The solution was evaporated to dryness, the residue was dissolved in ethanol, and 12 N HCl (0.05 ml) was added. The resulting crystalline solid was collected by filtration to give **11** as its bis-hydrochloride salt (0.248 g, 83% yield); mp: 178–180°C; <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 2.78 (t, 2H, NH-CH<sub>2</sub>-CH<sub>2</sub>), 3.6 (m, 2H, NH-CH<sub>2</sub>-CH<sub>2</sub>), 3.8 (s, 4H, N-CH<sub>2</sub>-CH<sub>2</sub>-N), 7.4–8.3 (m, 8H, 2 C<sub>6</sub>H<sub>4</sub>), 8.5 [br s, 2H, C(=N)NH<sub>2</sub><sup>+</sup>-CH<sub>2</sub>], 8.9 (t, 1H, NH-CH<sub>2</sub>), 10.3 (br s, 3H, NH<sub>2</sub><sup>+</sup>), 10.6 (s, 1H, NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix, glycerol) FAB > O: 352 [M+H-2HCl]<sup>+</sup>. This compound was sufficiently pure to be used without further purification for the preparation of **12**.

### 2-[2-[3-(3-(Guanidinoacetamido)benzene-1-carboxamido)-benzene-1-carboxamido]ethyl]imidazoline sulfate 12

To a solution of compound **11** (0.12 g, 0.28 mmol) and guanidinoacetic acid hydrochloride [29] (0.043 g, 0.28 mmol) in anhydrous DMF was added dicyclohexylcarbodiimide (DCC; 0.116 g, 0.56 mmol). After the mixture was stirred for 7 h at room temperature, the formed precipitate of dicyclohexylurea was removed by filtration, and the filtrate was evaporated. The resulting residue was dissolved in water (3 ml) at 60°C, and to the solution was added sodium sulfate (0.06 g, 0.42 mmol). By cooling to room temperature, the sulfate salt of **12** crystallized (0.041 g, 27 % yield); mp: 204–205°C; <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 2.8 (m, 2H, NH-CH<sub>2</sub>-CH<sub>2</sub>), 3.6 (m, 2H, NH-CH<sub>2</sub>-CH<sub>2</sub>), 3.7 (s, 4H, N-CH<sub>2</sub>-CH<sub>2</sub>-N), 4.1 (br s, 2H, NH-CH<sub>2</sub>-CO), 7.4–8.6 [m, 12H, 2 C<sub>6</sub>H<sub>4</sub> and (C=NH<sub>2</sub><sup>+</sup>)NH<sub>2</sub>], 8.9 [m, 1H, NH-CH<sub>2</sub>], 9.0–10.6 [very br s, 3H, NH<sub>2</sub>-C-NH and C(=N)NH<sub>2</sub><sup>+</sup>-CH<sub>2</sub>], 10.8 (br s, 2H, 2NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix, glycerol) FAB > O: 451 [M+H-H<sub>2</sub>SO<sub>4</sub>]<sup>+</sup>; HPLC retention time 9.7 min, spectrophotometric purity at 254 nm: > 97% (UV, λ<sub>max</sub> 224 nm) (same apparatus, column and solvents as described for the characterization of 7; linear gradient 0–100% B in 10 min; flow rate 3.0 ml/min).

### N<sup>1</sup>-[(3-Nitrobenzen-1-yl)carbonyl]-N<sup>4</sup>,N<sup>8</sup>-bis(tert-butoxycarbonyl)spermidine 13

To a solution of N<sup>4</sup>,N<sup>8</sup>-bis(tert)butoxycarbonyl)spermidine [32] (2.1 g, 6.15 mmol) in anhydrous pyridine (3 ml) was dropwise

added at 0°C a solution of the acyl chloride **2** (1.14 g, 6.15 mmol) in anhydrous acetonitrile (3 ml), and the mixture was stirred at ambient temperature for 3 h. After dilution with CH<sub>2</sub>Cl<sub>2</sub>, the mixture was poured into a saturated NaHCO<sub>3</sub> aqueous solution. The organic phase was separated, twice washed with water, dried over sodium sulfate, and evaporated. Chromatography of the residue on a silica gel column using as eluent a stepwise gradient of acetone (0–40%) in CH<sub>2</sub>Cl<sub>2</sub> led, after evaporation of the appropriate fractions, to the isolation of pure **13** (1.67 g, 55% yield); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.3–2.3 (m, 6H, 3CH<sub>2</sub>), 1.45 and 1.5 [2s, 2 × 9H, 2(CH<sub>3</sub>)<sub>3</sub>], 3.0–3.7 (m, 8H, 4 CH<sub>2</sub>), 4.75 (m, 1H, NHBOC), 7.7–8.7 (m, 4H, C<sub>6</sub>H<sub>4</sub>), 9.0 (m, 1H, CO-NH-CH<sub>2</sub>); MS (matrix, glycerol) FAB > O: 495 [M+H]<sup>+</sup>, 395 [M+2H-BOC]<sup>+</sup>, 295 [M+3H-2BOC]<sup>+</sup>.

*N*<sup>1</sup>-[[3-(3-Aminobenzene-1-yl)carbonyl]-N<sup>4</sup>,N<sup>8</sup>-bis(tert-butoxycarbonyl)spermidine **14**

Methanol (10 ml) cooled to –20°C was added to 10% palladium on charcoal (243 mg) and **13** (1.67 g, 3.38 mmol) was added. The solution was hydrogenated at atmospheric pressure and room temperature. After the calculated amount of hydrogen (≈ 250 ml) was taken up (≈ 4 h), the catalyst was removed by filtration, the methanol was removed *in vacuo*, and the evaporation was repeated with some methanol. The intermediate amine **14** (1.5 g, 96% yield) was not further purified but used directly for the synthesis of **15**; MS (matrix, glycerol) FAB > O: 465 [M+H]<sup>+</sup>, 365 [M+2H-BOC]<sup>+</sup>, 265 [M+3H-2BOC]<sup>+</sup>.

*N*<sup>1</sup>-[[3-(3-Nitrobenzene-1-carboxamido)-benzen-1-yl]-carbonyl]-N<sup>4</sup>,N<sup>8</sup>-bis(tert-butoxycarbonyl)spermidine **15**

To a solution of amine **14** (0.646 g, 1.39 mmol) in anhydrous pyridine (2 ml) was dropwise added at 0°C a solution of the acyl chloride **2** (0.258 g, 1.39 mmol) in anhydrous acetonitrile (1 ml), and the mixture was stirred at ambient temperature for 1 h. After dilution with CH<sub>2</sub>Cl<sub>2</sub>, the mixture was poured into a saturated NaHCO<sub>3</sub> aqueous solution. The organic phase was separated, twice washed with water, dried over sodium sulfate, and evaporated. Chromatography of the residue on a silica gel column using as eluent a stepwise gradient of acetone (0–50%) in CH<sub>2</sub>Cl<sub>2</sub> led to the isolation of pure **15** (0.79 g, 92% yield); <sup>1</sup>H NMR (Me<sub>2</sub>SO-*d*<sub>6</sub>) δ 1.4–1.6 (m, 6H, 3CH<sub>2</sub>), 1.42 and 1.44 [2s, 2 × 9H, 2(CH<sub>3</sub>)<sub>3</sub>], 3.1–3.3 (m, 8H, 4CH<sub>2</sub>), 4.7 (br s, 1H, NHBOC), 7.4–9.7 (m, 8H, 2 C<sub>6</sub>H<sub>4</sub>), 7.9 (br s, 1H, NH-CH<sub>2</sub>), 8.1 (br s, 1H, NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix, glycerol) FAB > O: 614 [M+H]<sup>+</sup>, 514 [M+2H-BOC]<sup>+</sup>.

*N*<sup>1</sup>-[[3-(3-Aminobenzene-1-carboxamido)-benzen-1-yl]-carbonyl]-N<sup>4</sup>,N<sup>8</sup>-bis(tert-butoxycarbonyl)spermidine **16**

This compound was synthesized (following a similar procedure as described for the preparation of **14**) from the nitro-derivative **15** (1.75 g, 2.85 mmol) by catalytic hydrogenation with 10% palladium on charcoal (243 mg) in methanol (10 ml). After a similar workup, the residue obtained by evaporation of methanol was chromatographed on a silica gel column using as eluent a stepwise gradient of methanol (0–20%) in CH<sub>2</sub>Cl<sub>2</sub> to afford pure **16** (1.58 g, 95% yield); <sup>1</sup>H NMR (Me<sub>2</sub>SO-*d*<sub>6</sub>) δ 1.1–1.4 [m, 22H, 2(CH<sub>3</sub>)<sub>3</sub>C and 2CH<sub>2</sub>], 1.7 (m, 2H, CH<sub>2</sub>), 2.9–3.3 (m, 8H, 4CH<sub>2</sub>), 5.28 (s, 2H, NH<sub>2</sub>), 6.7 (m, 1H, CH<sub>2</sub>-NHBOC), 6.7–8.2 (m, 8H, 2 C<sub>6</sub>H<sub>4</sub>), 8.37 (t, 1H, CO-NH-CH<sub>2</sub>), 10.16 (s, 1H, NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix, glycerol) FAB > O: 584 [M+H]<sup>+</sup>, 484 [M+2H-BOC]<sup>+</sup>, 384 [M+3H-2BOC]<sup>+</sup>.

*N*<sup>1</sup>-[[3-[3-(Guanidinoacetamido)benzene-1-carboxamido]-benzen-1-yl]carbonyl]-N<sup>4</sup>,N<sup>8</sup>-bis(tert-butoxycarbonyl)spermidine hydrochloride **17**

To a solution of compound **16** (0.583 g, 1.0 mmol) and guanidinoacetic acid hydrochloride [29] (0.154 g, 1.0 mmol) in anhydrous DMF (1 ml) was added a solution of DCC (0.309 g, 1.5 mmol) in DMF (1 ml). After the mixture was stirred for a night at room temperature, the formed precipitate of dicyclohexylurea was removed by filtration, and the filtrate was evaporated. The residue was chromatographed on a silica gel column using as eluent a stepwise gradient of methanol (0–30%) with 1% of water in CH<sub>2</sub>Cl<sub>2</sub>. After evaporation of the appropriate fractions, lyophilization from dioxane gave pure **17** (0.38 g, 53%); <sup>1</sup>H NMR (Me<sub>2</sub>SO-*d*<sub>6</sub>) δ 1.3–1.4 [m, 22H, 2(CH<sub>3</sub>)<sub>3</sub>C and 2CH<sub>2</sub>], 1.7 (m, 2H, CH<sub>2</sub>), 2.9 (m, 2H, CH<sub>2</sub>), 3.1–3.3 (m, 6H, 3CH<sub>2</sub>), 4.10 (d, 2H, NH-CH<sub>2</sub>-CO), 6.76 (t, 1H, CH<sub>2</sub>-NH-BOC), 7.3 [br s, 4H, C(=NH<sub>2</sub><sup>+</sup>)NH<sub>2</sub>], 7.64 [t, 1H, CH<sub>2</sub>-NH-C(=NH<sub>2</sub><sup>+</sup>)NH<sub>2</sub>], 7.4–8.2 (m, 8H, 2 C<sub>6</sub>H<sub>4</sub>), 8.43 [t, 1H, CO-NH-CH<sub>2</sub>], 10.39 and 10.50 (2s, 1H each, 2 CO-NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix glycerol) FAB > O: 683 [M+H-HCl]<sup>+</sup>, 483 [M+2H-HCl-BOC]<sup>+</sup>; HPLC retention time 15.5 min, spectrophotometric purity at 254 nm: 100% (UV, λ<sub>max</sub> 227 nm) (same apparatus, column and solvents as described for the characterization of **7**; linear gradient 0–100% B in 10 min, then 100% B for 10 min; flow rate 3.0 ml/min).

*N*<sup>1</sup>-[[3-[3-(Guanidinoacetamido)benzen-1-carboxamido]-benzen-1-yl]-carbonyl]-spermidine tris(trifluoroacetate) **18**

Compound **17** (0.05 g, 0.07 mmol) was dissolved in trifluoroacetic acid (TFA, 1 ml). After 20 min, the solution was evaporated to dryness, the resulting residue was dissolved in water and the aqueous layer was twice washed with ethyl ether. After evaporation of water, lyophilization from water gave pure **18** (0.054 g, 93% yield); <sup>1</sup>H NMR (Me<sub>2</sub>SO-*d*<sub>6</sub>) δ 1.6 and 1.9 (2 m, 4 and 2H, 3CH<sub>2</sub>), 2.8, 3.0 and 3.3 (3m, 2, 4 and 2H, 4 CH<sub>2</sub>-N), 4.08 (d, 2H, NH-CH<sub>2</sub>-CO), 7.5–8.6 (m, 8H, 4 C<sub>6</sub>H<sub>4</sub>), 7.4, 7.7, 7.9 and 8.6 [4 br s, 4, 1, 3 and 2H respectively, C(=NH<sub>2</sub><sup>+</sup>)NH<sub>2</sub>, NH, NH<sub>2</sub><sup>+</sup> and NH<sub>2</sub><sup>+</sup>], 8.63 (t, 1H, CO-NH-CH<sub>2</sub>), 10.39 and 10.41 (2s, 2H, 2 CO-NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix, glycerol) FAB > O: 597 [M+H 2CF<sub>3</sub>CO<sub>2</sub>H]<sup>+</sup>, 483 [M+H 3 CF<sub>3</sub>CO<sub>2</sub>H]<sup>+</sup>; HPLC retention, time 9.2 min, spectrophotometric purity at 254 nm: 100% (UV, λ<sub>max</sub> 227 nm) [same apparatus, column, solvents and conditions as described for the characterization of **17**].

*N*<sup>1</sup>-[[3-[3-(Succinamido)benzene-1-carboxamido]-benzen-1-yl]carbonyl]-N<sup>4</sup>,N<sup>8</sup>-bis(tert-butoxycarbonyl)spermidine triethylammonium **19**

Compound **16** (0.498 mg, 0.853 mmol) was first coevaporated 3 times with anhydrous pyridine and then dissolved in pyridine, and 4-dimethylaminopyridine (DMAP, 0.183 g, 1.5 mmol) was added to the solution.

The solution was evaporated to dryness, and after 3 coevaporations with anhydrous acetonitrile, the mixture of **16** and DMAP was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (4.5 ml). To the stirred solution succinic anhydride (0.150 g, 1.5 mmol) and triethylamine (0.21 ml, 1.5 mmol) were added. The reaction was stirred for 2 h at ambient temperature. Saturated aqueous NaHCO<sub>3</sub> (30 ml) was added, and the resulting mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layers were washed once with water, dried over sodium sulfate and evaporated to dryness. The residue was chromatographed on a silica gel column using as eluent a stepwise gradient of methanol (0–20%) with 1% of triethylamine in CH<sub>2</sub>Cl<sub>2</sub> to

afford **19** (0.57 g, 85% yield). This compound was sufficiently pure (TLC and mass spectrum analyses [matrix, glycerol; FAB < O: 682 [M-H-(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>N]-, 582 [M-H-(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>N-BOC]-] to be used directly for the preparation of **20**.

*N*<sup>1</sup>-[[3-{3-[N<sup>4</sup>,N<sup>8</sup>-bis(tert-butoxycarbonyl)spermidinosuccinamido]benzene-1-carboxamido}-benzen-1-yl]carbonyl]-N<sup>4</sup>,N<sup>8</sup>-bis(tert-butoxycarbonyl)spermidine **20**

To a solution of compound **19** (0.57 g, 0.73 mmol) and N<sup>4</sup>, N<sup>8</sup>-bis(tert-butoxycarbonyl)spermidine [32] (0.376 g, 1.09 mmol) in anhydrous acetonitrile (11 ml) was added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate [30, 31] (BOP, 0.321 g, 0.73 mmol). The mixture was stirred at room temperature for 2 h and then saturated NaCl solution (35 ml) was added. Compound **20** was extracted with ethyl acetate (3 x 10 ml). The combined organic layers were washed successively with 2 N HCl (50 ml), water (50 ml), 5% NaHCO<sub>3</sub> (50 ml) and water and then dried over sodium sulfate and filtered. The filtrate was evaporated to dryness. The resulting residue was chromatographed on a silica gel column using as eluent a stepwise gradient of methanol (0–10%) in CH<sub>2</sub>Cl<sub>2</sub>. After evaporation of the appropriate fractions, lyophilization from dioxane gave pure **20** (0.36 g, 49% yield); <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 1.3–1.4 [m, 44H, 4(CH<sub>3</sub>)<sub>3</sub>C and 4CH<sub>2</sub>], 1.6 and 1.7 (2m, 2H each, CH<sub>2</sub> each), 2.41 and 2.58 (2t, 2H each, CO-CH<sub>2</sub>-CH<sub>2</sub>-CO), 2.9 and 3.2 (2m, 4H and 12H respectively, 8CH<sub>2</sub>), 6.8 (br s, 2H, 2 CH<sub>2</sub>-NH-BOC), 7.8 [br s, 1H, CO-NH-(CH<sub>2</sub>)<sub>3</sub>], 7.4–8.2 (m, 8H, 4 C<sub>6</sub>H<sub>4</sub>), 8.4 (m, 1H, CO-NH-CH<sub>2</sub>), 10.12 and 10.35 (2s, 1H each, 2CO-NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix, glycerol) FAB > 0: 1011 [M+H]<sup>+</sup>, 911 [M+2H-BOC]<sup>+</sup>, 811 [M+3H-2BOC]<sup>+</sup>, 711 [m+4H-3BOC]<sup>+</sup>, 611 [M+5H-4BOC]<sup>+</sup>.

*N*<sup>1</sup>-[[3-{3-[N<sup>1</sup>-(spermidino)succinamido]benzene-1-carboxamido}-benzen-1-yl]carbonyl]-spermidine tetra(trifluoroacetate) **21**

Compound **20** (0.05 g, 0.049 mmol) was dissolved in TFA (2 ml). After 15 min, the solution was evaporated to dryness, the residue was dissolved in water and the aqueous layer was twice washed with ethyl ether and lyophilized from water to give pure **21** (0.051 g, 98% yield); <sup>1</sup>H NMR (Me<sub>2</sub>SO-d<sub>6</sub>) δ 1.6, 1.7 and 1.9 (3m, 8, 2 and 2H respectively, 6CH<sub>2</sub>), 2.44 and 2.61 (2t, 2H each, CO-CH<sub>2</sub>-CH<sub>2</sub>-CO), 2.8–3.0, 3.1 and 3.4 (3m, 12, 2 and 2H respectively, 8 CH<sub>2</sub>), 7.9 (br s, 6H, 2NH<sub>3</sub><sup>+</sup>), 8.09 (t, 1H, NH-CO), 7.4–8.3 (m, 8H, 2C<sub>6</sub>H<sub>4</sub>), 8.6 (m, 5H, 2NH<sub>3</sub><sup>+</sup> and CO-NH-CH<sub>2</sub>), 10.18 and 10.39 (2s, 1H each, 2CO-NH-C<sub>6</sub>H<sub>4</sub>); MS (matrix, glycerol) FAB > 0: 611 [M+H-4CF<sub>3</sub>CO<sub>2</sub>H]<sup>+</sup>; HPLC retention time 9.0 min, spectrophotometric purity at 254 nm: ≥ 99% (apparatus C; same column as described for the characterization of **9**; solvent A, 100% 0.1 M ammonium acetate buffer, pH 5.9; solvent B, 50% of acetonitrile in the same buffer; linear gradient 0–100% B in 10 min; flow rate 1.0 ml/min).

*Determination of association constant values to polynucleotides*

The binding constants (*K*<sub>app</sub>) of the test compounds to poly(dA-dT)·poly(dA-dT) and poly(dG-dC)·poly(dG-dC) were determined based on their ability to compete with the binding of ethidium bromide, according to Baguley *et al* [41] and as described previously for other netropsin analogues [6].

## Biological methods

### Cytostatic activity assays

The cytostatic activity assays were performed according to previously established procedures [42, 43].

### Anti-human immunodeficiency virus (HIV) assay

The method for determining the cytopathic effect of HIV in MT-4 cells has been previously described [46]. Briefly, MT-4 cells were seeded at 10<sup>6</sup> cells/ml and infected with HIV at 1000 CCID<sub>50</sub>/ml. Then, 10<sup>5</sup> cells were brought into wells of a microtiter tray containing 100 μl of various dilutions of the test compounds. After 5 d incubation at 37°C, the number of viable cells was determined microscopically in a hemacytometer by trypan blue exclusion.

### Antiviral activity assays

The antiviral assays were performed as previously reported [44, 45].

## Acknowledgments

The investigations were supported by grants from the French CNRS and INSERM (PNR-SIDA), the Belgian Fonds voor Geneeskundig Wetenschappelijk Onderzoek, the Belgian Geconcerteerde Onderzoeksacties, and the AIDS Basic Research Programme of the European Community. We thank JW Lown (Edmonton, Canada) for its interest and helpful discussions. We are indebted to A van Lierde, F De Meyer, L van Berckelaer and A Absillis for their excellent technical assistance with the biological assays. The assistance of C Duguet in typing this manuscript is also greatly appreciated.

## References

- 1 Portion of this work was presented at the NACON Meeting on Recognition Studies in Nucleic Acids, Sheffield University (UK), April 16–20th, 1989. This work is taken in part from the PhD Dissertation of F Debart, Université de Montpellier II, 1989
- 2 Lown JW (1988) *Anti-Cancer Drug Design* 3, 25–40
- 3 Cohen JS (1989) In: *Oligodeoxynucleotides, Antisense Inhibitors of Gene Expression, Topics in Molecular and Structural Biology*. MacMillan Press, London, vol 12, 255
- 4 Zimmer C, Wahnert U (1986) *Prog Biophys Mol Biol* 47, 31–112
- 5 Bailly C, Houssin R, Bernier JL, Henichart JP (1988) *Tetrahedron* 44, 5833–5843
- 6 Debart F, Périgaud C, Gosselin G, Mrani D, Rayner B, Le Ber P, Auclair C, Balzarini J, De Clercq E, Paoletti C, Imbach JL (1989) *J Med Chem* 32, 1074–1083
- 7 Rajagopalan M, Ayyer J, Sasisekharan V (1985) *J Biosci* 7, 27–32
- 8 Dasgupta D, Rajagopalan M, Sasisekharan V (1986) *Biochem Biophys Res Commun* 140, 626–631
- 9 Parrack P, Dasgupta D, Ayyer J, Sasisekharan V (1987) *FEBS Lett* 212, 297–301
- 10 Rajagopalan M, Rao KE, Ayyer J, Sasisekharan V (1987) *Indian J Chem* 26B, 1021–1024
- 11 Rao KE, Dasgupta D, Sasisekharan V (1988) *Biochemistry* 27, 3018–3024

- 12 Nishiwaki E, Tanaka S, Lee H, Shibuya M (1988) *Heterocycles* 27, 1945–1952
- 13 Leinsoo TA, Nikolaev VA, Grokhovskii SL, Strel'tsov SA, Zasedatelev AS, Zhuze AL, Gurskii GV (1988) *Mol Biol* 22 (part 2), 134–148
- 14 Arcamone FM, Animati F, Barbieri B, Configliacchi E, D'Alessio R, Geroni C, Giuliani FC, Lazzari E, Menozzi M, Mongelli N, Penco S, Verini MA (1989) *J Med Chem* 32, 774–778
- 15 Lown JW, Krowicki K, Balzarini J, Newman RA, De Clercq E (1989) *J Med Chem* 32, 2368–2375
- 16 Lown JW, Krowicki K, Bhat UG, Skorobogaty A, Ward B, Darbrowiak JC (1986) *Biochemistry* 25, 7408–7416
- 17 Kissinger K, Krowicki K, Dabrowiak JC, Lown JW (1987) *Biochemistry* 26, 5590–5595
- 18 Zakrzewska K, Lavery R, Pullman B (1987) *J Biomol Struct Dyn* 4, 833–843
- 19 Lee M, Chang DK, Hartley JA, Pon RT, Krowicki K, Lown JW (1988) *Biochemistry* 27, 445–455
- 20 Lee M, Pon RT, Krowicki K, Lown JW (1988) *J Biomol Struct Dyn* 5, 939–949
- 21 Lee M, Coulter DM, Pon RT, Krowicki K, Lown JW (1988) *J Biomol Struct Dyn* 5, 1059–1087
- 22 Krowicki K, Lee M, Hartley JA, Ward B, Kissinger K, Skorobogaty A, Dabrowiak JC, Lown JW (1988) In: *Structure and Expression, Vol 2: DNA and Its Drug Complexes* (Sarma RH, Sarma MH, eds), Adenine Press, Guiderland, NY, 251–271
- 23 Zimmer C, Luck G, Burckhardt G, Krowicki K, Lown JW (1988) In: *Structure and Expression, Vol 2: DNA and Its Drug Complexes* (Sarma RH, Sarma MH, eds), Adenine Press, Guiderland, NY, 291–303
- 24 Lee M, Hartley JA, Pon RT, Krowicki K, Lown JW (1988) *Nucleic Acids Res* 16, 665–684
- 25 Julia M, Gombert R (1968) *Bull Soc Chim Fr* 369–375
- 26 Julia M, Gombert R (1968) *Bull Soc Chim Fr* 376–382
- 27 Huffman CW (1958) *J Org Chem* 23, 727–729
- 28 Djerassi C, Scholz CR (1947) *J Am Chem Soc*, 69, 1688–1692
- 29 Julia M, Préau-Joseph N (1967) *Bull Soc Chim Fr* 4348–4356
- 30 Castro B, Dormoy JR, Dourtooglou B, Evin G, Selve C, Ziegler JC (1976) *Synthesis* 751–752
- 31 Dormoy JR, Castro B (1981) *Tetrahedron* 37, 3699–3706
- 32 Humora M, Quick J (1979) *J Org Chem* 44, 1166–1168
- 33 Bialer M, Yagen B, Mechoulam R (1980) *J Heterocycl Chem* 17, 1797–1798. For a discussion on attempted improvement during the introduction of the guanidinoacetyl moiety, see: Krowicki K, Lown JW (1987) *J Org Chem* 52, 3493–3501
- 34 Schabel FM, Laster WR, Brockman RW, Skipper HE (1953) *Proc Soc Exp Biol Med* 83, 1–3
- 35 Sugivra K (1959) *Gann* 50, 251–264
- 36 Werner GH, Maral R (1963) *Actual Pharmaceut Fr* 21, 133–167
- 37 Chirigos MA, Rusher RJ, Kamel IA, Fanning GR, Golding A (1963) *Cancer Res* 23, 1646–1650
- 38 Becker Y, Neuman S, Levitt-Hadar J (1972) *Isr J Med Sci* 8, 1225–1230
- 39 Becker Y, Asher Y, Zakay-Rones Z (1972) *Antimicrob Agents Chemother* 483–488
- 40 Kotler M, Becker Y (1972) *FEBS Lett* 21, 222–226
- 41 Baguley BC, Denny WA, Atwell GJ, Cain BF (1981) *J Med Chem* 24, 520–525
- 42 De Clercq E, Balzarini J, Torrence PF, Mertes MP, Schmidt CL, Shugar D, Barr PJ, Jones AS, Verhelst G, Walker RT (1981) *Mol Pharmacol* 19, 321–330
- 43 Balzarini J, De Clercq E, Torrence PF, Mertes MP, Park JS, Schmidt CL, Shugar D, Barr PJ, Jones AS, Verhelst G, Walker RT (1982) *Biochem Pharmacol* 31, 1089–1095
- 44 De Clercq E, Descamps J, Verhelst G, Walker RT, Jones AS, Torrence PF, Shugar D (1980) *J Infect Dis* 141, 563–574
- 45 De Clercq E, Montgomery JA (1983) *Antiviral Res* 3, 17–24
- 46 Balzarini J, Herdewijn P, Pauwels R, Broder S, De Clercq E (1988) *Biochem Pharmacol* 37, 2395–2403