

## The Ant and the Grasshopper: contrasting responses and behaviors to water stress of riparian trees along a hydroclimatic gradient

Pierre Lochin, Pauline Malherbe, Baptiste Marteau, Julien Godfroy, Michael Bliss Singer, John Stella, Hervé Piégay, Antoine Vernay

## ▶ To cite this version:

Pierre Lochin, Pauline Malherbe, Baptiste Marteau, Julien Godfroy, Michael Bliss Singer, et al.. The Ant and the Grasshopper: contrasting responses and behaviors to water stress of riparian trees along a hydroclimatic gradient. European Geosciences Union General Assembly 2024, Apr 2024, Vienne (Autriche), Austria. 10.5194/egusphere-egu24-12445. hal-04651472

## HAL Id: hal-04651472 https://hal.science/hal-04651472v1

Submitted on 17 Jul2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



EGU24-12445, updated on 17 Jul 2024 https://doi.org/10.5194/egusphere-egu24-12445 EGU General Assembly 2024 © Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



## The Ant and the Grasshopper: contrasting responses and behaviors to water stress of riparian trees along a hydroclimatic gradient

**Pierre Lochin**<sup>1</sup>, Pauline Malherbe<sup>1</sup>, Baptiste Marteau<sup>2</sup>, Julien Godfroy<sup>3</sup>, Michael Bliss Singer<sup>4,5,6</sup>, John Stella<sup>7</sup>, Hervé Piégay<sup>1</sup>, and Antoine Vernay<sup>8</sup> <sup>1</sup>ENS de Lyon, UMR 5600 Environnement Ville société, CNRS, Lyon, France <sup>2</sup>Univ Rennes 2, UMR6554 LETG, F-35043 Rennes, France

<sup>3</sup>Univ. Grenoble Alpes, INRAE, LESSEM, F-38402 St-Martind'Hères, France

<sup>4</sup>Earth Research Institute, University of California, Santa Barbara, CA 93106, USA

<sup>5</sup>Water Research Institute, Cardiff University, Cardiff CF10 3AX, UK

<sup>6</sup>School of Earth and Environmental Sciences, Cardiff University, Cardiff CF10 3AT, UK

<sup>7</sup>Department of Sustainable Resources Management, State University of New York College of Environmental Science and Forestry, Syracuse, NY 13210, USA

<sup>8</sup>Université Claude Bernard Lyon 1, LEHNA UMR 5023, CNRS, ENTPE, F-69622, Villeurbanne, France

The increasing threat of forest decline and mortality associated with more severe and frequent droughts calls for a better understanding of the tree response mechanisms to water stress. Among forest ecosystems, riparian forests, despite their privileged location in lowlands, are particularly vulnerable to drought because they are highly dependent on soil water availability for survival. At the scale of a large river and along a hydroclimatic gradient, riparian tree species may respond differently to water stress and, therefore, may not be equally vulnerable to drought events outside their normal range. To analyze the responses of white poplar (*Populus alba*) to seasonal changes in drought along a hydroclimatic gradient, we conducted a multi-tool analysis combining multispectral and thermal infrared remote sensing with in-situ ecophysiological measurements at different temporal scales.

Using this approach, we demonstrated that white poplars along the Rhône River (France) exhibit contrasting responses and behaviors during drought. We found that white poplars located downstream of the hydroclimatic gradient in a Mediterranean climate rapidly close their stomata during drought to reduce water loss and maintain high levels of minimum water potential ( $\Psi_m$ ), but at the expense of leaf density and greenness. Conversely, white poplars located upstream in a temperate climate show high levels of transpiration and stable greenness, but lower  $\Psi_m$  and water content. These results demonstrate that white poplars along an aridity gradient can have two opposing responses to drought, with isohydric regulation on the one hand and anisohydric regulation on the other. These results indicate that trees located upstream in an area unaccustomed to drought are at a high risk of hydraulic failure during more intense and prolonged droughts.

The combined use of different tools and metrics at different spatial and temporal scales is

therefore essential to consider the full range of tree responses to drought. These two distinct behaviors remind us of Aesop's fable, where upstream trees behave like the grasshopper, not paying attention to their water consumption and jeopardizing their long-term survival. In contrast, downstream white poplars respond quickly to drought and show greater long-term resilience, just like the ant. These results are important in light of future climate conditions because they show that the same species can have different levels of resilience to drought, but they also raise the question of the ability of trees to switch from grasshopper to ant to adapt to these future conditions.