






PROTEKER: Implementation of an underwater observatory at the Kerguelen Islands (Southern Ocean)

J.-P. Féral, T. Saucède, E. Poulin, C. Marschal, G. Marty,
J.-C. Roca, S. Motreuil, J.-P. Beurier









Antarctic Circumpolar Current

Kerguelen archipelago emerges in a very dynamic region




- Kerguelen archipelago emerges in a very dynamic region
- In the flow of ACC
 - At the confluence between several water masses AASW, PFSW and SASW
 - Close to the Antarctic convergence (PF)


Subantarctic Surface Water (SASW)
Polar Frontal Surface Water (PFSW)
Antarctic Surface Water (AASW)




Austral Ocean's hydrologic fronts and water masses



Position of the Polar Front




The 2°C isotherm corresponding to Park's PF (2014) is shown by a thick black line.




Position of the Polar Front

Kerguelen area is a major confluence where tropical and subtropical surface waters meet and interact with waters of Subantarctic and Antarctic origins




Subtropical Surface Water (**STSW**)
 Subantarctic Surface Water (**SASW**)
 Polar Frontal Surface Water (**PFSW**)
 Antarctic Surface Water (**AASW**)

Température de surface (°C)



Visiting the [predictive] models ...







Années 70

Années 2000


AIB 2100

AIB 2200

Température de surface (°C)**Amplitude absolue température de surface (°C)****Salinité de surface**



Maps drawn from data of WOCE (extent/old) and Bio-Oracle (future) in the framework of scenarios AIB (IPCC, 2007)



'Today' water masses shift


Doesn't take into account the south shift of the fronts (!) whose position is even not well defined ...

For 2100 and 2200, monthly mean temperature (SST, SAT) and average monthly mean salinity conditions over 10-year periods (2087–2096 and 2187–2196) were built with the R package “raster” (Hijmans and van Etten 2011). These grids represent environmental conditions at the end of the 21st and the 22nd century provided by the World Climate Research Programme Coupled Model Intercomparison Project (WCRP CMIP3) multi-model database (<http://esg.llnl.gov:8080/index.jsp>). This future scenario is based on A1B (720 ppm CO₂ stabilization) IPCC scenario represented by the UKMO-HadCM3 model.

Latitudinal shift of waters with other characteristics ($T^{\circ}C$, S°/oo , nutrients, ...)

→ environmental changes that may affect coastal biodiversity and ecosystem functioning.


Modern observations indicate that this highly dynamic region is already undergoing change in response to a warming climate.



Ampère glacier has retreated 5 km in 30 yrs

KERGUELEN # Glacier Ampère # Baie de la Table
Hélio do3


To be interpreted and to identify eventual trends, such changes must be recorded through the establishment of a long-term monitoring



A View of CHRISTMAS HARBOUR, in KERGUELEN's LAND.

To be interpreted and to identify eventual trends, such changes must be recorded through the establishment of a **long-term monitoring**.

The French Polar Institute (IPEV) program PROTEKER aims to implement such an **underwater observatory** that considers a multidisciplinary approach: oceanographic measurements, benthic mapping, genetic, ecophysiological, isotopic and environmental analyzes.



A View of CHRISTMAS HARBOUR, in KERGUELEN's LAND.

To be interpreted and to identify eventual trends, such changes must be recorded through the establishment of a **long-term monitoring**.


The French Polar Institute (IPEV) program PROTEKER aims to implement such an **underwater observatory** that considers a multidisciplinary approach: oceanographic measurements, benthic mapping, genetic, ecophysiological, isotopic and environmental analyzes.

In addition to the inventory and the monitoring of biodiversity, it also aims at providing **scientific criteria** to managers in charge of **protection and conservation policies** (French Southern Lands National Nature Reserve).

Thinking an
underwater
observatory
to track
warming,
acidification,
etc ...


OceanCube central node at Motobu Point, Okinawa

Thinking an
underwater
observatory
to track
warming,
acidification,
etc ...



OceanCube central node at Motobu Point, Okinawa

Thinking an
underwater
observatory
to track
warming,
acidification,
etc ...



Is this
possible in
Kerguelen?

There is a need to adapt practices to local conditions and circumstances

Between the roaring forties and the furious fifties

- ➔ weather and sea often harsh
- ➔ accessibility can be tough, risk of loss and degradation of gear
- ➔ implemented gear must be highly resistant, as inexpensive as possible, able of several years of functioning

Support vessel not always available

- ➔ possible gap in the observations and measurements
- ➔ great autonomy


Little / insufficient funding

- ➔ looking for "low cost"
- ➔ limited number of staff
- ➔ survey at sea difficult since it depends on a supporting vessel




The return of *La curieuse* to Kerguelen for summer campaigns

→PROTEKER (2011-2014)
phase 1



- To choose relevant sites

Not du Rendez-vous



Projection UTM - WGS84 datum



The cover of the magazine 'PROTEKER' is shown. The title 'PROTEKER' is written in large, stylized, block letters at the top. Below the title is a black and white photograph of a boat with a tall mast and a diver in the water. The word '100' is printed in a small box in the upper left corner.

OCÉAN INDIEN


49°5

PROTEKER 2011

- Campagne Nord
- Campagne Sud

- Campagne Morbihan, Norvégienne
- *Côte ouest / Ile de l'Ouest*

Not du Rendez-vous



Projection UTM - WGS84 datum



The cover of the magazine 'PROTEKER' is shown. The title 'PROTEKER' is written in large, stylized, block letters at the top. Below the title is a black and white photograph of a boat with a tall mast and a diver in the water. The word '100' is printed in a small box in the upper left corner.

OCÉAN INDIEN

49°S

done

PROTEKER 2011

- Campagne Nord
- Campagne Sud

Campagne Morbihan

Îles de Boynes

60°E

30°E

- 50° S



implemented
techniques

Beam Trawl



PROTEKER 2-27 II/2013
Station : Xmas
Latitude Sud :
Longitude Est :
Profondeur (m) : 100
Géotag 20 juillet 2013
Coll. J.-P. Féral, C. Marschal, G. Marty,
E. Poulin, T. Sauché

Provocator pulcher

Beam Trawl



Provocator pulcher

Remotely Operated Vehicle ROV




Beam Trawl



PROTEKER 2-27 II/2013
Station : Xmas
Latitude Sud :
Longitude Est :
Profondeur (m) : 100
Geste 20 profondeur (3)
Coll. J.-P. Féral, C. Marschal, G. Marty,
E. Poulin, T. Sauché

Provocator pulcher

Remotely Operated Vehicle ROV




Beam Trawl




Provocator pulcher

Remotely Operated Vehicle ROV



Labidiaster annulatus

Dot du Rendez-vous



Projection UTM - WGS84 datum



Océan Indien

- 49 -

The logo for PROTEKER features a stylized black silhouette of a scuba diver swimming towards the right. Above the diver, the word "PROTEKER" is written in a bold, sans-serif font, arched over the top. In the background, there is a white silhouette of a boat with a tall mast and a small cabin-like structure.

KERGUELEN monitored sites

Z = 5m

49°23'14,7" S

69°56'28,6" E

Photographs and GPS coordinates at each site

Ile Haute, thermoenregistreur 5m

Au pied de la paroi



Ile Haute

Z = 5m



49°23'14,7" S

69°56'28,6" E

Photographs and GPS coordinates at each site

Ile Haute, thermoenregistreur 5m

Au pied de la paroi





- To choose relevant sites
- To upgrade inventories (diving, towed gears, ROV)
 - guides (photos, video)
 - illustrated data bases (indexing, cataloguing, inter-operability, contextualization, traceability , international systems)


Shots / photographs & video



Field Identification Guide to Heard Island and McDonald Islands Benthic Invertebrates



A guide for scientific observers aboard fishing vessels



Ty Hibberd and Kirrily Moore

Pictorial guide to common HIMI taxa

Odontaster meridionalis



Five short arms with a mesh-like aboral surface.

Porania antarctica



Five armed orange/yellow coloured seastar with a mottled dorsal surface and soft, mucus covered oral surface. Marginal edge covered in conspicuous flattened spines.

Smilasterias triremis







Small, five arms, red/orange, tough, often arms missing. Four rows of suckered tube feet.



Photograph by Karen O'Regan
"Umbrella" shaped body with scaly surface.










- To choose relevant sites
- To upgrade inventories (diving, towed gears, ROV)
 - guides (photos, video)
 - illustrated data bases (indexing, cataloguing, inter-operability, contextualization, traceability , international systems)
- To setup sensors
- To estimate speed and quality of colonisation / recruitment (settlement plots)

Temperature logger




Temperature logger





-20 °C to +50 °C in water / waterproof to 120 m /
Accuracy: ±0.21°C over 0 to 50°C / 42000
measurements / logging intervals: 1 second to 18
hours, fixed / drift: 0.1°C per yr


Temperature logger



Settlement plots




-20 °C to +50 °C in water / waterproof to 120 m /
Accuracy: ±0.21°C over 0 to 50°C / 42000
measurements / logging intervals: 1 second to 18
hours, fixed / drift: 0.1°C per yr

An underwater photograph showing a kelp forest. In the foreground, a red, textured rock surface is visible. Several temperature loggers are deployed: one is attached to a vertical kelp stem on the left, another is suspended by a cable from a horizontal kelp stem in the center, and a third is mounted on a vertical kelp stem on the right. The water is a clear teal color.


Temperature loggers

T°C


Sea water temperature in the Baie du Morbihan at 15m depth , 2012-2015



Trois Bergers (15m)



Trois Bergers (15m)



[Proteker sites](#)[Campaign 2011-12](#)[Campaign 2013](#)[Campaign 2014](#)[Campaign 2016](#)[Home page](#) > [Thermorecorders](#) > Temperature monitoring, Ilot des Trois Bergers, Port Couvreux - Baleiniers sector

Rechercher dans le site

>>

**PROTEKER, IPEV program
1044 – Summer field campaigns**[Partners and participants](#)[Observation sites](#)[Thermorecorders](#)[Colonization plates](#)[What's up?](#)[Amsterdam](#)[Gallery](#)[Site Map](#)[Contact](#)[Member's logout](#)[Private area](#)

Jean-Pierre FERAL



Temperature monitoring, Ilot des Trois Bergers, Port Couvreux - Baleiniers sector

[\[Ilot des Trois Bergers 5m\]](#)[\[Ilot des Trois Bergers 15m\]](#)


2012

2013

2014

[\[Ilot des Trois Bergers 5m vs. 15m\]](#)**All data are copyright - ©PROTEKER**

Ilot des Trois Bergers, 15m depth



Publications

Derniers articles publiés

- ▶ Diaporama PROTEKER 1 (dec. 2011 - jan. 2012)
- ▶ Diaporama PROTEKER 2 (nov.-dec. 2013)
- ▶ Diaporama PROTEKER 4 (nov.-dec. 2015)
- ▶ pour voir la rubrique
- ▶ Iles du Prince de Monaco, 5m depth
- ▶ Iles du Prince de Monaco; 15m depth
- ▶ Iles du Prince de Monaco, 5m vs. 15m depth
- ▶ Fjord des Portes Noires, 15m depth
- ▶ Fjord des Portes Noires, 5m depth
- ▶ Fjord des Portes Noires, 5m vs. 15m depth
- ▶ Ilot des Trois Bergers, 5m vs. 15m depth
- ▶ Ilot des Trois Bergers, 15m depth

Statistics

Latest update


Saturday 16 April 2016

Publishing

58 Articles
Aucun album photo
No news item
No site
2 Authors


Visits

34 today
78 yesterday



Settlement plots






Settlement plots protocol

A_1

A_2




A_5

A_6


A_7

A_8



Ile Suhm, december 2013






Ile Suhm, december 2013



Ile Suhm, december 2014







Ile Suhm, december 2013




Ile Suhm, december 2014



A




B




C





D



E



F



H



Ile Suhm, december 2014



- To choose relevant sites
- To upgrade inventories (diving, towed gears, ROV)
 - guides (photos, video)
 - illustrated data bases (indexing, cataloguing, inter-operability, contextualization, traceability , international systems)
- To setup sensors
- To estimate speed and quality of colonisation / recruitment (settlement plots)
- To choose model species with large distribution area or endemic (population genetics, phylogeography → *connectivity, self recruitment*)
- To contribute to the necessary scientific bases to write a management plan of the coastal marine domain

Valuation



Model species
. *Abatus cordatus*.
. *Mytilus edulis*



Collaboration with Chile

Collaboration with labs of l'université de la Réunion and l'Institut de Chimie des Substances Naturelles du CNRS de Gif-sur-Yvette (active substances from Porifera)




Trophic webs (isotopic analyses)




First ROV exploration of île Amsterdam

Training of staff from the Nature Reserve




Phase 2

The overall scope of the project will depend on the support vessel availability (*La Curieuse* or equivalent)



Phase 2



The overall scope of the project will depend on the support vessel availability (*La Curieuse* or equivalent)

- At least the monitoring of the Baie du Morbihan [4 sites] will be ensured (small vessels such as *Le Commerson*)
- External (north and south) sites will depend on the route and schedule of the research and supply vessel *Marion Dufresne 2*

Phase 2



The overall scope of the project will depend on the support vessel availability (*La Curieuse* or equivalent)

- At least the monitoring of the Baie du Morbihan [4 sites] will be ensured (small vessels such as *Le Commerson*)
- External (north and south) sites will depend on the route and schedule of the research and supply vessel *Marion Dufresne 2*

In any case, the next step will be to adapt monitoring protocol produced by coastal research programs

Phase 2



The overall scope of the project will depend on the support vessel availability (*La Curieuse* or equivalent)

- At least the monitoring of the Baie du Mont-Saint-Michel [4 sites] will be ensured (small vessels such as Dufresne 2 and Emerson)
- External (north and south) and schedule of the Marion Dufresne 2

In any case, the protocol produced

Late Breaking News
La Curieuse will be in
Kerguelen next
(austral) summer !!

La Curieuse, Kerguelen,
PROTEKER 1 2011-12



La Curieuse, Kerguelen,
PROTEKER 3 2014



Thank you for attention

La Curieuse, Kerguelen,
PROTEKER 2 2013



Scientific divers:

Jean-Pierre Beurier [1]
Jean-Pierre Féral [1, 2, 3]
Christian Marschal [1, 2, 3]
Gilles Marty [2, 3]
Sébastien Motreuil [3]
Elie Poulin [2]
Jean-Claude Roca [1, 3]
Thomas Saucède [2]

Crew of La Curieuse for the 3 campaigns:

Baptiste Aspa (chief engineer) [1]
Vincent Benard (mechanic trainee) [3]
Mathias Cadet (mechanic trainee) [2]
Benoît Caffier (cook/deckhand) [1, 2, 3]
Laurent Conillen (first mate) [1]
Alix Daujat (captain) [1]
Corentin Hemmer (2nd engineer) [3]
Sébastien Laffont (deckhand) [3]
Pierre-Yves Le Bren (chief engineer) [2]
Arnaud Michelot (chief engineer) [3]
Yohann Mucherie (captain) [2, 3]
Jimmy Payet (deckhand) [1]
Xavier Payet (deckhand) [2]
Yann Rivière (mechanic trainee) [1]
Anthony Sautron (2nd engineer) [1, 2]
Pierre Samuel (first mate) [3]
Loïc Seguinneau (first mate) [2]

Romain Vergé (skipper of *Le Commerson*) [3]



An underwater photograph of a dense kelp forest. The scene is filled with long, brownish-yellow kelp fronds swaying in the current. Sunlight filters down from the surface in bright rays, illuminating the fronds and creating a dappled light effect. The water is a clear blue-green color. In the foreground, there are darker, more shadowed areas of the kelp and some small, colorful marine life or debris.

www.proteker.net

Mercí de
Votre
attention