



HAL
open science

Spiders in oyster shells: shellfish farming benefits an endangered terrestrial species

Julien Pétilon, Nela Glorikova

► **To cite this version:**

Julien Pétilon, Nela Glorikova. Spiders in oyster shells: shellfish farming benefits an endangered terrestrial species. *Frontiers in Ecology and the Environment*, 2024, 22 (4), pp.e2744. 10.1002/fee.2744 . hal-04653804

HAL Id: hal-04653804

<https://hal.science/hal-04653804>

Submitted on 19 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Spiders in oyster shells: shellfish farming benefits an endangered terrestrial species

Along the shore of Mont-Saint-Michel Bay in western France, large numbers of non-native Pacific oyster (*Crassostrea gigas*) and native blue mussel (*Mytilus edulis*) shells are deposited as waste from marine aquaculture operations. The resulting “shell beds” (above) cover hundreds of hectares and form a distinctive habitat that over time has become integrated into the salt marsh vegetation (mostly native *Atriplex portulacoides*) through sediment accretion. The shell beds appear in the form of banks, typically 1–2 m high, up to 200–300 m long, and 50 m wide. Both sexes of an endangered wolf spider (*Arctosa fulvolineata*) use the shells as a refuge and as breeding habitat (below). During May 2021, the species was found on several occasions near Vivier-sur-Mer along the center of the Bay, hiding under the shells during the day and hunting for amphipods at night. *A. fulvolineata* likely uses shells as a daytime shelter during high tides, as reported for intertidal spiders (*Desis* spp) on South African coasts. While the importance of anthropogenic resource subsidies for predatory spiders has been known for decades, examples documented in the scientific literature remain rare.

Julien Pétillon^{1,2} and Nela Gloriková³

¹University of Rennes, Rennes, France;

²Nelson Mandela University, Port Elizabeth, South Africa;

³Crop Research Institute, Prague, Czech Republic

doi:10.1002/fee.2744

Image credits: ©N Gloriková

