



HAL
open science

Preliminary assessment of blood mercury contamination in four African crocodile species

Jérémy Lemaire, Paco Bustamante, Matthew Shirley

► **To cite this version:**

Jérémy Lemaire, Paco Bustamante, Matthew Shirley. Preliminary assessment of blood mercury contamination in four African crocodile species. *Environment International*, 2024, 190, pp.108877. 10.1016/j.envint.2024.108877 . hal-04655055

HAL Id: hal-04655055

<https://hal.science/hal-04655055>

Submitted on 21 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Preliminary assessment of blood mercury contamination in four African crocodile species

Jérémy Lemaire¹, Paco Bustamante², Matthew H. Shirley^{3,4}

¹ Department of Behavioral and Cognitive Biology, University of Vienna, Djerassiplatz 1, 1030, Vienna, Austria

² Littoral Environnement et Sociétés (LIENSs), UMR 7266 CNRS-La Rochelle Université, 2 rue Olympe de Gouges, 17000 La Rochelle, France

³ Project *Mecistops*, Sarasota, Florida, USA

⁴ Global Forensics and Justice Center, Florida International University, Miami, Florida, USA

Corresponding author: jeremy.lemaire@univie.ac.at

Statement of author contributions:

JL: Conceptualization, Formal analysis, Methodology, Writing – original draft.

PB: Laboratory Methodology, Funding.

MHS: Conceptualization, Investigation, Project administration, Funding, Field Sampling Methodology, Writing – original draft.

Abstract: Heavy metal contamination in the environment is an increasingly pervasive threat to the long-term persistence of wildlife. As high trophic level consumers, crocodylians are at substantial risk from bioaccumulation of mercury (Hg). Despite that they are generally well-studied and the focal species of many conservation efforts around the world, little is known about Hg contamination levels in most crocodylians. Here we preliminarily evaluate blood Hg contamination in four African species – Central African slender-snouted crocodile (*Mecistops leptorhynchus*), African dwarf crocodile (*Osteolaemus tetraspis*), West African crocodile (*Crocodylus suchus*), and Nile crocodile (*Crocodylus niloticus*) – from a diversity of sites and habitats across 5 different countries representing varying degrees of environmental pollution. All of our sampled crocodiles were Hg contaminated and, worryingly, these African crocodiles generally showed the highest levels of Hg contamination of any crocodylian species examined to date. Of most concern was that Hg concentrations were not only highest in *M. leptorhynchus*, the most threatened amongst our study species, but also in individuals sampled in what are believed to be some of the most remote and pristine natural areas left in Africa – Gabon’s national parks. Our results underscore the need to better understand the impact of longstanding petroleum, mining, forestry, and agricultural industries on the entire aquatic food chain throughout much of Africa, including on the threatened species in these habitats and the human populations that depend on them for their subsistence and livelihoods.

Keywords: Environmental contaminant, Gabon, Topical ecosystems, Petroleum, Mining

INTRODUCTION

Mercury (Hg) is a major contaminant affecting ecosystems worldwide (Chen *et al.*, 2018). While it is naturally present in the environment, anthropogenic activities such as fossil fuel combustion, deforestation, and mining, have been shown to drastically increase the level of Hg in the environment, particularly in aquatic ecosystems (Pirrone *et al.*, 2010; Hsu-Kim *et al.*, 2018). Under anoxic conditions, Hg can be transformed by aquatic microorganisms to methylmercury (MeHg), the most toxic and bioavailable form of Hg (Campeau and Barta, 1985; Benoit *et al.*, 2003). In its methylated form, Hg biomagnifies through the food chain and bioaccumulates over time in the tissues of top predators (Eagles-Smith *et al.*, 2018). Mercury has been identified to have deleterious effects on humans and wildlife, causing neurobehavioral and neuronal dysfunction, and further impacting reproduction, embryonic development, hormonal synthesis, metabolic processes, and immune functions (Fingerman *et al.*, 1996; Tan *et al.*, 2009; Tartu *et al.*, 2013; Scheuhammer *et al.*, 2015). Because of this, it is crucial to evaluate Hg contamination in the environment and its impact on biodiversity.

As high trophic level consumers and potentially apex predators within aquatic ecosystems (Somaweera *et al.*, 2020), crocodylians bioaccumulate environmental contaminants that biomagnify through the food webs. As a result, they are excellent species in which to monitor Hg contamination using minimally invasive sampling methods (Lázaro *et al.*, 2015; Lemaire *et al.*, 2021a). Most research to date on Hg in crocodylians has focused on the American alligator (*Alligator mississippiensis*; Yanochko *et al.*, 1997; Elsey *et al.*, 1999; Burger *et al.*, 2000; Rumbold *et al.*, 2002; Campbell *et al.*, 2010; Horai *et al.*, 2014; Nilsen *et al.*, 2017a) and Morelet's crocodile (*Crocodylus moreletii*; Rainwater *et al.*, 2002; Trillanes *et al.*, 2014; Buenfil-Rojas *et al.*, 2015, 2020; Romero-Calderón *et al.*, 2022; Thirion *et al.*, 2022). More recent efforts to understand Hg contamination across the Crocodylia, notably on the

spectacled caiman (*Caiman crocodilus*; Schneider *et al.*, 2015; Marrugo-Negrete *et al.*, 2019; Lemaire *et al.*, 2021b) and the two dwarf caimans (*Paleosuchus sp.*; Lemaire *et al.*, 2021a, 2021c, 2022), is revealing the possibility that crocodylians across diverse geographies are Hg contaminated. And Hg contamination has been identified as a cause of detrimental effects on crocodylians, including DNA damage and physiological alterations (Marrugo-Negrete *et al.*, 2019; Lemaire *et al.*, 2021b). It is, therefore, crucial to evaluate Hg contamination and its impacts across the Order, especially in contexts where Hg contamination is a likely aggravating factor for their conservation - for example, in the face of anthropogenic activities, like gold mining and petroleum extraction.

Africa is rich in natural and mineral resources, and commercial and artisanal-scale extraction of these resources is expected to intensify across the continent after the Covid19 outbreak (Yoshimura *et al.*, 2021). Gold mining, oil extraction, and agriculture at both industrial and artisanal scales are among the most polluting activities in the region (UNEP, 2019). While it is likely that such extractive activities have led to increased Hg contamination, this is poorly documented across the continent, especially in wildlife species of conservation concern like crocodiles. Africa is home to seven crocodile species, all of which have significant cultural, subsistence, and financial importance to communities and governments across the continent (Thorbjarnarson *et al.*, 1992; Ross, 1998). However, to the best of our knowledge, only two studies have evaluated Hg concentrations in any African crocodile species (Almli *et al.* 2005; Du Preez *et al.* 2018). They evaluated Hg contamination in the eggs, liver, and kidneys of Nile crocodiles (*Crocodylus niloticus*) in Zambia and South Africa and found median concentrations up to 3.7 $\mu\text{g.g}^{-1}$ (wet weight) in liver, up to 2.7 $\mu\text{g.g}^{-1}$ (wet weight) in kidneys, and up to 0.37 $\mu\text{g.g}^{-1}$ (dry wet) in eggshell.

In this study, we evaluated Hg concentrations in the blood, which represent the recent contamination, of four African crocodile species (*Crocodylus niloticus*, *Crocodylus suchus*, *Mecistops leptorhynchus*, and *Osteolaemus tetraspis*) across a diversity of African ecosystems, land management regimes, and economic development contexts. Specifically, we aimed to: 1) evaluate Hg contamination in blood of three of these four African crocodile species for the first time, and 2) preliminarily describe sources of variation in Hg concentration related to species, sex, and/or site characteristics in three of them.

MATERIALS AND METHODS

Study species

We evaluated Hg levels in four of Africa's seven endemic crocodile species. Between them, they represent all three genera present on the continent, as well as a significant diversity of habitats, life history strategies, and socioeconomic and development contexts, which will certainly impact their exposure risk to Hg.

Two species represent the "true crocodiles" of the genus *Crocodylus*. The Nile crocodile (*C. niloticus*) is one of the largest crocodile species in the world and is widely distributed across East and southern Africa. It is a habitat and dietary generalist, capable of occupying most wetland types throughout its distribution, ranging from estuarine coastal environments (e.g., Gabon and South Africa), to tropical forested wetlands (e.g., Gabon and DR Congo), savannah river and lake systems (throughout), and even arid and desert wetlands (e.g., Namibia and Egypt). Nile crocodiles are considered Least Concern (LC) under IUCN Red List criteria and are generally thought to have stable populations globally (Isberg *et al.*, 2019). The West African crocodile (*C. suchus*) was only recently re-recognized as a relatively cryptic species, highly divergent from the Nile crocodile, to which it is not even a sister species (e.g., Hekkala *et al.*,

2011; Shirley *et al.*, 2015). It is widely distributed across West, Central, and parts of North and East Africa, where it occupies a staggering diversity of habitats, ranging from estuarine coastal environments, tropical forested wetlands, Guinea savannah, and even the Sahara Desert (e.g., Chad and Mauritania). It is generally a medium-bodied crocodile, dietary generalist, and generally not a threat to people, often living near rural communities throughout its distribution. West African crocodiles are currently being evaluated against IUCN Red List criteria, but minimally meet the criteria for Vulnerable and most populations are in decline (Kpera *et al.*, In Prep.).

The other two species in this study are habitat specialists, mostly occupying the freshwater, forested wetlands of the Lower Guinea and Congo Basin biomes of Central Africa. The African dwarf crocodile (*O. tetraspis*) is one of the smallest crocodylian species in the world. It is highly terrestrial and forages mostly on smaller invertebrates, including crustaceans, arachnids, millipedes and snails, as well as other small forest prey items like frogs and fish (Shirley *et al.*, 2017). It is widely distributed throughout forested habitats, including in coastal lagoon mangrove forests and throughout small stream networks within dense forest. Dwarf crocodiles are considered Vulnerable under IUCN Red List criteria, and though some populations are declining, it can be quite abundant within intact habitats and where hunting pressure is not too excessive (Crocodile Specialist Group, 1996). In contrast, the Central African slender-snouted crocodile (*M. leptorhynchus*) is one of the most aquatic crocodylian species in the world, occupying medium- and larger-bodied rivers, lakes, lagoons, and other wetlands throughout the forested wetlands of Central Africa. Like most crocodylians, it will eat whatever it can catch, but is much more specialized in its diet consuming more aquatic prey (e.g., fish, snakes, waterbirds, frogs, etc...) (Shirley *et al.*, 2018). Central African slender-snouted crocodiles are currently being evaluated against IUCN Red List

criteria, but minimally meet the criteria for Endangered and most populations are in decline (Shirley *et al.*, In Review).

Sample collection and preparation

We collected blood samples from wild-caught individuals of the four different crocodylian species from different sites and habitats throughout their ranges between 2006 and 2011 (Fig. 1, Table 1). Specifically, we analyzed *O. tetraspis* (n = 81) samples from 3 different sites, *M. leptorhynchus* (n = 124) samples from 8 different sites, *C. suchus* (n = 56) samples from 6 different sites, and *C. niloticus* (n = 9) samples from a single site (Table 2). To collect the samples, we captured crocodylians using standard methods, including by hand, with tongs, pole snaring, and research darting (Cherkiss *et al.*, 2004; Walsh, 1987; Webb *et al.*, 1983). For each captured individual, we drew a 0.5 ml blood sample from the cervical vessel posterior the occiput and spread it evenly on the lower half of a strip (2.5 cm x 10 cm) of lab grade, standard Whatman filter paper. We allowed the samples to air dry prior to storing them in paper coin envelopes at ambient environmental conditions. We took a standard series of measurements (e.g., head length, snout to vent length (SVL), mass) and determined their sex prior to releasing all captured animals at the site of capture.

Though these samples were originally collected for genetic research (e.g., Hekkala *et al.*, 2011; Shirley *et al.*, 2014), prior research on both humans and wildlife has demonstrated that dried blood spots (DBS) are appropriate for studies of Hg contamination because the obtained Hg concentrations do not vary from analysis of recently collected wet blood samples (Lehner *et al.*, 2013; Perkins and Basu, 2018; Barst *et al.*, 2020; Santa-Rios *et al.*, 2020). Though this method is fairly underrepresented for contaminant analysis to date, and the use of Whatman filter paper deserve more investigations, it holds extraordinary potential for adding

value to DBS samples originally collected for other purposes (e.g., genetics, health, etc.). To use DBS to quantify total mercury, we extracted two standardized 4 mm punches from the card using a biopsy punch. Punches were performed in the center of the DBS. To evaluate the mass of dried blood contained in the card, we weighed standardized 4 mm punches extracted from different cards without any blood to determine an average expected mass of the Whatman filter sample paper prior to soaking with blood (5.62 ± 0.21 mg, 3.8% of variation, $n = 9$). Though our procedure closely follows prior Hg analysis using DBS, we diverged in our use of Whatman filter paper which has not been validated so far, thus we additionally analyzed the blank punches for Hg as a form of control (see “Mercury analysis” section below). To evaluate the weight of dried blood in the 4mm punch, we used the following equation:

$$\text{Mass of dried blood} = \text{Mass of 4mm punch DBS} - \text{Mean of 4 mm blank punches mass}$$

Mercury Analysis

For all samples, we determined Hg concentrations by direct measurement of at least two replicated 4 mm punches from each individual’s DBS using an atomic absorption spectrometer AMA-254 (Advanced Mercury Analyser-254; Altec®). We validated inter-replicate reproducibility for each sample against a relative standard deviation (RSD) threshold of < 10 %. We analyzed Certified Reference Material (CRM) DOLT-5 (Lobster hepatopancreas from National Research Council of Canada; certified Hg concentration: 0.44 ± 0.18 $\mu\text{g}\cdot\text{g}^{-1}$ dry weight (dw)) at the beginning and at the end of the analytical cycle, as well as every 10 samples, to validate measurement consistency across samples. The measured value for the CRM was 0.436 ± 0.005 $\mu\text{g}\cdot\text{g}^{-1}$ dw ($n = 39$), giving a recovery of 99.98 ± 1.22 %. We analyzed the blank punches for Hg concentration and found that they were always under the limit of

quantification of the method. Blanks were included at the beginning of analytical runs and the limit of quantification was 0.1 ng.g^{-1} . We express Hg concentrations in $\mu\text{g.g}^{-1} \text{ dw}$.

Statistical Analysis

We performed all statistical analyses in R v.3.6.1 (R Development Core Team 2013). Prior to all analysis, we checked data for normality and homogeneity of variances.

For each species individually, we used paired t-tests to compare SVL between adult males and females. Depending on the results, we then used either a paired t-test or ANCOVA to compare Hg concentrations between the sexes. Ultimately, we only performed this inter-sex comparison for *M. leptorhynchus* and *O. tetraspis* due to the low number of sampled adults for the other species.

We used one-way ANOVAs to compare SVL between the four species and to compare Hg concentrations within each species. We implemented a post-hoc Tukey's honest significant difference (HSD) test to evaluate the variation of Hg concentration between species.

Also, for each species individually, we used paired t-tests to compare SVL between sites and then used ANCOVAs with site and SVL as explanatory variables predicting Hg concentration differences between sites. Finally, we used linear regressions to evaluate the relationships between Hg concentration and the body size (SVL) for each species and site individually.

RESULTS

We detected Hg in all the samples, with the highest average concentration in *M. leptorhynchus* ($3.400 \pm 2.008 \mu\text{g.g}^{-1}$) followed by *O. tetraspis* ($1.655 \pm 1.148 \mu\text{g.g}^{-1}$), *C. niloticus* ($1.580 \pm 0.296 \mu\text{g.g}^{-1}$), and *C. suchus* ($0.528 \pm 0.243 \mu\text{g.g}^{-1}$) (Table 1).

For *M. leptorhynchus*, we did not detect any difference in adult body size (SVL) between the sexes in our sample (*t*-test: $t = 2.134$, $p = 0.056$); however, for *O. tetraspis* adult females were smaller than adult males in our sample (*t*-test: $t = -2.071$, $p = 0.046$). We did not find any significant difference in Hg concentrations between adult males and females for either species (*M. leptorhynchus*, *t*-test: $t = 1.112$, $p = 0.290$; *O. tetraspis*, ANCOVA: $F_{1-32} = 0.006$, $p = 0.939$).

We found significant differences in body size (SVL) variation across sample sites for all three species sampled across multiple sites: *M. leptorhynchus* (ANOVA: $F_{1-116} = 6.233$, $p < 0.001$), *O. tetraspis* (ANOVA: $F_{2-78} = 4.396$, $p = 0.016$), and *C. suchus* (ANOVA: $F_{5-51} = 25.06$, $p < 0.001$). Likewise, we found that Hg concentrations were significantly different between sites for all three species: *M. leptorhynchus* (ANCOVA: $F_{5-115} = 5.218$, $p < 0.001$), *C. suchus* (ANCOVA: $F_{5-50} = 7.042$, $p < 0.001$) and *O. tetraspis* (ANCOVA: $F_{2-77} = 3.2013$, $p = 0.050$) (Table 2). However, we only found a significant relationship between Hg concentration and body size (SVL) at 4 sites for *M. leptorhynchus* and 1 site for *O. tetraspis* (Table 3).

DISCUSSION

We found that, at least for the localities examined as part of this study, that these four African crocodile species were contaminated at varying concentrations with Hg. Worryingly, *M. leptorhynchus* ($3.400 \pm 2.008 \mu\text{g.g}^{-1}$), *O. tetraspis* ($1.655 \pm 1.148 \mu\text{g.g}^{-1}$), and *C. niloticus* ($1.580 \pm 0.296 \mu\text{g.g}^{-1}$) showed levels of Hg contamination that were particularly high compared to blood Hg concentrations amongst crocodylians analyzed to date – which had average Hg concentrations ranging from 0.07 to $1.38 \mu\text{g.g}^{-1}$ dw (Table 4). Interestingly, *M. leptorhynchus* in Gabon had the highest Hg blood concentration both on average and as a single individual, with one individual having as much as $11 \mu\text{g.g}^{-1}$, of any species of crocodylian measured to

date. Here we discuss these findings both within the context of our study systems and for crocodylians globally.

Of particular note, we found high variability in Hg concentrations both between species and sampling sites for all species (Table 2). Several studies have shown that Hg concentration in the blood of vertebrates may represent a relatively recent exposure to Hg related to diet (e.g., Gilmour *et al.*, 2019; Chételat *et al.*, 2020; Seco *et al.*, 2021). It is also correlated with erythrocyte longevity, which ranges from up to two months in birds to four months in mammals, but may be as long as seventeen months in some reptiles (reviewed in Rodnan *et al.*, 1957; Monteiro and Furness, 2001). Because crocodylians are high level consumers, if not apex predators, within their ecosystems, we would expect individual Hg contamination to be closely related to both diet and individual position within the trophic chain. Mercury contamination in the trophic chain depends to some extent on the geological background of the environment but can also be increased by human activities. In this regard, mining, fossil fuel extraction, biomass burning, and agriculture are among the anthropogenic activities driving the largest increases in environmental Hg (Obrist *et al.*, 2018; Crespo-Lopez *et al.*, 2021).

The simultaneous assessment of Hg concentrations in four crocodylians, two of which are syntopic, provides an interesting preliminary opportunity to discuss the role of species ecology in Hg bioaccumulation in this group. Generally speaking, higher concentrations of Hg are to be found in higher trophic level species predominantly consuming prey from aquatic ecosystems (Lavoie *et al.*, 2013; Eagle-Smith *et al.*, 2018). Though *M. leptorhynchus* and *O. tetraspis* occupy the same freshwater wetlands habitats throughout Gabon, including sites sampled and analyzed as part of this study, they have quite divergent feeding ecologies. Slender-snouted crocodiles are predominantly piscivorous, though consume a wide variety of

prey accessible from within the wetlands (Shirley *et al.*, 2018 and references therein). In contrast, dwarf crocodiles are largely feeding on terrestrial and aquatic invertebrates (Pauwels *et al.*, 2007; Shirley *et al.*, 2017). This divergence in prey base is likely one of the most important factors explaining the higher Hg concentration in *M. leptorhynchus* despite both species being exposed to similar environmental conditions. While most invertebrates, like millipedes or even crustaceans, typically have low Hg concentrations, fishing spiders of the genus *Dolomedes* are known to have higher Hg concentrations likely because of their predation on fish (Ortega-Rodriguez *et al.*, 2019). These species are abundant throughout the wetlands also occupied by *M. leptorhynchus* and *O. tetraspis* and are almost certainly widely consumed by juveniles of both species. This may explain, at least in part, the higher concentrations of Hg in *O. tetraspis* than would be expected on the basis of observations in the ecologically homologous *Paleosuchus* spp. (Lemaire *et al.*, 2022).

In addition to likely explaining key interspecific differences, trophic level and foraging habitats are extremely important when considering intraspecific variability in Hg contamination between different study sites, as demonstrated in multiple species (e.g., caiman, Lemaire *et al.*, 2021d; birds, Carravieri *et al.*, 2016; fish, Goutte *et al.*, 2015). Across our sampling sites, the individuals with the highest Hg concentrations were often sampled at sites thought to represent among the most pristine environments in Central Africa (e.g., Ivindo and Loango National Parks in Gabon), even if they are relatively adjacent sites where intense and prolonged oil extraction and varying degrees of subsistence agriculture occur. Higher Hg concentrations in “pristine” environments can often be explained by the relative complexity and intactness of the food webs compared to degraded sites – pristine environments can effectively have more links in the food webs for bioaccumulation of Hg prior to consumption by species like crocodylians resulting in a higher degree of Hg biomagnification (De Almeida

Rodrigues *et al.*, 2019). Alternatively, the proximity of these sites, especially Loango National Park (Gabon), to long-term oil extraction concessions may mean they are not as pristine as believed and our results suggest that more in-depth investigation of the spatio-temporal scale and extent to which petroleum concessions drive environmental contamination is warranted. More interestingly, *C. suchus* had the lowest Hg concentrations of all our studied species in spite of its distribution amongst significant artisanal small-scale gold mining (ASGM) activities throughout West Africa, including Ghana and Cote d'Ivoire – countries with high reporting of Hg contamination due to prolific ASGM (World Health Organization, 2015). Aquatic habitats throughout West Africa are generally highly degraded and, thus, *C. suchus* are largely persisting in highly disrupted food webs, potentially limiting the degree of bioaccumulation despite potentially higher levels of exposure. Unfortunately, our preliminary data only allow an extrapolative inference to be made about the relationship between Hg concentrations in crocodylians and human activities, and we recommend that this issue be studied in greater depth, especially in West and Central Africa, which provides an interesting natural laboratory.

Mercury is known to affect physiology, reproduction, and behavior of many wildlife species (Tan *et al.*, 2009; Whitney and Cristol, 2017; Evers, 2018). In seabirds, Hg contamination reduces mating success (Tartu *et al.*, 2013), and hatchling and fledging rates (Hg concentrations in blood of *Diomedea exulans* were $6.2 \pm 3.0 \mu\text{g.g}^{-1}$ dw for males and $10.7 \pm 0.5 \mu\text{g.g}^{-1}$ dw for females, Goutte *et al.*, 2014). In freshwater turtles, Hg exposure has been related to increasing embryonic mortality (Hg concentration in eggs of *Chelydra serpentina* was $3.0 \pm 0.19 \mu\text{g.g}^{-1}$, Hopkins *et al.*, 2013a) and to a reduction of reproductive females in the population (Hg concentration in the claws of adult *Emys orbicularis* was $2.21 \pm 0.06 \mu\text{g.g}^{-1}$ dw, Beau *et al.*, 2019). However, little is known to date about the impact of Hg contamination in crocodylians. Nilsen *et al.* (2017a) showed that Hg contamination negatively impacted body

condition in *Alligator mississippiensis*, which had blood Hg concentrations ranging from 0.26 to 6.90 $\mu\text{g}\cdot\text{g}^{-1}$ dw. For *Caiman crocodilus*, Marrugo-Negrete *et al.* (2019) showed erythrocytes DNA damage related to Hg exposure (average $0.23 \pm 0.08 \mu\text{g}\cdot\text{g}^{-1}$ dw) and Lemaire *et al.* (2021b) found that blood Hg concentrations ranging from 0.169 to 1.532 $\mu\text{g}\cdot\text{g}^{-1}$ dw were related to disruption of osmoregulation, hepatic function, and endocrine processes. These studies highlight that Hg contamination has toxic effects in crocodylians even at relatively low levels and raise concerns regarding the Hg concentrations obtained in the species studied here. While we did not specifically study the linkage between Hg contamination and deleterious effects in African crocodylians, our results lay the framework for tentative discussion.

For example, in several species, Hg is transferred to the egg during vitellogenesis, including in birds (Ackerman *et al.*, 2016), snakes (Chin *et al.*, 2013; Cusaac *et al.*, 2016), turtles (Hopkins *et al.*, 2013b), and crocodylians (Du Preez *et al.*, 2018; Nilsen *et al.*, 2020; Lemaire *et al.*, 2021c). It has been shown that the quantity of Hg transferred to the egg by the mother is directly related to the body burden of Hg in the reproductive female (Akearok *et al.*, 2010; Nilsen *et al.*, 2020). While it is thought this is a potential detoxification pathway for the female, leading to a reduction of the Hg body burden, it has consequences for the next generation. In birds, Hg exposure during embryonic development reduces reproductive capacity of the next generation (Paris *et al.*, 2018), while in turtles it reduces hatchling success (Hopkins *et al.*, 2013a), and alters the behavior of hatchling snakes (Chin *et al.*, 2013). In crocodylians, maternally transferred Hg resulted in reduced body size of hatchling smooth-fronted caiman (*Paleosuchus trigonatus*) (Lemaire *et al.*, 2021c). We did not find any variation in Hg blood concentration between males and females for the individuals assessed here. While these results are consistent with several other crocodylian Hg studies (Yanochko *et al.*, 1997; Rumbold *et al.*, 2002; Campbell *et al.*, 2010; Schneider *et al.*, 2012; Lemaire *et al.*, 2021d), the

biological explanation is as yet clear and it may simply be that we did not sample any reproductively active females. However, the effects of Hg exposure from maternal transfer during embryonic development are concentration-dependent, and the high Hg concentrations found in both sexually mature females and juveniles in our study species is worrying and highlights the potential long-term consequences for reproductive success and population growth and sustainability in these species. In particular, further assessment of Hg contamination in *M. leptorhynchus* and *O. tetraspis* in and around the oil extraction concessions of Gabon, including maternal transfer and reproductive stability implications, should be further assessed and monitored.

Crocodylians are widely consumed as bushmeat throughout Africa. In particular, dwarf crocodiles (*Osteolaemus* spp.) are among the most widely consumed bushmeat species throughout Cameroon (Wilcox and Nambu, 2007; Smolensky, 2015; Dipita *et al.*, 2022), Congo (Behra, 1987; Thorbjarnarson and Eaton, 2004; Eaton, 2006; Eaton *et al.*, 2010; Mbetete *et al.*, 2011; Shirley *et al.*, 2019), Democratic Republic of Congo (Batumike *et al.*, 2021), Equatorial Guinea (Juste *et al.*, 1995; Fa *et al.*, 2002), Gabon (Behra, 1987; Thorbjarnarson and Eaton, 2004; Eaton, 2006; Abernethy and Ndong Obiang, 2010), and Nigeria (Dore, 1996; Akani *et al.*, 1998), where tens to hundreds of thousands of individuals are hunted and consumed annually. These diminutive species are also regularly transported for consumption outside of their natural range, including to destinations as close as Bioko Island (Equatorial Guinea; Cronin *et al.*, 2015) and as far away as western Europe (e.g., Chaber *et al.*, 2019). Similarly, *M. leptorhynchus* is readily consumed locally and nationally throughout its entire range (e.g., Shirley *et al.*, 2018 and references therein). While humans globally are exposed to Hg in the environment, Hg ingestion through the consumption of various food items (e.g., tuna and other fish) is the main exposure risk for people. Mercury toxicity through food consumption

can result in significant neurological and behavioral effects in humans (Yang *et al.*, 2020). As a result, the World Health Organization (WHO) has released safety guidelines for the consumption of fish recommending consumption of foods only at $0.5 \mu\text{g}\cdot\text{g}^{-1}$ of Hg or below. Though we did not measure Hg concentrations in commonly consumed crocodylian body parts (e.g., muscle tissue), previous work on *Caiman crocodilus*, *Melanoscuchus niger* and *Alligator mississippiensis* has shown that Hg concentration is higher in muscles than in blood (Eggins *et al.*, 2015; Nilsen *et al.*, 2017b). Though few people strictly depend on crocodile bushmeat for their protein, the volume of harvest and consumption of these species is sufficient to justify a more in-depth study of their potential consequences on human health. Further, the demographics most likely to be consuming crocodylians tend to be fishermen, their families, and others living in predominantly fishing communities suggesting that they may be bioaccumulating Hg at rates not dissimilar to the crocodiles. As previously recommended, further investigation of Hg concentrations in crocodylian and human prey (e.g., fish) throughout the rural areas of West and Central Africa is warranted.

CONCLUSION

Mercury contamination is understudied in crocodylians, including the extent to which it is an issue across the diversity of the Crocodylia, the habitats and geographies they occupy, and the specific and long-term ramifications of such contamination. However, all attempted studies to date have found that crocodylians are impacted by Hg contamination, almost regardless of species and study site. Our results reinforce this issue, with data from three crocodylian species that have never been studied, including from localities ranging from those thought to be heavily impacted by humans (most of West Africa) to those thought to be among the most pristine, unimpacted environments in the world today (most of Gabon).

Worryingly, individuals sampled in these latter sites and habitats have the highest individual Hg contamination levels for any species of crocodylian anywhere in the world. In consideration of the conservation status of these species, as well as their importance as protein and economic resources to local communities throughout their distributions, our preliminary results underscore the importance of further studies on Hg contamination in African crocodylians. Such future work should not only continue to map patterns and process of Hg contamination in these bioindicator species, but dig deep into the bioaccumulation pathways, the long-term ramifications for population dynamics, and the potential health consequences for people consuming crocodylians and their shared prey (e.g., fish and other species).

ACKNOWLEDGEMENTS: All crocodylians were captured and all samples exported/imported under authorization of research and CITES permits issued by the relevant national authority(ies). We thank Gabon's Centre National de la Recherche Scientifique et Technologique (N°AR0024/09/MENESRI/CENAREST/CG/CST/CSAR and N°AR00 13/11), Agence National des Parcs Nationaux, and Ministère des Eaux et Forêts (CITES certificate numbers 01019, 01081, 01157, 01158, 01159). We thank Cote d'Ivoire's DGRSIT (N°304/MESRS/DGRIST/tm), Office Ivorien des Parcs et Reserves, and the Direction de la Faune et Ressources Cynégétiques (for CITES export permit). We thank the Egyptian Environmental Affairs Agency (CITES certificate number 457). We thank Ghana's Wildlife Division (for both research and CITES export permits). We thank the Uganda Wildlife Authority (permit number NS281 and CITES certificate number 002158). We additionally thank the US Fish and Wildlife Service for CITES import permits (certificate numbers 07US146577/9, 08US164015/9, 09US164015/9, 10US164015/9). Logistical support for sample collection was provided by Wildlife Conservation Society, World Wildlife Fund, Smithsonian Institution, Projet Protections des Gorilles, Protection des Grandes Singes de Moukalaba-Doudou, Fondation Liambissi, Aspinnall Foundation. We thank R. Starkey, M. Starkey, R. Calaque, B. Huijbrechts, B. Verhage, L. Korte, M. Butler, N. Bout, K. Kombila, R. Beville, P. and S. du Plessis, M. Marti, for their assistance throughout. Sampling for this work was possible with the generous financial support of a National Science Foundation Doctoral Dissertation Improvement Grant (DDIG; agreement no. 1010574), Riverbanks Zoo and Gardens Conservation Support Fund, US Fish and Wildlife Service Wildlife Without Borders programme (agreement no. 96200-1-G003), Conservation, Food and Health Foundation, Columbus Zoo, Idea Wild Foundation, St. Augustine Alligator Farm Zoological Park, IUCN/SSC Crocodile Specialist Group, AZA Crocodylian Advisory Group, Minnesota Zoo, Fresno Chaffee Zoo, San Diego Zoological Society, Mohamed bin Zayed Species Conservation Fund, Cleveland Metroparks Zoo, Oklahoma City Zoo, Aspinnall Foundation, Wildlife Conservation Society, WWF, the Institut de Recherche pour le Développement (IRD), and Colin Stevenson. The Rare

Species Conservatory Foundation served as a zero-overhead funding fiduciary for all grants other than the DDIG. Sampling and animal handling methods were reviewed and approved by IFAS ARC at the University of Florida (approval # 011- 09WEC). We are grateful to the Plateforme Analyses Élémentaires of the LIENSs for the access to the mercury analyser. Thanks are also due to the CPER (Contrat de Projet Etat-Région) and the FEDER (Fonds Européen de Développement Régional) for funding the AMA of LIENSs laboratory. Paco Bustamante is an honorary member of the IUF (Institut Universitaire de France).

REFERENCES

- Abernethy, K., Ndong Obiang, A.M. 2010. Bushmeat in Gabon. Ministry of Water and Forests, Government of Gabon.
- Ackerman, J.T., Eagles-Smith, C.A., Herzog, M.P., Hartman, C.A. 2016. Maternal transfer of contaminants in birds: Mercury and selenium concentration in parents and their eggs. *Environ. Pollut.* 210, 145-154.
- Akani, G.C., Luiselli, L., Angelici, F.M., Politano, E. 1998. Bushmeat and herpetofauna: notes on amphibians and reptiles traded in bush-meat market of local people in the Niger Delta (Port Harcourt, River state, Nigeria). *Antropozoologica* 27, 21-26.
- Akearok, J.A., Hebert, C.E., Braune, B.M., Mallory, M.L. 2010. Inter- and intraclutch variation in egg mercury levels in marine birds species from the Canadian Arctic. *Sci. Total Environ.* 408, 836-840.
- Almli, B., Mwase, M., Sivertsen, T., Musonda, M.M., Flåøyen, A. 2005. Hepatic and renal concentrations of 10 trace element in crocodiles (*Crocodylus niloticus*) in Kague and Luangwa rivers in Zambia. *Sci. Total Environ.* 337, 75-82.
- Barst, B.D., Wooller, M.J., O'Brien, D.M., Santa-Rios, A., Basu, N., Köck, G., Johnson, J.J., Muir, D.C.G. 2020. Dried blood spot sampling of Landlocked Arctic Char (*Salvelinus alpinus*) for estimating mercury exposure and stable carbon isotope fingerprinting of essential amino acids. *Environ. Toxicol. Chem.* 39(4), 893-903.
- Batumike, R., Imani, G., Urom, C., Cuni-Sanchez, A. 2021. Bushmeat hunting around Lomami National Park, Democratic Republic of the Congo. *Oryx* 55(3), 421-431.
- Beau, F., Bustamante, P., Michaud, B., Brischoux, F. 2019. Environmental causes and reproductive correlates of mercury contamination in European pond turtles (*Emys orbicularis*). *Environ. Res.* 172, 338-344.

- Behra, O. 1987. Etude de répartition des populations de crocodile du Congo, du Gabon et de la R.C.A. Parc Zoologique de Paris, Muséum National d'Histoire Naturelle: Paris.
- Benoit, J.M., Gilmour, C.C., Heyes, A., Mason, R.P., Miller, C.L. 2003. Geochemical and biological controls over methylmercury production and degradation in aquatic ecosystems. *Am. Chem. Soc. Symp. Ser.* 835, 262-297.
- Buenfil-Rojas, A.M., Álvarez-Legorreta, T., Cedeño-Vázquez, J.R. 2015. Metals and metallothioneins in Morelet's Crocodile (*Crocodylus moreletii*) from transboundary river between Mexico and Belize. *Arch. Environ. Contam. Toxicol.* 68, 265-273.
- Buenfil-Rojas, A.M., Alvarez-Legorreta, T., Cedeño-Vazquez, J.R., Rendón-von Osten, J., González-Jáuregui, M. 2020. Distribution of metals in tissues of captive and wild Morelet's crocodiles and the potential of metallothioneins in blood fractions as a biomarker of metal exposure. *Chemosphere* 244, 125551.
- Burger, J., Gochfeld, M., Rooney, A.A., Orlando, E.F., Woodward, A.R., Guillette Jr., L.J. 2000. Metals and metalloids in tissues of American alligators in three Florida lakes. *Arch. Environ. Contam. Toxicol.* 38, 501-508.
- Campbell, J.W., Waters, M.N., Tarter, A., Jackson, J. 2010. Heavy metal and selenium concentrations in liver tissue from wild American alligator (*Alligator mississippiensis*) livers near Charleston, South Carolina. *J. Wildl. Dis.* 46(4), 1234-1241.
- Campeau, G.C., Bartha, R. 1985. Sulphate reducing bacteria: principal methylators of mercury in anoxic estuarine sediment. *Appl. Environ. Microbiol.* 50, 498-502.
- Carravieri, A., Cherel, Y., Jaeger, A., Churlaud, C., Bustamante, P. 2016. Penguins as bioindicators of mercury contamination in the southern Indian Ocean: geographical and temporal trends. *Environ. Pollut.* 213, 195-205.
- Chaber, A.L., Gaubert, P., Green, H., Garigliany, M., Renault, V., Busoni, V., Dieudonné, M., Saegerman, C. 2019. Report on the illegal importation of meat, including bushmeat, seized at Zaventem airport-2017/2018. Dead or alive: towards a sustainable wildlife trade, One World One Health recommendations, Brussels, 3-4.
- Chen, C.Y., Driscoll, T., Eagles-Smith, C.A., Eckley, C.S., Gay, D.A., Hsu-Kim, H., Keane, S.E., Kirk, J.L., Mason, R.P., Obrist, D., Selin, H., Selin, N.E., Thompson, M.R. 2018. A critical time for mercury science to inform global policy. *Environ. Sci. Technol.* 52(17), 9556-9561.
- Cherkiss, M.S., Fling, H. E., Mazzotti, F. J., Rice, K. G. 2004 Counting and capturing crocodilians. *Wildlife Ecology and Conservation Department*, 1-14.

- Chételat, J., Ackerman, J.T., Eagles-Smith, C.A., Hebert, C. 2020. Methylmercury exposure in wildlife: A review of the ecological and physiological processes affecting contaminant concentrations and their interpretation. *Sci. Total Environ.* 711, 135117.
- Chin, S.Y., Willson, J.D., Cristol, D.A., Drewett, D.V.V., Hopkins, W.A. 2013. Altered behavior of neonatal northern watersnakes (*Nerodia sipedon*) exposed to maternally transferred mercury. *Environ. Pollut.* 176, 144-150.
- CSG, Crocodile Specialist Group. 1996. *Osteolaemus tetraspis*. *The IUCN Red List of Threatened Species*: e.T15635A4931429.
- Crespo-Lopez, M.E., Augusto-Oliveira, M., Lopes-Araújo, A., Santos-Sacramento, L., Takeda, P.Y., de Matos Macchi, B., do Nascimento, J.L.M., Maia, C.S.F., Lima, R.R., Arrifano, G.P. 2021. Mercury: What can we learn from the Amazon? *Environ. Int.* 146, 106223.
- Cronin, D.T., Woloszynek, S., Morra, W.A., Honarvar, S., Linder, J.M., Gonder, M.K., O'Connor, M.P., Hearn, G.W. 2015. Long-Term Urban Market Dynamics Reveal Increased Bushmeat Carcass Volume despite Economic Growth and Proactive Environmental Legislation on Bioko Island, Equatorial Guinea. *PLoS ONE* 10(7): e0134464.
- Cusaac, J.P.W., Kremer, V., Wright, R., Henry, C., Otter, R.R., Bailey, F.C. 2016. Effects of maternally-transferred methylmercury on stress physiology in Northern water snake (*Nerodia sipedon*) neonates. *Bull. Environ. Contam. Toxicol.* 96, 725-731.
- De Almeida Rodrigues, P., Ferrari, R.G., Dos Santos, L.N., Junior, C.A.C. 2019. Mercury in aquatic fauna contamination: a systematic review on its dynamics and potential health risks. *J. Environ. Sci.* 84, 205-218.
- Dipita, A.D., Missoup, A.D., Tindo, M., Gaubert, P. 2022. DNA-typing improves illegal wildlife trade surveys: Tracing the Cameroonian bushmeat trade. *Biol. Conserv.*, 269, 109552.
- Dore, M.P.O. 1996. Status of crocodiles in Nigeria. Crocodile Specialist Group Newsletter 15, 15-16.
- Du Preez, M., Govender, D., Kylin, H., Bouwman, H. 2018. Metallic elements in Nil Crocodile eggs from Kruger National Park, South Africa. *Ecotox. Environ. Saf.* 148, 930-941.
- Eagles-Smith, C.A., Silbergeld, E.K., Basu, N., Bustamante, P., Diaz-Barriga, F., Hopkins, W.A., Kidd, K.A., Nyland, J.F. 2018. Modulators of mercury risk to wildlife and humans in the context of rapid global change. *Ambio* 47, 170-197.

- Eaton, M.J. 2006. Ecology, conservation and management of the Central African dwarf crocodile (*Osteolaemus tetraspis*), a progress report. Pp. 84-95 in Crocodiles. Proceedings of the 18th Working Meeting of the IUCN-SSC Crocodile Specialist Group. IUCN: Gland.
- Eaton, M.J., Meyers, G.L., Kolokotronis, S.O., Leslie, M.S., Martin, A.P., Amato, G. 2010. Barcoding bushmeat: molecular identification of Central African and South American harvested vertebrates. *Conserv. Genet.* 11, 1389-1404.
- Eggs, S., Schneider, L., Krikowa, F., Vogt, R., Da Silveira, R., Maher, W. 2015. Mercury concentrations in different tissues of turtle and caiman species from the Rio Purus, Amazonas, Brazil. *Environ. Toxicol. Chem.* 34(12), 2271-2781.
- Elsley, R.M., Lance, V.A., Campbell, L. 1999. Mercury levels in Alligator meat in South Louisiana. *Bull. Environ. Contam. Toxicol.* 63, 598-603.
- Evers, D. 2018. The effects of methylmercury on wildlife: A comprehensive review and Approach for Interpretation. In: Dominick A. DellaSala, and Michael I. Goldstein (eds.) *The Encyclopedia of the Anthropocene*, vol. 5, p. 181-194. Oxford: Elsevier.
- Fa, J.E., Juste, J., Burn, R.W., Broad, G. 2002. Bushmeat consumption and preferences of two ethnic groups in Bioko Island, West Africa. *Hum. Ecol.* 30, 397-416.
- Fingerman, M., Devis, M., Reddy, P.S., Katyayani, R. 1996. Impact of heavy metal exposure on the nervous system and endocrine-mediated processes in crustaceans. *Zool. Stud.* 35(1), 1-8.
- Gilmour, M.E., Lavers, J.L., Lamborg, C., Chastel, O., Kania, S.A., Shaffer, S.A. 2019. Mercury as an indicator of foraging ecology but not the breeding hormone prolactin in seabirds. *Ecol. Indic.* 103, 248-259.
- Goutte, A., Barbraud, C., Meillère, A., Carravieri, A., Bustamante, P., Labadie, P., Budzinski, H., Delord, K., Cherel, Y., Weimerskirch, H., Chastel, O. 2014. Demographic consequences of heavy metals and persistent organic pollutants in a vulnerable long-lived bird, the wandering albatross. *Proc. R. Soc. B.* 281, 20133313.
- Goutte, A., Cherel, Y., Churlaud, C., Ponthus, J., Massé, G., Bustamante, P. 2015. Trace elements in Antarctic fish species and the influence of foraging habitats and dietary habits on mercury levels. *Sci. Total Environ.* 538, 743-749.
- Hekkala, E., Shirley, M.H., Amato, G., Austin, J.D., Charter, S., Thorbjarnarson, J., Vliet, K.A., Houck, M.L., DeSalle, R., Blum, M. 2011. An ancient icon reveals new mysteries: mummy

- DNA resurrects a cryptic species within the Nile crocodile. *Molecular Ecology* 20(20), 4195-4215.
- Hopkins, B.C., Hepner, M.J., Hopkins, W.A. 2013b. Non-destructive techniques for biomonitoring of spatial, temporal, and demographic patterns of mercury bioaccumulation and maternal transfer in turtles. *Environ. Pollut.* 177, 164-170.
- Hopkins, B.C., Willson, J.D., Hopkins, W.A. 2013a. Mercury exposure is associated with negative effects on turtle reproduction. *Environ. Sci. Technol.* 47, 2416-2422.
- Horai, S., Itai, T., Noguchi, T., Yasuda, Y., Adachi, H., Hyobu, Y., Riyadi, A.S., Boggs, A.S.P., Lowers, R., Guillette Jr., L.J., Tanabe, S. 2014. Concentrations of trace elements in American alligators (*Alligator mississippiensis*) from Florida, USA. *Chemosphere* 108, 159-167.
- Hsu-Kim, H., Eckley, C.S., Achá, D., Feng, X., Gilmour, C.C., Jonsson, S., Mitchell, C.P. 2018. Challenges and opportunities for managing aquatic mercury pollution in altered landscapes. *Ambio* 47(2), 141-169.
- Isberg, S., Combrink, X., Lippai, C., Balaguera-Reina, S.A. 2019. *Crocodylus niloticus*. The IUCN Red List of Threatened Species 201 9: e.T45433088A3010181.
- Juste, J., Fa, J.E., Del Val, J.P., Castroviejo, J. 1995. Market dynamics of bushmeat species in Equatorial Guinea. *J. Appl. Ecol.* 454-467.
- Lavoie, R.A., Jardine, T.D., Chumchal, M.M., Kidd, K.A., Campbell, L.M. 2013. Biomagnification of mercury in aquatic food webs: A worldwide meta-analysis. *Environ. Sci. Technol.* 47, 13385-13394.
- Lázaro, W.L., de Oliveira, R.F., dos Santos-Filho, M., da Silva, C.J., Malm, O., Ignácio, Á.R.A., Díez, S. 2015. Non-lethal sampling for mercury evaluation in crocodylians. *Chemosphere* 138, 25-32.
- Lehner, A.F., Rumbelha, W., Shlosberg, A., Stuart, K., Johnson, M., Domenech, R., Langner, H. 2013. Diagnostic analysis of veterinary dried blood spots for toxic heavy metals exposure. *J. Anal. Toxicol.* 37, 406-422.
- Lemaire, J., Brischoux, F., Marquis, O., Mangione, R., Bustamante, P. 2021a. Variation of total mercury concentrations in different tissues of three neotropical caimans: Implications for minimally invasive biomonitoring. *Arch. Environ. Contam. Toxicol.* 81, 15-24.
- Lemaire, J., Brischoux, F., Marquis, O., Mangione, R., Caut, S., Brault-Favrou, M., Churlaud, C., Bustamante, P. 2022 Relationships between stable isotopes and trace element

- concentrations in the crocodylian community of French Guiana. *Sci. Total Environ.* 837, 155846.
- Lemaire, J., Bustamante, P., Mangione, R., Marquis, O., Churlaud, C., Brault-Favrou, M., Parenteau, C., Brischoux, F. 2021b. Lead, mercury, and selenium alter physiological functions in wild caimans (*Caiman crocodilus*). *Environ. Pollut.* 286, 117549.
- Lemaire, J., Bustamante, P., Marquis, O., Caut, S., Brischoux, F. 2021d. Influence of sex, size and trophic level on blood Hg concentrations in Black caiman, *Melanosuchus niger* (Spix, 1825) in French Guiana. *Chemosphere* 262, 127819.
- Lemaire, J., Marquis, O., Bustamante, P., Mangione, R., Brischoux, F. 2021c. I got it from my mother: Inter-nest variation of mercury concentration in neonate Smooth-fronted Caiman (*Paleosuchus trigonatus*) suggest maternal transfer and possible phenotypical effects. *Environ. Res.* 194, 110494.
- Marrugo-Negrete, J., Durango-Hernández, J., Calao-Ramos, C., Urango-Cárdenas, I., Díez, S. 2019. Mercury levels and genotoxic effect in caimans from tropical ecosystems impacted by gold mining. *Sci. Total Environ.* 664, 899-907.
- Mbete, R.A., Banga-Mboko, H., Racey, P., Mfoukou-Ntsakala, A., Nganga, I., Vermeulen, C., Doucet, J.L., Hornick, J.L., Leroy, P. 2011. Household bushmeat consumption in Brazzaville, the Republic of the Congo. *Trop. Conserv. Sci.* 4(2), 187-202.
- Monteiro, L.R., Furness, R.W. 2001. Kinetics, dose-response, and excretion of methylmercury in free-living cory's shearwaters. *Environ. Sci. Technol.* 35, 739-746.
- Nilsen, F.M., Dorsey, J.E., Lowers, R.H., Guillette Jr., L.J., Long, S.E., Bowden, J.A., Schock, T.B. 2017a. Evaluating mercury concentrations and body condition in American alligators (*Alligator mississippiensis*) at Merritt Island National Wildlife Refuge (MINWR), Florida. *Sci. Total Environ.* 607-608, 1056-1064.
- Nilsen, F.M., Kassim, B.L., Delaney, J.P., Lange, T.R., Brunell, A.M., Guillette Jr., L.J., Long, S.E., Schock, T.B. 2017b. Trace element biodistribution in the American alligator (*Alligator mississippiensis*). *Chemosphere* 181, 343-351.
- Nilsen, F.M., Rainwater, T.R., Wilkinson, P.M., Brunell, A.M., Lowers, R.H., Bowden, J.A., Guillette, L.J., Long, S.E., Schock, T.B. 2020. Examining maternal and environmental transfer of mercury into American alligator eggs. *Ecotox. Environ. Saf.* 189, 110057.

- Obrist, D., Kirk, J.L., Zhang, L., Sunderland, E.M., Jiskra, M., Selin, N.E. 2018. A review of global environmental mercury processes in response to human and natural perturbations: Changes of emissions, climate, and land use. *Ambio* 47, 116-140.
- Ortega-Rodriguez, C., Chumchal, M.M., Drenner, R.W., Kennedy, J.H., Nowlin, W.H., Barst, B.D., Polk, D.K., Hall, M.N., Williams, E.B., Lauck, K.C., Santa-Rios, A., Basu, N. 2019. Relationship Between Methylmercury Contamination and Proportion of Aquatic and Terrestrial Prey in Diets of Shoreline Spiders. *Environ. Toxicol.* 38(11), 2503-2508.
- Paris, O.J., Swaddle, J.P., Cristol, D.A. 2018. Exposure to dietary methyl-mercury solely during embryonic and juvenile development halves subsequent reproductive success in adult Zebra Finches. *Environ. Sci. Technol.* 52(5), 3117-3124.
- Pauwels, O.S., Barr, B., Sanchez, M.L., Burger, M. 2007. Diet records for the dwarf crocodile (*Osteolaemus tetraspis tetraspis*) in Rabi Oil Fields and Loango National Park, Southwestern Gabon. *Hamadryad*, 31(2), 258-264.
- Perkins, M., Basu, N. 2018. Dried blood spots for estimating mercury exposure in birds. *Environ. Pollut.* 236, 236-246.
- Pirrone, N., Cinnirella, S., Feng, X., Finkelman, R.B., Friedli, H.R., Leaner, J., Mason, R., Mukherjee, A.B., Stracher, G.B., Streets, D.G., Telmer, K. 2010. Global mercury emissions to the atmosphere from anthropogenic and natural sources. *Atmos. Chem. Phys.* 10, 5951-5964.
- Rainwater, T.R., Adair, B.M., Platt, S.G., Anderson, T.A., Cobb, G.P., McMurry, S.T. 2002. Mercury in Morelet's crocodile eggs from northern Belize. *Arch. Environ. Contam. Toxicol.* 42, 319-324.
- Rodnan, G.P., Ebaugh, F.G., Fox, M.R.S., Chambers, D.M. 1957. The life span of the red blood cell and the red blood cell volume in the chicken, pigeon and duck as estimated by the use of $\text{Na}_2\text{Cr}^{51}\text{O}_4$: with observation on red cell turnover rate in the mammal, bird and reptile. *Blood* 12(4), 355-366.
- Romero-Calderón, A.G., Alvarez-Legorreta, T., Rendón von Osten, J., González-Jáuregui, M., Cedeño-Vázquez, J.R. 2022. Stress responses in captive *Crocodylus moreletii* associated with metal exposure. *Environ. Pollut.* 308, 119685.
- Ross, J.P. 1998. Crocodiles: status survey and conservation action plan.

- Rumbold, D.G., Fink, L.E., Laine, K.A., Niemczyk, S.L., Chandrasekhar, T., Wankel, S.D., Kendall, C. 2002. Levels of mercury in alligators (*Alligator mississippiensis*) collected along a transect through the Florida Everglades. *Sci. Total Environ.* 297, 239-252.
- Santa-Rios, A., Barst, B.D., Basu, N. 2020. Mercury speciation in whole blood and dried blood spots from capillary and venous sources. *Anal. Chem.* 92, 3605-3612.
- Scheuhammer, A., Braune, B., Chan, H.M., Frouin, H., Krey, A., Letcher, R., Loseto, L., Noël, M., Ostertag, S., Ross, P., Wayland, M. 2015. Recent progress on our understanding of the biological effects of mercury in fish and wildlife in the Canadian Arctic. *Sci. Total Environ.* 509-510, 91-103.
- Schneider, L., Eggins, S., Maher, W., Vogt, R.C., Krikowa, F., Kinsley, L., Eggins, S.M., Da Silveira, R. 2015. An evaluation of the use of reptile dermal scutes as a non-invasive method to monitor mercury concentrations in the environment. *Chemosphere* 119, 163-170.
- Schneider, L., Peleja, R.P., Kluczkowski Jr., A., Freire, G.M., Marioni, B., Vogt, R.C., Da Silveira, R. 2012. Mercury concentration in the Spectacled Caiman and Black Caiman (Alligatoridae) of the Amazon: Implications for Human Health. *Arch. Environ. Contam. Toxicol.* 63, 270-279.
- Seco, J., Aparício, S., Brierley, A.S., Bustamante, P., Ceia, F.R., Coelho, J.P., Philips, R.A., Saunders, R.A., Fielding, S., Gregory, S., Matias, R., Pardal, M.A., Pereira, E., Stowasser, G., Tarling, G., Xavier, J.C. 2021. Mercury biomagnification in a Southern Ocean food web. *Environ. Pollut.* 275, 116620.
- Shirley, M.H. 2014. *Mecistops cataphractus*. *The IUCN Red List of Threatened Species*: e.T5660A3044332.
- Shirley, M.H., Burtner, B., Oslisly, R., Sebag, D., Testa, O. 2017. Diet and body condition of cave-dwelling dwarf crocodiles (*Osteolaemus tetraspis*, Cope 1861) in Gabon. *African Journal of Ecology* 55, 411 – 422.
- Shirley, M.H., Carr, A.N., Nestler, J.H., Vliet, K.A., Brochu, C.A. 2018. Systematic revision of the living African Slender-snouted Crocodiles (*Mecistops* Gray 1844). *Zootaxa* 4504(2): 151-193.
- Shirley, M.H., Evans, B., Madzoke, B., Bondeko, G., Mobongo, R. 2019. Sustainable Management of the Lac Tele Dwarf Crocodile Fishery: Laying the Foundation for Future Action. Report Presented to Wildlife Conservation Society – Congo. 29pp.

- Shirley, M.H., Villanova, V., Vliet, K.A., Austin, J.D. 2015. Genetic barcoding facilitates captive management of three cryptic African crocodile species complexes. *Animal Conservation* 18(4), 322 - 330.
- Smolensky, N.L 2015. Co-occurring cryptic species pose challenges for conservation: a case study of the African dwarf crocodile (*Osteolaemus* spp.) in Cameroon. *Oryx* 49(4), 584-590.
- Somaweera, R., Nifong, J., Rosenblatt, A., Brien, M.L., Combrink, X., Eelsey, R.M., Grigg, G., Magnusson, W.E., Mazzotti, F.J., Pearcy, A., Platt, S.G., Shirley, M.H., Tellez, M., van der Ploeg, J., Webb, G., Whitaker, R., Webber, B.L. 2020. The ecological importance of crocodylians: towards evidence-based justification for their conservation. *Biol. Rev.* 95, 936-959.
- Tan, S.W., Meiller, J.C., Mahaffey, K.R. 2009. The endocrine effects of mercury in humans and wildlife. *Crit. Rev. Toxicol.* 39(3), 228-269.
- Tartu, S., Goutte, A., Bustamante, P., Angelier, F., Moe, B., Clément-Chastel, C., Bech, C., Gabrielsen, G.W., Bustnes, J.O., Chastel, O. 2013. To breed or not to breed: endocrine response to mercury contamination by an Arctic seabird. *Biol. Lett.* 9: 20130317.
- Thirion, F., Tellez, M., Van Damme, R., Bervoets, L. 2022. Trace element concentrations in caudal scutes from *Crocodylus moreletii* and *Crocodylus acutus* in Belize in relation to biological variables and land use. *Ecotox. Environ. Safe.* 231, 113164.
- Thorbjarnarson, J. 1992. *Crocodiles: An Action Plan for their Conservation*. H. Messel, F.W., King, J.P. Ross (eds.). IUCN, Switzerland, 136pp.
- Thorbjarnarson, J.B., Eaton, M.J. 2004. Preliminary examination of crocodile bushmeat issues in the Republic of Congo and Gabon. Pp. 236-247 in *Crocodiles*. Proceedings of the 17th Working Meeting of the IUCN-SSC Crocodile Specialist Group. IUCN: Gland.
- Trillanes, C.E., Pérez-Jiménez, J.C., Rosiles-Martínez, R., González-Jáuregui, M. 2014. Metals in the Caudal Scutes of Morelet's Crocodile (*Crocodylus moreletii*) from the Southern Gulf of Mexico. *Bull. Environ. Contam. Toxicol.* 93, 423-428.
- UNEP, UN Environment, 2019. Global Mercury assessment 2018. UN Environment Programme, Chemicals and Health Branch Geneva, Switzerland.
- Walsh, B. 1987 Crocodile capture methods used in the Northern Territory of Australia. In *Wildlife Management: Crocodiles and Alligators* (eds G. J. W. Webb S. C. Manolis & P. J. Whitehead), pp. 249–252. Chipping Norton, New South Wales, Australia. 522 pp.: Surrey Beatty and Sons Pty Ltd.

- Webb, G. J. W., Buckworth, R. & Manolis, S. C. 1983 *Crocodylus johnstoni* in the McKinlay River, N.T. VI. Nesting biology. *Australian Wildlife Research* 10, 607–637.
- Whitney, M.C., Cristol, D.A. 2017. Impacts of sublethal mercury exposure on birds: A detailed review. In: de Voogt P. (eds) *Reviews of Environmental Contamination and Toxicology* volume 244. Springer, Cham.
- Wilcox, A.S., Nambu, D.M. 2007. Wildlife hunting practices and bushmeat dynamics of the Banyangi and Mbo people of Southwestern Cameroon. *Biol. Conserv.* 134(2), 251-261.
- World Health Organization. 2015. Chemicals of public health concern in the African Region and their management: Regional Assessment Report.
- Yang, L., Zhang, Y., Wang, F., Luo, Z., Guo, S., Strähle, U. 2020. Toxicity of mercury: Molecular evidence. *Chemosphere* 245, 125586.
- Yanochko, G.M., Jagoe, C.H., Brisbin Jr., I.L. 1997. Tissue mercury concentrations in Alligators (*Alligator mississippiensis*) from Florida Everglades and the Savannah River site, South Carolina. *Environ. Contam. Toxicol.* 32, 323-328.
- Yoshimura, A., Suemasu, K., Veiga, M.M. 2021. Estimation of mercury losses and gold production by artisanal and small-scale gold mining (ASGM). *J. Sustain. Metall.* 7, 1045-1059.




Figure 1. Geographic location of the study sites in Africa. The inset maps show more detailed sampling locations in Gabon (a), Cote d'Ivoire (b), and Ghana (c) relative to protected areas (light green polygons) in these countries. The purple dashed line in the Gabon inset (a) indicates coastal area where most of the country's on and offshore petroleum concessions are concentrated. Similarly, the red dashed line in the Ghana (c) and Cote d'Ivoire (b) inset indicates the general area of the major gold deposits where much of the artisanal gold extraction happens.

Table 1. Biometric data (cm) and Hg concentrations ($\mu\text{g}\cdot\text{g}^{-1}$ dw) in the blood of four African crocodylian species. Hg values marked with the same letter are not statistically different (Tukey's honesty significant difference test, $p < 0.05$).

Species	n	SVL (cm)		Hg concentration ($\mu\text{g}\cdot\text{g}^{-1}$)	
		Mean \pm SD	Min - Max	Mean \pm SD	Min - Max
<i>Mecistops leptorhynchus</i>	124	48.8 \pm 33.4	17.0 - 152.1	3.400 \pm 2.008 ^a	0.570 - 11.230
<i>Osteolaemus tetraspis</i>	81	48.6 \pm 18.5	20.8 - 92.6	1.655 \pm 1.148 ^b	0.390 - 5.180
<i>Crocodylus suchus</i>	56	50.1 \pm 31.9	20.0 - 139.5	0.528 \pm 0.243 ^c	0.207 - 1.291
<i>Crocodylus niloticus</i>	9	42.4 \pm 17.9	15.1 - 74.6	1.580 \pm 0.296 ^b	1.230 - 2.180

Table 2. Biometric data (cm) and Hg concentrations ($\mu\text{g}\cdot\text{g}^{-1}$ dw) in the blood of four African crocodylian species at each sampling site. All *M. leptorhynchus* and *O. tetraspis* individuals were sampled in Gabon. In the table, NP = national park. For Site Degradation, we qualitatively characterized sites based on our observations whilst in the field sampling as P = Pristine (e.g., habitat mostly or entirely intact, very little contemporary human disturbance), MD – Mildly Degraded (up to moderate habitat loss, some contemporary human presence), or HD = Heavily Degraded (considerable habitat loss, humans extensive in system contemporarily). For Proximity to Hg, we identified the most significant likely contributors of Hg to the habitat, where P = Petroleum, A = Industrial or Intensive Agriculture, and G = Gold mining, and estimated its relative proximity to the site, where 1 = < 50 km, 2 = 50 – 150 km, 3 = 150 – 300 km, and 4 = > 300 km. For both of these columns, the characterizations are applicable to the time of sampling, not after.

Species	Locality	Site Degradation	Proximity to Hg	N	SVL (cm)		Hg ($\mu\text{g}\cdot\text{g}^{-1}$)	
					Mean \pm SD	Min - Max	Mean \pm SD	Min - Max
<i>Mecistops leptorhynchus</i>	Ogooué River (middle stretch, Gabon)	MD	P3, A2	24	33.24 \pm 9.22	18.1 - 52.1	3.182 \pm 1.629	0.791 - 7.401
	Bongo River (Loango NP, Gabon)	P	P1	16	43.87 \pm 21.14	21.7 - 101.0	2.793 \pm 1.318	0.566 - 5.891
	Lac Divangui (Gabon)	MD	P1	1	26.2	-	1.823	-
	Lac Evaro (Gabon)	MD	P3, A2	2	46.55 \pm 5.02	43.0 - 50.1	3.064 \pm 0.232	2.900 - 3.227
	Akaka (Loango NP, Gabon)	P	P1	40	71.14 \pm 41.35	19.1 - 152.1	3.029 \pm 1.984	0.822 - 11.228
	Dji Dji River (Ivindo NP, Gabon)	P	G2	19	44.72 \pm 35.31	17.0 - 131.4	4.786 \pm 2.293	2.025 - 10.162
	Mpassa River (Plateau Bateke NP, Gabon)	P	-	16	26.69 \pm 11.82	17.1 - 61.0	3.919 \pm 2.440	0.928 - 8.608
	Nyanga River (Moukalaba-Doudou NP, Gabon)	MD	P2, A1	6	51.37 \pm 23.14	31.0 - 96.7	2.958 \pm 1.678	0.739 - 4.993

<i>Osteolaemus tetraspis</i>								
	Abanda (Fernan Vaz Lagoon area, Gabon)	MD	P1	32	43.1 ± 21.8	20.8 - 92.6	1.248 ± 0.620	0.535 - 3.014
	Akaka (Loango NP, Gabon)	P	P1	32	53.5 ± 15.3	21.7 - 81.8	2.279 ± 1.503	0.390 - 5.180
	Lope River (Lope NP, Gabon)	P	-	17	49.6 ± 15.1	25.1 - 76.9	1.245 ± 0.430	0.450 - 2.328
<i>Crocodylus suchus</i>								
	Abi Lagoon (Cote d'Ivoire)	HD	A1, G3	10	42.8 ± 32.0	20.0 - 100.2	0.578 ± 0.344	0.229 - 1.291
	Amansuri Wetlands (Ghana)	MD	A1, G2	6	24.5 ± 1.2	22.4 - 25.7	0.890 ± 0.164	0.667 - 1.106
	Bui National Park (Ghana)	MD	A1, G2	6	28.9 ± 11.4	22.7 - 52.0	0.397 ± 0.082	0.285 - 0.495
	Kidepo Valley NP (Uganda)	P	P4	19	85.3 ± 21.9	47.0 - 139.5	0.508 ± 0.172	0.253 - 1.002
	San Pedro River Dam (Cote d'Ivoire)	HD	A1, G2	8	26.3 ± 2.4	22.0 - 28.9	0.519 ± 0.186	0.289 - 0.815
	Tiatia Village (Ghana)	HD	A1, G2	7	32.7 ± 7.1	24.0 - 43.9	0.327 ± 0.080	0.207 - 0.404
<i>Crocodylus niloticus</i>								
	Lake Nasser (Egypt / Sudan)	P	-	9	42.4 ± 17.9	15.1 - 74.6	1.580 ± 0.296	1.230 - 2.180

Table 3. Summary statistics of linear regressions relating Hg concentration ($\mu\text{g.g}^{-1}$ dw) to Snout-Vent-Length (SVL in cm) for each sampling location for three African crocodylian species. All *M. leptorhynchus* and *O. tetraspis* individuals were sampled in Gabon. In the table, NP = national park.

Sites	n	R ²	p-value
<i>Mecistops leptorhynchus</i>			
Ogooué River (middle stretch)	24	0.405	<0.001
Bongo River (Loango NP)	16	0.105	0.222
Lac Divangui	1	-	-
Lac Evaro	2	-	-
Akaka (Loango NP)	40	0.352	<0.001
Dji Dji River (Ivindo NP)	19	0.297	0.016
Mpassa River (Plateau Bateke NP)	16	0.593	<0.001
Nyanga River (Moukalaba-Doudou NP)	6	0.190	0.387
<i>Osteolaemus tetraspis</i>			
Abanda (Fernan Vaz Lagoon area)	32	0.134	0.040
Akaka (Loango NP)	32	0.084	0.109
Lope River (Lope NP)	17	0.075	0.288
<i>Crocodilus suchus</i>			
Abi Lagoon (Cote d'Ivoire)	10	0.016	0.726
Amansuri Wetlands (Ghana)	6	0.034	0.728
Bui National Park (Ghana)	6	0.193	0.384
Kidepo Valley NP (Uganda)	19	0.091	0.211
San Pedro River Dam (Cote d'Ivoire)	8	0.145	0.351
Tiatia Village (Ghana)	7	0.014	0.804

Table 4. Review of Hg concentration ($\mu\text{g}\cdot\text{g}^{-1}$ dw) reported in blood of crocodylians. TL stands for Total Length and SVL for Snout-Vent-Length (cm). For *Caiman crocodilus* sites in Colombia, ^a: pristine area; ^b: gold mining area. ^ϕStudies did not report SVL values, only TL; using standard allometric relationships, the *Caiman crocodilus* individuals were ± 31 cm SVL and *Melanosuchus niger* were ± 78 cm SVL – both generally comparable to other species in the table. *Original data reported in wet weight, transformed to dry weight using a conversion factor of 3.6 following a mean moisture content of 72% in the blood of caimans from Lemaire *et al.* (2021a). These transformed values should be viewed cautiously until the universality of the conversion factor can be validated.

Species	Location	N	Tissue	Length (cm)		Hg ($\mu\text{g}\cdot\text{g}^{-1}$ dw)		Reference
				Mean \pm SD	Min-Max	Mean \pm SD	Min-Max	
<i>Mecistops leptorhynchus</i>	Gabon	124	Whole blood	48.8 \pm 33.4 (SVL)	17 - 152.1	3.400 \pm 2.008	0.57 - 11.23	Present study
<i>Osteolaemus tetraspis</i>	Gabon	81	Whole Blood	48.6 \pm 18.5 (SVL)	20.8 - 92.6	1.655 \pm 1.148	0.390 - 5.180	Present study
<i>Crocodylus suchus</i>	Cote d'Ivoire	18	Whole Blood	35.5 \pm 24.8 (SVL)	20.0 - 100.2	0.552 \pm 0.278	0.229 - 1.291	Present study
<i>Crocodylus suchus</i>	Ghana	19	Whole Blood	28.9 \pm 8.1 (SVL)	22.4 - 52.0	0.527 \pm 0.277	0.207 - 1.106	Present study
<i>Crocodylus suchus</i>	Uganda	19	Whole Blood	85.3 \pm 21.9 (SVL)	47.0 - 139.5	0.508 \pm 0.172	0.253 - 1.002	Present study
<i>Crocodylus niloticus</i>	Egypt	9	Whole Blood	42.4 \pm 17.9 (SVL)	15.1 - 74.6	1.580 \pm 0.296	1.230 - 2.180	Present study
<i>Paleosuchus trigonatus</i>	French Guiana	24	Whole Blood	33.3 \pm 19.8 (SVL)	10.9 - 81	0.300 \pm 0.178	0.032 - 0.738	Lemaire <i>et al.</i> , 2021a
<i>Paleosuchus palpebrosus</i>	French Guiana	7	Whole blood	38.1 \pm 15.3 (SVL)	16.5 - 62.0	1.376 \pm 0.986	0.540 - 3.415	Lemaire <i>et al.</i> , 2021a
<i>Caiman crocodilus</i>	French Guiana	40	Whole blood	32.7 \pm 13.6 (SVL)	14.5 - 103.0	0.605 \pm 0.380	0.089 - 1.532	Lemaire <i>et al.</i> , 2021a
<i>Caiman crocodilus</i>	Colombia ^a	23	Whole blood	57.5 \pm 6.8 (TL) ^ϕ	-	0.05 \pm 0.02*	-	Marrugo-Negrete <i>et al.</i> , 2019
<i>Caiman crocodilus</i>	Colombia ^b	23	Whole blood	57.2 \pm 3.5 (TL) ^ϕ	-	0.23 \pm 0.08*	-	Marrugo-Negrete <i>et al.</i> , 2019
<i>Melanosuchus niger</i>	French Guiana	72	Whole blood	143.2 \pm 61.3 (TL) ^ϕ	46 - 326	1.284 \pm 0.672	0.300 - 3.410	Lemaire <i>et al.</i> , 2021d
<i>Alligator mississippiensis</i>	USA / Florida	37	Whole blood	92.1 \pm 31.6 (SVL)	43.9 - 153.5	0.694*	0.280 - 6.900	Nilsen <i>et al.</i> , 2017b