



HAL
open science

A roadmap for Ocean Negative Carbon Emission eco-engineering in sea-farming fields

Nianzhi Jiao, Chenba Zhu, Jihua Liu, Tingwei Luo, Mindong Bai, Zhiming Yu,
Quanrui Chen, Buki Rinkevich, Markus Weinbauer, Helmuth Thomas, et al.

► **To cite this version:**

Nianzhi Jiao, Chenba Zhu, Jihua Liu, Tingwei Luo, Mindong Bai, et al.. A roadmap for Ocean Negative Carbon Emission eco-engineering in sea-farming fields. *The Innovation Geoscience*, 2023, 1 (2), pp.100029. 10.59717/j.xinn-geo.2023.100029 . hal-04660302

HAL Id: hal-04660302

<https://hal.science/hal-04660302>

Submitted on 24 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



A roadmap for Ocean Negative Carbon Emission eco-engineering in sea-farming fields

Nianzhi Jiao,^{1,2,*} Chenba Zhu,^{1,2} Jihua Liu,^{2,3} Tingwei Luo,^{1,2} Mindong Bai,⁴ Zhiming Yu,⁵ Quanrui Chen,^{1,2} Buki Rinkevich,^{2,6} Markus Weinbauer,^{2,7} Helmuth Thomas,^{2,8} Mar Fernández-Méndez,⁹ Celeste López-Abbate,^{2,10} Camila Negrão Signori,^{2,11} Ramaiah Nagappa,^{2,12} Michal Koblížek,^{2,13} Hermann Kaartokallio,^{2,14} Jung-Ho Hyun,^{2,15} Fanglue Jiao,^{2,16} Feng Chen,^{2,16} and Wei-Jun Cai^{2,17}

¹Carbon Neutral Innovation Research Center and Fujian Key Laboratory of Marine Carbon Sequestration, Xiamen University, Xiamen 361005, China

²Global Ocean Negative Carbon Emissions Program Alliance

³Institute of Marine Science and Technology, Shandong University, Qingdao 266237, China

⁴School of Environmental Science and Engineering, Tianjin University, Tianjin 300072, China

⁵CAS Key Laboratory of Marine Ecology and Environmental Sciences, Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China

⁶Israel Oceanographic & Limnological Research, National Institute of Oceanography, Haifa Tel Shikmona 3109701, Israel

⁷Laboratoire d'Océanographie de Villefranche, Sorbonne Université-CNRS UMR7093, Institut de la Mer de Villefranche, Villefranche BP28, France

⁸Institute of Carbon Cycles, Helmholtz-Zentrum Hereon, Geesthacht 21502, Germany

⁹Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven 27570, Germany

¹⁰Instituto Argentino de Oceanografía, Consejo Nacional de Investigaciones Científicas y Técnicas, Bahía Blanca 8000, Argentina

¹¹Universidade de São Paulo, Praça do Oceanográfico, São Paulo-SP, Brazil

¹²Biological Oceanography Division, CSIR-National Institute of Oceanography, Dona Paula, Goa, 403004, India

¹³Institute of Microbiology of the Czech Academy of Sciences, Novohradská 237, Třeboň 37981, Czech Republic

¹⁴Marine Research Centre, Finnish Environment Institute, Helsinki 00790, Finland

¹⁵Department of Marine Science and Convergence Technology, Hanyang University, Gyeonggi-do 15588, South Korea

¹⁶Institute of Marine and Environmental Technology, University of Maryland Center for Environmental Science, Baltimore, MD 21613, USA

¹⁷School of Marine Science and Policy, University of Delaware, Newark, DE 19716, USA

*Correspondence: jjiao@xmu.edu.cn (N.J.)

Received: August 8, 2023; Accepted: September 13, 2023; Published Online: September 14, 2023; <https://doi.org/10.59717/j.xinn-geo.2023.100029>

© 2023 The Author(s). This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Citation: Jiao N., Zhu C., Liu J., et al., (2023). A roadmap for Ocean Negative Carbon Emission eco-engineering in sea-farming fields. *The Innovation Geoscience* **1**(2), 100029.

Carbon neutralization has become a significant, inevitable, and urgent strategy for both adaptation and mitigation of global warming caused by anthropogenic CO₂ emissions, and its environmental consequences such as

ocean acidification. However, the reduction of anthropogenic CO₂ emissions often conflicts with economic development. In contrast, environmentally-friendly negative carbon emissions can be a way of killing two birds with one




Figure 1. An illustration of the BCMS ecoengineering approaches POC: Particulate Organic Carbon, RDOC: Refractory Dissolved Organic Carbon, N,P: Nitrogen and phosphorus.

stone, capturing carbon dioxide and ensuring economic development, and therefore become imperative to achieve carbon-neutral goals.

The ocean, as the largest active carbon reservoir, has great potential for

negative carbon emissions. However, since the ocean has stored 93% of the world's CO₂, further negative carbon emissions remain a formidable challenge. On top of that, both the approaches and implementation fields should

be appropriate and must comply with the UN Conventions on the Law of the Sea, and the International Maritime Organization London Convention and Protocol.

Sea-farming, being a low cost and high productivity manner, is a well-developed marine industry, and is further promoted nowadays by the UN given that the world food needs are under pressure of 8 billion world population. However, rapid and over-intensive sea-farming developments entail a series of environmental risks and problems, such as eutrophication, deoxygenation, acidification in the bottom water, and potential harmful microalgal blooms in the surface water.

Under such circumstances, the Global Ocean Negative Carbon Emissions (ONCE) program (<https://www.global-once.org>), endorsed by the United Nations Decade of Ocean Science for Sustainable Development, has proposed a comprehensive eco-engineering approach, which integrates the well-known biological carbon pump (BCP) and the carbonate counter pump (CCP), and the newly established microbial carbon pump (MCP),¹ as well as the “unmanageable” solubility carbon pump (SCP) into a comprehensive “BCP-CCP-MCP-SCP (BCMS)” approach, just like the Business-Continuity-Management-System (BCMS). We here propose a road map for the BCMS approach. As illustrated in Figure 1, this ONCE roadmap is based on systematic measures, starting with solar energy-driven artificial upwelling as an internal ecosystem regulation solution, followed by further scientific intervention measures to simultaneously increase carbon sequestration capabilities and mitigate potential environmental impacts (Figure 1).

BCMS STEP 1, Intelligent artificial upwelling to enhance carbon fixation

Artificial upwelling can bring up nutrients from deep water to the surface water for enhancement of photosynthesis, and the consequent downwelling can drag high oxygen surface water down to the subsurface for mitigation of hypoxia and acidification. Ideally, solar energy powered artificial upwelling would enhance nutrient supply during the daytime fitting the photosynthesis rhythm and the growth of primary producers, and would stop during the night to avoid excessive dissolved inorganic carbon (DIC).

Artificial upwelling should be implemented in legal sites. Grounded in the United Nations Convention on the Law of the Sea and the London Convention, seaweeds culture fields would be the right place for BCMS practice. An intelligent operation of the artificial upwelling can maintain a nutrient-DIC balance, ensuring optimal and efficient seaweed growth without release of CO₂, which can be achieved through solar radiation based automatic adjustment of upwelling flux and rhythm.

BCMS STEP 2, Controlling algal blooms and enhancing carbon sequestration


Once the microalgal bloom occurs, the microalgal cells can be treated by inactivation processing at proper scales. E.g., bloom cells can be exposed to hydroxyl radicals (•OH) that effectively breaks the DNA strands of the microalgal cells while keeping their cell walls intact,² ensuring no intracellular organic matter and no extra dissolved organic carbon is released during the BCP process. Meanwhile MCP is well developed in seaweeds culture area,³ favoring the aggregation of the dead cells and therefore enhancing BCP efficiency.

In the controlled areas of intentional upwelling, on-site floating photobioreactors can be applied for cultivation of specific microalgae such as marine *Chlorella sp.* for high population densities to generate “Green Water”.⁴ Continuous supply of large quantities of microalgae not only produces sufficient dissolved oxygen and food for aquaculture animals but also offers a counter-measure to bead harmful blooms. Up on needed, flocculants can be introduced to aggregate and settle the microalgal cells down to the seafloor for long term burial.

In case microalgal bloom spreads, the cells can be flocculated and settled with modified clay which can also reduce the release rate of dissolved inorganic phosphorus (P).⁵ Dissolved organic carbon could be immobilized for uptake due to the enlarged C/P ratio, leading to the accumulation of recalcitrant dissolved organic matter through the MCP. Consequently, carbon sequestration efficiency is enhanced for both the BCP and the MCP.

BCMS STEP 3, Integrating the unmanageable SCP with BCP, CCP and MCP toward maximum complex carbon sequestration

Enhancing seawater alkalinity simultaneously with clay addition can secure both organic and inorganic carbon sequestration in the water column and sediment. Olivine with the main ingredients of Mg₂SiO₄, an effective alkaline mineral for CO₂ removal (Eq. 1), can chelate 4 moles of CO₂ per mole of olivine. More importantly, the CO₂ is captured and stored in the form of bicarbonate (Eq. 1), which is a major component of total alkalinity. This makes the alkaline addition an efficient approach for the enhancement of ocean alkalinity, which could significantly decrease surface seawater pCO₂ and promote CO₂ uptake from the atmosphere, enhancing the SCP efficiency.



Moreover, the use of olivine minerals could also supply extra nutrients including silicate and iron to promote the growth of micro/macroalgae. Consequently, the elevated seawater pH resulting from the enhanced micro/macroalgal photosynthesis drives the SCP to accelerates CO₂ absorption from the atmosphere into the sea, making the SCP a “manageable” approach for ONCE.

In the seabed, the metabolic activities of anaerobic methanotrophic archaea, nitrogen reduction and sulfate-reduction bacteria can not only contribute to the MCP but also enhance the concentration of bicarbonate alkalinity, promoting the combination of microbial mats/biofilm with Mg²⁺ and/or Ca²⁺ ions to induce carbonate deposition. The effective formation of carbonates-microbial mats/biofilm precipitation would then turn the CCP counter pump to be an effective pump for ONCE.

Closing remark

A comprehensive BCMS-based ONCE eco-engineering roadmap is proposed here towards achieving the twin goals of enhancement of carbon sink alongside remediation of the ecosystem. BCMS would be best implemented in sea-farming fields. This ONCE roadmap is theoretically riskless, environmentally friendly, sustainable, and efficient by using specialized measures, including photosynthesis rhythm-fit artificial upwelling, specifically designed DNA-inactivation hydroxyl radical treatment, modified clay sedimentation of organic matter, as well as natural minerals mediated water alkalinity enhancement. BCMS could become a forward-looking best practice for MRV (Measurement, Reporting, and Verification) based ONCE eco-engineering.

REFERENCES

- Jiao, N. Z., Herndl, G. J., Hansell, D. A., et al. (2010). Microbial production of recalcitrant dissolved organic matter: long-term carbon storage in the global ocean. *Nat. Rev. Microbiol.* **8**, 593–599.
- Bai, M. D., Zheng, Q. L., Zheng, W., et al. (2019). •OH Inactivation of Cyanobacterial Blooms and Degradation of Toxins in Drinking Water Treatment System. *Water Res.* **154**, 144–152.
- Li, H. M., Zhang, Z. H., Xiong, T. Q., et al. (2022). Carbon Sequestration in the Form of Recalcitrant Dissolved Organic Carbon in a Seaweed (Kelp) Farming Environment. *Environ. Sci. Technol.* **56**, 9112–9122.
- Zhai, X., Zhu, C., Zhang, Y., et al. (2020). Seawater supplemented with bicarbonate for efficient marine microalgae production in floating photobioreactor on ocean: A case study of *Chlorella sp.* *Sci. Total. Environ.* **738**, 139439.
- Lu, G. Y., Song, X. X., Yu, Z. M., et al. (2017). Application of PAC-modified kaolin to mitigate *Prorocentrum donghaiense*: effects on cell removal and phosphorus cycling in a laboratory setting. *J. Appl. Psychol.* **29**, 917–928.

ACKNOWLEDGMENTS

This study is supported by the National Natural Science Foundation of China (Grant No. 42188102), the Chinese Academy of Sciences (Grant No. 2020-ZW09-A-021 and XK2022DXA001), the Marine Economic Development Special Fund Project of Fujian Province of China (Grant No. FJHJF-L-2022-11), and the MOST ONCE project.

DECLARATION OF INTERESTS

The authors declare no competing interests.