



HAL
open science

Medication adherence in pediatric transplantation and assessment methods: a systematic review

Delphine Hoegy, Nathalie Bleyzac, Philip Robinson, Yves Bertrand, Claude Dussart, Audrey Janoly-Dumenil

► To cite this version:

Delphine Hoegy, Nathalie Bleyzac, Philip Robinson, Yves Bertrand, Claude Dussart, et al.. Medication adherence in pediatric transplantation and assessment methods: a systematic review. Patient Preference and Adherence, 2019, Volume 13, pp.705-719. 10.2147/PPA.S200209 . hal-04665713

HAL Id: hal-04665713

<https://hal.science/hal-04665713v1>

Submitted on 31 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Medication adherence in pediatric transplantation and assessment methods: a systematic review

This article was published in the following Dove Press journal:
Patient Preference and Adherence

Delphine Hoegy^{1,2}
Nathalie Bleyzac^{3,4}
Philip Robinson⁵
Yves Bertrand^{3,6}
Claude Dussart^{1,7}
Audrey Janoly-Dumenil^{1,2}

¹EA 4129 P2S Parcours Santé Systémique – Univ Claude Bernard Lyon I, Univ Lyon I, Lyon, France; ²Pharmacy, Hôpital Edouard Herriot, Hospices Civils de Lyon, Lyon, France; ³Institut d'Hématologie et d'Oncologie Pédiatrique, Hospices Civils de Lyon, Lyon, France; ⁴EMR 3738, PK/PD Modeling in Oncology, université Lyon-Sud, Lyon, France; ⁵Direction de la Recherche Clinique et de l'Innovation, Hospices Civils de Lyon, Lyon, France; ⁶UI111-CNRS UMR 5308, University of Lyon I, ENS Lyon, Lyon, France; ⁷Central Pharmacy, Hospices Civils de Lyon, Lyon, France

Background: Medication adherence is a major concern in public health. It is fully established that immunosuppressive therapy (IT) and concomitant medications affect transplant outcomes in the pediatric population, showing interest in adherence to this therapy. The aim of the present review was to report on medication adherence in pediatric population post-transplantation. This will enable us to know the situation in this particular population.

Methods: A literature search was performed using the MEDLINE database. Studies that were published from January 1999 to January 2016 in English language and which investigated medication adherence in pediatric transplantation were included. The type of organ and the methods used to assess medication adherence were studied.

Results: A total of 281 records were identified, from which 34 studies were selected: 38% (n=13) on kidney transplantation, 32% (n=11) on liver transplantation, and 23% (n=10) on the transplantation of other organs. Medication adherence was found to be lower than 80% in two-thirds of the studies (64%), and varied from 22% to 97%. This wide range was explained in part by the important heterogeneity of assessment methods among studies. The methods used were objective, non-objective, or combined both types. Most studies did not fully describe the data collected: the time since transplantation, the period over which adherence was assessed, the population, the medications, and the threshold discriminating adherence and non-adherence.

Conclusion: The present study found poor medication adherence in the pediatric population post-transplantation. There was a wide range of medication adherence, explained largely by the heterogeneity of assessment methods. Future studies must consider the characteristics of each methodology, but also the threshold defining adherence should be chosen on the basis of clinical outcomes, and describe all data collected to gain precision. To improve adherence in this population, it is essential to identify factors influencing medication (IT and concomitant medications) adherence.

Keywords: medication adherence, patient compliance, child, transplantation

Introduction

Adherence is defined “as the extent to which a person’s behavior [. . .] corresponds with agreed recommendations from a health care provider”. Medication adherence averages 50% among adults suffering from chronic diseases, and as poor medication adherence compromises the therapeutic approach it is a major concern in public health.¹

In adults, immunosuppressive therapy (IT) adherence averages 48% after renal transplantation, which is the most frequent transplant procedure.² In adults and pediatric populations, IT is recognized as essential after solid organ transplantation (SOT)^{3,4} and hematopoietic stem cell transplantation (HSCT).⁵ IT is part of treatment regimen post-transplantation that includes concomitant medications to prevent infection. After SOT in pediatric patients, the risk of biopsy-proven acute rejection is doubled, the risk of

Correspondence: Delphine Hoegy
P2S, EA 4129, universités Lyon I & III,
5 place d'Arsonval, 69003, Lyon, France
Email delphine.hoegy@chu-lyon.fr

hospitalization is increased by 60%, and that of organ loss by 80% when patients considered non-adherent to IT.⁶ Medication adherence is therefore a major concern in pediatrics. In 2009, Dew et al⁴⁹ published a meta-analysis that investigated medical regimen adherence after SOT in a pediatric population; the authors included studies that reported non-adherence to IT and/or clinical appointments. The authors concluded that there was a wide range of IT adherence rates due to the heterogeneity of methodological aspects and analysis of data in published studies. However, there is no recent systematic review that has investigated adherence to both IT and concomitant medications in a pediatric population after both SOT and HSCT. The aim of the present review was therefore to report on medication adherence in pediatric populations after both SOT and HSCT. This will provide an overview in this particular population, and a special focus will be made on the assessment methods used, which can be objective methods such as drug assays, and non-objective methods such as questionnaires.^{1,7}

Methods

The MEDLINE database was searched for relevant studies published from January 1, 1999 to January 1, 2016 using the following search strategy: (“patient compliance” [Mesh] OR “medication adherence” [Mesh] OR medication compliance) AND (pediatr* OR child) AND (transplan*). The inclusion

criteria were: studies in English language, reporting on pediatric transplant patients (age <21 years at the time of transplantation), measuring medication adherence, and describing the assessment method of medication adherence. Case reports, editorials, letters, and syllabi were excluded. Systematic and non-systematic reviews, as well as meta-analysis were excluded but checked for additional studies.^{11,49} The selection was independently reviewed by two of the authors (DH and AJD), and disagreements were resolved by consensus.

The analysis of selected studies consisted of description of organs, mean time since transplantation, population, medications; the methods used to assess medication adherence were also investigated.

Results

The first search strategy found 281 records; 221 were excluded because they did not talk about medication adherence in pediatric transplantation. A further 19 studies were excluded because a measure of medication adherence was not reported, and 7 because results obtained in pediatric subjects were not reported separately from other populations (Figure 1). Thus, 34 studies were included.

Thirteen studies concerned kidney transplantation (Table 1), and 11 liver transplantation (Table 2). Other studies reported IT adherence in heart transplantation (Table 3), HSCT (Table 4), or combined results of




Figure 1 Review inclusion and exclusion flowchart followed PRISMA guidelines.

Table 1 Medication adherence and assessment methods in kidney transplantation

References	Setting & Design	Population	Medications	Mean time since transplantation	Methods	Medication adherence (regarding methods)
Penkower et al, 2003 ⁸	USA Prospective, longitudinal	n=22 adolescents. Age 13–18 yr (mean 16)	Tac, CsA	4.3 yr	Interview (patient): ≤3 missed doses/month	87%
Shaw et al, 2003 ¹²	USA Retrospective	n=112. 48 child/64 adolescents Age 9–22 yr (mean 15) n ₁ =7 n ₂ =6. Age 2–21 yr (mean 11)	Tac, CsA	5 yr	Drug assay by 2 consecutive levels (Tac<2 ng/mL, CsA<50 ng/mL)	68% Child: 81% Ado: 65%
Gerson et al, 2004 ¹³	USA Prospective, comparative (methods)	n=79 adolescents. Age >12 yr	Tac, CsA	2 m	Gp 1: Combined: MEMS over 3 m, drug assay (with gradual physician's opinion) on 6 consecutive levels (Tac 4–12, CsA 100–250 ng/mL) Gp 2: drug assay only	Gp 1: 60% Gp 2: 80%
Feinstein et al, 2005 ¹⁴	Israel Retrospective	n=79 adolescents. Age >12 yr	Tac, CsA	ND	Combined: drug assay by levels (Tac>2, CsA>20 ng/mL) and/or questionnaire: ≤1 missed, frequency, "drug holiday"	84%
Takemoto et al, 2007 ¹⁵	USA Retrospective	n=91 adolescents. Age <18 yr (n=7046 with adults) n=150 adolescents. Age 10–22 yr (mean 17)	Tac, CsA	ND	MPR over 36 m: poor <69%, low 69–81%, medium 81–98%, high >98%. high defining adherent patients	56% (high)
Delucchi et al, 2008 ¹⁶	Chile Prospective	n=27 adolescents. Age 13–16 yr (mean 14) n=56 adolescents. Age 11–18 yr (mean 14) n=877.	IS	5.8 yr	Combined: BM (High creatininemia), drug assay by levels; and if suspected: questionnaire (10 patients/5 caregivers)	90%
Wu et al, 2008 ⁹	China Retrospective	n=27 adolescents. Age 13–16 yr (mean 14) n=56 adolescents. Age 11–18 yr (mean 14) n=877.	Tac, CsA, Aza, pred, MMF	ND (range 5.9–14.7)	Questionnaire	70%
Zelikovsky et al, 2008 ¹⁷	USA Prospective	n=23 adolescents. Age 11–18 yr (mean 15)	IS	ND	Questionnaire: MAM >90% or MAM >80%	62% 78%
Chisholm-Burns et al, 2009 ¹⁸	USA Retrospective	Age mean 12 yr n=23 families: 23 adolescents/22 parents	Tac, CsA	ND	MPR over 36 m: Interquartile 4= adherent Interquartile 1–3= non-adherent	92%
Dobbels et al, 2010 ¹⁹	Belgium Prospective	Age 11–18 yr (mean 15)	Tac (14), CsA (7), other IS (2)	3.8 yr	Combined: Questionnaire BAASIS (patient/caregiver); drug assay by levels (Tac>5, CsA>100 ng/mL); Gradual physician's opinion	22% 26/27% 63% 52%
Ingerski et al, 2011 ²⁰	USA Prospective	n=13 adolescents. Age 14–18 yr (mean 16)	IS	65.5 m	MAMS over 1 m	74%

(Continued)

Table 1 (Continued).

References	Setting & Design	Population	Medications	Mean time since trans-plantation	Methods	Medication adherence (regarding methods)
Pai et al, 2012 ²¹	USA Prospective	n=48 adolescents. 2 non-transplanted patients mean age: 16	Tac, SrL	3.9 yr	MEMS over 1 m	79%
Akchurin et al, 2014 ²²	USA Retrospective	n=46 adolescents. Age ND n ₁ =25 "child to adult care transition already done" n ₂ =22 "graft loss before transition"	Tac	ND	Questionnaire MAM (patient) Drug assay by SD (Tac): ND Drug assay by <10% level/patient ≤2.5 ng/mL during 2 yr (T1=before transition. T2=after transition)	97% (SD: 2.17) 59% n ₁ =68% n ₂ =32% (T1=T2)

Abbreviations: Ado, adolescents; Aza, Azathioprine; BAA/SIS, Basel Adherence Assessment Scale for immunosuppressive medication; CsA, cyclosporin; IS, immunosuppressant; m, month; MACS, multidimensional adherence classification system; MAM, medication adherence measure; MEMS, medication event monitoring system; MMF, mycophenolate mofetil; MPR, medication possession ratio; ND, not documented; Pred, prednisone; SrL, sirolimus; Tac, tacrolimus; T1, time 1; T2, time 2; yr, year.

different organs in the same study (multi-organ studies) (Table 5).

Among all types of organ, medication adherence ranged from 22%¹⁹ to 97%.²¹ Medication adherence was found to be lower than 80% in 22 studies, between 80% and 90% in 5 studies,^{8,12,14,23,24} and 90% or greater in 7 studies.^{16,18,21,32,34,37,42} Among these 7 studies, 3 focused on kidneys,^{16,18,21} 1 on liver,³² 1 on heart,³⁴ and 2 were multi-organ.^{37,42}

Twenty studies reported the mean time of medication adherence measurement since transplantation, which ranged from 27 days³⁵ to 9.4 years.^{31,33} The majority of these studies (n=12) examined medication adherence 5 years or later after transplantation, and among these medication adherence ranged from 27%³¹ to 97%.³⁷

The majority of studies (n=23) reported data for adolescents (≥10 years of age) only. Two studies reported data for children, in whom adherence was 30%⁴¹ and 91%.³⁴ One study stratified according to the patient's age, and concluded that medication adherence was poorer among adolescents (65%) than in children (81%).¹²

All included studies concerned IT. Twenty-two studies measured adherence to cyclosporine (CsA), 26 to tacrolimus (Tac), and 21 to both CsA and Tac. Three studies measured also concomitant medications: antibiotics, antivirals, antihypertensives for instance, without separated IT and concomitant medication adherence measures.^{35,41,42} One study compared IT to concomitant medications: IT adherence was poorer (60% vs 80%).⁴¹

Assessment methodologies

Period over which adherence was assessed

The majority of studies (n=24) reported the period over which adherence was assessed; this ranged from 7 days^{17,42} to 36 months.^{15,18} The period was related to the method used: medication event monitoring systems (MEMS) generally examined adherence over a 1–6-month period,^{13,20,21,35,37} medication possession ratio (MPR) over a 36-month period,^{15,18} and medication adherence measure (MAM) questionnaire over a 7-day period.^{17,42} Five studies^{13,20,21,35,37} examined medication adherence in a 3–6-month period: medication adherence ranged from 60%¹³ to 73%.³⁵ Eight studies^{15,18,22,23,29–31,38} examined medication adherence over a 1–2.5-year period: medication adherence ranged from 27%³⁸ to 92%.¹⁸

Table 2 Medication adherence and assessment methods in liver transplantation

References	Setting and design	Population	Medications	Mean time since transplantation	Methods	Medication adherence (regarding methods)
Shemesh et al, 2000 ²³	USA Retrospective	n=19. Age 9–20 yr (mean 14)	Tac	ND	Combined: physician's or nurse's opinion; drug assay by SD during 1 yr	84%
Falkenstein et al, 2004 ²⁴	USA Retrospective	n=234 (n=40 with non-compliance). n=28<10 yr & n=12>10 yr	Tac (11), CsA (29)	ND	Combined: BM (low AST/ALT); drug assay by levels (low levels C0 and C2h)	83%
Rumbo et al, 2004 ²⁵	USA Retrospective	n=14 adolescents. Age 12–17 yr	Aza	ND	Drug assay by SD (metabolite <50 pmol/8X 10 ⁸ erythrocytes) during 25m	71%
Shemesh et al, 2004 ²⁶	USA Prospective	n=81. Age 0.5–21 yr (30>8 yr)	Tac, CsA	ND	Drug assay by SD Questionnaire Likert-like scale (patient/care-giver)	(SD 2.48) 70/70%
Berquist et al, 2006 ²⁷	USA Retrospective	n=97 adolescents. Age 12–18 yr (mean 15)	Tac, CsA	ND	Nurse's opinion	52%
Berquist et al, 2008 ²⁸	USA Retrospective	n=111 adolescents. Age 12–21 yr (mean 15)	Tac, CsA	Non adherent: 12.4 yr Adherent: 9.6 yr	Physician's opinion	61%
Fredericks et al, 2008 ²⁹	USA Retrospective	n=38 adolescents. Age 12–18 yr (mean 15)	Tac	7.5 yr	RCR	62%
Stuber et al, 2008 ³⁰	USA Retrospective	n=68 adolescents. Age <21 yr	Tac	ND	RCR	50%
Fredericks et al, 2010 ³¹	USA Retrospective	n=71 adolescents. Age 11–20 yr (mean 12)	Tac, CsA	9.4 yr	Combined: questionnaire MAM (patient)	42%
Masuda et al, 2010 ³²	Japan Retrospective	n=108. Age 9–21 yr (mean 18)	Tac, CsA	8 yr	Drug assay by SD (<2) during 1 yr Drug assay by SD (≤3) during 2 yr or more	50%
Bilharz et al, 2015 ³³	USA Retrospective	n=48 adolescents. Age 11–20 yr (mean 16)	Tac	9.4 yr	Drug assay by SD (Tac<2, CsA <30) and by levels (<50% out of interval) during 1 yr	30%
					Combined: interview; and if suspected: Drug assay and BM (AST/ALT)	27%
					Drug assay by SD (≤2) during 6 m	94%
						67%

Abbreviations: ALT, Alanine Transaminase; AST, Aspartate Transaminase; Aza, Azathioprin; BM, Biological Markers; C0, Residual concentration; C2h, Concentration 2hours after administration; CsA, Cyclosporin; m, month; ND, Not Documented; RCR, retrospective chart review; SD, Standard Deviation; Tac, Tacrolimus; yr, year.

Table 3 Medication adherence and assessment methods in heart transplantation

References	Setting & Design	Population	Medications	Mean time since transplantation	Methods	Medication adherence (regarding methods)
Oliva et al, 2013 ³⁴	USA Retrospective	n=2030 children. Mean age: 6 yr	IS	ND	RCR	91%

Abbreviations: IS, immunosuppressant; ND, not documented; RCR, retrospective chart review; yr, year.

Table 4 Medication adherence and assessment methods in hematopoietic stem cell transplantation

References	Setting and design	Population	Medications	Mean time since transplant	Methods	Medication adherence (regarding methods)
Mc Grady et al, 2014 ³⁵	USA Prospective	n=6 adolescents. Age 12–18 yr (mean 15) malignant hematology (4), Non-malignant (1), oncology (1)	Tac, CsA, FLC, VRC, ACV, SXT	27 days	MEMS over 3 m	73%

Abbreviations: ACV, aciclovir; CsA, cyclosporin; FLC, fluconazole; m, month; MEMS, medication event monitoring system; ND, not documented; SXT, cotrimoxazole; Tac, Tacrolimus; VRC, voriconazole; yr, year.

Objective methods

Thirty-one studies assessed medication adherence using objective methods: drug assays (n=21), MEMS (n=6), MPR (n=2). A total of 21 studies used drug assays (Table 6) among which 2 did not mention the type of drug assay (level or SD).^{32,36} The 21 other studies considered either drug or metabolite levels in plasma, serum, or blood (n=10), and/or the SD that reports drug or metabolite levels over a period of time (n=11). SD studies (performed only for Tac) found medication adherence from 27%³⁸ to 90%.⁴¹ Studies using levels reported medication adherence that ranged from 27%³¹ to 84%.¹⁴ Six other studies used MEMS,^{13,20,21,35,37,40} which is a specific medication device recording date and timing of each use, and reported medication adherence ranged from 69%³⁷ to 79%²¹ (Table 6). Two studies used MPR: medication adherence reported was 56%¹⁵ and 92%.¹⁸

Non-objective methods

Twenty studies assessed medication adherence by non-objective methods: questionnaires or interviews (n=16), the opinion of physicians and/or nurses (n=4), retrospective chart review (RCR; n=3). RCR evaluated the number of patient records documenting medication non-adherence: the medication adherence reported in the 3 studies was 50%,²⁸ 62%,²⁷ and 91%.³⁴ Among the 4 studies using the opinion of physicians and/or nurses,^{13,19,23,26} 2 reported medication adherence by this method: 52%¹⁹ and 61%.²⁶ The medication

adherence, reported by the 16 studies using questionnaires and interviews ranged from 26%¹⁹ to 97%.²¹ Six studies used the MAM questionnaire,^{17,21,38,39,42,43} which is a specific and validated pediatric tool for adherence post-transplantation that evaluates the number of missed or late doses during the previous 7 days. The medication adherence from MAM questionnaire-studies ranged from 65%¹⁷ to 97%.²¹ Furthermore, a total of 7 studies reported caregiver or parent's medication adherence. Among these, 3 studies compared patient and caregiver reports and found results that were close: 26% and 27%,¹⁹ 70% and 70%,²⁶ and 86% and 88%.³⁹

Combined methods

Sixteen studies used several assessment methods and a total of 10 studies combined these to calculate medication adherence.^{13,14,16,19,23,24,32,38,41,43} With this methodology, medication adherence ranged from 22%¹⁹ to 94%,³² and 5 studies reported medication adherence greater than 80%.^{14,16,23,24,32} Four studies detailed precisely the combination of techniques.^{19,38,41,43}

One combination technique defined non-adherence as below a threshold for at least one item used.^{19,41} Two studies used a combination of drug assays, nurse/physician's opinion, and basal adherence assessment scale for immunosuppressive medication, a validated questionnaire on a 6-point scale. Medication adherence obtained with this combination technique was 22%¹⁹ and 30%.⁴¹

Table 5 Medication adherence and assessment methods in multi-organ studies

References	Setting and design	Population	Medications	Mean time since transplantation	Methods	Medication adherence (regarding methods)
Wray et al, 2006 ³⁶	UK Prospective	n=40 adolescents. Age 12–25 yr (mean 18) Heart (33), Heart-lungs (7) n=70.	Tac, CsA	8.8 yr	Questionnaire (patient): BMQ and if intentional: drug assay	72% 82%
Maikranz et al, 2007 ³⁷	USA Prospective	Age 7–18 yr (mean 13) heart (42), kidney (28)	IS	6 yr	MEMS (n=28) over 3 m Questionnaire (caregiver): missed doses past 3 days each month	69% 97%
Simons et al, 2009 ³⁸	USA Prospective	n=68 families: 71 adolescents/78 parents Age 11–21 yr (mean 16) kidney (47), liver (20), heart (14), lung (1)	Tac, CsA, SrL	4.8 yr	Combined: MACS: Questionnaire MAM >90%; drug assay (by SD <3 only for Tac) during 1 yr	All organs: 27% kidney: 28% liver: 25% heart: 29%
Simons et al, 2010 ³⁹	USA Prospective	n=66 families: 51 adolescents/62 parents Age 11–20 yr (mean 16) kidney (39), liver (16), heart (10), lungs (1)	Tac, CsA, SrL	ND	Questionnaire MAM >90% (caregiver/patient) Drug assay by levels during 16 m (<1 CsA <150, SrL<5, Tac <4ng/mL or SD<3 only for Tac)	86/88% 57%
Wu et al, 2010 ⁴⁰	USA Prospective	n=55 adolescents. Mean age: 13 yr liver (32), kidney (23) n=10 children.	Tac, CsA, MMF, SrL, Pred, SXT	6.2 yr	MEMS over 87 days	70%
Claes et al, 2014 ⁴¹	Belgium Prospective	Age 2–11 yr (mean 6) kidney (5), liver (5)	Tac (9), CsA (1) comedications	5.5 yr	Combined: Questionnaire BAASIS IS/comedications (caregiver) Drug assay (by SD<2 only for Tac) during 6–12 m Nurse's opinion	30% 60/80% 90% 30%
McCormick King et al, 2014 ⁴²	USA Prospective	n=72 adolescents. Age 12–21 yr (mean 18) kidney (37), liver (27), heart (5), heart & kidney (2), kidney & liver (1)	IS and other medications (antihypertensive)	6.7 yr	Questionnaire MAM (patient)	96%
Loiselle et al, 2015 ⁴³	USA Prospective Longitudinal: T1: since transplant. T2: 18 m after T1	N=66 adolescents. Age: ND (mean 15.8) kidney (39), liver (16), heart (10), heart & lung (1)	CsA, Tac, SrL	3.4 yr	Combined: MACS: Questionnaire MAM >90% (patient/caregiver); drug assay by levels (<1 level CsA <150, SrL <5, Tac <4 ng/mL or SD<3 only for Tac)	65% 82/73% 33%

Abbreviations: BMQ, Belief about Medication Questionnaire; CsA, cyclosporin; IS, immunosuppressant; MACS, multidimensional adherence classification system; m, month; MAM, medication adherence measure; MEMS, medication event monitoring system; MMF, mycophenolate mofetil; ND, not documented; Pred, prednisone; SrL, sirolimus; SXT, cotrimoxazole; Tac, tacrolimus; T, time; yr, year.

Table 6 Methods used to assess medication adherence

Methods	Number of studies concerned	References concerned	Medication adherence prevalence
Drug assays	21	12–14,16,19,21–26,29–33,36,38,39,41,43	27–90%
Questionnaire/interview	16	8,9,14,16,17,19,21,26,32,36–39,41–43	26–97%
Combined	10	13,14,16,19,23,24,32,38,41,43	22–94%
Medication event monitoring system	6	13,20,21,35,37,40	69–79%
Physician or nurse's opinion	4	13,19,23,26	30–61%
Retrospective chart review	3	27,28,34	50–91%

Another combination technique created balance scores of different methods, such as the multidimensional adherence classification system (MACS),^{38,43} which is a validated tool combining the MAM questionnaire and drug assay. Two studies used this technique: medication adherence was 27%³⁸ and 65%.⁴³

Discussion

To the best of our knowledge, this is the first time a systematic review has focused on medication adherence in a pediatric population after both SOT and HSCT. A total of 34 studies investigated medication adherence after transplantation over the last 16 years in pediatric populations, which is low considering the number of studies in other chronic diseases.⁴⁴ Irrespective of the transplanted organ, medication adherence ranged widely among studies. It remained poor with two-thirds studies (64%) describing medication adherence lower than 80%, highlighting the importance of considering medication adherence in the pediatric population post-transplantation.

Data for HSCT, a relatively frequent procedure, is reported only in one study,³⁵ whereas two studies reported data for heart-lung transplantation which is a rare procedure.^{34,36} This may be related to the short period required of IT after HSCT, as compared to the indefinite period required after SOT. Nevertheless, IT is essential for good prognosis after HSCT,⁵ and it is therefore of great interest to evaluate medication adherence after such transplantations.

All studies in this review investigated adherence to IT, underlining the importance of this concern in the post-transplantation period. Despite the need for an effective prevention of infections after transplantation, for instance, only 3 studies reported adherence to concomitant

medications such as antibiotics.^{35,41,42} In adults, Russel et al have considered that adherence to a medication – measured by MEMS – allowed the extrapolation of results to the comedications.⁴⁵ However in pediatrics, it is more appropriate to evaluate medication adherence for individual drugs^{41,46} as aspects such as medication flavor or form (syrup or pill) could cause different behaviors in children.⁷ It would be interesting that the 3 studies found in this review did compare IT adherence to concomitant medication adherence.

Most studies reported data for adolescents, which reflects a concern regarding medication adherence in this specific population after SOT and HSCT. Only one study compared data for children and adolescents, which is regrettable. This study revealed that medication adherence in children (81%) is higher than in adolescents (65%).¹² The adolescence period of life is known to decrease medication adherence by several reasons;⁷ for instance, Dobbels et al enunciate the transition to adulthood health care system as an important factor of non-adherence during adolescence.¹⁰ This is confirmed by a study included in this present review which found that IT adherence after transition (32%) is lower than before (68%).³² In addition, adolescence is a time when difficulties in the parent-patient relationship are important.⁷ Regarding it, the unfrequent comparison of adolescent and caregiver reports on medication adherence (3 studies in this review using subjective methods) would be interested to consider more.

An important aspect to consider when interpreting data is the measurement method employed. At the time of writing no “gold standard” method exists,^{1,7} which explains the heterogeneity of those used in this review. Surprisingly, 16 studies used only 1 method whereas it is known that no single method is optimal.^{1,7} For example,

Table 7 Assessment methods and definition of medication adherence or non-adherence by authors

References	Definition of medication adherence (MA) or medication non-adherence (MNA) by authors	MA or MNA	Methods	A priori threshold of MA
Penkower et al, 2003 ⁸	How often patients missed taking prescribed medication (seven options ranging from missing medication every day to never missing medications). Nonadherent if they missed taking medications at least 3 times in a month.	MNA	Interview (patient): ≤3 missed doses/month	Not documented
Shaw et al, 2003 ¹²	Non-adherence if the patient had consistently and significantly low serum IS levels (Tac<2 ng/mL, CsA<50 ng/mL) on a minimum of 2 consecutive levels without maintenance dose change, unaccompanied by an explanation of lack of absorption	MNA	Drug assay by 2 consecutive levels (Tac<2 ng/mL, CsA<50 ng/mL)	Not documented
Gerson et al, 2004 ¹³	"Medication adherence reported as a percentage and calculated by dividing the number of recorded presumptive medication doses by the number of prescribed dose and then multiplying the quotient by 100. Also, 2 nephrologists classify as 'probably adherent' to 'possibly non-adherent' regarding the target range"	MA	Gp 1: Combined: MEMS over 3m, drug assay (with gradual physician's opinion) on 6 consecutive levels (Tac 4–12, CsA 100–250 ng/mL) Gp 2: drug assay only	Not documented
Feinstein et al, 2005 ¹⁴	"Noncompliant behavior was defined as discontinuation of 1 or more IS medications, deviation from prescribed dose or frequency, and/or 'drug holidays'. Patients self reports and/or levels served as indicator (Tac<2, CsA<20 ng/mL) of non-compliance"	MNA	Combined: drug assay by levels (Tac>2, CsA>20 ng/mL) and/or questionnaire: ≤1 missed, frequency, "drug holiday"	Not documented
Takemoto et al, 2007 ¹⁵	We modeled compliance as a time-varying variable by splitting patients into quartiles based on overall 1–3 MPR values: poor compliance when MPR fell below 69%, low for MPR between 69 and 81%, medium for MPR 81 and 98% and high when the MPR exceeded 98%	MA	MPR over 36 m: poor <69%, low 69–81%, medium 81–98%, high >98%	>98%
Delucchi et al, 2008 ¹⁶	Not documented	MNA	Combined: BM (High creatinemia), drug assay by levels; and if suspected: questionnaire (10 patients/5 caregivers)	Not documented
Wu et al, 2008 ⁹	Not documented	MNA	Questionnaire	Not documented
Zelkowsky et al, 2008 ¹⁷	"Classifying patient as adherent or not adherent is also possible by selecting cut off with regard to percent adherence"	MA	Questionnaire: MAM >90% or MAM >80%	>90% >80%
Chisholm-Burns et al, 2009 ¹⁸	We also calculated MPR quartiles and used these data to classify pediatric renal transplant recipient into binary (adherent and nonadherent) groups. MPR Quartile 4 (the highest quartile) was classified as the adherent group and MPR quartile 1 to 3 were collapsed and classified as the nonadherent group.	MA	MPR over 36 m: Interquartile 4= adherent Interquartile 1–3= non-adherent	>75%
Dobbels et al, 2010 ¹⁹	Non-adherence is defined as any self-reported non-adherence (response score 1–5) on any of the four items [BAASIS questionnaire]. The four trough blood levels for cyclosporine or tacrolimus preceding the data collection were retrieved from the medical charts, with a tacrolimus trough level below 5 ng/mL, and a cyclosporine level below 100 ng/mL on at least one measurement being indicative for non-adherence	MNA	Combined: Questionnaire BAASIS (patient/caregiver); drug assay by levels (Tac>5, CsA>100 ng/mL); Gradual physician's opinion	Not documented

(Continued)

Table 7 (Continued).

References	Definition of medication adherence (MA) or medication non-adherence (MNA) by authors	MA or MNA	Methods	A priori threshold of MA
Ingerski et al, 2011 ²⁰	"Adherence was defined as the number of doses of oral medication taken divided by the number prescribed across the 1-month period"	MA	MEMS over 1 m	Not documented
Pai et al, 2012 ²¹	[MEMS] Adherence was defined as the number of times that doses of oral medication were taken out of the number of doses prescribed within a 30-day period. [MAM] Adherence was defined as the number of times that doses of oral medication were taken out of those prescribed during the designated seven-day time period. [drug assay] A larger SDTACRO or SDSIRO is considered indicative of poorer adherence.	MNA	MEMS over 1 m Questionnaire MAM (patient) Drug assay by SD (Tac): ND	Not documented
Akhurin et al, 2014 ²²	"Based on the percent of low tacrolimus levels, we categorized patients as nonadherent if they had more than 10% of low levels during the period of observation and as adherent otherwise"	MNA	Drug assay by <10% level/patient ≤2.5 ng/mL during 2 yr (T1=before transition, T2=after transition)	>90%
Shemesh et al, 2000 ²³	"higher SD, and therefore, increased variability between individual levels overtime, is a fairly specific but not a sensitive measure of nonadherence in our population"	MNA	Combined: physician's or nurse's opinion; drug assay by SD during 1 yr	Not documented
Falkenstein et al, 2004 ²⁴	Not documented	MNA	Combined: BM (Low AST/ALT); drug assay by levels (low levels C0 and C2h)	Not documented
Rumbo et al, 2004 ²⁵	Not documented	MNA	Drug assay by SD (metabolite <50 pmol/8X10 ⁸ erythrocytes) during 25 m	Not documented
Shemesh et al, 2004 ²⁶	"[SD] higher SD means a higher degree of difference between individual levels, which suggests less consistent medication taking and, therefore, less adherence [questionnaire] scale, 1 "I always take my medications as prescribed," 2 "I mostly take my medications as prescribed," 3 "I sometimes take my medications as prescribed," and 4 "I rarely take my medications as prescribed".	MA	Drug assay by SD Questionnaire Likert-like scale (patient/caregiver) Nurse's opinion Physician's opinion	Not documented
Berquist et al, 2006 ²⁷	"non-adherence was defined prior to data collection as any documented report by a patient, parent or health care provider of non-adherence to immunosuppressive medication taking by the patient that was formally recorded in the patient's legal medical record. Any patient with at least one episode of chart-documented non-adherence to taking the immunosuppressive medication after 1 yr post-transplantation was considered to be non-adherent in our study."	MNA	RCR	Not documented

(Continued)

Table 7 (Continued).

References	Definition of medication adherence (MA) or medication non-adherence (MNA) by authors	MA or MNA	Methods	A priori threshold of MA
Berquist et al, 2008 ²⁸	"non-adherence was defined as patient admission of not having taken the immunosuppressive(s) medication or not having attended an annual clinic visit in 2005. Any patient with at least one episode of not taking the immunosuppressive medication as prescribed by the healthcare provider after the six month post-transplantation visit was considered to be non-adherent. This definition included patients who changed the dose, those who took fewer doses, and a subgroup who made the independent decision to stop their immunosuppression completely without the healthcare provider's knowledge"	MNA	RCR	Not documented
Fredericks et al, 2008 ²⁹	"Adherence was defined as s.d. <2"	MA	Drug assay by SD (<2) during 1 yr	SD <2 for Tac
Stuber et al, 2008 ³⁰	Not documented	MNA	Drug assay by SD (≤ 3) during 2 yr or more	Not documented
Fredericks et al, 2010 ³¹	"adherence was defined as a tacrolimus SD <2 and cyclosporine SD <30, and/or <50% of immunosuppressant blood values out of the target range »	MA	Drug assay by SD (Tac<2, CsA <30) and by levels (<50% out of interval) during 1 yr	SD <2 for Tac
Masuda et al, 2010 ³²	"Noncompliance was defined as not taking the immunosuppressant medication"	NMA	Combined: Interview; and if suspected: Drug assay and BM (AST/ALT)	Not documented
Bilhartz et al, 2015 ³³	Not documented	MA	Drug assay by SD (≤ 2) during 6 m	SD <2 for Tac
Oliva et al, 2013 ³⁴	"Patient was defined to be NA when first reported to be so, irrespective of subsequent reports of adherence."	MNA	RCR	Not documented
Mc Grady et al, 2014 ³⁵	"Daily adherence was calculated by dividing the number of pill bottle openings per day by the number of prescribed doses for that day. Weekly and monthly adherence values were calculated by computing the mean of daily adherence values across the specified time frame"	MA	MEMS over 3 m	Not documented
Wray et al, 2006 ³⁶	Those patients who reported forgetting to take their medications regularly in response to a specific question on the BMQ questionnaire were considered to be unintentionally non-adherent. The second response was non-adherence which had been specified in the medical notes and which had been documented by the consultant cardiologist as intentional non-adherence	MNA	Questionnaire (patient); BMQ and if intentional: drug assay	Not documented

(Continued)

Table 7 (Continued).

References	Definition of medication adherence (MA) or medication nonadherence (MNA) by authors	MA or MNA	Methods	A priori threshold of MA
Maikranz et al, 2007 ³⁷	Adherence as assessed by MEMS was calculated as the number of recorded doses divided by the number of prescribed doses over the 3-month study period. Self-reported adherence was calculated as the percent of prescribed dose that were successfully administered for each medication.	MA	MEMS (n=28) over 3 m Questionnaire (caregiver): missed doses past 3 days each month	Not documented
Simons et al, 2009 ³⁸	"MACS places each patient into one of four categories based on parent/self report and immunosuppressant levels [Table 3]"	MNA	Combined: MACS: Questionnaire MAM >90%; drug assay (by SD <3 only for Tac) during 1 yr Questionnaire MAM >90% (caregiver/patient) Drug assay by levels during 16 m (<1 CsA <150, SrL<5, Tac <4 ng/mL or SD<3 only for Tac)	>90% SD<3 for Tac
Simons et al, 2010 ³⁹	"The final dichotomous categorization of drug levels, as "adherent" or "non-adherent" was determined by the presence of one or more out-of-range blood levels of tacrolimus with SD>3"	MNA	MEMS over 87 days	>90% SD<3 for Tac
Wu et al, 2010 ⁴⁰	"For each day of monitoring, a continuous measure of medication adherence was calculated by dividing the number of bottle openings by the number of prescribed dosage per day"	MA		Not documented
Claes et al, 2014 ⁴¹	When combining those 3 elements, the patient was considered nonadherent if he or she scored non-adherent in at least 1 of 3 of the following measures. (.) A score of 1 or higher on at least 1 of 4 questions [BAASIS] is considered as non-adherence. (.) A standard deviation of at least 2 is indicative of non-adherence (.) Average or bad [adherence; as scored by nurses] were therefore considered as medication nonadherent	MNA	Combined: Questionnaire BAASIS IS/comedications (caregiver) Drug assay (by SD<2, only for Tac) during 6–12 m Nurse's opinion	SD<2 for Tac
McCormick King et al, 2014 ⁴²	"Percentage of missed and late doses were calculated by dividing the total number of missed or late doses by the total number of prescribed doses in the previous 7 days"		Questionnaire MAM (patient)	Not documented
Loiselle et al, 2015 ⁴³	"Participants who missed or were late in taking 10% or more of their prescribed doses in the past week were classified as "non-adherent" (.) Based on these empirically derived out-of-target ranges, participants were coded as "non-adherent" if they had at least one serum levels that fell out of the target ranges (high or low levels for CsA, SrL, Tac, or >3 s.d. for Tac) and adherent if all levels were within the specified ranges."	MNA	Combined: MACS: Questionnaire MAM >90% (patient/caregiver); drug assay by levels (<1 level CsA <150, SrL <5, Tac <4 ng/mL or SD<3 only for Tac)	>90% SD<3 for Tac

Abbreviations: Ado, adolescents; ALT, alanine transaminase; AST, aspartate transaminase; Aza, azathioprine; BAASIS, Basel Adherence Assessment Scale for immunosuppressive medication; BM, biological markers; C0, Residual concentration; C2h, concentration 2 hrs after administration; CsA, cyclosporin; IS, immunosuppressant; m, month; MACS, multidimensional adherence classification system; MAM, medication adherence measure; MEMS, medication event monitoring system; MMF, mycophenolate mofetil; MPR, medication possession ratio; ND, not documented; Pred, prednisone; RCR, retrospective chart review; SrL, sirolimus; Tac, tacrolimus; T1, time 1; T2, time 2; yr, year.

drug assays will not detect patients who take their medications only before clinical visits. Conversely, MEMS is a longitudinal method taking into account “drug holidays” (when a patient discontinues medication of his/her own accord). Nevertheless, patients cannot be blinded to MEMS; it interferes with daily life and causes anxiety.⁴⁵ In pediatric patients, all these methods have additional drawbacks. For instance, MEMS-based studies may be too difficult to be feasible in pediatric patients who are unable to take pills,^{20,21,35,40} and drug assay methods are invasive so this may be a limiting factor in pediatric patients.⁴⁶ Yet, 23 of the 34 studies used this method, which could indicate that this is not a drawback too hard to surpass. In this review, only 10 studies (32%) used combined methods – 4 were published before 2005 and 6 after 2008 – suggesting that this is yet to become the preferred method. Among these studies, only a minority described precisely the combination method although this is essential to interpret data as the different combination techniques have different aims. One of the combination techniques has a high sensitivity: it underestimates medication adherence but misses less frequently non-adherent patients.^{19,41} Whereas the other combination technique used a well-balanced score (MACS),^{38,43} which allows categorization of patients according to their adherence.⁴⁷

An additional difficulty for the interpretation of data in the present review is the definition of medication adherence which differed between each selected study, as synthesized in the Supplementary material (Table 7). It is interesting to note that many studies did not define a threshold discriminating medication adherence and non-adherence, and when they did so this was rarely based on clinical outcomes. Generally, 80% of taken medication is the arbitrary threshold discriminating medication adherence and non-adherence.¹ To be relevant this threshold should be correlated to clinical outcomes, which was a posteriori investigated in 2 studies included herein; these found a close relationship between IT adherence and graft survival ($p=0.017$;¹⁸ $p=0.02^{48}$) and also mortality ($p=0.009$).⁴⁸ Also, Stuber et al³⁰ constructed ROC curve to try answering about the appropriate SD cut-off for Tac.

Limitations

This review included studies from the MEDLINE database only. Given the topic of this review, studies are essentially published in PubMed and research on other database is unlikely to have added additional relevant articles. An additional point is that this review is not a meta-analysis. Dew Ma et al published a meta-analysis

in 2009 focusing on medication adherence after pediatric SOT only.⁴⁹ The authors recognized that there was a lack of studies in certain transplantation types (lung, intestine) and noted the heterogeneity in published articles concerning methodological aspects and the analysis of data. This still seems to be the case 10 years later.

Conclusion

This review revealed the poor medication adherence in the pediatric population post-transplantation, and underlined also the wide range of medication adherence reported in the literature. The heterogeneity of assessment methods explained at least in part this wide range found. Future studies must weigh the advantages and disadvantages of each methodology and describe all data collected to gain precision. It is also essential that the threshold discriminating adherence and non-adherence must be chosen on the basis of clinical outcomes. Beyond the observation of poor medication adherence, there is an obvious need to improve this situation. For that it is essential to identify factors of medication non-adherence to propose adapted interventions and so try to improve medication adherence in adolescents and children after SOT and HSCT.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Sabaté E. *Adherence to Long-Term Therapies: Evidence for Action*. Geneva: World Health Organization; 2003. http://www.who.int/chp/knowledge/publications/adherence_report/en/. Accessed June 20, 2015.
2. Chisholm MA, Vollenweider LJ, Mulloy LL, et al. Renal transplant patient compliance with free immunosuppressive medications. *Transplantation*. 2000;70:1240–1244.
3. Neu AM. Special issues in pediatric kidney transplantation. *Adv Chronic Kidney Dis*. 2006;13:62–69. doi:10.1053/j.ackd.2005.10.001
4. Kerker N, Emre S. Issues unique to pediatric liver transplantation. *Clin Liver Dis*. 2007;11:323–335. doi:10.1016/j.cld.2007.04.001
5. Locatelli F, Zecca M, Rondelli R, et al. Graft versus host disease prophylaxis with low-dose cyclosporine-A reduces the risk of relapse in children with acute leukemia given HLA-identical sibling bone marrow transplantation: results of a randomized trial. *Blood*. 2000;95:1572–1579.
6. Connelly J, Pilch N, Oliver M, et al. Prediction of medication non-adherence and associated outcomes in pediatric kidney transplant recipients. *Pediatr Transplant*. 2015;19:1–8. doi:10.1111/ptr.12479
7. Osterberg L, Blaschke T. Adherence to medication. *N Engl J Med*. 2005;353:487–497. doi:10.1056/NEJMra050100
8. Penkower L, Dew MA, Ellis D, Sereika SM, Kitutu JMM, Shapiro R. Psychological distress and adherence to the medical regimen among adolescent renal transplant recipients. *Am J Transplant*. 2003;3:1418–1425. doi:10.1046/j.1600-6135.2003.00226.x

9. Wu ZX, Yang SL, Wu WZ, et al. The long-term outcomes of pediatric kidney transplantation: a single-centre experience in China. *Pediatr Transplant*. 2008;12:215–218. doi:10.1111/j.1399-3046.2007.00814.x
10. Dobbels F, Van Damme-Lombaert R, Vanhaecke J, De Geest S. Growing pains: non-adherence with the immunosuppressive regimen in adolescent transplant recipients. *Pediatr Transplant*. 2005;9:381–390. doi:10.1111/j.1399-3046.2005.00356.x
11. Dobbels F, Ruppert T, De Geest S, Decorte A, Van Damme-Lombaerts R, Fine RN. Adherence to the immunosuppressive regimen in pediatric kidney transplant recipients: a systematic review. *Pediatr Transplant*. 2010;14:603–613. doi:10.1111/j.1399-3046.2010.01299.x
12. Shaw RJ, Palmer L, Blasey C, Sarwal M. A typology of non-adherence in pediatric renal transplant recipients. *Pediatr Transplant*. 2003;7:489–493.
13. Gerson AC, Furth SL, Neu AM, Fivush BA. Assessing associations between medication adherence and potentially modifiable psychosocial variables in pediatric kidney transplant recipients and their families. *Pediatr Transplant*. 2004;8:543–550. doi:10.1111/j.1399-3046.2004.00215.x
14. Feinstein S, Keich R, Becker-Cohen R, Rinat C, Schwartz SB, Frishberg Y. Is noncompliance among adolescent renal transplant recipients inevitable? *Pediatrics*. 2005;115:969–973. doi:10.1542/peds.2004-0211
15. Takemoto SK, Pinsky BW, Schnitzler MA, et al. A retrospective analysis of immunosuppression compliance, dose reduction and discontinuation in kidney transplant recipients. *Am J Transplant*. 2007;7:2704–2711. doi:10.1111/j.1600-6143.2007.01966.x
16. Delucchi A, Gutierrez H, Arrellano P, Slater C, Meneses M, López I. Factors that influence nonadherence in immunosuppressant treatment in pediatric transplant recipients: a proposal for an educational strategy. *Transplant Proc*. 2008;40:3241–3243. doi:10.1016/j.transproceed.2008.03.125
17. Zelikovsky N, Schast AP, Palmer J, Meyers KEC. Perceived barriers to adherence among adolescent renal transplant candidates. *Pediatr Transplant*. 2008;12:300–308. doi:10.1111/j.1399-3046.2007.00886.x
18. Chisholm-Burns MA, Spivey CA, Rehfeld R, Zawaideh M, Roe DJ, Gruessner R. Immunosuppressant therapy adherence and graft failure among pediatric renal transplant recipients. *Am J Transplant*. 2009;9:2497–2504. doi:10.1111/j.1600-6143.2009.02793.x
19. Dobbels F, Decorte A, Roskams A, Van Damme-Lombaerts R. Health-related quality of life, treatment adherence, symptom experience and depression in adolescent renal transplant patients. *Pediatr Transplant*. 2010;14:216–223. doi:10.1111/j.1399-3046.2009.01197.x
20. Ingerski L, Perrazo L, Goebel J, Pai ALH. Family strategies for achieving medication adherence in pediatric kidney transplantation. *Nurs Res*. 2011;60:190–196. doi:10.1097/NNR.0b013e318215fdfe
21. Pai AL, Rausch J, Tackett A, Marsolo K, Drotar D, Goebel J. System for integrated adherence monitoring: real-time non-adherence risk assessment in pediatric kidney transplantation. *Pediatr Transplant*. 2012;16:329–334. doi:10.1111/j.1399-3046.2012.01657.x
22. Akchurin OM, Melamed ML, Hashim BL, Kaskel FJ, Del Rio M. Medication adherence in the transition of adolescent kidney transplant recipients to the adult care. *Pediatr Transplant*. 2014;18:538–548. doi:10.1111/petr.12289
23. Shemesh E, Lurie S, Stuber ML, et al. A pilot study of posttraumatic stress and nonadherence in pediatric liver transplant recipients. *Pediatrics*. 2000;105:E29. doi:10.1542/peds.105.2.e29
24. Falkenstein K, Flynn L, Kirkpatrick B, Casa-Melley A, Dunn S. Non-compliance in children post-liver transplant. Who are the culprits? *Pediatr Transplant*. 2004;8:233–236. doi:10.1111/j.1399-3046.2004.00136.x
25. Rumbo C, Shneider BL, Emre SH. Utility of azathioprine metabolite measurements in post-transplant recurrent autoimmune and immune-mediated hepatitis. *Pediatr Transplant*. 2004;8:571–575. doi:10.1111/j.1399-3046.2004.00230.x
26. Shemesh E, Shneider BL, Savitzky JK, et al. Medication adherence in pediatric and adolescent liver transplant recipients. *Pediatrics*. 2004;113:825–832.
27. Berquist RK, Berquist WE, Esquivel CO, et al. Adolescent non-adherence: prevalence and consequences in liver transplant recipients. *Pediatr Transplant*. 2006;10:304–310. doi:10.1111/j.1399-3046.2005.00451.x
28. Berquist RK, Berquist WE, Esquivel CO, Cox KL, Wayman KI, Litt IF. Non-adherence to post-transplant care: prevalence, risk factors and outcomes in adolescent liver transplant recipients. *Pediatr Transplant*. 2008;12:194–200. doi:10.1111/j.1399-3046.2007.00809.x
29. Fredericks EM, Magee JC, Opiari-Arrigan L, Shieck V, Well A, Lopez MJ. Adherence and health-related quality of life in adolescent liver transplant recipients. *Pediatr Transplant*. 2008;12:289–299. doi:10.1111/j.1399-3046.2008.00901.x
30. Stuber ML, Shemesh E, Seacord D, Washington J, Hellemann G, McDiarmid S. Evaluating non-adherence to immunosuppressant medications in pediatric liver transplant recipients. *Pediatr Transplant*. 2008;12:284–288. doi:10.1111/j.1399-3046.2008.00923.x
31. Fredericks EM, Dore-Stites D, Well A, et al. Assessment of transition readiness skills and adherence in pediatric liver transplant recipients. *Pediatr Transplant*. 2010;14:944–953. doi:10.1111/j.1399-3046.2010.01349.x
32. Masuda Y, Mita A, Ohno Y, et al. Noncompliance with medications in pediatric patients after living-donor liver transplantation. *Transplant Proc*. 2010;42:4191–4192. doi:10.1016/j.transproceed.2010.09.087
33. Bilhartz JL, Lopez MJ, Magee JC, Shieck VL, Eder SJ, Fredericks EM. Assessing allocation of responsibility for health management in pediatric liver transplant recipients. *Pediatr Transplant*. 2015;19:538–546. doi:10.1111/petr.12466
34. Oliva M, Singh TP, Gauvreau K, Vanderpluym CJ, Bastardi HJ, Almond CS. Impact of medication non-adherence on survival after pediatric heart transplantation in the U.S.A. *J Heart Lung Transplant*. 2013;32:881–888. doi:10.1016/j.healun.2013.03.008
35. McGrady ME, Williams SN, Davies SM, Pai ALH. Adherence to outpatient oral medication regimens in adolescent hematopoietic stem cell transplant recipients. *Eur J Oncol Nurs*. 2014;18:140–144. doi:10.1016/j.ejon.2013.11.007
36. Wray J, Waters S, Radley-Smith R, Sensky T. Adherence in adolescents and young adults following heart or heart-lung transplantation. *Pediatr Transplant*. 2006;10:694–700. doi:10.1111/j.1399-3046.2006.00554.x
37. Maikranz JM, Steele RG, Dreyer ML, Stratman AC, Bovaird JA. The relationship of hope and illness-related uncertainty to emotional adjustment and adherence among pediatric renal and liver transplant recipients. *J Pediatr Psychol*. 2007;32:571–581. doi:10.1093/jpepsy/jsl046
38. Simons LE, McCormick ML, Mee LL, Blount RL. Parent and patient perspectives on barriers to medication adherence in adolescent transplant recipients. *Pediatr Transplant*. 2009;13:338–347. doi:10.1111/j.1399-3046.2008.00940.x
39. Simons LE, McCormick ML, Devine K, Blount RL. Medication barriers predict adolescent transplant recipients' adherence and clinical outcomes at 18-month follow-up. *J Pediatr Psychol*. 2010;35:1038–1048. doi:10.1093/jpepsy/jsq025
40. Wu YP, Aylward BS, Steele RG. Associations between internalizing symptoms and trajectories of medication adherence among pediatric renal and liver transplant recipients. *J Pediatr Psychol*. 2010;35:1016–1027. doi:10.1093/jpepsy/jsq014
41. Claes A, Decorte A, Levchenko E, Knops N, Dobbels F. Facilitators and barriers of medication adherence in pediatric liver and kidney transplant recipients: a mixed-methods study. *Prog Transplant*. 2014;24(4):311–321. doi:10.7182/pit2014873

42. McCormick King ML, Mee LL, Gutiérrez-Colina AM, Eaton CK, Lee JL, Blount RL. Emotional functioning, barriers, and medication adherence in pediatric transplant recipients. *J Pediatr Psychol*. 2014;39:283–293. doi:10.1093/jpepsy/jst074
43. Loiselle KA, Gutierrez-Colina AM, Eaton CK, et al. Longitudinal stability of medication adherence among adolescent solid organ transplant recipients. *Pediatr Transplant*. 2015;19:428–435. doi:10.1111/ptr.12480
44. Kim SH, Gerver SM, Fidler S, et al. Adherence to antiretroviral therapy in adolescents living with HIV: systematic review and meta-analysis. *Aids*. 2014;28:1945–1956. doi:10.1097/QAD.0000000000000316
45. Russell CL, Conn VS, Ashbaugh C, Madsen R, Hayes K, Ross G. Intrasubject medication adherence patterns. *Clin Nurs Res*. 2007;16:153–163. doi:10.1177/1054773806296429
46. Chisholm MA, Mulloy LL, DiPiro JT. Comparing renal transplant patients' adherence to free cyclosporine and free tacrolimus immunosuppressant therapy. *Clin Transplant*. 2005;19:77–82. doi:10.1111/j.1399-0012.2004.00301.x
47. Simons LE, Gilleland J, Blount RL, et al. Multidimensional Adherence Classification System (MACS): initial development with adolescent transplant recipients. *Pediatr Transplant*. 2009;13:590–598. doi:10.1111/j.1399-3046.2008.01038.x
48. Chartrand C, Servando ES, Chartrand S. Risk factors for acute rejection after pediatric heart transplantation. *Transplant Proc*. 2001;33:1732–1734.
49. Dew MA, Dabbs AD, Myaskovsky L, et al. Meta-analysis of medical regimen adherence outcomes in pediatric solid organ transplantation. *Transplantation*. 2009;88(5):736–746. doi:10.1097/TP.0b013e3181b2a0e0

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease

states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>