

Combined InSAR-geophysical-seismological data for high-resolution site-effect modeling and ShakeMap generation, case study of Grenoble, France

Valentin Schindelholz, Aya Cheaib, Emeline Maufroy, Cécile Cornou, Erwan Pathier, Antoine Schlupp, Léna Pellorce, Véronique Mendel, Marc Grunberg

▶ To cite this version:

Valentin Schindelholz, Aya Cheaib, Emeline Maufroy, Cécile Cornou, Erwan Pathier, et al.. Combined InSAR-geophysical-seismological data for high-resolution site-effect modeling and ShakeMap generation, case study of Grenoble, France. 1ères Rencontres Epos-France, Oct 2023, Saint-Jean-Cap-Ferrat, France. hal-04668784

HAL Id: hal-04668784 https://hal.science/hal-04668784v1

Submitted on 7 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Combined InSAR-geophysical-seismological data for high-resolution siteeffect modeling and ShakeMap generation, case study of Grenoble, France Valentin Schindelholz¹, Aya Cheaib¹, Emeline Maufroy¹, Cécile Cornou¹, Erwan Pathier¹, Antoine Schlupp², Léna Pellorce³,

Véronique Mendel³, Marc Grunberq³ 1:ISTerre, Grenoble, France 2:ITES, Strasbourg, France 3:EOST, Strasbourg, France Contact: schindev@univ-grenoble-alpes.fr

Abstract: In the urbanized valley of Grenoble, a well-known sedimentary basin prone to complex 3D site amplifications of seismic waves, multiple geophysical and seismological data from previous temporary campaigns and from permanent networks are gathered to characterize the basin response to earthquakes. Our goal is to find physical correlations between the parameters available at the highest spatial resolutions and the seismic response that is recorded by seismological stations only at a few sparse points in the basin. If such correlations exist, it would allow us to map the seismic response at a finer spatial scale [1]. We process InSAR data to retrieve the subsidence rate of the sediments at millimetric precision and at 354291 points in the basin. We then test multiple correlations between the subsidence rates and all the available siteeffect parameters (as for example the depth of the bedrock below the sedimental resonance period), in order to determine a model of aggravation of the ground motion based on the subsidence rates. The obtained model can be applied at a fine spatial scale to predict the Grenoble basin response to earthquakes in high resolution. The modeled response can in turn be used as an input to the ShakeMaps, or to predict earthquake-damage potential at building scale.

Input Subsidence rates from InSAR (354291 points)



We investigate the correlation between the **subsidence rates** of the sediments and **5 parameters** of the site response: V_{S10}, f₀, T₀, log₁₀(*f*₀) and H_{bed} [2, 3, 4]. The best linear fits are obtained with the bedrock depth (**R²=0.67**) and the resonance period (**R²=0.53**) (shown below). The amplification factors are available **only** at 7 RAP stations (black triangles on the right), using the standard spectral ratios (SSR) from an earthquake and ground-motion catalog [5]. To estimate the amplification in all of the Grenoble basin, we use the link between observed the amplifications at low frequency and the bedrock depths at the stations to extrapolate these values. The subsidence rate is used to retrieve the bedrock depth at a fine spatial scale, using the correlation found, hence we finaly map at high resolution the amplification factor (see right). The misfits between the amplification factors obtained with this method and from the bedrock depth model are shown on the right. The largest misfits are located on the edges of the basin, where the depth of the basin is less well constrained.

