



HAL
open science

Serum and Salivary IgG and IgA Response After COVID-19 Messenger RNA Vaccination

Guy Gorochov, Jacques Ropers, Odile Launay, Karim Dorgham, Omaira da Mata-Jardin, Said Lebbah, Christine Durier, Rebecca Bauer, Anne Radenne, Corinne Desaint, et al.

► **To cite this version:**

Guy Gorochov, Jacques Ropers, Odile Launay, Karim Dorgham, Omaira da Mata-Jardin, et al.. Serum and Salivary IgG and IgA Response After COVID-19 Messenger RNA Vaccination. JAMA Network Open, 2024, 7 (4), pp.e248051. 10.1001/jamanetworkopen.2024.8051 . hal-04669127

HAL Id: hal-04669127

<https://hal.science/hal-04669127>

Submitted on 7 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Serum and Salivary IgG and IgA Response After COVID-19 Messenger RNA Vaccination

Guy Gorochov, MD, PhD¹; Jacques Ropers, PharmD²; Odile Launay, MD, PhD³; et al Karim Dorgham, PhD¹; Omaira da Mata-Jardin, PhD¹; Said Lebbah, MD²; Christine Durier, PhD⁴; Rebecca Bauer, PhD⁴; Anne Radenne, MSc⁵; Corinne Desaint, PhD³; Louis-Victorien Vieillard, MSc³; Claire Rekacewicz, MSc³; Marie Lachatre, MD³; Béatrice Parfait, MD, PhD⁶; Frédéric Batteux, MD, PhD⁷; Philippe Hupé, PhD^{8,9}; Lăétitia Ninove, MD¹⁰; Maeva Lefebvre, MD¹¹; Anne Conrad, MD, PhD¹²; Bertrand Dussol, MD, PhD¹³; Zoha Maakaroun-Vermesse, MD¹⁴; Giovanna Melica, MD¹⁵; Jean-François Nicolas, MD, PhD¹⁶; Renaud Verdon, MD, PhD¹⁷; Jean-Jacques Kiladjian, MD, PhD¹⁸; Paul Loubet, MD, PhD¹⁹; Catherine Schmidt-Mutter, MD, PhD²⁰; Christian Dualé, MD, PhD²¹; Séverine Ansart, MD, PhD²²; Elisabeth Botelho-Nevers, MD, PhD²³; Jean-Daniel Lelièvre, MD, PhD²⁴; Xavier de Lamballerie, MD, PhD¹⁰; Marie-Paule Kieny, PhD²⁵; Eric Tartour, MD, PhD²⁶; Stéphane Paul, PhD²⁷

Author Affiliations

- ¹Sorbonne Université, Institut National de la Santé et de la Recherche Médicale (INSERM), Centre d'Immunologie et des Maladies Infectieuses (CIMI), Département d'Immunologie, Assistance Publique–Hôpitaux de Paris (AP-HP), Hôpital Pitié-Salpêtrière, Paris, France
- ²INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique, AP-HP, Hôpital Pitié-Salpêtrière, Département de Santé Publique, Unité de Recherche Clinique Paris Sciences et Lettres (PSL)–CFX, Sorbonne Université, Paris, France
- ³Université Paris Cité, INSERM, Centre d'Investigation Clinique (CIC) 1417 Cochin Pasteur, French Clinical Research Infrastructure Network, Innovative Clinical Research Network in Vaccinology, APHP, Hôpital Cochin, Paris, France
- ⁴INSERM SC10-US019, Villejuif, France
- ⁵AP-HP, Hôpitaux Universitaires Pitié-Salpêtrière–Charles Foix, Unité de Recherche Clinique des Hôpitaux Universitaires Pitié-Salpêtrière, Paris, France
- ⁶AP-HP, Hôpital Cochin, Fédération des Centres de Ressources Biologiques–Plateforme de Ressources Biologiques Centre de Ressources Biologique Cochin, Paris, France
- ⁷AP-HP, Hôpital Cochin, Service d'Immunologie Biologique et Plateforme d'Immunomonitoring Vaccinal, Paris, France
- ⁸Institut Curie, PSL Research University, INSERM U900, MINES ParisTech, PSL, Paris, France
- ⁹Centre National de la Recherche Scientifique (CNRS), Unité Mixte de Recherche (UMR) 144, Paris, France
- ¹⁰Research Institute for Sustainable Development 190, INSERM 1207, Institut Hospitalier Universitaire Méditerranée Infection, Unité des Virus Émergents, Aix Marseille Université, Marseille, France
- ¹¹Centre Hospitalier Universitaire (CHU) de Nantes, INSERM CIC 1413, Maladies Infectieuses et Tropicales, Centre de Prévention des Maladies Infectieuses et Transmissibles, Nantes, France
- ¹²Département des Maladies Infectieuses et Tropicales, Hôpital de la Croix-Rousse, Hospices Civils de Lyon, Centre International de Recherche en Infectiologie (CIRI), INSERM U1111, Université Claude Bernard Lyon I, CNRS, UMR5308, École Normale Supérieure de Lyon, Université Lyon, Lyon, France

- ¹³CIC 1409, INSERM–Hôpitaux Universitaires de Marseille–Aix Marseille Université, Hôpital de la Conception, Marseille, France
- ¹⁴Centre de Vaccination CHU de Tours, CIC 1415, INSERM, Centre Hospitalier Régional et Universitaire de Tours, Tours, France
- ¹⁵Service d’Immunologie Clinique et Maladies Infectieuses, AP-HP, Hôpital Henri Mondor, Créteil, Centre d’Investigation Clinique 1430 INSERM, AP-HP, Hôpital Henri Mondor, Créteil, France
- ¹⁶CIRI, INSERM U1111, Université Claude Bernard Lyon I, Lyon, CHU Lyon-Sud, Pierre-Bénite, France
- ¹⁷Service de Maladies Infectieuses, CHU de Caen, Dynamicure INSERM UMR 1311, Normandie Université, University of Caen Normandy, Caen, France
- ¹⁸Université Paris Cité, AP-HP, Hôpital Saint-Louis, Centre d’Investigations Cliniques, INSERM, CIC 1427, Paris, France
- ¹⁹Virulence Bactérienne et Maladies Infectieuses, INSERM U1047, Department of Infectious and Tropical Diseases, CHU 37 Nîmes, Université de Montpellier, Nîmes, France
- ²⁰INSERM CIC 1434, CHU Strasbourg, Strasbourg, France
- ²¹CIC, INSERM CIC1405, CHU Clermont-Ferrand, Clermont-Ferrand, France
- ²²CIC 1412, INSERM, CHU Brest, Brest, France
- ²³INSERM CIC 1408, Axe Vaccinologie, CHU de Saint-Étienne, Service d’Infectiologie, Saint-Étienne, France
- ²⁴INSERM U955, Vaccine Research Institute, Créteil, France
- ²⁵INSERM 101, Paris, France
- ²⁶AP-HP, Hôpital Européen Georges Pompidou, INSERM U970, Paris Cardiovascular Research Center, Université Paris Cité, Paris, France
- ²⁷INSERM, U1111, CNRS, UMR 5308, CIRI-GIMAP, Université Claude Bernard Lyon 1, Université Jean Monnet, Immunology and Immunomonitoring Laboratory, iBiothera, CIC 1408, Saint-Étienne, France

Article Information

JAMA Netw Open. 2024;7(4):e248051. doi:10.1001/jamanetworkopen.2024.8051

Key Points

Question Can intramuscularly administered messenger RNA vaccines induce mucosal immunity?

Findings In this cohort study of 427 vaccinated individuals, a modest increase in spike-specific IgA levels in saliva of SARS-CoV-2–naïve individuals having received the Moderna or Pfizer-BioNTech messenger RNA vaccines was observed using ultrasensitive digital enzyme-linked immunosorbent assay. This increase persisted after Moderna vaccination, and previously infected individuals demonstrated a stronger mucosal IgA response to vaccination.

Meaning Specific salivary IgA can be detected after messenger RNA vaccination solely, but to much lower levels than in previously infected individuals.

Abstract

Importance

There is still considerable controversy in the literature regarding the capacity of intramuscular messenger RNA (mRNA) vaccination to induce a mucosal immune response.

Objective

To compare serum and salivary IgG and IgA levels among mRNA-vaccinated individuals with or without previous SARS-CoV-2 infection.

Design, Setting, and Participants

In this cohort study, SARS-CoV-2-naïve participants and those with previous infection were consecutively included in the CoviCompare P and CoviCompare M mRNA vaccination trials and followed up to day 180 after vaccination with either the BNT162b2 (Pfizer-BioNTech) vaccine or the mRNA-1273 (Moderna) vaccine at the beginning of the COVID-19 vaccination campaign (from February 19 to June 8, 2021) in France. Data were analyzed from October 25, 2022, to July 13, 2023.

Main Outcomes and Measures

An ultrasensitive digital enzyme-linked immunosorbent assay was used for the comparison of SARS-CoV-2 spike-specific serum and salivary IgG and IgA levels. Spike-specific secretory IgA level was also quantified at selected times.

Results

A total of 427 individuals were included in 3 groups: participants with SARS-CoV-2 prior to vaccination who received 1 single dose of BNT162b2 (Pfizer-BioNTech) ($n = 120$) and SARS-CoV-2-naïve individuals who received 2 doses of mRNA-1273 (Moderna) ($n = 172$) or 2 doses of BNT162b2 (Pfizer-BioNTech) ($n = 135$). The median age was 68 (IQR, 39-75) years, and 228 (53.4%) were men. SARS-CoV-2 spike-specific IgG saliva levels increased after 1 or 2 vaccine injections in individuals with previous infection and SARS-CoV-2-naïve individuals. After vaccination, SARS-CoV-2-specific saliva IgA levels, normalized with respect to total IgA levels, were significantly higher in participants with previous infection, as compared with the most responsive mRNA-1273 (Moderna) recipients (median normalized levels, 155×10^{-5} vs 37×10^{-5} at day 29; 107×10^{-5} vs 54×10^{-5} at day 57; and 104×10^{-5} vs 70×10^{-5} at day 180 [$P < .001$]). In contrast, compared with day 1, spike-specific IgA levels in the BNT162b2-vaccinated SARS-CoV-2-naïve group increased only at day 57 (36×10^{-5} vs 49×10^{-5} [$P = .01$]). Bona fide multimeric secretory IgA levels were significantly higher in individuals with previous infection compared with SARS-CoV-2-naïve individuals after 2 antigenic stimulations (median optical density, 0.36 [IQR, 0.16-0.63] vs 0.16 [IQR, 0.10-0.22]; $P < .001$).

Conclusions and Relevance

The findings of this cohort study suggest that mRNA vaccination was associated with mucosal immunity in individuals without prior SARS-CoV-2 infection, but at much lower levels than in previously infected individuals. Further studies are needed to determine the association between specific saliva IgA levels and prevention of infection or transmission.

Introduction

IgA is monomeric in human serum but is also produced locally in mucosal tissues mostly under dimeric or even polymeric forms and released associated with the secretory component as secretory IgA (SIgA).¹ Monomeric IgA is also present in saliva following passive transport from blood.¹ Saliva represents an easily accessible body fluid permitting probing of IgA mucosal immunity. Sterlin et al² previously showed that potent SARS-CoV-2–neutralizing IgA antibodies are rapidly produced following infection, even before IgG antibodies, although their levels decline and fail to be detected in saliva around 6 months after symptom onset. Recent studies^{3,4} suggest that SARS-CoV-2 spike-specific mucosal IgA antibodies could provide protection against breakthrough infection. However, it remains unclear whether messenger RNA (mRNA)–based vaccines can induce such SIgA responses. Several studies conclude that humoral mucosal immunity is largely driven by previous infection, with little impact of vaccination,⁴⁻⁶ while others suggest that mRNA vaccination can by itself elicit long-lasting anti–SARS-CoV-2 mucosal IgA responses.⁷⁻¹¹ Herein, we compared individuals with previous SARS-CoV-2 infection and SARS-CoV-2–naïve individuals included in the CoviCompare P and CoviCompare M vaccination trials¹² for their saliva and serum responses after 2 different modes of antigenic challenge: infection and vaccination or vaccination alone.

Methods

The CoviCompare P and CoviCompare M trials received ethical approval from Comité De Protection des Personnes Ile De France 1. All participants provided written informed consent. The study followed the Strengthening the Reporting of Observational Studies in Epidemiology ([STROBE](#)) reporting guideline.

Participants

From February 19 to June 8, 2021, participants without prior SARS-CoV-2 infection received either 2 doses of the mRNA-1273 (Moderna) vaccine in the CoviCompare M trial or the BNT162b2 (Pfizer-BioNTech) vaccine in the CoviCompare P trial at 28 days apart, whereas participants with a documented prior SARS-CoV-2 infection and positive serologic findings received only 1 dose of BNT162b2 (Pfizer-BioNTech) in accordance with the French guidelines at the time. In each trial, participants were healthy adults in stable medical condition. Samples were collected at multiple times: before the first vaccine administration (day 1), before the second vaccine administration (day 29), 28 days after the second vaccine dose (day 57), and 6 months after the first dose (day 180). Saliva was collected in a cup and serum was collected by venipuncture. All biospecimen were stored at –80°C prior to analysis.

Overall, 180 and 267 patients were included in the CoviCompare M and P trials, respectively. For the purpose of mucosal antibody analyses, 20 patients were excluded because (1) the second vaccine dose was delayed by more than 14 days after the planned date (n = 2); (2) anti–nucleocapsid levels were either missing before day 180 or were above the positivity threshold (n = 13); or (3) a SARS-CoV-2 infection was documented (using polymerase chain reaction analysis) at any time until day 180 (n = 5).

After these exclusions, 427 participants were included in these analyses, among whom 120 had a documented SARS-CoV-2 infection at least 5 months prior to their single dose of BNT162b2 vaccine (Pfizer-BioNTech) at the beginning of the pandemic, therefore at a period

when only ancestral wild-type SARS-CoV-2 (D614G) was circulating in France (Table). The remaining 307 patients had no prior SARS-CoV-2 infection and received either 2 doses of mRNA-1273 (Moderna) (n = 172) or 2 doses of BNT162b2 (Pfizer-BioNTech) (n = 135).

Quantification of Total Saliva IgA and IgG by Enzyme-Linked Immunosorbent Assay

For quantification of total IgA levels, high-binding 96-half-well plates (2310M; Nunc) were coated overnight at 4°C with 4 µg/mL of polyclonal goat anti-human IgA (I1261; Sigma-Aldrich) in phosphate-buffered solution (PBS). Serial dilutions of saliva in 5% skim milk in PBS, supplemented with 0.1% polysorbate 20 (Tween 20; Sigma-Aldrich), were added and plates were subsequently incubated for 1.5 hours at room temperature. Plates were then washed and incubated for 1 hour at room temperature with horseradish peroxidase (HRP)-conjugated polyclonal goat anti-human IgA (A0295; Sigma-Aldrich) diluted at 1:10 000 in PBS 5% milk. Bound antibodies were detected using tetramethylbenzidine substrate (T0440; Sigma-Aldrich). The color reaction was stopped with 0.5M sulfuric acid (H₂SO₄) after a 10-minute incubation, and the absorbance was measured at 450 nm in an enzyme-linked immunosorbent assay (ELISA) plate reader (Varioskan; Thermo Fisher Scientific Inc). Serial dilutions of prequantified saliva pool and monoclonal IgA were used for the generation of standard curves and measurement of concentration.

For measurement of total IgG, goat antihuman IgG Fc fragment antibody (A80-104A; Bethyl Laboratories) was coated overnight at 4°C (100 µL per well at 1 µg/mL in PBS) in 96-well plates (Immuno Maxisorp plates; Nunc). Plates were blocked for 1 hour in combined PBS and 3% bovine serum albumin, and a 500-fold dilution of saliva in combined PBS-3% bovine serum albumin was added and incubated for 1 hour at room temperature. Plates were washed 5 times with PBS-0.5% polysorbate 20 (Sigma-Aldrich) and incubated for 1 hour at room temperature with goat anti-human IgG Fc fragment HRP conjugated (A80-104P; Bethyl Laboratories) diluted to 1:150 000. After washing, 100 µL of tetramethylbenzidine (002023; Life Technologies) was added to each well, the reaction was stopped with 50 µL of 0.5M H₂SO₄ after 10 minutes of incubation, and the absorbance was measured at 450 nm in a microplate reader. Calculation of individual IgG concentrations was based on standard curves prepared with a commercial standard (N Protein Standard SL, model OQIM13; Siemens Healthcare Diagnostics AG).

SARS-CoV-2 Spike- and Nucleocapsid-Specific Serum Antibody Quantification by ELISA

Serum samples were tested for anti-SARS-CoV-2 IgG and IgA antibodies directed against the S1 domain of the spike protein of the virus using a commercial ELISA kit (EUROIMMUN). The kit is based on the detection of the Wuhan spike protein. Quantitative results were expressed in standardized units (binding antibody units [BAU] per milliliter). Serum IgG levels against SARS-CoV-2 nucleocapsid protein were detected using the V-PLEX COVID-19 serology kit (Meso Scale Discovery) and interpreted according to the manufacturer's instructions.

SARS-CoV-2 Spike-Specific Saliva Antibody Quantification by Digital ELISA

SARS-CoV-2 spike IgG antibody levels were measured as previously described^{13,14} by using a SARS-CoV-2 spike IgG assay (Simoa Advantage kit; Quanterix Corporation) on the SR-X

Platform (Quanterix Corporation) according to the manufacturer's instructions. Saliva samples were diluted 100-fold in sample diluent provided in the kit. We used an ultrasensitive digital ELISA for the measurement of specific anti-SARS-CoV-2 spike IgA mucosal responses, which allowed a better discrimination from background signals than classical ELISA performed in parallel on the same samples. The SARS-CoV-2 spike IgA antibody assay was developed using reagents provided in the SARS-CoV-2 spike IgG assay kit (Quanterix Corporation). Saliva samples were diluted 100-fold in sample diluent plus 0.5% polysorbate 20. Biotinylated antihuman IgA (clone REA1014; Miltenyi Biotec) diluted in sample diluent at 100 ng/mL was used as detector reagent. Streptavidin- β -galactosidase (S β G) was diluted to 25 pM in S β G diluent (SR-X 103158; Quanterix Corporation). Calibrator reagent consisted of recombinant anti-SARS-CoV-2 spike IgA (clone CR3022; Diaclone) diluted in sample diluent in a 4-fold dilution series (range, 0.05-200 ng/mL). Signals are expressed in BAU per milliliter. The lower limit of quantification was estimated to be 10 BAU/mL (eMethods in Supplement 1).

To avoid biases related to the fact that total IgA concentration within saliva is variable in samples from the same individual collected at different times, depending on various factors such as mode of sample collection and stress,¹ we normalized SARS-CoV-2 spike-specific IgA and IgG titers to the total IgA and IgG concentrations, respectively, within each saliva sample. Normalization of SARS-CoV-2 spike-specific saliva antibodies was calculated as follows: Normalized IgG = Anti-spike IgG/Total IgG and Normalized IgA = Anti-spike IgA/Total IgA.

Secretory SARS-CoV-2 Specific IgA Quantification by ELISA

To compare the best response in saliva samples among SARS-CoV-2-naïve participants (ie, the group receiving mRNA-1273 vaccine) with individuals who had a previous infection, for secretory piece-bound SIgA only, we selected groups exposed to 2 antigenic stimulations: previous infection and SARS-CoV-2-naïve individuals after a single or a second injection, respectively. High-binding 96-half-well plates (2310M) were coated overnight at 4°C with 1 μ g/mL of S1 spike protein (Wuhan strain) in carbonate coating buffer. Saturation of the coated plates were performed with 5% skim milk in PBS during 90 minutes at 37°C. Serial dilutions of saliva in 5% skim milk in PBS supplemented with 0.1% polysorbate 20 were added, and plates were subsequently incubated overnight at 4°C. Plates were then washed and incubated for 1 hour at room temperature with mouse antihuman secretory piece (1:1000 dilution; MilliporeSigma). After washing, plates were then revealed with HRP-conjugated polyclonal goat antimouse IgG (Invitrogen). Bound antibodies were detected using tetramethylbenzidine substrate (T0440; Sigma-Aldrich). The color reaction was stopped with 0.5M H₂SO₄ after 10-minute incubation, and the absorbance was measured at 450 nm in an ELISA plate reader (Varioskan).

Statistical Analysis

Data were analyzed from October 25, 2022, to July 13, 2023. Data were expressed as medians (IQRs) for continuous data and as frequencies and proportions for categorical data. For specific IgG and IgA levels, we computed a cutoff arbitrarily set at the mean plus 1 SD of measurements performed at baseline in SARS-CoV-2-naïve participants. Comparisons between groups were performed using the Wilcoxon rank sum test (with stratification on age as sensitivity analysis) and Fisher exact test for categorical data. Within groups, comparisons between different

times were performed using Wilcoxon rank sum test for paired data or McNemar test for proportions of participants above threshold. Fold changes of the geometric means were estimated with their 95% CI. All tests were performed on log₁₀-transformed data at the 2-sided 5% significance level ($P < .05$). Correlations between salivary and serum antibody levels were estimated using the Spearman correlation coefficient. All calculations were performed using R software, version 4.1.1 (R Program for Statistical Computing).

Results

We included 427 participants at the beginning of the pandemic, a period when only ancestral wild-type SARS-CoV-2 (D614G) was circulating in France. Of these, 120 participants had a documented SARS-CoV-2 infection prior to their single dose of BNT162b2 vaccine. The median age was 68 (IQR, 39-75) years; 228 participants (53.4%) were men and 199 (46.6%) were women. Race and ethnicity data were not collected, in conformity to French law.

Serum baseline SARS-CoV-2 spike-specific IgG and IgA levels (Figure 1 and eTable 1 in Supplement 1) were elevated in individuals with previous infection and increased to higher levels following vaccination, compared with SARS-CoV-2-naïve individuals. Among SARS-CoV-2-naïve patients, at all times following vaccination, serum anti-spike IgG and IgA levels were significantly higher in the mRNA-1273 compared with the BNT162b2 groups (IgG: 285.4 vs 117.5 BAU/mL at day 29, 2090.9 vs 1292.4 BAU/mL at day 57, and 395.4 vs 247.9 BAU/mL at day 180 [$P < .001$ for all comparisons]; IgA: 217.1 vs 74.4 BAU/mL at day 29, 603.9 vs 249.4 BAU/mL at day 57, and 119.2 vs 63.0 BAU/mL at day 180 [$P < .001$ for all comparisons]) (Figure 1).

A significant increase in specific IgG saliva levels was mostly detected after vaccination in previously infected individuals, but also in SARS-CoV-2-naïve individuals after each injection (Figure 2). Levels of salivary and serum anti-spike IgG antibodies were correlated in all 3 groups and at all times measured (eFigure 1 in Supplement 1), consistently with the known predominant systemic origin of saliva IgG.

The highest levels of SARS-CoV-2 spike-specific saliva IgA were detected in patients with previous infection 1 month after vaccination and slightly declined afterward (medians of normalized levels: 155×10^{-5} [IQR, 69×10^{-5} to 387×10^{-5}] at day 29; 107×10^{-5} [IQR, 63×10^{-5} to 217×10^{-5}] at day 57; and 104×10^{-5} [61×10^{-5} to 208×10^{-5}] at day 180) (Figures 1 and 3 and eTable 2 in Supplement 1). Levels were lower in SARS-CoV-2-naïve recipients of the mRNA-1273 vaccine (37×10^{-5} [IQR, 23×10^{-5} to 71×10^{-5}] at day 29; 54×10^{-5} [IQR, 31×10^{-5} to 90×10^{-5}] at day 57; and 70×10^{-5} [IQR, 41×10^{-5} to 118×10^{-5}] at day 180 [$P < .001$]). Levels of serum anti-spike IgA were better correlated with salivary-specific IgA among individuals with previous infection (remaining at day 180), rather than that of individuals without a prior infection (eFigure 1 in Supplement 1). In the latter groups, the correlation between blood and saliva IgA levels remained weak, with correlation coefficients substantially smaller than in individuals with previous infection (eFigure 1 in Supplement 1).

SARS-CoV-2 spike-specific saliva IgA levels remained more elevated 180 days after vaccination in individuals with previous infection compared with mRNA-1273-vaccinated individuals (104×10^{-5} [IQR, 61×10^{-5} to 208×10^{-5}] vs 70×10^{-5} [IQR, 41×10^{-5} to 118×10^{-5}];

$P < .001$) (Figure 1 and eTable 2 in Supplement 1). In individuals without prior infection, compared with day 1, increases of specific salivary IgA levels were substantially smaller, particularly after BNT162b2 vaccination, reaching statistical significance only at day 57 (36×10^{-5} [IQR, 19×10^{-5} to 79×10^{-5}] vs 49×10^{-5} [IQR, 25×10^{-5} to 103×10^{-5}]; $P = .01$) (Figure 3). In mRNA-1273-vaccinated individuals, specific saliva IgA levels increased significantly, albeit modestly, after the first (day 29 vs day 1, 1.4-fold [95% CI, 1.3- to 1.6-fold] change of geometric mean; $P < .001$) (eTable 3 in Supplement 1) and second vaccine doses (day 57 vs day 29, 1.3-fold [95% CI 1.1- to 1.5-fold] change of geometric mean; $P < .001$) (Figure 3 and eTable 3 in Supplement 1). In parallel, specific serum IgA levels increased in the same participants by 64.5-fold at day 29 vs day 1 (95% CI, 54.8- to 75.8-fold) and 182.6-fold at day 57 vs day 1 (95% CI, 154.3- to 216.1-fold) (eTable 1 in Supplement 1). Similar to IgG, salivary anti-spike IgA levels were significantly higher at day 180 in the mRNA-1273 group, compared with the BNT162b2 group (70×10^{-5} [IQR, 41×10^{-5} to 118×10^{-5}] vs 43×10^{-5} [IQR, 24×10^{-5} to 84×10^{-5}]; $P < .001$) (Figure 1). By definition, anti-nucleocapsid IgG levels were below the positivity threshold in individuals assigned to the mRNA-1273 group. Nevertheless, to verify whether asymptomatic infections might explain the late increase in anti-spike saliva IgA in the mRNA-1273 group, we split individuals in 2 groups according to anti-nucleocapsid IgG levels (SARS-CoV-2 nucleocapsid is not encoded by mRNA vaccines). As shown, anti-spike saliva IgA levels were not significantly elevated in individuals with anti-nucleocapsid serum IgG levels at day 180 above the 90th percentile of the distribution, compared with the others (eFigure 2 in Supplement 1). In conclusion, we did not find evidence that asymptomatic infections might explain the late increase in anti-spike saliva IgA in the mRNA-1273 group.

Finally, we verified whether bona fide mucosal secretory IgA could be locally induced at similar levels in SARS-CoV-2-naïve individuals and those with previous infections by comparing SIgA levels at their peak of response in both groups (ie, days 57 and 29, respectively). As shown (Figure 4 and eTable 4 in Supplement 1), spike-specific SIgA levels were significantly more elevated in previously infected individuals (median optical density, 0.36 [IQR, 0.16-0.63] vs 0.16 [IQR, 0.10-0.22]; $P < .001$). Age-stratified Wilcoxon rank sum tests as sensitivity analyses for the comparisons between individuals with previous infection and SARS-CoV-2-naïve individuals produced similar results as nonstratified tests, suggesting that differences between these groups were not confounded by age.

Discussion

In this cohort study, we compared individuals with previous SARS-CoV-2 infection and SARS-CoV-2-naïve individuals included in 2 large vaccination trials for their saliva and serum IgA and IgG antibody responses. Importantly, we limited the study to individuals vaccinated at the beginning of the pandemic to select a substantial number of confirmed SARS-CoV-2-naïve participants to be included in the study.

Since SARS-CoV-2 infection became extremely prevalent in Europe and in the US, we speculate that some of the previous inconsistencies reported in the literature³⁻¹¹ on saliva IgA responses might relate to difficulties recruiting participants who were truly SARS-CoV-2 naïve. Many SARS-CoV-2 infections are indeed asymptomatic and even more so in the recent phases of virus circulation.¹⁵ Furthermore, infections do not always leave a lasting serologic signature.¹⁵ A recent study¹¹ including a proportion of individuals with a history of SARS-CoV-

2 infection concluded that mRNA vaccination induces by itself a robust mucosal antibody production. In this study, more robust responses were induced primarily in participants with previous infection (Figures 1 and 3).

However, a modest but significant increase in specific saliva IgA levels was observed in SARS-CoV-2-naïve individuals receiving mRNA-1273 (Moderna) after the first and second administrations (Figure 1 and eTable 3 in Supplement 1). In parallel, specific serum IgA levels increased by 64.5-fold in the same participants (eTable 1 in Supplement 1). The correlations observed between serum and saliva IgA levels (eFigure 1 in Supplement 1) do not imply a blood origin for salivary IgA. Since the digital ELISA used in the study detects total anti-spike IgA, encompassing SIgA and possibly monomeric blood-borne IgA, we also measured electively secretory piece-bound SIgA to verify that bona fide SIgA levels were detected both in SARS-CoV-2-naïve individuals and those with previous infection. However, specific SIgA levels were significantly more elevated in individuals with previous infection compared with SARS-CoV-2-naïve individuals who received 2 antigenic stimulations (Figure 4).

Interestingly, we noted that anti-spike saliva IgA levels significantly increased between days 57 and 180 in the mRNA-1273 group (Figures 1 and 3), corresponding to an increase from 40 of 164 individuals (24%) to 55 of 159 (35%) with specific IgA levels above threshold (Figure 3). We reasoned that such a late variation could be related to asymptomatic SARS-CoV-2 infections. Data pertaining to anti-nucleocapsid saliva levels, by definition always below detection threshold in infection-naïve individuals, were available to test that hypothesis. We found no correlation between anti-spike saliva IgA and serum anti-nucleocapsid IgG levels (eFigure 2 in Supplement 1) that could have pointed in the direction of asymptomatic infections to explain a late increase in anti-spike saliva IgA levels in the mRNA-1273 vaccine group. It appears unlikely as well that late seronegative SARS-CoV-2 infections would have only occurred in the same latter group. Finally, our observation that specific serum and saliva IgA levels were significantly higher in SARS-CoV-2-naïve individuals at day 180 after vaccination with mRNA-1273 (Moderna) compared with BNT162b2 (Pfizer-BioNTech) (Figure 1) coincides with reports of more elevated antibody responses, which are associated with slightly lower hospitalization rates in the former group.^{16,17}

Limitations

This study has some limitations. We cannot determine whether the discrete increase in specific SIgA observed in some SARS-CoV-2-naïve vaccinations is the result of asymptomatic or seronegative exposures to respiratory virus, mucosal seeding by rare antibody-secreting cells trafficking from vaccine-draining lymph nodes, or both. In addition, we did not include data on neutralizing antibodies; however, prior studies^{2,18} suggest that specific antibody levels are strongly correlated. The number of vaccine breakthrough cases reported in our cohort was too limited to address the association between SIgA levels and prevention of infection or transmission of SARS-CoV-2.

Conclusion

The findings of this cohort study suggest that mRNA vaccination was associated with mucosal immunity in individuals with prior SARS-CoV-2 infection. Further studies are needed to determine the association between SIgA levels and prevention of infection or transmission.
Back to top

Article Information

Accepted for Publication: February 26, 2024.

Published: April 23, 2024. doi:10.1001/jamanetworkopen.2024.8051

Open Access: This is an open access article distributed under the terms of the CC-BY License. © 2024 Gorochov G et al. JAMA Network Open.

Corresponding Authors: Guy Gorochov, MD, PhD, Sorbonne Université, Centre d'Immunologie et des Maladies Infectieuses, Institut National de la Santé et de la Recherche Médicale (INSERM), Assistance Publique–Hôpitaux de Paris, Hôpital Pitié-Salpêtrière, 83 Boulevard de l'Hôpital, 75013 Paris, France (guy.gorochov@sorbonne-universite.fr); Stéphane Paul, PhD, Laboratoire d'Immunologie et d'Immunomonitoring, Centre d'Investigation Clinique, 1408 INSERM/Agence Nationale de Recherches sur le Sida Vaccinologie, Faculté de Médecine, 10 Rue de la Marandière, 42270 Saint-Priest-en-Jarez, France (stephane.paul@chu-st-etienne.fr).

Author Contributions: Drs Gorochov and Ropers had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Gorochov, Ropers, Launay, Durier, Desaint, Vieillard, Lachatre, Parfait, Hupé, Ninove, Kieny, Tartour, Paul.

Acquisition, analysis, or interpretation of data: Gorochov, Ropers, Launay, Dorgham, da Mata-Jardin, Lebbah, Bauer, Radenne, Vieillard, Rekacewicz, Batteux, Ninove, Lefebvre, Conrad, Dussol, Maakaroun-Vermesse, Melica, Nicolas, Verdon, Kiladjian, Loubet, Schmidt-Mutter, Dualé, Ansart, Botelho-Nevers, Lelièvre, de Lamballerie, Tartour, Paul.

Drafting of the manuscript: Gorochov, Ropers, Lebbah, Desaint, Parfait, Tartour, Paul.

Critical review of the manuscript for important intellectual content: Gorochov, Ropers, Launay, Dorgham, da Mata-Jardin, Durier, Bauer, Radenne, Vieillard, Rekacewicz, Lachatre, Batteux, Hupé, Ninove, Lefebvre, Conrad, Dussol, Maakaroun-Vermesse, Melica, Nicolas, Verdon, Kiladjian, Loubet, Schmidt-Mutter, Dualé, Ansart, Botelho-Nevers, Lelièvre, de Lamballerie, Kieny, Tartour, Paul.

Statistical analysis: Ropers, Lebbah, Durier, Bauer, Tartour, Paul.

Obtained funding: Gorochov, Launay, Radenne, de Lamballerie, Kieny, Tartour.

Administrative, technical, or material support: Gorochov, Ropers, Launay, da Mata-Jardin, Radenne, Vieillard, Rekacewicz, Parfait, Batteux, Hupé, Ninove, Lefebvre, Nicolas, Botelho-Nevers, Lelièvre, de Lamballerie, Paul.

Supervision: Gorochov, Ropers, Launay, Vieillard, Kiladjian, Paul.

Conflict of Interest Disclosures: Dr Launay reported receiving nonfinancial support from Pfizer Inc, MSD, Sanofi SA, and AstraZeneca, grant funding from Moderna Inc, Pfizer Inc, MSD, Sanofi SA, and AstraZeneca, and personal fees from Moderna Inc, Pfizer Inc, MSD, Sanofi SA, and AstraZeneca outside the submitted work. Dr Conrad reported consulting for Pfizer Inc and Moderna Inc outside the submitted work. Dr Schmidt-Mutter reported receiving financial support for personal and material investigation from Assistance Publique–Hôpitaux de Paris (AP-HP) outside the submitted work. Dr Dualé reported receiving grant funding from INSERM during the conduct of the study. Dr Botelho-Nevers reported participating on the scientific advisory boards for Pfizer Inc, Moderna Inc, Sanofi SA, and Janssen Global Services LLC outside the submitted work. Dr de Lamballerie reported receiving grant funding from the French Ministry of Research during the conduct of the study. No other disclosures were reported.

Funding/Support: The CoviCompare M and CoviCompare P trials have been supported by the Ministère des Solidarités et de la Santé and Ministère de l'Enseignement Supérieur, de la Recherche et de l'Innovation. In addition, the CoviCompare P trial was supported by ANRS Maladies Infectieuses Émergentes and the CoviCompare M trial was supported by AP-HP.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Disclaimer: The views expressed in this article are those of the authors and do not necessarily reflect the opinions of the French research and public health institutions with which they are affiliated.

Data Sharing Statement: See Supplement 2.

Additional Contributions: The authors are grateful to the volunteers included in CoviCompare M and CoviCompare P trials and clinical participating centers. This study has been labeled as a National Research Priority by the National Orientation Committee for Therapeutic Trials and other research on COVID-19 (CAPNET). The investigators would like to acknowledge ANRS Maladies Infectieuses Émergentes for their scientific support, the French Ministry of Health and Prevention, and the French Ministry of Higher Education, Research and Innovation for their funding and support.

References

1.

Brandtzaeg P. Induction of secretory immunity and memory at mucosal surfaces. *Vaccine*. 2007;25(30):5467-5484. doi:10.1016/j.vaccine.2006.12.001

2.

Sterlin D, Mathian A, Miyara M, et al. IgA dominates the early neutralizing antibody response to SARS-CoV-2. *Sci Transl Med.* 2021;13(577):eabd2223. doi:10.1126/scitranslmed.abd2223

3.

Havervall S, Marking U, Svensson J, et al. Anti-spike mucosal IgA protection against SARS-CoV-2 Omicron infection. *N Engl J Med.* 2022;387(14):1333-1336. doi:10.1056/NEJMc2209651

4.

Sheikh-Mohamed S, Isho B, Chao GYC, et al. Systemic and mucosal IgA responses are variably induced in response to SARS-CoV-2 mRNA vaccination and are associated with protection against subsequent infection. *Mucosal Immunol.* 2022;15(5):799-808. doi:10.1038/s41385-022-00511-0

5.

Sano K, Bhavsar D, Singh G, et al; PARIS Study Group. SARS-CoV-2 vaccination induces mucosal antibody responses in previously infected individuals. *Nat Commun.* 2022;13(1):5135. doi:10.1038/s41467-022-32389-8

6.

Thomas SN, Karger AB, Altawallbeh G, et al. Ultrasensitive detection of salivary SARS-CoV-2 IgG antibodies in individuals with natural and COVID-19 vaccine-induced immunity. *Sci Rep.* 2022;12(1):8890. doi:10.1038/s41598-022-12869-z

7.

Azzi L, Dalla Gasperina D, Veronesi G, et al. Mucosal immune response after the booster dose of the BNT162b2 COVID-19 vaccine. *EBioMedicine.* 2023;88:104435. doi:10.1016/j.ebiom.2022.104435

8.

Zuo F, Marcotte H, Hammarström L, Pan-Hammarström Q. Mucosal IgA against SARS-CoV-2 Omicron infection. *N Engl J Med.* 2022;387(21):e55. doi:10.1056/NEJMc221315

9.

Soffritti I, D'Accolti M, Gallenga C, et al. Evaluation of anti-SARS-CoV-2 IgA response in tears of vaccinated COVID-19 subjects. *Viruses.* 2023;15(2):399.

10.

Stolovich-Rain M, Kumari S, Friedman A, et al. Intramuscular mRNA BNT162b2 vaccine against SARS-CoV-2 induces neutralizing salivary IgA. *Front Immunol.* 2023;13:933347. doi:10.3389/fimmu.2022.933347

11.

Cao KT, Cobos-Uribe C, Knight N, et al. SARS-CoV-2 mRNA vaccination induces an intranasal mucosal response characterized by neutralizing antibodies. *J Allergy Clin Immunol Glob.* 2023;2(4):100129. doi:10.1016/j.jacig.2023.100129

12.

Molino D, Durier C, Radenne A, et al. A comparison of SARS-CoV-2 vaccine platforms: the CoviCompare project. *Nat Med.* 2022;28(5):882-884. doi:10.1038/s41591-022-01785-4

13.

Norman M, Gilboa T, Ogata AF, et al. Ultrasensitive high-resolution profiling of early seroconversion in patients with COVID-19. *Nat Biomed Eng.* 2020;4(12):1180-1187. doi:10.1038/s41551-020-00611-x

14.

Shan D, Johnson JM, Fernandes SC, et al. N-protein presents early in blood, dried blood and saliva during asymptomatic and symptomatic SARS-CoV-2 infection. *Nat Commun.* 2021;12(1):1931. doi:10.1038/s41467-021-22072-9

15.

Doll MK, Waghmare A, Heit A, et al. Acute and postacute COVID-19 outcomes among immunologically naive adults during Delta vs Omicron waves. *JAMA Netw Open.* 2023;6(2):e231181. doi:10.1001/jamanetworkopen.2023.1181

16.

Bajema KL, Dahl RM, Evener SL, et al; SUPERNOVA COVID-19; Surveillance Group; Surveillance Platform for Enteric and Respiratory Infectious Organisms at the VA (SUPERNOVA) COVID-19 Surveillance Group. Comparative effectiveness and antibody responses to Moderna and Pfizer-BioNTech COVID-19 vaccines among hospitalized veterans—five Veterans Affairs Medical Centers, United States, February 1-September 30, 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70(49):1700-1705. doi:10.15585/mmwr.mm7049a2

17.

Self WH, Tenforde MW, Rhoads JP, et al; IVY Network. Comparative Effectiveness of Moderna, Pfizer-BioNTech, and Janssen (Johnson & Johnson) vaccines in preventing COVID-19 hospitalizations among adults without immunocompromising conditions—United States, March-August 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70(38):1337-1343. doi:10.15585/mmwr.mm7038e1

18.

Marking U, Havervall S, Norin NG, et al. Correlates of protection and viral load trajectories in Omicron breakthrough infections in triple vaccinated healthcare workers. *Nat Commun.* 2023;14(1):1577. doi:10.1038/s41467-023-36984-1

Table. Participant Characteristics

Characteristic	Participant group				P value ^a
	All (n = 427)	Infection naive with mRNA-1273 (Moderna) (n = 172)	Infection naive with BNT162b2 (Pfizer-BioNTech) (n = 135)	Previous infection BNT162b2 (Pfizer-BioNTech) (n = 120)	
Age, median (IQR), y	68 (39-75)	69 (40-75)	68 (39-75)	67 (37-70)	.05
Age group, No. (%)					
18-45	152 (36)	55 (32)	47 (35)	50 (42)	.20
65-74	168 (39)	69 (40)	50 (37)	49 (41)	
>74	107 (25)	48 (28)	38 (28)	21 (18)	
Sex, No. (%)					
Men	228 (53)	98 (57)	68 (50)	62 (52)	.49
Women	199 (47)	74 (43)	67 (50)	58 (48)	
Body mass index, median (IQR) ^b	25 (22-27)	25 (22-27)	24 (22-27)	25 (22-29)	.11
First dose injection date, range	February 19 to June 8, 2021	February 19 to April 16, 2021	March 10 to May 4, 2021	March 15 to June 8, 2021	NA


Abbreviation: NA, not applicable.

^a Calculated using Kruskal-Wallis rank sum test or Fisher exact test.

^b Calculated as weight in kilograms divided by height in meters squared.

Figure 1. Longitudinal Evolution of Serum and Salivary Anti-Spike IgG and IgA Following Messenger RNA (mRNA) Vaccination in Individuals With or Without Previous Exposure to SARS-CoV-2


A Salivary anti-spike IgG



No. at risk


	1	29	57	180
Previous infection, BNT162b2	117	118	110	116
Infection naive, mRNA-1273	167	171	169	159
Infection naive, BNT162b2	135	135	134	133

B Salivary anti-spike IgA



	1	29	57	180
Previous infection, BNT162b2	113	114	100	116
Infection naive, mRNA-1273	167	170	164	159
Infection naive, BNT162b2	129	127	121	133


C Serum anti-spike IgG



No. at risk

	1	29	57	180
Previous infection, BNT162b2	120	120	116	115
Infection naive, mRNA-1273	172	172	172	172
Infection naive, BNT162b2	135	135	135	134

D Serum anti-spike IgA



	1	29	57	180
Previous infection, BNT162b2	120	120	116	115
Infection naive, mRNA-1273	172	172	172	172
Infection naive, BNT162b2	134	135	135	134


Evolution over time in individuals with previous SARS-CoV-2 infection receiving the BNT162b2 (Pfizer-BioNTech) vaccine, SARS-CoV-2-naive individuals receiving the mRNA-1273 (Moderna) vaccine, and SARS-CoV-2-naive individuals receiving BNT162b2 (Pfizer-BioNTech) of median \log_{10} levels of salivary and serum anti-spike IgG or IgA at days 1, 29, 57, and 180 after vaccination. Error bars indicate 95% CI. Dotted lines indicate threshold values computed as mean plus 1 SD of measurements made at day 1 among patients without prior infection. Results of comparison between groups at each time (Wilcoxon rank sum test) are shown.

^a $P \leq .001$, previous infection vs infection naive with mRNA-1273 vaccine.

^b $P \leq .001$, previous infection vs infection naive with BNT162b2 vaccine.


^c $P \leq .001$, infection naive with mRNA-1273 vaccine vs infection naive with BNT162b2 vaccine.

Figure 2. Salivary Anti-Spike IgG Response Following Messenger RNA (mRNA) Vaccination




Evolution over time in SARS-CoV-2-naive individuals receiving the mRNA-1273 (Moderna) vaccine or BNT162b2 (Pfizer-BioNTech) vaccine and previously infected individuals receiving BNT162b2 (Pfizer-BioNTech) of median \log_{10} salivary anti-spike IgG normalized on total salivary IgG levels at days 1, 29, 57, and 180 after vaccination, measured using digital enzyme-linked immunosorbent assay. Dotted lines indicate threshold values computed as mean plus 1 SD of measurements made at day 1 among patients without prior infection. Horizontal blue bars indicate median values. Results of the comparison between time points (Wilcoxon rank sum test) are provided in eTable 5 in Supplement 1.

Figure 3. Salivary Anti-Spike IgA Response Following Messenger RNA (mRNA) Vaccination



Evolution over time in SARS-CoV-2-naive individuals receiving the mRNA-1273 (Moderna) vaccine or the BNT162b2 (Pfizer-BioNTech) vaccine and previously infected individuals receiving BNT162b2 (Pfizer-BioNTech) of median \log_{10} salivary anti-spike IgA levels normalized on total salivary IgA at days 1, 29, 57, and 180 after vaccination, measured using digital enzyme-linked immunosorbent assay (taking into account both monomeric and multimeric secretory IgA). Dotted lines indicate threshold values computed as mean plus 1 SD of measurements made at day 1 among patients without prior infection. Horizontal blue bars indicate median values. Results of the comparison between time points (Wilcoxon rank sum test) are provided in eTable 5 in Supplement 1.

Figure 4. Multimeric Secretory Anti-Spike IgA After Vaccination in the Saliva of Individuals With Previous Infection Compared With SARS-CoV-2-Naive Individuals



Measure of anti-spike secretory IgA by classic enzyme-linked immunosorbent assay, using antisecretory piece detection antibodies. SARS-CoV-2-naive individuals received the mRNA-1273 (Moderna) vaccine and previously infected individuals received the BNT162b2 (Pfizer-BioNTech) vaccine. Horizontal orange bars indicate median values. Results of the comparison between groups (Wilcoxon rank sum test) are shown.

^a $P \leq .01$, days 1 vs 57, for infection-naive participants receiving mRNA-1273 vaccine.

^b $P \leq .001$, day 57 for infection-naive participants receiving mRNA-1273 vaccine vs day 29 for previously infected individuals.

Supplementary Online Content

Gorochov G, Ropers J, Launay O, et al. Serum and salivary IgG and IgA response after COVID-19 messenger RNA vaccination. *JAMA Netw Open*. 2024;7(4):e248051. doi:10.1001/jamanetworkopen.2024.8051

eTable 1. Serum SARS-CoV-2 Spike-Specific IgG and IgA

eTable 2. Salivary SARS-CoV-2 Spike-Specific IgG and IgA

eTable 3. Intragroup Longitudinal Variations of Salivary SARS-CoV-2 Spike-Specific IgA

eTable 4. SARS-CoV-2 Spike-Specific Secretory IgA (OD)

eFigure 1. Correlations Between Serum and Saliva IgG and IgA

eFigure 2. No Serological Evidence for Asymptomatic Infection in Moderna-Vaccinated Individuals With Late Increase in Specific Saliva IgA

eMethods. Digital ELISA Assay for SARS-CoV-2 Spike-Specific IgA

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable 1. Serum SARS-CoV-2 Spike-Specific IgG and IgA

Abbreviation: IQR: interquartile range

	D1	D29	D57	D180
mRNA-1273, N = 172				
IgG, Median (IQR)	0.3 (0.2 – 0.7)	285.4 (152.4 – 493.6)	2,090.9 (1,250.6 – 3,051.4)	395.4 (201.4 – 621.3)
Missing	0	0	0	0
IgA, Median (IQR)	2.9 (1.9 – 5.5)	217.1 (111.5 – 474.1)	603.9 (315.3 – 1,209.1)	119.2 (53.5 – 260.0)
Missing	0	0	0	0
BNT162b2, N = 135				
IgG, Median (IQR)	0.3 (0.2 – 0.6)	117.5 (57.4 – 243.0)	1,292.4 (794.2 – 2,031.6)	247.9 (144.5 – 380.0)
Missing	0	0	0	1
IgA, Median (IQR)	3.4 (2.0 – 6.7)	74.4 (40.0 – 164.0)	249.4 (144.3 – 483.3)	63.0 (33.0 – 151.8)
Missing	1	0	0	1
Previously Infected, N = 120				
IgG, Median (IQR)	132.7 (38.8 – 276.2)	6,888.8 (4,000.2 – 12,696.6)	4,859.1 (2,850.4 – 7,970.2)	2,110.9 (698.8 – 4,493.4)
Missing	0	0	4	5
IgA, Median (IQR)	103.2 (44.8 – 215.0)	2,211.5 (858.9 – 4,497.9)	1,438.6 (705.1 – 2,753.1)	813.4 (409.1 – 1,880.7)
Missing	0	0	4	5

Quantitative results expressed in standardized binding antibody units per mL (BAU/mL).

Table 2. Salivary SARS-CoV-2 Spike-Specific IgG and IgA

	D1	D29	D57	D180
mRNA-1273, N = 172				
IgG saliva, normalized*, Median (IQR)	0.00002 (0.00001 – 0.00007)	0.00060 (0.00030 – 0.00109)	0.00440 (0.00278 – 0.00725)	0.00081 (0.00040 – 0.00137)
Missing	5	1	3	13
IgA-saliva, normalized**, Median (IQR)	0.00026 (0.00015 – 0.00048)	0.00037 (0.00023 – 0.00071)	0.00054 (0.00031 – 0.00090)	0.00070 (0.00041 – 0.00118)
Missing	5	2	8	13
BNT162b2, N = 135				
IgG saliva, normalized*, Median (IQR)	0.00003 (0.00001 – 0.00009)	0.00028 (0.00011 – 0.00061)	0.00292 (0.00140 – 0.00544)	0.00045 (0.00025 – 0.00073)
Missing	0	0	1	2
IgA saliva, normalized**, Median (IQR)	0.00036 (0.00019 – 0.00079)	0.00044 (0.00021 – 0.00070)	0.00049 (0.00025 – 0.00103)	0.00043 (0.00024 – 0.00084)
Missing	6	8	14	2
Previously Infected, N = 120				
IgG saliva, normalized*, Median (IQR)	0.00021 (0.00009 – 0.00059)	0.00990 (0.00340 – 0.01915)	0.00935 (0.00387 – 0.01900)	0.00382 (0.00152 – 0.01085)
Missing	3	2	10	4
IgA saliva, normalized**, Median (IQR)	0.00052 (0.00030 – 0.00105)	0.00155 (0.00069 – 0.00387)	0.00107 (0.00063 – 0.00217)	0.00104 (0.00061 – 0.00208)
Missing	7	6	20	4

Abbreviation: IQR: interquartile range

*Normalized IgG = ng (specific IgG)/ng (total IgG)

**Normalized IgA = AU (specific IgA)/ng (total IgA)

eTable 3. Intragroup Longitudinal Variations of Salivary SARS-CoV-2 Spike-Specific IgA

		Missing	p-value ¹	Fold change (CI 95%)
mRNA-1273, N = 172				
D1	D29	6	<0.001	1.41 (1.25 - 1.59)
D1	D57	12	<0.001	1.89 (1.64 - 2.17)
D1	D180	18	<0.001	2.45 (2.09 - 2.87)
D29	D57	9	<0.001	1.30 (1.14 - 1.48)
D57	D180	21	0.002	1.27 (1.08 - 1.50)
BNT162b2, N = 135				
D1	D29	10	0.25	1.11 (0.94 - 1.31)
D1	D57	17	0.01	1.29 (1.08 - 1.54)
D1	D180	8	0.41	1.15 (0.93 - 1.42)
D29	D57	15	0.07	1.17 (0.96 - 1.42)
D57	D180	15	0.16	0.86 (0.68 - 1.09)
Previously Infected, N = 120				
D1	D29	10	<0.001	2.86 (2.31 - 3.54)
D1	D57	23	<0.001	1.89 (1.52 - 2.36)
D1	D180	11	<0.001	1.80 (1.45 - 2.25)
D29	D57	21	<0.001	0.64 (0.53 - 0.79)
D57	D180	21	0.28	0.93 (0.76 - 1.15)

Abbreviation: CI: confidence interval

¹Wilcoxon rank sum test for paired data


eTable 4. SARS-CoV-2 Spike-Specific Secretory IgA (OD)

	mRNA-1273 D57, N = 172	Previously infected D29, N = 120	p-value¹
OD, Median (IQR)	0.16 (0.10 – 0.22)	0.36 (0.16 – 0.63)	<0.001
Missing	2	2	

Abbreviation: IQR: interquartile range; OD: optical density


¹Wilcoxon rank sum test; ²Wilcoxon rank sum test stratified on age

eFigure 1. Correlations Between Serum and Saliva IgG and IgA



Correlations between serum and saliva IgG (left column), serum and saliva IgA (right) at indicated times (day 29, 57 and 180) after vaccination in BNT162b2-vaccinated previously infected individuals (PI, blue), mRNA-1273-vaccinated naïve individuals (M, red) and BNT162b2-vaccinated naïve individuals (P, green). Regression lines are standard linear regressions. Spearman rank correlations are shown on top of each graph along with 95% confidence intervals and p-values.

eFigure 2. No Serological Evidence for Asymptomatic Infection in Moderna-Vaccinated Individuals With Late Increase in Specific Saliva IgA



By definition all individuals tested here had antinucleocapsid antibodies below detection threshold. Levels of salivary anti-spike IgA at day 180 (measured by digital Elisa) among SARS-CoV-2-naïve individuals following 2 mRNA-1273 injections split in two groups according to antinucleocapsid signals recorded. Groups were formed according to serum Elisa signals against nucleocapsid measured by classical Elisa at day 180 in naïve individuals (being lower or greater than the 90th percentile of the anti-N distribution). Dotted line indicates threshold value for specific IgA positivity computed arbitrarily as the mean + 1 standard deviation of measurements performed at baseline in naïve individuals. Green plain lines indicate median values of antinucleocapsid signals. Results of comparison between groups (Wilcoxon rank sum test) are shown with $p > 0.05$: ns.

eMethods. Digital ELISA Assay for SARS-CoV-2 Spike-Specific IgA

In the first step of the assay, 25 μ L of SARS-CoV-2 spike-coupled beads (Bead Reagent from the SARS-CoV-2 spike IgG Advantage Assay kit) were incubated with 100 μ L of diluted saliva in a 96-well conical bottom plate for 30 min at 30°C at 800 rpm. The plate was washed on a Simoa microplate washer equipped with a magnet, and the beads were incubated with 100 μ L of detector reagent for 10 minutes at 30°C at 800 rpm. After a new washing step, beads were incubated with the SBG reagent and incubated at 30°C at 800 rpm for 10 minutes. The beads were then washed twice in buffer B at 800 rpm for 1 minute. Buffer was then removed and the beads were dried for 10 minutes on the magnet. The plate was then transferred to the SR-X instrument with RGP reagent and read according to a home-brew analysis protocol. The concentration of each sample was calculated based on the four-parameter logistic fitting model generated with the calibrators. Signals are expressed in binding antibody units (BAU)/mL.

Limit of Detection: Lower Limit of Detection (LOD) is calculated as 2.5 Standard Deviations above the background (mean of calibrator blanks).

Mean LOD: 0.074 BAU/mL (Range: 0.030 – 0.134 BAU/mL)

Limit of Quantification: Lower Limit of Quantification (LLOQ) is the highest concentration of calibrator with $\leq 20\%$ pooled CV.

Mean LLOQ: 0.093 (Range: 0.067 – 0.148 BAU/mL)

Functional LLOQ is the Analytical LLOQ multiplied by the Minimum Required Dilution (1:100): 9.25 BAU/mL

Data normalization: To minimize the impact of variations between runs, all sample concentrations were normalized. Six saliva samples diluted 1:100 were used as controls in each run performed on the SR-X analyser (Quanterix®). The average concentration of controls and the corresponding AEB values (AEB: average enzymes per bead is the unit of measurement for Simoa) were used to construct a new calibration curve for each analysis. The mean concentration corrected for MRD (BAU/mL) and the coefficient of variation (% CV) based on concentration are reported in the table below.

Experiments	Control 1	Control 2	Control 3	Control 4	Control 5	Control 6
Mean (BAU/mL)	16	95	146	241	1248	1976
Range	13 - 20	76 - 123	121 - 163	203 - 280	893 - 1665	1398 - 2843
CV	14%	17%	11%	12%	18%	22%

Anti-SARS-CoV-2 spike IgA binding capacity in samples is related to antibody concentration but also to affinity and avidity (dimer vs monomer) effects. IgA are mostly dimeric in saliva. For that reason, we chose here to express anti-Spike IgA data as BAU (binding antibody units) and not in concentration (ng/ml). Although useful, available monomeric recombinant anti-Spike IgA antibodies cannot be used as calibrators to measure accurate concentrations. Furthermore, after vaccination or infection, the anti-Spike IgA response is polyclonal and the overall binding capacity of this heterogeneous population of antibodies may not be comparable to that of a monoclonal IgA used as a reference.