



HAL
open science

Full stereochemical understanding in a new (2 R ,3 R ,4 R)-4-hydroxyisoleucine synthesis

Marc Rolland, Tarek Kassem, Valérie Rolland, Jean Martinez

► **To cite this version:**

Marc Rolland, Tarek Kassem, Valérie Rolland, Jean Martinez. Full stereochemical understanding in a new (2 R ,3 R ,4 R)-4-hydroxyisoleucine synthesis. *Acta Crystallographica Section C: Crystal Structure Communications* [1968-2013], 2001, 57 (12), pp.1415-1417. 10.1107/s0108270101014597 . hal-04671408

HAL Id: hal-04671408

<https://hal.science/hal-04671408v1>

Submitted on 8 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Full stereochemical understanding in a new (2R,3R,4R)-4-hydroxyisoleucine synthesis

Marc Rolland,* Tarek Kassem,* Valérie Rolland* and Jean Martinez*

*Laboratoire des Aminoacides, Peptides et Protéines, Université Montpellier 1 et II, Faculté de Pharmacie, UMR CNRS 5810, 15 Avenue Charles Flahault, 34060 Montpellier CEDEX 2, France

*Correspondence e-mail: rolland@colombes.pharma.univ-montp1.fr (Received 21 June 2001; accepted 6 September 2001)


We present the crystal and molecular structures of 2,3,6,7,8,8a-hexahydro-6,8-methano-7,7,8a-trimethyl-3-(1-methyl-2-oxopropylidene)-5H-1,4-benzoxazin-2-one, C16H25NO3, (III), and 2,3,6,7,8,8a-hexahydro-3-(2-hydroxy-1-methylpropyl)-6,8-methano-7,7,8a-trimethyl-5H-1,4-benzoxazin-2-one, C16H25NO3, (V). These compounds are two of the four key intermediates in our synthetic route to (2R,3R,4R)-4-hydroxyisoleucine. The two structures provide a full understanding of the stereochemistry in successive steps. This synthesis was based on a new optically pure chiral oxazinone auxiliary derived from (1R,2R,5R)-2-hydroxypipnan-3-one.

Reuse permissions | PowerPoint slides

Comment

For the synthesis of gamma-hydroxy-alpha-amino acids (Jacob et al., 1997), and in particular 4-hydroxyisoleucine, we have explored a new strategy using the oxazinone (El Achkar et al., 1998) derived from (1R,2R,5R)-2-hydroxypipnan-3-one as the starting material. By this route, enantiomerically pure isomers of (2R,3R,4R)-4-hydroxyisoleucine have been prepared (Kassem et al., 2001). The strategy used is outlined in the reaction Scheme below. We have recently described the molecular structures of compounds (II) and (IV) (Kassem et al., 2000). At that time, we did not possess crystals of compounds (III) and (V) of sufficient quality to be able to determine their structure and stereochemistry, and without this information it is difficult to obtain a full stereochemical understanding of the different steps in the synthetic route; a particular requirement is the structure of the key dihydro intermediate, (III).

During the enantioselective synthesis of 4-hydroxyisoleucine from the oxazinone, (I), three stereogenic centers were created, the key intermediate being the chiral dihydro amino acid derivative, (II), where hydrogenation of the double bond would allow control of the configurations of atoms C2 and C3. The configuration of atom C3 controls the stereochemistry of atom C4 after reduction of the ketone. Hence, it is necessary to determine the geometry of the double bond of (III) and the configuration of atom C4 in compound (IV).





For these intermediates, we found the R configuration for C2 and S for C3 in compound (II), and the R configuration for both C2 and C3 in compound (IV). To determine the stereochemistry unambiguously during the hydrogenation of (II) to (III), and the reduction step between (IV) and (V), the structures of two key intermediates, (III) and (V), were determined, and their structures are presented here.

The C2=C3 double bond was found to have the E configuration in compound (III). From the known R configuration of C7, C10 and C12, corresponding to the chiral auxiliary oxazinone, we found the R configuration for C2, C3 and C4 in compound (V).

Compounds (III) and (V) have different structures. The carbonyl group on C4 and the C2=C3 double bond in compound (III) were reduced in two steps, producing three asymmetric C atoms, viz. C2, C3 and C4, in compound (V). The six-membered C1-C2-N1-C8-C7-O3 ring of the oxazinone shares one side (C7-C8) with the bicyclo system of the (1R,2R,5R)-2-hydroxypipnan-3-one. On atom C2 is found the future amino acid side chain (C3-C4-C5).

The C1-C2-N1-C8-C7-O3 ring can be described as having a boat conformation, with slight distortion differences between (III) and (V). Atoms C2 and C7 are 0.2915 (17) and 0.5533 (17) Å, respectively, above the mean plane of the other four atoms (r.m.s. deviation = 0.065 Å) in (III). However, in (V), atoms C2 and C7 are 0.3347 (17) and 0.5163 (18) Å, respectively, out of the same mean plane (r.m.s. deviation = 0.066 Å). This difference is better indicated by the C8-N1-C2-C3 torsion angle, which is -158.28 (19)° in (III) and 155.44 (17)° in (V). In compound (III), a partial conjugation of the C2=C3 double bond with C8-N1 can be observed, but not with the carbonyl group of the ketone.

The bicyclo system, including the six-membered C7-C8-C9-C10-C11-C12 ring, is bridged by atom C13 between C12 and C10.



Experimental

The enantioselective condensation of butane-2,3-dione with (1R,2R,5R)-oxazinone (I) resulted in alcohol (II). After a dehydration step, a stereoselective hydrogenation of the double bond of compound (III) gave the second optically pure intermediate, (IV). After a reduction step, the final cleavage of the chiral auxiliary produced one pure isomer of 4-hydroxyisoleucine. Crystals of (III) (m.p. 375-377 K) and (V) (m.p. 382-384 K) suitable for single-crystal X-ray diffraction were grown from solutions in diethyl ether.

Compound (III)

Crystal data

- C16H25NO3
Mr = 275.34
Orthorhombic, P212121
a = 8.4160 (3) Å
b = 12.3127 (5) Å
c = 14.7039 (4) Å
V = 1523.72 (9) Å³
Z = 4
Dx = 1.200 Mg m⁻³
Mo Kα radiation
Cell parameters from 10 878 reflections
θ = 1.0-26.4°
μ = 0.08 mm⁻¹
T = 293 (2) K
Prism, colourless
0.4 × 0.3 × 0.2 mm

Data collection

- Nonius KappaCCD area-detector diffractometer
φ scans
10 878 measured reflections
1775 independent reflections
1668 reflections with I > 2σ(I)
Rint = 0.033
θmax = 26.4°
h = -10 → 10
k = -15 → 15
l = -17 → 17

Refinement

- Refinement on F²
R[F² > 2σ(F²)] = 0.038
wR(F²) = 0.100
S = 1.06
1775 reflections
181 parameters
H-atom parameters constrained
w = 1/[σ²(Fo²) + (0.0551P)² + 0.1091P] where P = (Fo² + 2Fc²)/3
Δσmax = 0.001
Δρmax = 0.13 e Å⁻³
Δρmin = -0.16 e Å⁻³

Table 1

Selected geometric parameters (Å, °) for (III)

Table with 2 columns: Bond/angle and Value. Rows include O1-C4, O2-C1, O3-C1, O3-C7, N1-C8, N1-C2, C1-C2, C2-C3, C3-C4, C3-C6, C4-C5, C7-C8, C1-O3-C7, C8-N1-C2, O2-C1-O3, O2-C1-C2, O3-C1-C2, C3-C2-N1, C3-C2-C1, N1-C2-C1, C2-C3-C4, C2-C3-C6, C4-C3-C6, O1-C4-C5, O1-C4-C3, C5-C4-C3, O3-C7-C8, N1-C8-C7, C8-N1-C2-C3, O3-C1-C2-C3, C1-C2-C3-C4, C2-C3-C4-O1, C2-C3-C4-C5.

Compound (V)

Crystal data

- C16H25NO3
Mr = 279.37
Orthorhombic, P212121
a = 10.2905 (3) Å
b = 11.8864 (5) Å
c = 12.8317 (5) Å
V = 1569.50 (10) Å³
Z = 4
Dx = 1.182 Mg m⁻³
Mo Kα radiation
Cell parameters from 11 088 reflections
θ = 1.0-26.3°
μ = 0.08 mm⁻¹
T = 293 (2) K
Prism, colourless
0.4 × 0.3 × 0.2 mm

Data collection

- Nonius KappaCCD area-detector diffractometer
φ scans
11 088 measured reflections
1791 independent reflections
1638 reflections with I > 2σ(I)
Rint = 0.036
θmax = 26.3°
h = -12 → 12
k = -14 → 14
l = -16 → 16

Refinement

- Refinement on F²
R[F² > 2σ(F²)] = 0.038
wR(F²) = 0.100
S = 1.06
1791 reflections
181 parameters
H-atom parameters constrained
w = 1/[σ²(Fo²) + (0.0677P)² + 0.1278P] where P = (Fo² + 2Fc²)/3
Δσmax = 0.001
Δρmax = 0.13 e Å⁻³
Δρmin = -0.12 e Å⁻³

Table 2

Selected geometric parameters (Å, °) for (V)

Table with 2 columns: Bond/angle and Value. Rows include O1-C4, O2-C1, O3-C1, O3-C7, N1-C8, N1-C2, C1-C2, C2-C3, C3-C4, C3-C6, C4-C5, C7-C8, C1-O3-C7, C8-N1-C2, O2-C1-O3, O2-C1-C2, O3-C1-C2, N1-C2-C1, N1-C2-C3, C1-C2-C3, C6-C3-C4, C6-C3-C2, C4-C3-C2, O1-C4-C5, O1-C4-C3, C5-C4-C3, O3-C7-C8, N1-C8-C7, C8-N1-C2-C3, O3-C1-C2-C3, C1-C2-C3-C4, C2-C3-C4-O1, C2-C3-C4-C5.


For both structures, the H atoms were introduced at calculated positions and refined as riding (O-H = 0.82 Å, and C-H = 0.96, 0.97 or 0.98 Å), with displacement parameters equal to 1.2 (OH, CH and CH2) or 1.5 (CH3) times that of the parent atom.

For both compounds, data collection: KappaCCD Reference Manual (Nonius, 1998b); data reduction: DENZO and SCALEPACK (Otwinowski & Minor, 1997); program(s) used to solve structure: SIR92 (Altomare et al., 1994); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: ORTEPII (Johnson, 1976); software used to prepare material for publication: maxus (Mackay et al., 1999).

Supporting information

Crystal structure: contains datablocks global, I. DOI: 10.1107/S0108270101014597/gd11665sup1.cif
Structure factors: contains datablock III. DOI: 10.1107/S0108270101014597/gd11665IIIsup2.hkl
Structure factors: contains datablock V. DOI: 10.1107/S0108270101014597/gd11665Vsup3.hkl

3D view



Supplementary crystallographic information

(III) 2,3,5,6,7,8-Hexahydro-6,8-methano-7,7,8a-trimethyl-3-(1-methyl-2-oxopropylidene)-5H-1,4-benzoxazin-2-one
(V) 2,3,5,6,7,8-Hexahydro-3-(2-hydroxy-1-methylpropyl)-6,8-methano-7,7,8a-trimethyl-5H-1,4-benzoxazin-2-one

Acknowledgements

We thank Professor M. Pierrot and M. Giorgi, Laboratoire de Bio-inorganique Structurale (UMR CNRS 6517), Faculté des Sciences de St. Jérôme, Marseille, France, for data collection and kind help with using MAXUS.

References

Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). J. Appl. Cryst. 27, 435.
El Achkar, A., Boumezbra, M., Roumestant, M. L. & Viallefont, P. (1988). Tetrahedron, 44, 5319-5332.
Jacob, M., Roumestant, M. L., Viallefont, P. & Martinez, J. (1997). Synlett, pp. 691-692.
Johnson, C. K. (1976). ORTEPII. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
Kassem, T., Rolland, V., Martinez, J. & Rolland, M. (2000). Acta Cryst. C56, 1037-1039.
Kassem, T., Rolland, V., Rolland, M., Roumestant, M. L. & Martinez, J. (2001). Tetrahedron Asymmetry, Accepted.
Mackay, S., Gilmore, C. J., Edwards, C., Stewart, N. & Shankland, K. (1999). maxus. Nonius BV, Delft, The Netherlands, MacScience Co. Ltd, Japan, and The University of Glasgow, Scotland.
Nonius (1998). KappaCCD Reference Manual. Nonius BV, Delft, The Netherlands.
Otwinowski, Z. & Minor, W. (1997). Methods in Enzymology, Vol. 276, Macromolecular Crystallography, Part A, edited by C. W. Carter & R. M. Sweet, pp. 307-326. London: Academic Press.
Sheldrick, G. M. (1997). SHELXL97. University of Göttingen, Germany.

© International Union of Crystallography. Prior permission is not required to reproduce short quotations, tables and figures from this article, provided the original authors and source are cited. For more information, click here.

We recommend

- Reexamination of some reactions of 3-(D-galactosylidenehydrazino)-1,2,4-triazolo[5,6-b]indole
Shaban, M. A. E., Die Pharmazie - An International Journal of Pharmaceutical Sciences, 2003
Synthesis and anxiolytic activity of 1-phenyl-2-(4-aryl-1,3,4,5-tetrahydropyrido[2,3-b][1,4]diazepin-2-ylidene)ethanone
Liszekiewicz, H., Die Pharmazie - An International Journal of Pharmaceutical Sciences, 2006
Synthesis and anti-inflammatory activity of some heterocyclic derivatives of phenothiazine
Sharma, S., Die Pharmazie - An International Journal of Pharmaceutical Sciences
Synthesis, chemical behavior, structure elucidation and iNOS inhibitory activity of 1-substituted 3-methylsulfanyl-5,6,7,8-tetrahydro-1H-[1,2,4]triazolo[1,2-a]p...
Schutz, U., Die Pharmazie - An International Journal of Pharmaceutical Sciences, 2017
Synthesis and antimicrobial testing of a new S-substituted-thiopyridines, thienopyridines, pyridothienopyridines and pyridothienotriazines
Abdel-Rahman, A. E., Die Pharmazie - An International Journal of Pharmaceutical Sciences, 2003