



**HAL**  
open science

## The interplay between abiotic and biotic factors in dispersal decisions in metacommunities

Mélanie Thierry, Julien Cote, Elvire Bestion, Delphine Legrand, Jean Clobert,  
Staffan Jacob

► **To cite this version:**

Mélanie Thierry, Julien Cote, Elvire Bestion, Delphine Legrand, Jean Clobert, et al.. The interplay between abiotic and biotic factors in dispersal decisions in metacommunities. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 2024, 379 (1907), 10.1098/rstb.2023.0137 . hal-04676422

**HAL Id: hal-04676422**

**<https://hal.science/hal-04676422v1>**

Submitted on 12 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# The interplay between abiotic and biotic factors on dispersal decisions in metacommunities

Mélanie Thierry<sup>1\*</sup>, Julien Cote<sup>2</sup>, Elvire Bestion<sup>1</sup>, Delphine Legrand<sup>1</sup>, Jean Clobert<sup>1†</sup> & Staffan Jacob<sup>1†</sup>

*Philosophical Transactions of the Royal Society B*, 2024, 379(1907):20230137, doi:10.1098/rstb.2023.0137

*Part of the Special Issue: “Diversity-dependence of dispersal interspecific interactions determine spatial dynamics”*

<sup>1</sup> Station d'Ecologie Théorique et Expérimentale, UAR2029, CNRS, 09200, Moulis, France.

<sup>2</sup> Centre de Recherche sur la Biodiversité et l'Environnement (CRBE), UMR 5300 CNRS-IRD-TINP-UT3  
Université Toulouse III – Paul Sabatier, Bât. 4R1, 118 route de Narbonne 31062 Toulouse cedex 9, France

\* Corresponding author: [melanie.thierry34@gmail.com](mailto:melanie.thierry34@gmail.com)

† Jean Clobert and Staffan Jacob share last authorship

**Key words:** Habitat choice, plasticity, biotic context, *Tetrahymena*, competition.

## Abstract

Suitable conditions for species to survive and reproduce constitute their ecological niche, which is built by abiotic conditions and interactions with conspecifics and heterospecifics. Organisms should ideally assess and use information about all these environmental dimensions to adjust their dispersal decisions depending on their own internal conditions. Dispersal plasticity is often considered through its dependence to abiotic conditions or conspecifics density, and to a lesser extent with the effects of interactions with heterospecifics, potentially leading to misinterpretation of dispersal drivers. Here, we first review the evidence for effects of, and the potential interplays between abiotic factors, biotic interactions with conspecifics and heterospecifics, and phenotype on dispersal decisions. We then present an experimental test of these potential interplays, investigating the effects of density and interactions with conspecifics and heterospecifics on temperature-dependent dispersal in microcosms of *Tetrahymena* ciliates. We found significant differences in dispersal rates depending on temperature, density, and presence of another strain or species. However, the presence and density of conspecifics and heterospecifics had no effects on the thermal-dependency of dispersal. We discuss the causes and consequences of the (lack of) interplay between the different environmental dimensions and the phenotype for metacommunity assembly and dynamics.

## Introduction

Dispersal is a major ecological process consisting in the emigration of individuals from one patch, their transition between patches, and their immigration to another one. Dispersal can lead to gene flow between habitats and changes in population demography and meta-population dynamics [1–3]. It is thus essential for ecological and evolutionary dynamics of populations, communities and ecosystems [4–7]. To leave unsuitable conditions and try to settle in more suitable ones, individuals can adjust their dispersal by behavioral changes at any of its steps based on ecological cues (named context-dependent dispersal, habitat choice or dispersal plasticity) [2,8–10], which can be associated with phenotypic changes [11,12]. Dispersal plasticity is observed in many taxa [10,13], with variability within and between species [14–16].

In spatially and temporally fluctuating environments, dispersal plasticity should allow organisms to track environmental conditions within which their fitness is maximized (*i.e.*, their ecological niche) [9,10,17–19]. Benefits of such adaptive dispersal plasticity should exceed potential costs induced by both dispersal [20] and plasticity of associated traits [21] (but see [14,22]). Trait plasticity is expected to evolve to adaptively respond to fluctuating and somewhat predictable environmental conditions when cues are available and reliable [23–28]. How cues are used should depend on how organisms perceive and integrate information in their environment [8,29]. Adaptive plasticity cannot be viewed as a unidimensional response, as organisms are exposed to a multitude of environmental factors [8,9,30–32]. Indeed, species ecological niche is constrained by the suite of abiotic conditions within which organisms can survive and reproduce (*i.e.*, the fundamental niche), biotic interactions, and dispersal barriers [33–35]. All the elements composing organism's ecological niche and their interplay are therefore important for the expression and adaptiveness of dispersal plasticity, but which cues are prioritized in dispersal decisions remains poorly understood. Disentangling effects of abiotic conditions and biotic interactions on the evolution of dispersal plasticity requires an integrative approach [36] in a metacommunity context [37], which has been rarely investigated so far.

Biotic interactions modify the fitness landscape in which species live, influencing species' evolutionary responses to environmental changes [38–41]. Thus, organisms cannot track environmental changes based solely on their expected performance in given abiotic conditions [42,43], but should modify their dispersal decisions

accounting for the effects of conspecifics and heterospecifics and their interactions with abiotic conditions on their fitness. For instance, optimal abiotic conditions may increase species abundances and thus strengthen competitive interactions. In that case, an adaptive strategy could be to avoid such competitive habitats by choosing sub-optimal conditions, potentially resulting in niche partitioning [44]. For instance, in the ciliate *Tetrahymena thermophila*, an organism able to differentiate kin from non-kin [45], the most generalist genotypes have been found to disperse toward sub-optimal habitats, which has been hypothesized as a way to avoid competition with specialists [22]. Further, dispersal plasticity of one focal species can have cascading effects on the population dynamics of the other trophic levels [46] and strongly affects metacommunity diversity at both local and regional scales that depend on both consumer and resource dispersal [47].

To fully understand the causes and consequences of dispersal plasticity in metacommunities, we need to understand which cues are used for dispersal decisions, whether organisms change their use of abiotic cues depending on their biotic context (both intra- and interspecific), and what is the subsequent interplay between abiotic and biotic factors and the organism's phenotype on dispersal decisions. To address this knowledge gap, we first review existing evidence for the role of abiotic, intraspecific and interspecific contexts and their interplay on dispersal. We do not aim to be exhaustive but rather to highlight how the different dimensions of an organism's environment, and their interactions, may affect its dispersal decisions. We then experimentally explore these interplays in actively dispersing ciliates of the genus *Tetrahymena* [22]. Finally, we propose key perspectives for future research in metacommunity ecology, aiming to better understand the effects of the multidimensionality of the environment on dispersal decisions for metacommunity assembly and dynamics. We emphasize the importance of individual variation to gain insight into eco-evolutionary dynamics in metacommunities.

## **The evidence for the role of abiotic, intraspecific and interspecific contexts, and their interplay on dispersal**

### **Use of information in dispersal decisions**

In the case of informed dispersal, organisms need to perceive and process information about their environment to decide whether or not to disperse and to settle in another habitat [29,48,49], potentially in relation with their

phenotype (physiology, behavior, morphology and life-history traits) [9,50,51]. The external information acquired can be abiotic (*e.g.*, temperature, humidity, light), or biotic (*e.g.*, conspecifics density, presence and density of heterospecifics like competitors, resource/prey or predators/parasites) [52], and can be acquired via visual, auditory, olfactory, chemical, or haptic cues. Information can also be transmitted by ascendants [53,54], or other (unrelated) individuals. Abiotic conditions can drastically influence organism's fitness. Adaptive dispersal plasticity following changes in abiotic conditions should allow organisms to track abiotic conditions in which they live and thrive (*i.e.*, their fundamental niche) [14]. Population density, involving either kin or non-kin, is an important biotic factor for ecological processes, including dispersal [55]. Conspecific density-dependent dispersal has been observed in many taxa including plants [56], birds [57], lizards [58], mammals [59,60], and ciliates [16]. For the information coming from heterospecific presence and density, how organisms perceive and use it likely depends on the type and strength of biotic interactions [37,61]. In the case of mutualistic interactions, dispersers may prefer to join an occupied patch rather than to disperse toward an empty habitat [62]. In the case of competitive or trophic interactions, dispersers may avoid strong levels of competition or predation. Trophic interactions lead to fluctuations between prey and predator densities [63], making disentangling density-dependent dispersal from direct effects of interactions with heterospecifics on dispersal decisions difficult [64]. We highlight in Table 1 a few empirical evidences of dispersal plasticity along these three key environmental dimensions: abiotic and biotic context, both intra- and interspecific. We did not aim to review the literature exhaustively here, as it has previously been done [8,60]. We however aimed to illustrate the environmental dimensions experienced by an organism that can modify dispersal, thus presenting effect of the different types of biotic interactions with very different underlying mechanisms.

### **The relative importance and interplay between abiotic and biotic dimensions of the environment**

Organisms likely integrate multiple cues to fine-tune their dispersal decisions in temporally and spatially fluctuating environment [65]. However, little is known on how they process these multiple information sources, whether, some cues are prioritized, and under which conditions they have or not interactive effects on dispersal decisions [66,67]. Whether effects of abiotic conditions and interactions with conspecifics and heterospecifics on dispersal decisions have interactive, additive or compensatory effects thus remains an open question.

Optimality in information use should result from a trade-off between the cost of information acquisition and the benefits resulting from informed dispersal [5,68,69]. Cues do not have to be the direct environmental conditions selecting for adaptive dispersal plasticity, but should correlate with the selective environment [70], at least partially [71]. Expressed dispersal plasticity may not be seen as adaptive when considering one dimension of the environment if the fitness responses to abiotic context and biotic interactions go in different directions. For instance, the presence of competitors constrained the adaptive response of the algae *Chlamydomonas reinhardtii* to an increase in CO<sub>2</sub> level [72], changed both the plastic and evolutionary responses of *Arabidopsis thaliana* to drought [40], and hindered evolutionary changes in cell shape and velocity of the protist *Paramecium aurelia* – two traits linked to dispersal propensity – in response to change in salinity [73]. While these studies looked at the plasticity of other traits than dispersal, they show that an integration of the different abiotic and biotic elements of the environment in studies of dispersal plasticity may be essential.

Densities of prey, predators and competitors are not independent. Hence, effects of intra- and interspecific contexts on dispersal decisions likely interact [74]. However, this topic still requires experimental validation. After detailed literature searching, among the studies investigating interspecific effect on dispersal decisions, 5 out of 95 for host-parasite, 22 out of 208 for prey-predator, 4 out of 91 for resource-consumer, and 20 out of 108 investigated the interplay between intra- and interspecific interactions (Bestion *et al.* this issue). The relative proportion of their effects, when additive, may also be crucial to understand metacommunity dynamics. Indeed, the balance between competitive intra- and interspecific interactions is important for species coexistence in community [75] but less considered in the metacommunity framework (but see [76]). We could expect stronger positive density-dependence of dispersal with conspecifics than heterospecifics because intraspecific competition may be stronger than interspecific competition for species to coexist [75]. However, the reverse could also hold if the competitive species is a better competitor, with high heterospecific densities, or if the niche overlap between heterospecifics is strong. We may predict that the relative importance of intra- vs. interspecific interactions and the sign of their relationship with dispersal depend of niche overlap.

Effects of interactions with conspecifics and heterospecifics on dispersal decisions should be tightly coupled with abiotic conditions. Indeed, the abiotic context can change species carrying capacity [77], the way species interact within a habitat because it changes their niche [78,79], and the outcome of species interactions

[80–82]. The abiotic context may thus change the effect of conspecifics and heterospecifics presence and density on dispersal decisions, and *vice versa*. For instance, inferior competitors may avoid species with whom they would strongly compete (*i.e.*, with a strong niche overlap) by choosing a sub-optimal habitat that would maximize their fitness when considering both the abiotic and biotic dimensions of the habitat [22]. As conspecific density can be used as proxy for abiotic conditions, disentangling density- and abiotic-dependency of dispersal and which cues may be prioritized can be challenging. Moreover, the dispersal decisions resulting from the interplay between one biotic and one abiotic factor could change depending on the variation of another factor [83–86].

Some phenotypes influence the competitiveness and optimality of organisms in a given environment. Therefore, in addition to interactive effects of abiotic and biotic factors on dispersal decisions, the environment and the phenotype can also interact [87] (Fig.1). An individual's phenotype likely plays a major role in how it uses information of its environment, both biotic and abiotic [9,50,51]. Further, phenotypic differences between dispersers and residents are widespread across taxa [8], and can be affected by environmental factors depending on the associated costs of dispersal [88].

We illustrate the different interplays between these three axes of external conditions (abiotic, conspecifics, and heterospecifics) and the internal conditions with selected evidence in Table 1.

### **The interplay between all dimensions of the environment**

Studies integrating the interplay of two or more dimensions of the environment on dispersal plasticity are scarce although interactive effects between all dimensions of the environment and the phenotype is probably the rule rather than the exception. It would require very large experimental designs to have enough statistic power to test more than two-way interactions between environmental factors. Short-distance dispersal of the predator *Orius minutus* was affected by both weather conditions and intra- and inter-trophic densities in the field, but the interactive effects between abiotic and biotic factors were not tested [89]. Emigration of the stream fish *Rivulus hartii* was positively correlated to water level and body size, but only when the predator *Hoplias malabaricus* was present [90]. The mesocosm experiment using butterflies mentioned above [67] presents ten dimensions including weather conditions, conspecifics density, habitat quality, and several phenotypic traits. However, they only tested two-way interactions between the most significant environmental factors and phenotypic factors as higher order interactions require statistical power that even this highly controlled mesocosm experiment was not able to meet.

We can expect, for two hypothetical co-occurring species having partial niche overlap in a two-dimensional space (Fig. 2a), that their dispersal decisions along these environmental axes would depend on their optimum and performance relative to the other species (Fig. 2b) [22]. This would translate into an interactive effect between the abiotic factors and the biotic context on dispersal (Fig. 2c). An organism's dispersal decision should thus depend on its niche and its competitive abilities compared to other organisms with whom it shares its habitat.

### **Case study: interplay of abiotic and biotic conditions on dispersal in *Tetrahymena***

To illustrate such dependency of dispersal decisions to an organism's performance and biotic interactions, we present below a case study using microcosm with *Tetrahymena* species. We investigated the dependence of dispersal decisions to thermal conditions, biotic context, and their potential interplay in light of their thermal niche and competitive abilities. We manipulated the biotic context as conspecific population density and presence/absence of conspecifics or heterospecifics, and quantified the consequences for the decision to leave a habitat (emigration step of dispersal) depending on thermal local conditions.

We used two strains for three *Tetrahymena* species (D2 and D3 for *T. thermophila*, A3 and A5 for *T. americanis*, P2 and P3 for *T. pyriformis*; Table S1), 20 to 50  $\mu\text{m}$  ciliates naturally living in freshwater ponds and streams [91]. Evidence for thermal-dependent dispersal and important intra- and interspecific variations in the direction and magnitude of the response has been provided in previous studies [14,22,92,93]. We created a design comprising in total six monocultures, three intraspecific cocultures (D2/D3, A3/A5, P2/P3) and three interspecific cocultures (D3/A5, D3/P3 and A5/P3, *i.e.*, one focal strain per species for logistical reasons). Cells were maintained in the laboratory in 24-wells plates in 2mL of axenic liquid growth media (0.6% Difco proteose peptone, 0.03% yeast extract, 0.006% glucose, 0.0009% sequestrene Fe-EDTA) at 23°C. All manipulations were performed in sterile conditions under a laminar flow hood. We prepared cultures in Erlenmeyer flasks with 0.5 mL of the stock culture in 50 mL of media five days before the start of the experiments to reach densities close to asymptotic density.

We first measured the thermal niche of each strain by following their growth at eight temperatures (11, 15, 19, 23, 27, 31, 35 and 39°C) for eleven days, with six replicates per strain and temperature. The thermal optimum corresponds to the temperature for which population growth is maximal (Fig. S1). *T. americanis* strains



(A3 and A5) have their thermal optimum close to 23°C (22.7 and 22.2°C respectively), *T. pyriformis* strains (P2 and P3) have their optimum slightly below it (21.6 and 21.2 respectively), and *T. thermophila* strains (D2 and D3) above 23°C (27.6 and 25°C respectively). For all of them, 31°C is above their thermal optimum, but either within (D2, D3) or outside (A3, A5, P2, P3) the 80% performance zone (Fig. S1). Based solely on their thermal niche, we could therefore expect *T. americanis* and *pyriformis* strains to avoid 31°C, a temperature that negatively impact their fitness, while *T. thermophila* strains should be less impacted by an increase in temperature from 23 to 31°C.

Then, we assessed potential intra- and interspecific interactions (either competition or facilitation) for each pair of strains. To do so, we inoculated 24-well plates by diluting the five-days old cultures to constitute monocultures of each strain with ~10,000 cells/mL, and co-cultures of each of the six pairs with ~10,000 cells/ml of each strain (~20,000 cells/mL in total). We placed these 24-well plates at two temperatures: 23°C, the standard temperature at which strains are maintained at the laboratory, and 31°C, resulting in 24 experimental conditions, each replicated six times (N = 144). We assessed biotic interactions for each pair in each temperature by comparing the sum of the monocultures with the abundance of cells in cocultures measured with flow cytometry (Supplement Text 1) after five days following [94]. If the abundance of cells in coculture is inferior to the sum of the monocultures this indicates competitive interaction, if it is superior facilitation, and if it is equal neutral interaction. Error rates to discriminate between strains and species are presented in Table S2. Despite intra- and interspecific differences in performance at 23 and 31°C (Fig. S1), all strains decreased in abundance when in co-culture (Fig. S2). Biotic interactions at 23°C were always competitive while biotic interactions at 31°C were either competitive or neutral (Fig. S2). There were no marked differences between intra- and interspecific competition, reflecting an overall similar effect on fitness. The type of biotic interaction is thus expected to have no significant effect on dispersal. D3 and P3 strains had significant lower abundances in cocultures of conspecifics than in monocultures at 23°C while their conspecific strain had no significant decrease in abundance. We could therefore expect dispersal of these strains to be affected by presence of their conspecific strain (D2 and P2 respectively). A5 and D3 paired with P2 had significant lower abundances in cocultures of heterospecifics than in monocultures at 23°C, meaning that P2 strain seems to be a stronger heterospecific competitor than these two strains at 23°C. We could therefore expect this decrease in fitness of A5 and D3 to be reflected on their dispersal decisions when paired with P2. However, these declines in abundance of D3 and P3 with conspecific strains and of A5 and D3 paired with P2

became non-significant at 31°C. At 31°C, the effect of biotic interaction on dispersal may thus become non-significant if dispersal decisions depend on the fitness consequences of biotic interactions.

Finally, we tested the effect of density and the presence of conspecifics vs. heterospecifics on thermal dispersal plasticity. We used standard 1.5 mL two-patches systems where two 1.5mL Eppendorf tubes are connected by a corridor (4 mm internal diameter, 2.5 cm long silicone tube), the whole system being filled with growth medium [16]. As for to the biotic interaction assessment experiment, we inoculated the start patches of the systems with monocultures of each strain with either 10,000 cells (low density) or 20,000 cells (high density) and co-cultures of the six pairs with 10,000 cells of each strain (20,000 cells in total). Each combination of strains and density was placed at either 23 or 31°C. We replicated each of the resulting 36 conditions six times in three block days (N = 216). After a 30 min acclimation at the corresponding temperature, we allowed cells to disperse for three hours to remain below the minimum generation time at the tested temperatures, but still letting sufficient time for cells to disperse. We then closed the corridors to prevent further movements and sampled residents (start patch) and dispersers (immigration patch). We estimated total and relative abundance of each strain in samples with flow cytometry (Supplement Text 1). Error rates to discriminate between strains and species are presented in Table S3. They were low enough to not change the estimated abundances. We calculated dispersal rate as  $\frac{N_{disp}}{N_{resi} + N_{disp}}$ , where  $N_{disp}$  and  $N_{resi}$  are the abundance of dispersers and residents, respectively. Statistical analyses are described in Supplement Text 2.

In monocultures, we found evidence for thermal dispersal plasticity ( $\chi^2_{(1)} = 6.30$ , p-value = 0.012; Fig. 3a&d) and density-dependent dispersal ( $\chi^2_{(1)} = 5.86$ , p-value = 0.016; Fig. 3a&d), as previously described [14,16,22,92,93,95]. Thermal dispersal plasticity varied between strains in terms of magnitude and direction (temperature x strain:  $\chi^2_{(5)} = 27.46$ , p-value < 0.0001; Fig. 3d; Table S4&5), but the direction of the effect did not follow what we could have expected according to their thermal niche. Indeed, strains of a same species have very similar thermal niche (Fig. S1) while their dispersal response to temperature were in opposite direction (Fig. 3a&d). Density-dependent dispersal was always positive when significant (density x strain:  $\chi^2_{(5)} = 5.51$ , p-value = 0.36; Fig. 3d; Table S5). However, we found no significant interactive effect between temperature and

monoculture density ( $\chi^2_{(1)} = 0.90$ , p-value = 0.342; Table S4), meaning that the density dimension of the biotic context had no significant effect on thermal dispersal plasticity (Fig. 3a&g).

Looking at dispersal response in monocultures at high density and cocultures, we found that the effects for both temperature and the presence of another strain (either conspecific or heterospecific) varied depending on the strain (temperature x strain:  $\chi^2_{(5)} = 20.38$ , p-value = 0.0011; presence/absence of another strain x strain:  $\chi^2_{(5)} = 39.33$ , p-value < 0.0001; Fig. 3b&e; Table S6&7). Three out of the six strains tested showed a decrease in dispersal rate with presence of another strain (A3, D3 and P2), two showed an increase in dispersal with presence of another strain (D2 and P3), and one strain was not significantly affected by the presence of another strain nor by temperature (A5) (Fig. 3e). Interestingly, the strain P3, that showed no dispersal plasticity to temperature despite very low performance at 31°C compared to 23°C (Fig. S1), had significant changes in its dispersal rate between monoculture and coculture (Fig. 3e; Table S7). Further, despite change in biotic interactions with temperature (Fig. S2), temperature did not modify the effect of presence or absence of another strain on dispersal ( $\chi^2_{(1)} = 0.08$ , p-value = 0.781; Fig. 3h; Table S6).

We found that conspecific vs. heterospecific biotic interactions had no significant effect on dispersal rate and its thermal plasticity (interaction type:  $\chi^2_{(1)} = 0.36$ , p-value = 0.55; strain x interaction type:  $\chi^2_{(2)} = 0.51$ , p-value = 0.77; Fig. 3c&f&i; Table S8&9). This was in line with the similar overall effect of interaction type observed on fitness (Fig. S2). However, effect of competitors on dispersal rates varied depending on their identity for A5 and D3 (but not for P2; Fig. S3; Table S10), without an interactive effect with temperature (Table S10). Surprisingly, effects of P2 strain compared to the other heterospecific strain and conspecifics on dispersal were opposite between A5 and D3 despite that both had a decrease in fitness in coculture of P2 compared to monocultures (Fig. S2).

Overall, our results show that temperature and the biotic context independently affect dispersal decisions in *Tetrahymena*, with both intra- and interspecific variations in their effect. Further, the direction of the effects of temperature and the biotic context could not have been predicted by effect on fitness from their thermal niche and the biotic interaction assessment.

## Discussion

Our literature overview and case study highlight that, to understand how environmental changes affect dispersal decisions, we need a deeper consideration of both biotic and abiotic environmental factors and their incorporation in the context of metacommunities. In the literature, we found several studies investigating the additive and interactive effects of the multiple dimensions of the environment on dispersal decisions. Integrating effects of more than two environmental axes is however technically challenging, which explains the scarcity of evidence. In our study case using microcosms of *Tetrahymena* species, effects of temperature and the biotic context on dispersal did not interact significantly. Further, we observed important intra- and interspecific variations in terms of magnitude and direction of dispersal response to temperature and the biotic context despite similar direction in their effect on fitness. Below, we propose perspectives on the causes and consequences of presence or absence of interplay between the multiple dimensions of the environment and the phenotype on dispersal plasticity expression and evolution for metacommunity dynamics.

### **Dispersal plasticity evolution and expression along multiple environmental axes**

We highlighted in the first sections the importance of considering interactions between multiple environmental factors to understand the evolution and expression of dispersal plasticity by reviewing the different interplays possible between the three external environmental context types (abiotic, conspecifics, heterospecifics). Based on our literature survey, we expected in our case study that the biotic factors (conspecific density, presence of another strain, and biotic interaction type) would interact with thermal effect on dispersal. This was not the case. Both temperature and biotic interactions (presence of another strain) significantly changed dispersal decisions, but their effects were additive. This was surprising considering the decrease in competition observed with an increase in temperature. The expected interaction between the abiotic and biotic context on dispersal decisions may simply not hold in all cases and/or for all organisms, and likely depends on the degree of niche overlap between organisms. A large experimental study on 21 species ranging from protozoa to vertebrates found general effects of resource availability and predation risk on emigration decisions [46]. Effects of resource availability in this study could be similar than effects of density or competition. We could hope for similar generality of the abiotic- and biotic-dependency of dispersal, which may help forecasting metacommunity dynamics in a changing world. Expanding our comprehension of the interplay (or lack of) between abiotic conditions and interactions with conspecific and

heterospecifics (both within and between trophic levels) on dispersal plasticity, and the consequences for metacommunities, will require further systematic investigation across environmental contexts and taxa.

Here, we investigated effects and interplay of abiotic and biotic factors over a single generation. The interplay between different environmental dimensions may not act on dispersal plasticity, but on its evolution. Indeed, organisms may have previously adapted their dispersal decisions based on the fitness landscape combining both abiotic and biotic conditions, and what is observed may be the result of niche partitioning [44]. We may expect organisms free of competition or predation (*e.g.*, during range expansion or invasion) to evolve their dispersal decisions without the biotic pressure exerted by other species (illustrated by the differences in dispersal reaction norms on Fig. 2c&d). This is often observed in the realized niche shifts of invasive species free of their native competitors [96] and natural enemies [97]. Experimental evolution integrating dispersal with a factorial design combining effects of abiotic factors and biotic context could tell us under which conditions interactive effects could emerge as drivers of dispersal plasticity evolution [36].

### **High intra- and interspecific variations in dispersal response**

Dispersal rates were always positively correlated with conspecific density independently of the abiotic conditions. Another study found a similar positive effect of density of conspecifics or heterospecifics on nematodes' dispersal in laboratory independent of the level of salinity [98]. However, dispersal response to the other external factors studied (temperature, presence of another strain, and competitor identity) varied significantly in terms of both magnitude and direction across strains and species despite a same direction of their effect on fitness. Therefore, even with a lack of interplay between the abiotic and biotic contexts, it might be difficult to predict how organisms would disperse depending on the fitness landscape. These variations in dispersal response may be linked to high intra- and interspecific variations observed in phenotypic traits and depending on the dispersal status (morphological and movement traits; Fig. S4). This highlights the importance of the interplay between external factors and the phenotype for dispersal decisions. The studies discussed in the previous section about the interplay between external and internal conditions found generalities in this interplay when considering personality (*e.g.*, for boldness [99,100]), starvation [101], and life stage [102]. These generalities become harder to make when talking about phenotypic traits that are also plastic to these external factors as in our case study.

### **Importance of intra- vs. interspecific interactions in metacommunities**

We did not find differences between the effect of conspecifics and heterospecifics presence and density on dispersal rates nor on thermal dispersal plasticity in our case study. They are however able to differentiate themselves from other clonal lineages [45], and another study found that *T. thermophila* is able to change its dispersal decisions based on biotic cues coming from kin or non-kin [103]. Therefore, in our experiment, cells likely differentiate cues coming from conspecifics and heterospecifics, but they translated into a similar dispersal response because intra- and interspecific interactions had similar effects on fitness. Such generic response to potential competitors has been observed in the propensity to move in flour beetles [104]. We tested the effects of biotic interactions during the emigration phase only. Effects in the settlement phase may be different because of different associated costs [106] or different cues potentially prioritized [62].

## **Conclusion and perspectives**

We have discussed the causes and consequences of the interplay (or lack of) between the environmental dimensions and the phenotype in metacommunities. Comparative and integrative approaches are essential to build our knowledge on the multi-cause and consequences of dispersal decisions. How and why organisms decide to disperse based on the interplay between the different dimensions of the environment and their internal conditions is particularly important to forecast the impact of ongoing climate change, and to advance our understanding on biological invasions and disease biogeography. In this section, we discuss timely questions considering dispersal decisions that integrate the multiple dimensions of the environment.

### **How important individual variations are in dispersal plasticity for metacommunity dynamics?**

Ecological and evolutionary processes such as dispersal and niche partitioning are often looked at the population or community scales, but looking at the individual scale allows to merge ecological and evolutionary processes [107]. The ecological niche an individual occupies can be mediated by its phenotype and makes explicit the intraspecific context as a whole dimension of the niche [108]. Each individual can thus occupy a specific niche space that can be different from another individual of a same species, leading to intra-specific variations in performance along environmental axes, and therefore differences in dispersal decisions [109]. Individual variations can be determinant in metacommunity dynamics because they impact both dispersal decisions [110]

and the strength and outcome of biotic interactions [111]. If individuals adaptively disperse depending on their fitness landscape, we could expect a spatial sorting of the metacommunity that would maximize niche partitioning. At the landscape scale, this would translate into a spatial homogenization of fitness among individuals and of trophic and competitive impacts on other species, which should stabilize the metacommunity [112]. Although not possible with our experimental design, methods exist to individually track dispersers for our biological system and others [113], both in the laboratory and in the field. This would allow for a deeper understanding of the link between phenotype and dispersal decisions and its consequences for populations and communities. Such methodological and conceptual advances could bring a deeper understanding on how individual variations in dispersal decisions would scale up to metacommunity dynamics and ecosystem functioning [87].

### **Dispersal syndromes: a key driver of the interplay between abiotic conditions and biotic interactions on dispersal decisions?**

Dispersal is often associated with a suite of traits (*i.e.*, dispersal syndrome) [9,114]. Dispersers are generally larger than residents [11,12], which may be linked to the level of energy reserves required to overcome costs of dispersal [20]. Dispersers also often have a more locomotion-oriented morphology than residents to move faster and farther (*e.g.*, bigger wings, longer legs, elongated body shape). This dispersal syndrome was observed to be stronger under predation risk across taxa [12]. Dispersal decisions likely depend on the balance between dispersal costs and local predation and competition strength [12]. This balance could be dictated for instance by the relationship between dispersal and body size. Indeed, both dispersal propensity and interaction strengths can be mediated by body size [115]. In our dispersal experiment, we could not characterize phenotypic traits at the strain level in competitive treatments with two strains. However, the relationship between dispersal status and cell size was either positive, negative or null depending on the strains and thermal conditions at the community scale and in monocultures (Figure S4). These intra- and inter-specific variations were found in previous studies on *Tetrahymena* species [14,95] and other species [11]. This variability in dispersal syndromes can have important implications for metapopulation dynamics [95]. The lack of a univocal relationship between dispersal and phenotypes suggests the existence of a range of dispersal strategies (*e.g.*, [116]) and would require further investigation.

### **What would be the consequences of dispersal plasticity for metacommunities under global change?**

With the ongoing global change, abiotic conditions are changing and induce numerous phenotypic changes including life-history traits. These individual changes alter community composition and biotic interactions, with range shifts, reshufflings of entire species communities [117–119] and desynchronization of previously interacting species [120–122], which feedback to individual evolutionary responses [123]. These abiotic and biotic changes modify the fitness landscape in which dispersal plasticity has evolved. By overlooking unpredictable novel biotic interactions and differences in dispersal response among individuals we are at risk of underestimating the impact of climate change on biodiversity and ecosystem functioning [41,118,124,125]. Moreover, environmental factors impacting organisms' fitness (*e.g.*, temperature) are increasingly variable and fluctuate in an unpredictable manner [126]. Future climate variability may therefore prevent organisms to adaptively track favorable conditions through dispersal plasticity, which would lead to a destabilization of the whole meta-community. Landscape fragmentation is also increasing [127], preventing free movement between habitats [128], and therefore limiting the potential for spatial sorting through adaptive dispersal and so its potential stabilizing effects. Further, dispersal plasticity could become maladaptive with changing climate if the cues used for adaptive dispersal behavior became decoupled with the selective conditions [129]. This is especially true when plastic capacity depends on developmental conditions [130], which would induce a mismatch between adult dispersal and environmental conditions [131]. With climate change, cues to detect environmental changes may not be reliable anymore, and bet-hedging (or other unconditional strategies) may be better evolutionary strategy than plasticity in some cases [132].

**Data accessibility.** All raw data used for this study are available from the Zenodo database: [10.5281/zenodo.8406074](https://doi.org/10.5281/zenodo.8406074).

**Authors' contributions.** MT, JCl and SJ conceived the project; MT performed the experiments; MT analyzed the data and led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

**Competing Interest.** The authors declare no competing interests.



**Acknowledgments.** We thank Michèle Huet for the maintenance of cell cultures and technical support, Pablo Molina Vila for technical support, Amélie Cocchiara for her help in the lab to perform the experiments, and Léonard Dupont for fruitful discussions and his help to fit the thermal niche.

**Funding.** This work benefitted from financial support by the Agence Nationale de la Recherche for the project CHOOSE (ANR-19-CE02-0016), and is part of TULIP (Laboratory of Excellence Grant ANR-10 LABX-41). JCo is supported by a funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement no. 817779).

## References

1. Clobert J, Danchin E, Dhondt AA, Nichols JD, Clobert J, Danchin E, Dhondt AA, Nichols JD, editors. 2001 *Dispersal*. Oxford, UK: Oxford University Press.
2. Clobert J, Ims RA, Rousset F. 2004 Causes, Mechanisms and Consequences of Dispersal. In *Ecology, Genetics and Evolution of Metapopulations* (eds I Hanski, OE Gaggiotti), pp. 307–335. Burlington: Academic Press. (doi:10.1016/B978-012323448-3/50015-5)
3. Ronce O. 2007 How does it feel to be like a rolling stone? Ten questions about dispersal evolution. *Annual review of ecology, evolution, and systematics* **38**, 231–253. (doi:10.1146/ANNUREV.ECOLSYS.38.091206.095611)
4. Clobert J, Baguette M, Benton TG, Bullock JM. 2012 *Dispersal Ecology and Evolution*. Oxford, UK: Oxford University Press.
5. Delgado MM, Bartoń KA, Bonte D, Travis JMJ. 2014 Prospecting and dispersal: their eco-evolutionary dynamics and implications for population patterns. *Proceedings of the Royal Society B: Biological Sciences* **281**, 20132851. (doi:10.1098/rspb.2013.2851)
6. Hanski I, Mononen T. 2011 Eco-evolutionary dynamics of dispersal in spatially heterogeneous environments. *Ecology Letters* **14**, 1025–1034. (doi:10.1111/j.1461-0248.2011.01671.x)
7. Kubisch A, Holt RD, Poethke H-J, Fronhofer EA. 2014 Where am I and why? Synthesizing range biology and the eco-evolutionary dynamics of dispersal. *Oikos* **123**, 5–22. (doi:10.1111/j.1600-0706.2013.00706.x)
8. Bowler DE, Benton TG. 2005 Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. *Biological Reviews* **80**, 205–225. (doi:10.1017/S1464793104006645)
9. Clobert J, Le Galliard JF, Cote J, Meylan S, Massot M. 2009 Informed dispersal, heterogeneity in animal dispersal syndromes and the dynamics of spatially structured populations. *Ecology Letters* **12**, 197–209. (doi:10.1111/J.1461-0248.2008.01267.X)
10. Edelaar P, Siepielski AM, Clobert J. 2008 Matching habitat choice causes directed gene flow: a neglected dimension in evolution and ecology. *Evolution* **62**, 2462–2472. (doi:10.1111/J.1558-5646.2008.00459.X)
11. Stevens VM *et al.* 2014 A comparative analysis of dispersal syndromes in terrestrial and semi-terrestrial animals. *Ecology Letters* **17**, 1039–1052. (doi:10.1111/ele.12303)

12. Cote J *et al.* 2022 Dispersal syndromes in challenging environments: A cross-species experiment. *Ecology Letters* **25**. (doi:10.1111/ele.14124)
13. Arendt JD. 2015 Effects of dispersal plasticity on population divergence and speciation. *Heredity* **115**, 306–311. (doi:10.1038/hdy.2015.21)
14. Campana JLM, Raffard A, Chainé AS, Huet M, Legrand D, Jacob S. 2022 Dispersal plasticity driven by variation in fitness across species and environmental gradients. *Ecology Letters* **25**, 2410–2421. (doi:10.1111/ele.14101)
15. De Meester N, Bonte D. 2010 Information use and density-dependent emigration in an agrobiont spider. *Behavioral Ecology* **21**, 992–998. (doi:10.1093/beheco/arq088)
16. Pennekamp F, Mitchell KA, Chainé A, Schtickzelle N. 2014 Dispersal propensity in *tetrahymena thermophila* ciliates - a reaction norm perspective. *Evolution* **68**, 2319–2330. (doi:10.1111/EVO.12428)
17. Fronhofer EA, Nitsche N, Altermatt F. 2017 Information use shapes the dynamics of range expansions into environmental gradients. *Global Ecology and Biogeography* **26**, 400–411. (doi:10.1111/geb.12547)
18. Holt RD. 1987 Population dynamics and evolutionary processes: the manifold roles of habitat selection. *Evolutionary Ecology 1987 1:4* **1**, 331–347. (doi:10.1007/BF02071557)
19. Pellerin F, Cote J, Bestion E, Aguilée R. 2019 Matching habitat choice promotes species persistence under climate change. *Oikos* **128**, 221–234. (doi:10.1111/oik.05309)
20. Bonte D *et al.* 2012 Costs of dispersal. *Biological Reviews* **87**, 290–312. (doi:10.1111/j.1469-185X.2011.00201.x)
21. Moran NA. 1992 The evolutionary maintenance of alternative phenotypes. *The American Naturalist* **139**, 971–989. (doi:10.1086/285369)
22. Jacob S *et al.* 2018 Habitat choice meets thermal specialization: competition with specialists may drive suboptimal habitat preferences in generalists. *Proceedings of the National Academy of Sciences of the United States of America* **115**, 11988–11993. (doi:10.1073/PNAS.1805574115)
23. Botero CA, Weissing FJ, Wright J, Rubenstein DR. 2015 Evolutionary tipping points in the capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 184–189. (doi:10.1073/PNAS.1408589111)
24. Fryxell JM. 1997 Evolutionary dynamics of habitat use. *Evolutionary Ecology* **11**, 687–701. (doi:10.1023/A:1018434302138)
25. Lande R. 2014 Evolution of phenotypic plasticity and environmental tolerance of a labile quantitative character in a fluctuating environment. *Journal of Evolutionary Biology* **27**, 866–875. (doi:10.1111/JEB.12360)
26. Leung C, Rescan M, Grulois D, Chevin LM. 2020 Reduced phenotypic plasticity evolves in less predictable environments. *Ecology Letters* **23**, 1664–1672. (doi:10.1111/ELE.13598)
27. Reed TE, Robin SW, Schindler DE, Hard JJ, Kinnison MT. 2010 Phenotypic plasticity and population viability: the importance of environmental predictability. *Proceedings of the Royal Society B: Biological Sciences* **277**, 3391–3400. (doi:10.1098/RSPB.2010.0771)

28. Tufto J. 2015 Genetic evolution, plasticity, and bet-hedging as adaptive responses to temporally autocorrelated fluctuating selection: A quantitative genetic model. *Evolution* **69**, 2034–2049. (doi:10.1111/evo.12716)
29. Enfjäll K, Leimar O. 2009 The evolution of dispersal – the importance of information about population density and habitat characteristics. *Oikos* **118**, 291–299. (doi:10.1111/j.1600-0706.2008.16863.x)
30. Matthysen E. 2012 Multicausality of dispersal: a review. In *Dispersal Ecology and Evolution*, pp. 3–12. Oxford, UK: Oxford University Press.
31. Sgrò CM, Terblanche JS, Hoffmann AA. 2016 What can plasticity contribute to insect responses to climate change? *Annual Review of Entomology* **61**, 433–451. (doi:10.1146/annurev-ento-010715-023859)
32. Singh P, van Bergen E, Brattström O, Osbaldeston D, Brakefield PM, Oostra V. 2020 Complex multi-trait responses to multivariate environmental cues in a seasonal butterfly. *Evol Ecol* **34**, 713–734. (doi:10.1007/s10682-020-10062-0)
33. Hutchinson GE. 1957 Concluding Remarks. *Cold Spring Harb Symp Quant Biol* **22**, 415–427. (doi:10.1101/SQB.1957.022.01.039)
34. Soberón J, Arroyo-Peña B. 2017 Are fundamental niches larger than the realized? Testing a 50-year-old prediction by Hutchinson. *PLOS ONE* **12**, e0175138. (doi:10.1371/journal.pone.0175138)
35. Soberón J, Peterson AT. 2005 Interpretation of models of fundamental ecological niches and species' distributional areas. *Biodiversity Informatics* **2**. (doi:10.17161/bi.v2i0.4)
36. Lustenhouwer N *et al.* 2023 Experimental evolution of dispersal: Unifying theory, experiments and natural systems. *Journal of Animal Ecology* **92**, 1113–1123. (doi:10.1111/1365-2656.13930)
37. Fronhofer EA, Klecka J, Melián CJ, Altermatt F. 2015 Condition-dependent movement and dispersal in experimental metacommunities. *Ecology Letters* **18**, 954–963. (doi:10.1111/ele.12475)
38. Kleynhans EJ, Otto SP, Reich PB, Vellend M. 2016 Adaptation to elevated CO<sub>2</sub> in different biodiversity contexts. *Nat Commun* **7**, 12358. (doi:10.1038/ncomms12358)
39. Lawrence D, Fiegna F, Behrends V, Bundy JG, Phillimore AB, Bell T, Barraclough TG. 2012 Species interactions alter evolutionary responses to a novel environment. *PLOS Biology* **10**, e1001330. (doi:10.1371/journal.pbio.1001330)
40. Lorts CM, Lasky JR. 2020 Competition × drought interactions change phenotypic plasticity and the direction of selection on Arabidopsis traits. *New Phytologist* **227**, 1060–1072. (doi:10.1111/NPH.16593)
41. Alexander JM, Diez JM, Levine JM. 2015 Novel competitors shape species' responses to climate change. *Nature* **525**, 515–518. (doi:10.1038/nature14952)
42. Davis AJ, Jenkinson LS, Lawton JH, Shorrocks B, Wood S. 1998 Making mistakes when predicting shifts in species range in response to global warming. *Nature* **391**, 783–786. (doi:10.1038/35842)
43. Thompson PL, Hürlemann S, Altermatt F. 2021 Species interactions limit the predictability of community responses to environmental change. *The American Naturalist* (doi:10.1086/716724)
44. Paterson JE, Blouin-Demers G. 2017 Do ectotherms partition thermal resources? We still do not know. *Oecologia* **183**, 337–345. (doi:10.1007/s00442-016-3762-7)

45. Chaine AS, Schtickzelle N, Polard T, Huet M, Clobert J. 2010 Kin-based recognition and social aggregation in a ciliate. *Evolution* **64**, 1290–1300. (doi:10.1111/j.1558-5646.2009.00902.x)
46. Fronhofer EA *et al.* 2018 Bottom-up and top-down control of dispersal across major organismal groups. *Nature Ecology & Evolution* 2018 2:12 **2**, 1859–1863. (doi:10.1038/s41559-018-0686-0)
47. Haegeman B, Loreau M. 2014 General relationships between consumer dispersal, resource dispersal and metacommunity diversity. *Ecology Letters* **17**, 175–184. (doi:10.1111/ele.12214)
48. Dall SRX, Giraldeau L-A, Olsson O, McNamara JM, Stephens DW. 2005 Information and its use by animals in evolutionary ecology. *Trends in Ecology & Evolution* **20**, 187–193. (doi:10.1016/j.tree.2005.01.010)
49. Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, Smouse PE. 2008 A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences* **105**, 19052–19059. (doi:10.1073/pnas.0800375105)
50. Bonte D, De La Peña E. 2009 Evolution of body condition-dependent dispersal in metapopulations. *Journal of Evolutionary Biology* **22**, 1242–1251. (doi:10.1111/j.1420-9101.2009.01737.x)
51. Reim E, Baguette M, Günter F, Fischer K. 2018 Emigration propensity and flight performance are decoupled in a butterfly. *Ecosphere* **9**, e02502. (doi:10.1002/ecs2.2502)
52. Usinowicz J, O'Connor MI. 2023 The fitness value of ecological information in a variable world. *Ecology Letters* **26**, 621–639. (doi:10.1111/ele.14166)
53. Bestion E, Teyssier A, Aubret F, Clobert J, Cote J. 2014 Maternal exposure to predator scents: offspring phenotypic adjustment and dispersal. *Proceedings of the Royal Society B: Biological Sciences* **281**, 20140701. (doi:10.1098/rspb.2014.0701)
54. Mousseau TA, Fox CW. 1998 The adaptive significance of maternal effects. *Trends in Ecology & Evolution* **13**, 403–407. (doi:10.1016/S0169-5347(98)01472-4)
55. Antonovics J, Levin DA. 1980 The ecological and genetic consequences of density-dependent regulation in plants. *Annual Review of Ecology and Systematics* **11**, 411–452. (doi:10.1146/annurev.es.11.110180.002211)
56. Hamilton WD, May RM. 1977 Dispersal in stable habitats. *Nature* **269**, 578–581. (doi:10.1038/269578a0)
57. Kim S-Y, Torres R, Drummond H. 2009 Simultaneous positive and negative density-dependent dispersal in a colonial bird species. *Ecology* **90**, 230–239. (doi:10.1890/08-0133.1)
58. Le Galliard J-F, Ferrière null, Clobert J. 2003 Mother–offspring interactions affect natal dispersal in a lizard. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **270**, 1163–1169. (doi:10.1098/rspb.2003.2360)
59. Lambin X. 2001 Dispersal, intraspecific competition, kin competition and kin facilitation : a review of the empirical evidence. *Dispersal*
60. Matthysen E. 2005 Density-dependent dispersal in birds and mammals. *Ecography* **28**, 403–416. (doi:10.1111/j.0906-7590.2005.04073.x)
61. Ramachandran R. 1987 Influence of host-plants on the wind dispersal and the survival of an Australian geometrid caterpillar. *Entomologia Experimentalis et Applicata* **44**, 289–294. (doi:10.1111/j.1570-7458.1987.tb00557.x)

62. Jacob S, Bestion E, Legrand D, Clobert J, Cote J. 2015 Habitat matching and spatial heterogeneity of phenotypes: implications for metapopulation and metacommunity functioning. *Evol Ecol* **29**, 851–871. (doi:10.1007/s10682-015-9776-5)
63. Lotka AJ. 1920 Analytical note on certain rhythmic relations in organic systems. *Proceedings of the National Academy of Sciences* **6**, 410–415. (doi:10.1073/pnas.6.7.410)
64. Hauzy C, Hulot FD, Gins A, Loreau M. 2007 Intra- and interspecific density-dependent dispersal in an aquatic prey-predator system. *Journal of Animal Ecology* **76**, 552–558. (doi:10.1111/j.1365-2656.2007.01227.x)
65. Munoz NE, Blumstein DT. 2020 Optimal multisensory integration. *Behavioral Ecology* **31**, 184–193. (doi:10.1093/beheco/arz175)
66. Budaev S, Jørgensen C, Mangel M, Eliassen S, Giske J. 2019 Decision-making from the animal perspective: bridging ecology and subjective cognition. *Frontiers in Ecology and Evolution* **7**. (doi:10.3389/fevo.2019.00164)
67. Legrand D, Trochet A, Moulherat S, Calvez O, Stevens VM, Ducatez S, Clobert J, Baguette M. 2015 Ranking the ecological causes of dispersal in a butterfly. *Ecography* **38**, 822–831. (doi:10.1111/ecog.01283)
68. DeWitt TJ, Sih A, Wilson DS. 1998 Costs and limits of phenotypic plasticity. *Trends in Ecology & Evolution* **13**, 77–81. (doi:10.1016/S0169-5347(97)01274-3)
69. Schjørring S. 2002 The evolution of informed natal dispersal: inherent versus acquired information. *Evol Ecol Res* **4**, 227–238.
70. Chevin L-M, Lande R. 2015 Evolution of environmental cues for phenotypic plasticity. *Evolution* **69**, 2767–2775. (doi:10.1111/evo.12755)
71. Tufto J. 2000 The evolution of plasticity and nonplastic spatial and temporal adaptations in the presence of imperfect environmental cues. *The American Naturalist* **156**, 121–130. (doi:10.1086/303381)
72. Collins S. 2010 Competition limits adaptation and productivity in a photosynthetic alga at elevated CO<sub>2</sub>. *Proceedings of the Royal Society B: Biological Sciences* **278**, 247–255. (doi:10.1098/rspb.2010.1173)
73. Govaert L, Gilarranz LJ, Altermatt F. 2021 Competition alters species' plastic and genetic response to environmental change. *Sci Rep* **11**, 23518. (doi:10.1038/s41598-021-02841-8)
74. Kreuzinger-Janik B, Gansfort B, Traunspurger W, Ptatscheck C. 2022 It's all about food: Environmental factors cause species-specific dispersal. *Ecosphere* **13**, e4251. (doi:10.1002/ecs2.4251)
75. Barabás G, J. Michalska-Smith M, Allesina S. 2016 The effect of intra- and interspecific competition on coexistence in multispecies communities. *The American Naturalist* **188**, E1–E12. (doi:10.1086/686901)
76. Thompson PL, Guzman LM, De Meester L, Horváth Z, Ptacnik R, Vanschoenwinkel B, Viana DS, Chase JM. 2020 A process-based metacommunity framework linking local and regional scale community ecology. *Ecology Letters*, e13568. (doi:10.1111/ele.13568)
77. Bernhardt JR, Sunday JM, O'Connor MI. 2018 Metabolic theory and the temperature-size rule explain the temperature dependence of population carrying capacity. *The American Naturalist* **192**, 687–697. (doi:10.1086/700114)

78. Barton BT, Schmitz OJ. 2009 Experimental warming transforms multiple predator effects in a grassland food web. *Ecology Letters* **12**, 1317–1325. (doi:10.1111/j.1461-0248.2009.01386.x)
79. Schmitz OJ, Barton BT. 2014 Climate change effects on behavioral and physiological ecology of predator-prey interactions: Implications for conservation biological control. *Biological Control* **75**, 87–96. (doi:10.1016/j.biocontrol.2013.10.001)
80. Bestion E, García-Carreras B, Schaum C-E, Pawar S, Yvon-Durocher G. 2018 Metabolic traits predict the effects of warming on phytoplankton competition. *Ecology Letters* **21**, 655–664. (doi:10.1111/ele.12932)
81. Sentis A, Gémard C, Jaugeon B, Boukal DS. 2017 Predator diversity and environmental change modify the strengths of trophic and nontrophic interactions. *Global Change Biology* **23**, 2629–2640. (doi:10.1111/gcb.13560)
82. Thierry M, Pardikes NA, Rosenbaum B, Ximénez-Embún MG, Hrček J. 2022 The presence of multiple parasitoids decreases host survival under warming, but parasitoid performance also decreases. *Proceedings of the Royal Society B* **289**, 20220121. (doi:10.1098/RSPB.2022.0121)
83. Pärn H, Ringsby TH, Jensen H, Sæther B-E. 2011 Spatial heterogeneity in the effects of climate and density-dependence on dispersal in a house sparrow metapopulation. *Proceedings of the Royal Society B: Biological Sciences* **279**, 144–152. (doi:10.1098/rspb.2011.0673)
84. Holomuzki JR, Pillsbury RW, Khandwala SB. 1999 Interplay between dispersal determinants of larval hydropsychid caddisflies. *Can. J. Fish. Aquat. Sci.* **56**, 2041–2050. (doi:10.1139/f99-141)
85. Cortesero AM, Monge JP, Huignard J. 1997 Dispersal and Parasitizing Abilities of *Eupelmus vuilleti* (Hymenoptera: Eupelmidae) within a Column of Cowpea Seeds. *Environmental Entomology* **26**, 1025–1030. (doi:10.1093/ee/26.5.1025)
86. Berry JS, Holtzer TO. 1990 Ambulatory dispersal behavior of *Neoseiulus fallacis* (Acarina: Phytoseiidae) in relation to prey density and temperature. *Exp Appl Acarol* **8**, 253–274. (doi:10.1007/BF01202136)
87. Shaw AK. 2020 Causes and consequences of individual variation in animal movement. *Mov Ecol* **8**, 12. (doi:10.1186/s40462-020-0197-x)
88. Winandy L, Cote J, Di Gesu L, Pellerin F, Trochet A, Legrand D. 2019 Local predation risk and matrix permeability interact to shape movement strategy. *Oikos* **128**, 1402–1412. (doi:10.1111/oik.06403)
89. Tuda M, Shima K. 2002 Relative importance of weather and density dependence on the dispersal and on-plant activity of the predator *Orius minutus*. *Popul Ecol* **44**, 0251–0257. (doi:10.1007/s101440200028)
90. Gilliam JF, Fraser DF. 2001 Movement in corridors: Enhancement by predation threat, disturbance, and habitat structure. *Ecology* **82**, 258–273. (doi:10.1890/0012-9658(2001)082[0258:MICEBP]2.0.CO;2)
91. Doerder FP, Brunk C. 2012 Natural populations and inbred strains of *Tetrahymena*. *Methods in Cell Biology* **109**, 277–300. (doi:10.1016/B978-0-12-385967-9.00009-8)
92. Jacob S, Legrand D, Chaine AS, Bonte D, Schtickzelle N, Huet M, Clobert J. 2017 Gene flow favours local adaptation under habitat choice in ciliate microcosms. *Nature Ecology & Evolution* **2017 1:9 1**, 1407–1410. (doi:10.1038/s41559-017-0269-5)
93. Laurent E, Schtickzelle N, Jacob S. 2020 Fragmentation mediates thermal habitat choice in ciliate microcosms. *Proceedings of the Royal Society B* **287**. (doi:10.1098/RSPB.2019.2818)

94. Mitri S, Richard Foster K. 2013 The genotypic view of social interactions in microbial communities. *Annual Review of Genetics* **47**, 247–273. (doi:10.1146/annurev-genet-111212-133307)
95. Jacob S, Chaine AS, Huet M, Clobert J, Legrand D. 2019 Variability in dispersal syndromes is a key driver of metapopulation dynamics in experimental microcosms. *American Naturalist* **194**, 613–626. (doi:10.1086/705410)
96. Tingley R, Vallinoto M, Sequeira F, Kearney MR. 2014 Realized niche shift during a global biological invasion. *Proceedings of the National Academy of Sciences* **111**, 10233–10238. (doi:10.1073/pnas.1405766111)
97. Torchin ME, Lafferty KD, Dobson AP, McKenzie VJ, Kuris AM. 2003 Introduced species and their missing parasites. *Nature* **421**, 628–630. (doi:10.1038/nature01346)
98. De Meester N, Derycke S, Rigaux A, Moens T. 2015 Active dispersal is differentially affected by inter- and intraspecific competition in closely related nematode species. *Oikos* **124**, 561–570. (doi:10.1111/oik.01779)
99. Cote J, Fogarty S, Tymen B, Sih A, Brodin T. 2013 Personality-dependent dispersal cancelled under predation risk. *Proceedings of the Royal Society B: Biological Sciences* **280**, 20132349. (doi:10.1098/rspb.2013.2349)
100. Gillies N, Weimerskirch H, Thorley J, Clay TA, Martín López LM, Joo R, Basille M, Patrick SC. 2023 Boldness predicts plasticity in flight responses to winds. *Journal of Animal Ecology* **92**, 1730–1742. (doi:10.1111/1365-2656.13968)
101. Achiano KA, Giliomee JH. 2008 Food-, temperature- and crowding-mediated laboratory dispersal of *Carcinops pumilio* (Erichson) (Coleoptera: Histeridae), a predator of house fly (Diptera: Muscidae) eggs and larvae. *African Entomology* **16**, 115–121. (doi:10.10520/EJC32752)
102. Nielsen ME, Nylin S, Wiklund C, Gotthard K. 2023 Evolution of butterfly seasonal plasticity driven by climate change varies across life stages. *Ecology Letters* **26**, 1548–1558. (doi:10.1111/ele.14280)
103. Jacob S, Chaine AS, Schtickzelle N, Huet M, Clobert J. 2015 Social information from immigrants: multiple immigrant-based sources of information for dispersal decisions in a ciliate. *Journal of Animal Ecology* **84**, 1373–1383. (doi:10.1111/1365-2656.12380)
104. Cronin JT, Goddard J, Krivchenia A, Shivaji R. 2023 Density-dependent within-patch movement behavior of two competing species. *Ecol Evol* **13**, e10753. (doi:10.1002/ece3.10753)
105. Cayuela H, Grolet O, Joly P. 2018 Context-dependent dispersal, public information, and heterospecific attraction in newts. *Oecologia* **188**, 1069–1080. (doi:10.1007/s00442-018-4267-3)
106. Travis JMJ *et al.* 2012 Modelling dispersal: an eco-evolutionary framework incorporating emigration, movement, settlement behaviour and the multiple costs involved. *Methods in Ecology and Evolution* **3**, 628–641. (doi:10.1111/j.2041-210X.2012.00193.x)
107. Hrčák J. 2022 Individual-based eco-evolutionary framework: towards unifying ecology and evolution. *EcoEvoRxiv*
108. Takola E, Schielzeth H. 2022 Hutchinson's ecological niche for individuals. *Biol Philos* **37**, 25. (doi:10.1007/s10539-022-09849-y)
109. Bestion E, Clobert J, Cote J. 2015 Dispersal response to climate change: scaling down to intraspecific variation. *Ecology Letters* **18**, 1226–1233. (doi:10.1111/ele.12502)

110. Bocedi G, Heinonen J, Travis JMJ. 2012 Uncertainty and the role of information acquisition in the evolution of context-dependent emigration. *The American Naturalist* **179**, 606–620. (doi:10.1086/665004)
111. Forsman A, Wennersten L. 2016 Inter-individual variation promotes ecological success of populations and species: evidence from experimental and comparative studies. *Ecography* **39**, 630–648. (doi:10.1111/ecog.01357)
112. Luo M, Wang S, Saavedra S, Ebert D, Altermatt F. 2022 Multispecies coexistence in fragmented landscapes. *Proceedings of the National Academy of Sciences* **119**, e2201503119. (doi:10.1073/pnas.2201503119)
113. Costa-Pereira R, Moll RJ, Jesmer BR, Jetz W. 2022 Animal tracking moves community ecology: Opportunities and challenges. *Journal of Animal Ecology* **91**, 1334–1344. (doi:10.1111/1365-2656.13698)
114. Belichon S, Clobert J, Massot M. 1996 Are there differences in fitness components between philopatric and dispersing individuals? *Acta oecologica* **17**, 503–517.
115. Brown JH, Gillooly JF, Allen AP, Savage VM, West GB. 2004 Toward a metabolic theory of ecology. *Ecology* **85**, 1771–1789. (doi:10.1890/03-9000)
116. Junker AD, Jacob S, Philippe H, Legrand D, Pearson CG. 2021 Plastic cell morphology changes during dispersal. *iScience* **24**, 102915. (doi:10.1016/j.isci.2021.102915)
117. Gilman SE, Urban MC, Tewksbury J, Gilchrist GW, Holt RD. 2010 A framework for community interactions under climate change. *Trends in Ecology & Evolution* **25**, 325–331. (doi:10.1016/j.tree.2010.03.002)
118. Urban MC, Tewksbury JJ, Sheldon KS. 2012 On a collision course: competition and dispersal differences create no-analogue communities and cause extinctions during climate change. *Proceedings of the Royal Society B: Biological Sciences* **279**, 2072–2080. (doi:10.1098/rspb.2011.2367)
119. Williams JW, Jackson ST. 2007 Novel climates, no-analog communities, and ecological surprises. *Frontiers in Ecology and the Environment* **5**, 475–482. (doi:10.1890/070037)
120. Hughes L. 2000 Biological consequences of global warming: is the signal already apparent? *Trends in Ecology & Evolution* **15**, 56–61. (doi:10.1016/S0169-5347(99)01764-4)
121. Koelle K, Vandermeer J. 2005 Dispersal-induced desynchronization: from metapopulations to metacommunities. *Ecology Letters* **8**, 167–175. (doi:10.1111/j.1461-0248.2004.00703.x)
122. Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin J-M, Hoegh-Guldberg O, Bairlein F. 2002 Ecological responses to recent climate change. *Nature* **416**, 389–395. (doi:10.1038/416389a)
123. Boukal DS, Bideault A, Carreira BM, Sentis A. 2019 Species interactions under climate change: connecting kinetic effects of temperature on individuals to community dynamics. *Current Opinion in Insect Science* **35**, 88–95. (doi:10.1016/j.cois.2019.06.014)
124. Carscadden KA, Emery NC, Arnillas CA, Cadotte MW, Afkhami ME, Gravel D, Livingstone SW, Wiens JJ. 2020 Niche breadth: Causes and consequences for ecology, evolution, and conservation. *The Quarterly Review of Biology* **95**, 179–214. (doi:10.1086/710388)
125. Westley PAH, Dittman AH, Ward EJ, Quinn TP. 2015 Signals of climate, conspecific density, and watershed features in patterns of homing and dispersal by Pacific salmon. *Ecology* **96**, 2823–2833. (doi:10.1890/14-1630.1)



126. Salinger MJ. 2005 Climate variability and change: Past, present and future – An overview. *Climatic Change* **70**, 9–29. (doi:10.1007/s10584-005-5936-x)
127. Newbold T *et al.* 2015 Global effects of land use on local terrestrial biodiversity. *Nature* **520**, 45–50. (doi:10.1038/nature14324)
128. Fahrig L. 2003 Effects of Habitat Fragmentation on Biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **34**, 487–515. (doi:10.1146/annurev.ecolsys.34.011802.132419)
129. Bonamour S, Chevin L-M, Charmantier A, Teplitsky C. 2019 Phenotypic plasticity in response to climate change: the importance of cue variation. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**, 20180178. (doi:10.1098/rstb.2018.0178)
130. Seebacher F, Beaman J, Little AG. 2014 Regulation of thermal acclimation varies between generations of the short-lived mosquitofish that developed in different environmental conditions. *Functional Ecology* **28**, 137–148. (doi:10.1111/1365-2435.12156)
131. Bonte D, Travis JMJ, De Clercq N, Zwertvaegher I, Lens L. 2008 Thermal conditions during juvenile development affect adult dispersal in a spider. *Proceedings of the National Academy of Sciences* **105**, 17000–17005. (doi:10.1073/pnas.0806830105)
132. Ketola T, Kronholm I. 2023 Experimental evolution of evolutionary potential in fluctuating environments. *Journal of Evolutionary Biology* **36**, 945–949. (doi:10.1111/jeb.14178)
133. Imbert E, Ronce O. 2001 Phenotypic plasticity for dispersal ability in the seed heteromorphic *Crepisancta* (Asteraceae). *Oikos* **93**, 126–134. (doi:10.1034/j.1600-0706.2001.930114.x)
134. Harman RR, Goddard J, Shivaji R, Cronin JT. 2020 Frequency of occurrence and population-dynamic consequences of different forms of density-dependent emigration. *The American Naturalist* **195**, 851–867. (doi:10.1086/708156)
135. Travis JMJ, Murrell DJ, Dytham C. 1999 The evolution of density-dependent dispersal. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **266**, 1837–1842. (doi:10.1098/rspb.1999.0854)
136. Kuussaari M, Saccheri I, Camara M, Hanski I. 1998 Allee effect and population dynamics in the Glanville fritillary butterfly. *Oikos* **82**, 384–392. (doi:10.2307/3546980)
137. Kenward RE. 1978 Hawks and doves: factors affecting success and selection in goshawk attacks on woodpigeons. *Journal of Animal Ecology* **47**, 449–460. (doi:10.2307/3793)
138. Doligez B, Danchin E, Clobert J. 2002 Public information and breeding habitat selection in a wild bird population. *Science* **297**, 1168–1170. (doi:10.1126/science.1072838)
139. Cote J, Clobert J. 2007 Social information and emigration: lessons from immigrants. *Ecology Letters* **10**, 411–417. (doi:10.1111/j.1461-0248.2007.01032.x)
140. Chiatante G. 2018 Heterospecific social attraction in migrant birds: habitat niche overlap between two threatened shrikes. *Wildl. Res.* **46**, 25–36. (doi:10.1071/WR18031)
141. Forsman JT, Hjernquist MB, Gustafsson L. 2009 Experimental evidence for the use of density based interspecific social information in forest birds. *Ecography* **32**, 539–545. (doi:10.1111/j.1600-0587.2008.05635.x)

142. Hämäläinen R, Kajanus MH, Forsman JT, Kivelä SM, Seppänen J-T, Loukola OJ. 2023 Ecological and evolutionary consequences of selective interspecific information use. *Ecology Letters* **26**, 490–503. (doi:10.1111/ele.14184)
143. Hodge MA, Storfer-Isser A. 1997 Conspecific and heterospecific attraction: a mechanism of web-site selection leading to aggregation formation by web-building spiders. *Ethology* **103**, 815–826. (doi:10.1111/j.1439-0310.1997.tb00123.x)
144. Mönkkönen M, Forsman JT. 2002 Heterospecific attraction among forest birds: a review. *Ornithological Science* **1**, 41–51. (doi:10.2326/osj.1.41)
145. Seppänen J-T, Forsman JT, Mönkkönen M, Thomson RL. 2007 Social information use is a process across time, space, and ecology, reaching heterospecifics. *Ecology* **88**, 1622–1633. (doi:10.1890/06-1757.1)
146. Parejo D, White J, Clobert J, Dreiss A, Danchin E. 2007 Blue Tits use fledgling quantity and quality as public information in breeding site choice. *Ecology* **88**, 2373–2382. (doi:10.1890/06-2000.1)
147. Sloggett JJ, Weisser WW. 2002 Parasitoids induce production of the dispersal morph of the pea aphid, *Acyrtosiphon pisum*. *Oikos* **98**, 323–333. (doi:10.1034/j.1600-0706.2002.980213.x)
148. Gols R, Veenemans C, Potting RPJ, Smid HM, Dicke M, Harvey JA, Bukovinszky T. 2012 Variation in the specificity of plant volatiles and their use by a specialist and a generalist parasitoid. *Animal Behaviour* **83**, 1231–1242. (doi:10.1016/j.anbehav.2012.02.015)
149. Brown CR, Brown MB. 1992 Ectoparasitism as a cause of natal dispersal in Cliff swallows. *Ecology* **73**, 1718–1723. (doi:10.2307/1940023)
150. Cote J, Clobert J. 2010 Risky dispersal: avoiding kin competition despite uncertainty. *Ecology* **91**, 1485–1493. (doi:10.1890/09-0387.1)
151. Hammill E, Fitzjohn RG, Srivastava DS. 2015 Conspecific density modulates the effect of predation on dispersal rates. *Oecologia* **178**, 1149–1158. (doi:10.1007/s00442-015-3303-9)
152. French DR, Travis JMJ. 2001 Density-dependent dispersal in host-parasitoid assemblages. *Oikos* **95**, 125–135. (doi:10.1034/j.1600-0706.2001.950114.x)
153. Baines CB, McCauley SJ, Rowe L. 2014 The interactive effects of competition and predation risk on dispersal in an insect. *Biology Letters* **10**, 20140287. (doi:10.1098/rsbl.2014.0287)
154. Matsumura M, Suzuki Y. 2003 Direct and feeding-induced interactions between two rice planthoppers, *Sogatella furcifera* and *Nilaparvata lugens*: effects on dispersal capability and performance. *Ecological Entomology* **28**, 174–182. (doi:10.1046/j.1365-2311.2003.00498.x)
155. Pitcher KA, Yee DA. 2014 Investigating habitat use, prey consumption, and dispersal response as potential coexistence mechanisms using morphologically similar species of predaceous diving beetles (Coleoptera: Dytiscidae). *Annals of the Entomological Society of America* **107**, 582–591. (doi:10.1603/AN13170)
156. Byers JE. 2000 Effects of body size and resource availability on dispersal in a native and a non-native estuarine snail. *Journal of Experimental Marine Biology and Ecology* **248**, 133–150. (doi:10.1016/S0022-0981(00)00163-5)
157. Baines CB, McCauley SJ, Rowe L. 2015 Dispersal depends on body condition and predation risk in the semi-aquatic insect, *Notonecta undulata*. *Ecology and Evolution* **5**, 2307–2316. (doi:10.1002/ece3.1508)

158. Baines CB, Ferzoco IMC, McCauley SJ. 2019 Phenotype-by-environment interactions influence dispersal. *Journal of Animal Ecology* **88**, 1263–1274. (doi:10.1111/1365-2656.13008)
159. Debeffe L, Morellet N, Verheyden-Tixier H, Hoste H, Gaillard J-M, Cargnelutti B, Picot D, Sevila J, Hewison AJM. 2014 Parasite abundance contributes to condition-dependent dispersal in a wild population of large herbivore. *Oikos* **123**, 1121–1125. (doi:10.1111/oik.01396)
160. Terui A, Ooue K, Urabe H, Nakamura F. 2017 Parasite infection induces size-dependent host dispersal: consequences for parasite persistence. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20171491. (doi:10.1098/rspb.2017.1491)
161. De Bona S, Bruneaux M, Lee AEG, Reznick DN, Bentzen P, López-Sepulcre A. 2019 Spatio-temporal dynamics of density-dependent dispersal during a population colonisation. *Ecology Letters* **22**, 634–644. (doi:10.1111/ele.13205)

## Figure captions

**Figure 1. Overview of the cause of dispersal decisions depending on the different dimensions of the environment and their interplay.** Dispersal decisions can depend on the abiotic and biotic (both intra- and interspecific) conditions and are influenced by the individual internal conditions (plain arrows). However, abiotic and biotic factors are not independent (dashed arrows) and can thus have interactive effects on dispersal decisions.

**Figure 2. Predictions about dispersal decisions along two environmental gradients depending on ecological niche and biotic interactions.** a) Hypothetical ecological niche of individuals of two species (species A in orange and species B in blue) along two environmental gradients with their performance curves. Along the environmental gradient 1, species A is a specialist while species B is a generalist. Their optimum along the gradient do not overlap. Along the environmental gradient 2, species A is a generalist while species B is a specialist and they have the same optimum. b) Dispersal reaction norms of the two species in isolation along each environmental gradient following their performance curves. We hypothesize that, for a given environmental axis, the generalist species has never a null dispersal rate while the specialist species does not disperse at its optimum. c) In presence of the other species, dispersal decisions of species B along the environmental gradient 1 change slightly due to the small niche overlap with the superior competitor at the beginning of the gradient. Along the environmental gradient 2, species A avoids the superior competitor species B by dispersing toward sub-optimal habitats [22]. More complex dispersal decisions would appear in a three-dimensional space combining environmental gradients 1 and 2 and biotic interactions.

**Figure 3. a-c) Reactions norms of dispersal rates along the thermal axis depending on a) monoculture density, b) presence of another strain in cultures at high density and c) biotic interaction type in cocultures.** Small points represent raw data and big point the means with their standard errors for each strain (color) and species (shape). Lines between factor levels of the x-axis represent the reaction norms. **d-f) Dispersal plasticity along the abiotic (x axis) and biotic (y axis) axes in d) monocultures, e) cultures at high density, and f) cocultures. g-i) Thermal effect on dispersal depending on g) monoculture density, h) presence of another strain in cultures at high density and i) biotic interaction type in cocultures.** Points represent the effects size on dispersal rates (contrasts  $\pm$  95% CIs) recovered from the models for each strain (color) and species (shape). Horizontal and vertical dot lines situate the effect size values corresponding to no plasticity.

**Table 1. Evidence of the effects and interplays between abiotic conditions, biotic context (decomposed in conspecifics and heterospecifics), and the phenotype on dispersal decisions (presented with the biological system, the observed effects, the type of study, and references).**

<b>FACTORS</b>	<b>Species</b>	<b>EFFECTS ON DISPERSAL DECISIONS</b>	<b>TYPE OF STUDY</b>	<b>REFERE NCES</b>
<b>Abiotic</b>	Five <i>Tetrahymena</i> species	Dispersal plasticity correlates with fitness sensitivity to the environment along three abiotic gradients (temperature, salinity and nutrients)	Microcosm experiment	[14]
	Money spiders ( <i>Erigone atra</i> )	Thermal conditions during the development determine adult dispersal strategy	Microcosm experiment	[131]
	<i>Crepis sancta</i> (Asteraceae)	A greater proportion of seed dispersal morphs are produced when nutrients are depleted	Mesocosm experiment	[133]
<b>Conspecifics</b>	Various	High conspecific densities can reflect a good habitat quality or a crowded competitive habitat and lead to negative or positive density-dependent dispersal respectively	Various	[60,134,135 ]
	Glanville Fritillary butterfly ( <i>Melitaea cinxia</i> )	Low conspecific densities can lead to a scarcity of mates and thus enhance dispersal	Mesocosm experiment	[136]
	Wood pigeons ( <i>Columba palumbus</i> )	Low conspecific densities can reduce the effectiveness of antipredator strategies such as flocking	Field experiment	[137]
	Collared flycatcher ( <i>Ficedula albicollis</i> )	Fitness of conspecifics is used as information sources about habitat quality	Field experiment	[138]
	Common lizard ( <i>Zootoca vivipara</i> )	Informed emigration decisions with the intra-specific competitive level that immigrants experienced (probably cued by differences in behavior or physiology)	Mesocosm experiment	[139]
<b>Heterospecifics of a same trophic level</b>	Various	Densities of competitive heterospecifics can be cues for either habitat quality or competitive habitats	Various	[140–145]
	Blue Tits ( <i>Cyanistes caeruleus</i> )	Condition and reproductive success of heterospecifics represent cues about habitat quality	Field experiment	[146]
<b>Heterospecifics</b>	21 species ranging from protozoa to	Cues about the presence of	Microcosm	[46]

<b>of a different trophic level</b>	vertebrates	resources and predators affect dispersal decisions	and mesocosm experiments	
	<i>Tetrahymena pyriformis</i> and its prey <i>Dileptus</i> sp.	A decrease in the prey density led to an increase in predator dispersal while an increase in the predator density led to an increase in prey dispersal	Microcosm experiment	[64]
	Pea aphid ( <i>Acyrtosiphon pisum</i> )	Specialized dispersal morphs production in the presence of natural enemies	Mesocosm experiment	[147]
	Parasitoids ( <i>Diadegma semiclausum</i> and <i>D. fenestrata</i> ) and their host caterpillars of the diamondback moth ( <i>Plutella xylostella</i> )	Natural enemies locate and disperse toward herbivores via the volatile organic compounds produced by the plant due to herbivory	Mesocosm experiment	[148]
<b>Abiotic x conspecifics</b>	Cliff swallows ( <i>Petrochelidon pyrrhonota</i> )	Natal dispersal induced by ectoparasitism	Field observations	[149]
	Common lizard ( <i>Zootoca vivipara</i> )	Biotic information about kin competition prioritized over cues about suitable habitats	Mesocosm experiment	[150]
	Large white butterfly ( <i>Pieris brassicae</i> )	Weather conditions and habitat quality prioritized over population density and sex ratio	Mesocosm experiment	[67]
<b>Abiotic x heterospecifics</b>	House sparrow ( <i>Passer domesticus</i> )	Temperature- and density-dependent dispersal vary depending on resource availability	Field observations	[83]
	Larval hydroptychid caddisflies	Interactive effects of predator presence, algal cover and substrate size	Experiments in real and artificial streams	[84]
	Parasitoid ( <i>Eupelmus vuilleti</i> )	Independent effect of host detection via olfactory cues and interactive effect of light and gravity	Microcosm experiment	[85]
<b>Conspecifics x heterospecifics</b>	Predatory spider ( <i>Neoseiulus fallacis</i> )	Negative effect of prey density, but only for temperatures above 23°C	Microcosm experiment	[86]
	Ciliate ( <i>Paramecium aurelia</i> ) and its predator ( <i>Stenostomum virginianum</i> )	Positive effect of predation risk at low conspecific density and negative effect at high conspecific density	Microcosm experiment	[151]
	Cowpea bruchid ( <i>Callosobruchus chinensis</i> ) and its parasitoid ( <i>Anisopteromalus calandrae</i> )	Negative effect of parasitoid:host ratio, potentially due to increase in intraspecific competition	Microcosm experiment	[152]
	Backswimmers ( <i>Notonecta undulata</i> )	Positive effect of predation risk at intermediate conspecific density but almost no effects of predation at high and low conspecific densities	Mesocosm experiment	[153]
	Predatory bug ( <i>Orius minutes</i> ) and its prey	Importance of the ratio between prey and predator	Field experiment	[89]

		densities		
	Nematods ( <i>Litoditis marina</i> )	Positive effect of density, either with conspecifics or heterospecifics	Microcosm experiment	[98]
	Wing-dimorphic planthoppers ( <i>Sogatella furcifera</i> and <i>Nilaparvata lugens</i> )	Intraspecific effects stronger for <i>S. furcifera</i> and interspecific stronger for <i>N. lugens</i>	Microcosm experiment	[154]
	Diving beetles ( <i>Laccophilus fasciatus rufus</i> and <i>Laccophilus Proximus</i> )	Effect of unknown intraspecific interactions rather than interspecific interactions	Field and microcosm experiments	[155]
	Nematode ( <i>Caenorhabditis elegans</i> )	Increase in conspecifics competition (with decreasing food availability and increasing population density) increases dispersal. The effect was amplified by the presence of a predator for certain food-supply levels	Microcosm experiment	[74]
<b>Environment x phenotype</b>	Poultryhouse pill beetle ( <i>Carcinops pumilio</i> )	Effect of temperature but only when individuals were starved	Microcosm experiment	[101]
	Mosquitofish ( <i>Gambusia affinis</i> )	Bold and social mosquitofish dispersed more than shy and less social ones without predation risk but no differences under predation risk	Mesocosm experiment	[99]
	Common toad ( <i>Bufo bufo</i> )	Interactive effect of matrix permeability and predation risk on movement syndrome	Mesocosm experiment	[88]
	Common lizard ( <i>Zootoca vivipara</i> )	Individuals with low preferred temperatures dispersed more from warmer habitats, whereas individuals with higher preferred temperatures dispersed more from cooler habitats.	Mesocosm experiment	[109]
	Salt marsh snails ( <i>Cerithidea californica</i> Haldeman and <i>Batillaria attramentaria</i> Sowerby)	Dispersal of large snails for both species increased as resource levels decreased, supporting that competition influences dispersal rates. However, small snails always dispersed at relatively higher rates than larger individuals and were not influenced by variation in resource levels because competition reduces with body size.	Field experiment	[156]
	Backswimmers ( <i>Notonecta undulata</i> )	Additive effects between body condition and	Field experiment	[157,158]

		predation risk but interactive effect between sex, body mass and population density		
	Stream fish ( <i>Rivulus hartii</i> )	Emigration positively correlated to water level and body size, but only when presence of the predator <i>Hoplias malabaricus</i>	Field observations	[90]
	Roe deer ( <i>Capreolus capreolus</i> )	Natal dispersal propensity depends on both endoparasite abundance and body mass	Field observations	[159]
	Salmonid fish ( <i>Oncorhynchus masou masou</i> )	Large fish infested by larval parasites of the freshwater mussel <i>Margaritifera laevis</i> become dispersed more, whereas small infested fish tended to stay in the patch	Field observations	[160]
	Wild guppies ( <i>Poecilia reticulata</i> )	Natal dispersal is positively influenced by conspecific density at both the local and landscape scale because of intra-specific competition for resources while adult dispersal switched from conspecific attraction to conspecific avoidance along a predation-risk gradient (from high to low) at the local scale	Field observations	[161]