



The KdV equation and solutions in terms of Wronskians

+

Pierre Gaillard

► To cite this version:

| Pierre Gaillard. The KdV equation and solutions in terms of Wronskians +. 2024. hal-04677152

HAL Id: hal-04677152

<https://hal.science/hal-04677152v1>

Preprint submitted on 25 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The KdV equation and solutions in terms of Wronskians

⁺Pierre Gaillard, ⁺ Université de Bourgogne Franche Comté, Dijon, France :
e-mail: Pierre.Gaillard@u-bourgogne.fr,

Abstract

A method to construct solutions to the Korteweg-de-Vries (KdV) equation in terms of wronskians is given.

For this, a particular type of polynomials is considered and we obtain for each positive integer n , rational solutions in terms of determinants of order n .

Explicit solutions can be easily constructed and rational solutions from order 1 until order 10 are given.

1 Introduction

In the following, the Korteweg-de-Vries (KdV) equation

$$u_t + \frac{3}{2}uu_x + u_{xxx} = 0, \quad (1)$$

is considered, with the usual notations, where the subscripts x and t denote partial derivatives.

Korteweg and de Vries [77] introduced this equation (1) for the first time in 1895. This KdV equation is the basis of the most common tool for the 2-dimensional modelling of shallow water waves.

Numerical modelling done by Zabusky and Kruskal revealed the presence of soliton solutions of this equation [?]. The inverse scattering technique developed in a work published in 1967 by Gardner et al. for the KdV equation enabled the derivation of analytic solutions for given initial conditions with zeros at infinity [74]. Being the historic first among the integrable nonlinear evolution equations, the KdV equation attracted significant attention from both physicists and mathematicians. Zakharov and Faddeev in 1971 [87] proved that this equation is a completely integrable.

Solutions were built by Hirota in 1971 by using the bilinear method [75].

Solutions in terms of Riemann theta functions [76] were constructed by Its and Matveev in 1975. In the same year the expressions of periodic and almost periodic solutions [79] were given by Lax. Among different works realized, we can mention for example Airault et al. in 1977 [3], Adler and Moser in 1978 [2],

Ablowitz and Cornille in 1979 [1], Freeman and Nimmo in 1984 [5], Matveev in 1992 [84], Ma in 2004 [81], Kovalyov in 2005 [78] and more recently Ma in 2015 [82].

Considering certain polynomials, we construct rational solutions in terms of wronskians of order n . So we get a very efficient method to construct rational solutions to the KdV equation. We give explicit solutions in the simplest cases for orders $n = 1$ until 10.

2 Solutions to the KdV equation

Let p_n be the polynomials defined by

$$\begin{cases} p_{3k}(x, t) = \sum_{l=0}^k \frac{x^{3l}}{(3l)!} \frac{(-t)^{k-l}}{(k-l)!}, & k \geq 0, \\ p_{3k+1}(x, t) = \sum_{l=0}^k \frac{x^{3l+1}}{(3l+1)!} \frac{(-t)^{k-l}}{(k-l)!}, & k \geq 0, \\ p_{3k+2}(x, t) = \sum_{l=0}^k \frac{x^{3l+2}}{(3l+2)!} \frac{(-t)^{k-l}}{(k-l)!}, & k \geq 0, \\ p_n(x, t) = 0, & n < 0. \end{cases} \quad (2)$$

Let $W_n(x, t) = W(p_{2n}, \dots, p_{n+1})$ be the classical wronskian

$$W_n(x, t) = \det(\partial_x^{i-1}(p_{2n+1-j})_{\{1 \leq i \leq n, 1 \leq j \leq n\}}) \quad (3)$$

with $\partial_x^0(p_{n+j})$ meaning p_{n+j} .

We get the following statement

Theorem 2.1 *The function $v_n(x, t)$ expressed as*

$$v_n(x, t) = 4\partial_x^2(\ln(W_n(x, t))) \quad (4)$$

is a rational solution to the (KdV) equation (1)

$$u_t + \frac{3}{2}uu_x + u_{xxx} = 0. \quad (5)$$

3 Rational solutions to the KdV equation

We construct explicitly rational solutions to the KdV equation.

We call rational solution to the KdV equation of order k , the following function

$$v_k(x, t) = 4\partial_x^2(\ln(W_k(x, t))).$$

In the following, we give some examples of these solutions.

3.1 Solutions of order 1

Example 3.1 The function $v_k(x, t)$ expressed as

$$v_k(x, t) = -\frac{8}{x^2} \quad (6)$$

is a rational solution to the KdV equation.

3.2 Solutions of order 2

Example 3.2 The function $v_k(x, t)$ expressed as

$$v_k(x, t) = 24 \frac{(-x^3 + 24t)x}{x^6 + 24tx^3 + 144t^2} \quad (7)$$

is a rational solution to the KdV equation.

3.3 Solutions of order 3

Example 3.3 The function $v_k(x, t)$ expressed as

$$v_k(x, t) = -48 \frac{(x^9 + 5400t^2x^3 + 43200t^3)x}{x^{12} + 120tx^9 + 2160t^2x^6 - 86400t^3x^3 + 518400t^4} \quad (8)$$

is a rational solution to the KdV equation.

3.4 Solutions of order 4

Example 3.4 The function $v_k(x, t)$ expressed as

$$v_k(x, t) = \frac{n(x, t)}{d(x, t)} \quad (9)$$

with $n(x, t) = -80(9144576000t^6 - 152409600t^4x^6 - 2116800t^3x^9 + 22680t^2x^{12} + 144tx^{15} + x^{18})$

and

$d(x, t) = (x^{18} + 360tx^{15} + 32400t^2x^{12} + 604800t^3x^9 + 108864000t^4x^6 + 91445760000t^6)x^2$
is a rational solution to the KdV equation.

3.5 Solutions of order 5

Example 3.5 The function $v_k(x, t)$ expressed as

$$v_k(x, t) = \frac{n(x, t)}{d(x, t)} \quad (10)$$

with

$$\begin{aligned} n(x, t) = & 120(-x^{27} - 504tx^{24} - 120960t^2x^{21} - 1447891200t^4x^{15} \\ & - 512096256000t^5x^{12} - 17667320832000t^6x^9 - 184354652160000t^7x^6) \end{aligned}$$

$$\begin{aligned}
& - 19357238476800000 t^8 x^3 + 154857907814400000 t^9) x \\
& \text{and} \\
d(x, t) & = x^{30} + 840 t x^{27} + 226800 t^2 x^{24} + 25401600 t^3 x^{21} \\
& + 1905120000 t^4 x^{18} - 109734912000 t^5 x^{15} - 9601804800000 t^6 x^{12} \\
& - 1152216576000000 t^7 x^9 + 58071715430400000 t^8 x^6 \\
& + 774289539072000000 t^9 x^3 + 2322868617216000000 t^{10}
\end{aligned}$$

is a rational solution to the KdV equation.

3.6 Solutions of order 6

Example 3.6 The function $v_k(x, t)$ expressed as

$$v_k(x, t) = \frac{n(x, t)}{d(x, t)} \quad (11)$$

with

$$\begin{aligned}
n(x, t) & = -168(x^{39} + 1200 t x^{36} + 604800 t^2 x^{33} + 120960000 t^3 x^{30} + 23278752000 t^4 x^{27} + \\
& 2112397056000 t^5 x^{24} - 1594448271360000 t^6 x^{21} - 294045670195200000 t^7 x^{18} \\
& - 17292927259238400000 t^8 x^{15} - 749512273821696000000 t^9 x^{12} \\
& + 66893970438586368000000 t^{10} x^9 - 404736627863715840000000 t^{11} x^6 \\
& + 70828909876150272000000000 t^{12} x^3 + 339978767405521305600000000 t^{13}) x \\
& \text{and} \\
d(x, t) & = x^{42} + 1680 t x^{39} + 1038240 t^2 x^{36} + 313286400 t^3 x^{33} + 53317958400 t^4 x^{30} \\
& + 3990083328000 t^5 x^{27} + 441683020800000 t^6 x^{24} + 93283453992960000 t^7 x^{21} \\
& + 18418412410675200000 t^8 x^{18} - 534879213590937600000 t^9 x^{15} \\
& + 81739423771213824000000 t^{10} x^{12} + 10152143748914872320000000 t^{11} x^9 \\
& + 138824663357254533120000000 t^{12} x^6 - 2379851371838649139200000000 t^{13} x^3 \\
& + 7139554115515947417600000000 t^{14}
\end{aligned}$$

is a rational solution to the KdV equation.

3.7 Solutions of order 7

Example 3.7 The function $v_k(x, t)$ defined by

$$v_k(x, t) = \frac{n(x, t)}{d(x, t)} \quad (12)$$

with

$$\begin{aligned}
n(x, t) & = -224 x^{54} - 532224 t x^{51} - 533433600 t^2 x^{48} - 274743705600 t^3 x^{45} - 90973899878400 t^4 x^{42} - \\
& 17237159976960000 t^5 x^{39} - 341160574132224000 t^6 x^{36} - 1094458263478272000000 t^7 x^{33} \\
& - 741477502246327418880000 t^8 x^{30} - 154020874806534655180800000 t^9 x^{27} \\
& - 16034390275582650875904000000 t^{10} x^{24} - 177938087284701740924928000000 t^{11} x^{21} \\
& - 93996037692962389591326720000000 t^{12} x^{18} - 4714367526984320747757895680000000 t^{13} x^{15} \\
& - 86292038044598141254568509440000000 t^{14} x^{12} \\
& + 20603060246307153105741938688000000000 t^{15} x^9
\end{aligned}$$

$$\begin{aligned}
& + 1483420337734115023613419585536000000000 t^{16}x^6 \\
& - 29668406754682300472268391710720000000000 t^{18} \\
& \text{and} \\
d(x, t) = & (x^{54} + 3024tx^{51} + 3646944t^2x^{48} + 2336947200t^3x^{45} + 895482604800t^4x^{42} \\
& + 211662185011200t^5x^{39} + 34736257188864000t^6x^{36} + 3672731815796736000t^7x^{33} \\
& - 240312372794081280000t^8x^{30} - 285039517734390988800000t^9x^{27} \\
& - 26606879994975849676800000t^{10}x^{24} - 5005337403127787421696000000t^{11}x^{21} - \\
& 341169543038359143972864000000t^{12}x^{18} + 130784494154255779579822080000000t^{13}x^{15} \\
& + 11439814652602935286431744000000000t^{14}x^{12} \\
& + 73582358022525546806221209600000000t^{15}x^9 \\
& + 13244824444054598425119817728000000000t^{16}x^6 \\
& + 3708550844335287559033548963840000000000t^{18})x^2
\end{aligned}$$

is a rational solution to the KdV equation.

3.8 Solutions of order 8

Example 3.8 The function $v_k(x, t)$ expressed as

$$v_k(x, t) = \frac{n(x, t)}{d(x, t)} \quad (13)$$

with

$$\begin{aligned}
n(x, t) = & 288(-x^{69} - 4200tx^{66} - 7623000t^2x^{63} - 7730553600t^3x^{60} - 5000876496000t^4x^{57} \\
& - 2135257886361600t^5x^{54} - 609838468081920000t^6x^{51} - 163518036732610560000t^7x^{48} \\
& - 25169090600103690240000t^8x^{45} + 24130294695297087897600000t^9x^{42} \\
& + 1726822502597498322722560000t^{10}x^{39} + 5035052987586864132194304000000t^{11}x^{36} \\
& + 705713595558650622946050048000000t^{12}x^{33} + 83184077095173887007853117440000000t^{13}x^{30} \\
& + 3466115954227539946436186603520000000t^{14}x^{27} \\
& - 51516112586940473202237562159104000000t^{15}x^{24} \\
& - 28979499285932207292849826257960960000000t^{16}x^{21} \\
& - 2969807516143698277274066010243072000000000t^{17}x^{18} \\
& + 219447562532189704845715806256889856000000000t^{18}x^{15} \\
& - 216142591024938360620006524225490780160000000000t^{19}x^{12} \\
& - 5465663450786023746477963825911452139520000000000t^{20}x^9 \\
& - 36692565823458620955376540369755203174400000000000t^{21}x^6 \\
& - 2407949632164472000196585461765185208320000000000000t^{22}x^3 \\
& + 1155815823438946560094361021647288899936000000000000t^{23})x
\end{aligned}$$

and

$$\begin{aligned}
d(x, t) = & x^{72} + 5040tx^{69} + 10674720t^2x^{66} + 12666931200t^3x^{63} + 9461359353600t^4x^{60} \\
& + 4710504608409600t^5x^{57} + 1632339334112256000t^6x^{54} + 397315117938106368000t^7x^{51} \\
& + 65431718836877352960000t^8x^{48} + 9437611212912434872320000t^9x^{45} \\
& + 4685020977200251685437440000t^{10}x^{42} + 1631982341617690935243571200000t^{11}x^{39} \\
& + 419453485567536573208697241600000t^{12}x^{36} + 56299529140380549326208761856000000t^{13}x^{33} \\
& - 4653435919389780491390674796544000000t^{14}x^{30} \\
& + 1234383299752144727965990775685120000000t^{15}x^{27}
\end{aligned}$$

$$\begin{aligned}
& + 7237700541575855582398849996554240000000000 t^{16}x^{24} \\
& + 630659838963943927023985589175189504000000000 t^{17}x^{21} \\
& + 4632051208750985398754036099976265728000000000 t^{18}x^{18} \\
& - 66958128775409005273214599142943227904000000000 t^{19}x^{15} \\
& - 5232860239595519238522449791368497725440000000000 t^{20}x^{12} \\
& - 614600477542931901002557051193399653171200000000000 t^{21}x^9 \\
& + 150256057047063052812266932814147556999168000000000000 t^{22}x^6 \\
& + 138697898812673587211323322597674667999232000000000000 t^{23}x^3 \\
& + 277395797625347174422646645195349335998464000000000000 t^{24}
\end{aligned}$$

is a rational solution to the KdV equation.

3.9 Solutions of order 9

Example 3.9 The function $v_k(x, t)$ defined by

$$v_k(x, t) = \frac{n(x, t)}{d(x, t)} \quad (14)$$

with

$$\begin{aligned}
n(x, t) = & -360(x^{87} + 6864tx^{84} + 20845440t^2x^{81} + 36883123200t^3x^{78} + 42793643692800t^4x^{75} \\
& + 34479450061056000t^5x^{72} + 20002051134793728000t^6x^{69} + 8731536922920075264000t^7x^{66} \\
& + 2751355428529975787520000t^8x^{63} + 497883367475949628293120000t^9x^{60} \\
& + 217830264215804373294120960000t^{10}x^{57} + 325706259771597584319475875840000t^{11}x^{54} \\
& + 23613668450824898355530712678400000t^{12}x^{51} + 90842180055466965502594778136576000000t^{13}x^{48} \\
& + 22464586263799027916853721618710528000000t^{14}x^{45} + 3471147967080811533200910965060665344000000t^{15}x^{42} \\
& + 305398496769968754512043060254019158016000000t^{16}x^{39} \\
& + 1750666438551366314018316164380741337088000000t^{17}x^{36} \\
& + 23112992363226111560405467570860512929382400000000t^{18}x^{33} \\
& + 389754549604414714891502174564268060652339200000000t^{19}x^{30} \\
& + 35573513199032343174082894988317881256346910720000000t^{20}x^{27} \\
& + 9378985656695796563336893013692224467033294028800000000t^{21}x^{24} \\
& - 628602036005341304797250525826869047108513352908000000000t^{22}x^{21} \\
& - 162329335579032153111484405548792940444728667668480000000000t^{23}x^{18} \\
& - 75986637119012033976830106990619719139025700740136960000000000t^{24}x^{15} \\
& - 2856556091043632926211314732491165804870086303730892800000000000t^{25}x^{12} \\
& + 13087464265664955125039994390898846051632948803209789440000000000t^{26}x^9 \\
& - 634543721971634187880727000770853141897355093488959488000000000000t^{27}x^6 \\
& + 58298704456143891011541793195822132411819499214298152960000000000000t^{28}x^3 \\
& + 199881272421064769182429005242818739697666854449022238720000000000000t^{29})_x
\end{aligned}$$

and

$$\begin{aligned}
d(x, t) = & x^{90} + 7920tx^{87} + 27324000t^2x^{84} + 54752544000t^3x^{81} + 71622115488000t^4x^{78} + 65066476598323200t^5x^{75} \\
& + 42677705283333120000t^6x^{72} + 20691357050667663360000t^7x^{69} + 7527123164747777310720000t^8x^{66} \\
& + 2107511386632847215820800000t^9x^{63} + 469311169848026334959370240000t^{10}x^{60} \\
& + 40479537955403752535044915200000t^{11}x^{57} - 38641983187081149702891896832000000t^{12}x^{54} \\
& - 28229717733062389116522121396224000000t^{13}x^{51} - 975524760175268043556099819831296000000t^{14}x^{48} \\
& - 1605415983686251630919982642723028992000000t^{15}x^{45}
\end{aligned}$$

$$\begin{aligned}
& -371454265116593208604981692729309265920000000 t^{16} x^{42} \\
& -84128558269428776137058782348652314624000000000 t^{17} x^{39} \\
& +28352032914692269976628454317964798407475200000000 t^{18} x^{36} \\
& +13128049960975410746073137017106342394986496000000000 t^{19} x^{33} \\
& +2430885024143419353984724033937664218012162457600000000 t^{20} x^{30} \\
& +146410665507920500533143394324200504477481959424000000000 t^{21} x^{27} \\
& +6602294506252984121623391996311398594267033108480000000000 t^{22} x^{24} \\
& +6340693439092960095436410881602153532582910335385600000000000 t^{23} x^{21} \\
& +195086521542338342335825547841625228435453001072640000000000000 t^{24} x^{18} \\
& -29476806265415574162291514021822848249977034812424192000000000000 t^{25} x^{15} \\
& +202801615686956949251652805644093688816626272777011200000000000000 t^{26} x^{12} \\
& +25778338705097638882654534406315908889580050672988979200000000000000 t^{27} x^9 \\
& +2748367495789640576258398822088757670842919248674055782400000000000000 t^{28} x^6 \\
& -29982190863159715377364350786422810954650028167353358080000000000000000 t^{29} x^3 \\
& +59964381726319430754728701572845621909300056334706671616000000000000000 t^{30}
\end{aligned}$$

is a rational solution to the KdV equation.

3.10 Solutions of order 10

Example 3.10 The function $v_k(x, t)$ expressed as

$$v_k(x, t) = \frac{n(x, t)}{d(x, t)} \quad (15)$$

with

$$\begin{aligned}
n(x, t) = & -440 x^{108} - 4656960 t x^{105} - 22239360000 t^2 x^{102} - 63527473152000 t^3 x^{99} - 121936711792896000 t^4 x^{96} \\
& - 167286869336530944000 t^5 x^{93} - 170457089665294688256000 t^6 x^{90} - 132572493457040495738880000 t^7 x^{87} \\
& - 79773272960436964977868800000 t^8 x^{84} - 37509353281280939031016243200000 t^9 x^{81} \\
& - 14763698113159334472421736448000000 t^{10} x^{78} - 4978722500676502355093319843840000000 t^{11} x^{75} \\
& + 307544827106787687424323357769728000000 t^{12} x^{72} + 3062181416347097524628843272549171200000000 t^{13} x^{69} + \\
& 2999970822894313674432005462201582223360000000 t^{14} x^{66} \\
& + 1673514712689379266786652586236034049638400000000 t^{15} x^{63} \\
& + 61418906352803836665076046329111551179489280000000 t^{16} x^{60} \\
& + 155794693835149410545599807873777332894931353600000000 t^{17} x^{57} \\
& + 2902768310737004959527251853307297196031344640000000000 t^{18} x^{54} \\
& + 521861007163420135519099414763336936986835209420800000000 t^{19} x^{51} \\
& + 7878664545429697476399924229707963182514752716800000000000 t^{20} x^{48} \\
& - 378545538586374130684752374182777086090607685208637440000000000 t^{21} x^{45} \\
& - 1313294750805698415322003375543919305004529356936891596800000000000 t^{22} x^{42} \\
& - 3513566821938187068323458047568782732506359185178361856000000000000 t^{23} x^{39} \\
& - 1213277153327325812628585554718526296390804347854945202995200000000000 t^{24} x^{36} \\
& + 259542378662438531031176273828288551677962322567714166328524800000000000 t^{25} x^{33} \\
& - 161332608981727230241305201619521450571485883696858596226419916800000000000 t^{26} x^{30} \\
& - 38619657275477595326725531332288937493194588220016949924227710976000000000000 t^{27} x^{27} \\
& - 3949639858292160818760942549919713001064650682276256769927135887360000000000000 t^{28} x^{24} \\
& - 22398063092453206574593483244714613369097180484635029213910275194880000000000000 t^{29} x^{21}
\end{aligned}$$

$$\begin{aligned}
& -9599260625185284618515095398119217595753908235129788366990376823685120000000000000000 t^{30} x^{18} \\
& -2443740619574356759064843703785097208419752494714286388156930420690124800000000000000 t^{31} x^{15} \\
& -46085411092170352959601147874341981236415924513805933760867375334922649600000000000000 t^{32} x^{12} \\
& +5199379712962809051852437196079608139493078663096054065328626960863068160000000000000000 t^{33} x^9 \\
& +37435533933332225173337547811773178604350166374291589270366114118214090752000000000000000 t^{34} x^6 \\
& -374355339333322251733375478117731786043501663742915892703661141182140907520000000000000000 t^{36}
\end{aligned}$$

and

$$\begin{aligned}
d(x, t) = & (x^{108} + 11880 t x^{105} + 63082800 t^2 x^{102} + 199635321600 t^3 x^{99} + 423349568928000 t^4 x^{96} \\
& + 640877607943372800 t^5 x^{93} + 720356494667217408000 t^6 x^{90} + 617087988787839528960000 t^7 x^{87} \\
& + 410285748104906240286720000 t^8 x^{84} + 214683708353169921004339200000 t^9 x^{81} \\
& + 8918844995327630048644300800000 t^{10} x^{78} + 29191715419424489706841178112000000 t^{11} x^{75} \\
& + 775168403895240598967724461260800000 t^{12} x^{72} + 2539364289397734716586557530177536000000 t^{13} x^{69} \\
& + 1689948975066768235981089928013414400000000 t^{14} x^{66} \\
& + 1130347213654414430863954455878457578520000000 t^{15} x^{63} \\
& + 50655611168547447404944610107264472186800000000 t^{16} x^{60} \\
& + 1702055990512445735572775413812531031965696000000000 t^{17} x^{57} \\
& + 42383814923944123214485620080346613723378483200000000 t^{18} x^{54} + \\
& 1773594944261910353764201873309567045620203520000000000 t^{19} x^{51} \\
& - 200783526308251655278234458571207724313002115072000000000 t^{20} x^{48} \\
& + 1570726919413516781314315564980913434623903111577600000000000 t^{21} x^{45} \\
& + 980600167365812967816911712883777312427743463677624320000000000 t^{22} x^{42} \\
& + 235480271406556214857047367906374310739240351979892899840000000000 t^{23} x^{39} \\
& + 3314466802974799576312045027363982121858353580325666816000000000000 t^{24} x^{36} \\
& + 3296619340170051451908628366879682725845760352684222164500480000000000 t^{25} x^{33} \\
& + 283516231803993134625581454811050981056415186227296565880422400000000000 t^{26} x^{30} \\
& - 6892888005774141197866666184603837337804670060290450959368192000000000000 t^{27} x^{27} \\
& - 5137727874828207926581210072659589786905078681099262312719581184000000000000 t^{28} x^{24} \\
& - 63154005936950859882279738645114023793800141576489849722667458560000000000000 t^{29} x^{21} \\
& - 706399143837922099139889865939402543693176476409406132348522762076160000000000000 t^{30} x^{18} \\
& + 65932146650332942670872712529373985285363388094360016735129733326438400000000000000 t^{31} x^{15} \\
& + 5725198244302650719380471063365452626127627381183172848266189152976896000000000000000 t^{32} x^{12} \\
& + 18569213260581460899472989985998600498189566653914478804745096288796672000000000000000 t^{33} x^9 \\
& + 33424583869046629619051381974797480896741219977046061848541173319834009600000000000000 t^{34} x^6 \\
& + 4679441741666528146667193476471647325543770796786448658795764264776761344000000000000000 t^{36})_x^2
\end{aligned}$$

is a rational solution to the KdV equation.

4 Conclusion

New representation of the solutions to the KdV equation in terms of wronskians have been constructed. Precisely, these solutions can be written as the second derivative with respect to x of a logarithm of a wronskian involving particular polynomials. Surprisingly, although the representation of the solutions is different from this given in [62], we recover the same solutions.

In particular, rational solutions appear as the quotient of two polynomials in x and t ; the numerator is a polynomial of degree $n(n + 1) - 2$ in x and the denominator is a polynomial of degree $n(n + 1)$ in x have been construced. We get a very efficient method to construct rational solutions to the KdV equation. Other recent works as [83], [85] or [80] can be quoted.

The solutions presented in this paper are different from these presented in a previous work of the author [52].

References

- [1] M.J. Ablowitz, H. Cornille, On solutions of the KdV equation, Phys. Lett., V. 72A, N. 4, 277-280, 1979
- [2] M. Adler, J. Moser, On a class of polynomials connected with the Korteweg-de-Vries equation, Commun. Math. Phys., V. 61, 1-30, 1978
- [3] H. Airault, H.P. McKean, J. Moser, Rational and elliptic solutions of the KdV equation and a related many-body problem, Comm. Pure and Appl. Math., V. XXX, 2, 95-148, 1977
- [4] L. Cheng, KdV-type Wronskian rational solutions to the 4+1-dimensional Fokas equation, Par. Diff. Eq. in Appl. Math., V. 5, 100222-1-6, 2022
- [5] N. C Freeman, J.J.C. Nimmo Rational solutions of the KdV equation in wronskian form, Phys. Letters, V. 96 A, N. 9, 443-446, 1983
- [6] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, Max-Planck-Institut für Mathematik, MPI 02-31, V. 161, 2002
- [7] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, Lett. Math. Phys., V. 68, 77-90, 2004
- [8] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, Lett. Math. Phys., V. 89, 1-12, 2009
- [9] P. Gaillard, V.B. Matveev, Wronskian and Casorati determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, J. Phys A : Math. Theor., V. 42, 404409-1-16, 2009
- [10] P. Dubard, P. Gaillard, C. Klein, V. B. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, Eur. Phys. J. Spe. Top., V. 185, 247-258, 2010
- [11] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, J. Phys. A : Meth. Theor., V. 44, 435204-1-15, 2011
- [12] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, J. Math. Sciences : Adv. Appl., V. 13, N. 2, 71-153, 2012

- [13] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *J. Math. Phys.*, V. 54, 013504-1-32, 2013
- [14] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. 4, N. 4, 246-266, 2013
- [15] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. 2013, ID 645752, 1-10, 2013
- [16] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. 7, N. 45, 1-6, 2013
- [17] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. 88, 042903-1-9, 2013
- [18] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. 2013, 1-111, 2013
- [19] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. 54, 073519-1-22, 2013
- [20] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. 61, 365-369, 2014
- [21] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. 89, 015004-1-7, 2014
- [22] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. 482, 012016-1-7, 2014
- [23] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. 26, N. 2, 1550016-1-14, 2014
- [24] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. 5, 093506-1-12, 2014
- [25] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. 574, 012031-1-5, 2015
- [26] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. 355, 293-298, 2015

- [27] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation Phys. Lett. A, V. 379, 1309-1313, 2015
- [28] P. Gaillard, Other $2N-2$ parameters solutions to the NLS equation and $2N+1$ highest amplitude of the modulus of the N -th order AP breather, J. Phys. A: Math. Theor., V. 48, 145203-1-23, 2015
- [29] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, Adv. Res., V. 4, N. 5, 346-364, 2015
- [30] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, Jour. Phys. : Conf. Ser., V. 633, 012106-1-6, 2016
- [31] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, Journal of Theoretical and Applied Physics, V. 10, 1-7, 2016
- [32] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, Commun. Theor. Phys, V. 65, N. 2, 136-144, 2016
- [33] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, Annals Of Physics, V. 367, 1-5, 2016
- [34] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, Adv. Res., V. 10, 83-89, 2016
- [35] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, Theor. And Math. Phys., V. 189, N. 1, 1440-1449, 2016
- [36] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, Jour. of Math. Phys., V. 57, 063505-1-13, doi: 10.1063/1.4953383, 2016
- [37] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, Jour. Of Bas. And Appl. Res. Int., V. 21, N. 3, 130-139, 2017
- [38] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on $2N - 2$ parameters, Int. Jour. of Appl. Sci. And Math., V. 4, N. 3, 60-70, 2017
- [39] P. Gaillard, Families of rational solutions of order 5 to the KPI equation depending on 8 parameters, New Hor. in Math. Phys., V. 1, N. 1, 26-31, 2017

- [40] P. Gaillard, 6-th order rational solutions to the KPI Equation depending on 10 parameters, Jour. Of Bas. And Appl. Res. Int., V. 21, N. 2, 92-98, 2017
- [41] P. Gaillard, N -Order rational solutions to the Johnson equation depending on $2N - 2$ parameters, Int. Jour. of Adv. Res. in Phys. Sci., V. 4, N. 9, 19-37, 2017
- [42] P. Gaillard, Families of rational solutions to the KPI equation of order 7 depending on 12 parameters, Int. Jour. of Adv. Res. in Phys. Sci., V. 4, N. 11, 24-30, 2017
- [43] P. Gaillard, Rational solutions to the Johnson equation and rogue waves, Int. Jour. of Inn. In Sci. and Math., V. 6, N. 1, 14-19, 2018
- [44] P. Gaillard, Multiparametric families of solutions of the KPI equation, the structure of their rational representations and multi-rogue waves, Theo. And Mat. Phys., V. 196, N. 2, 1174-1199, 2018
- [45] P. Gaillard, The Johnson equation, Fredholm and wronskian representations of solutions and the case of order three, Adv. In Math. Phys., V. 2018, 1-18, 2018
- [46] P. Gaillard, Families of solutions of order 5 to the Johnson equation depending on 8 parameters, NHIMP, V. 2, N. 4, 53-61, 2018
- [47] P. Gaillard, Multiparametric families of solutions to the Johnson equation, J. Phys. : Conf. Series, V. 1141, 012102-1-10, 2018
- [48] P. Gaillard, Rational solutions to the Boussinesq equation, Fund. Jour. Of Math. And Appl., V. 2, 109-112, 2019
- [49] P. Gaillard, Differential relations for the solutions to the NLS equation and their different representations, Comm. In Adv. Math. Sci., V. 2, N. 4, 1-4, 2019
- [50] P. Gaillard, Multi-parametric families of solutions of order N to the Boussinesq and KP equations and the degenerate rational case, UJMA, V. 3, N. 2, 44-52, 2020
- [51] P. Gaillard, The mKdV equation and multi-parameters rational solutions, Wave Motion, V. 100, 102667-1-9, 2021
- [52] P. Gaillard, Degenerate Riemann theta functions, Fredholm and wronskian representations of the solutions to the KdV equation and the degenerate rational case, Jour. Of Geom. And Phys., V. 161, 104059-1-12, 2021
- [53] P. Gaillard, $2N$ parameter solutions to the Burgers equation, Jour. Of Appl. Nonlin. Dyn., V. 1, N. 1, 1-6, 2021

- [54] P. Gaillard, Multi-Parametric rational Solutions to the KdV equation, *Asi. Jour. Of Res. and Rev. In Phys.*, V. 3, N. 3, 14-21, 2021
- [55] P. Gaillard, Multiparametric solutions to the Gardner equation and the degenerate rational case, *Jour. Of Appl. Ana. And Appl.*, V. 11, N.4, 2102-2113, 2021
- [56] P. Gaillard, Other families of rational solutions to the KPI equation, *Asi. Res. Jour Of Math.*, V. 17, N. 6, 27-34, 2021
- [57] P. Gaillard, Rational solutions to the mKdV equation associated to particular polynomials, *Wave Motion*, V. 107, 102824-1-11, 2021
- [58] P. Gaillard, Multiparametric rational solutions of order N to the KPI equation and the explicit case of order 3, *Arch. Of cur. Res. Int.*, V. 21, N . 6, 58-71, 2021
- [59] P. Gaillard, Rational solutions to the KPI equation from particular polynomials, *Wave Motion*, V. 108, 102828-1-9, 2022
- [60] P. Gaillard, Rogue Waves of the Lakshmanan Porsezian Daniel Equation Depending on Multi-parameters, *As. Jour. Of Adv. Res. And Rep.*, V 16, N. 3, 32-40, 2022
- [61] P. Gaillard, Quasi-rational and rational solutions to the defocusing NLS equation, *Jour. Of Math. Sci. And Mod. Res. And Rep.*, V 5, N. 1, 24-34, 2022
- [62] P. Gaillard, Rational solutions to the KdV equation in terms of particular polynomials, *Jour. Of Nonlin. Mod. and Ana.*, V. 4, N. 4, 615-626, 2022
- [63] P. Gaillard, Rational solutions to the Painlev II equation from particular polynomials, *Int. J. Of Appl. Nonlin. Sci.*, V. 3, N. 3, 189-196, 2022
- [64] P. Gaillard, Multi-lump solutions to the KPI equation with a zero degree of derivation, *Jour. Of Theor. And Appl. Phys.*, V. 16, N. 4, 1-8, 2022
- [65] P. Gaillard, Families of solutions to the KPI equation given by an extended Darboux transformation, *Part. Diff. Equ. And Appl.*, V. 3, N. 80, 1-14, 2022
- [66] P. Gaillard, Rogue waves of the Hirota equation in terms of quasi-rational solutions depending on multi-parameters, *WSEAS Trans. On Math.*, DOI: 10.37394/23206.2023.22.24, V. 22, 190-203, 2023
- [67] P. Gaillard, Different representations of the solutions to the cylindrical nonlinear Schrödinger equation, *Dis. Nonlin. And Compl.*, V. 12, N. 3, 539-553, 2023
- [68] P. Gaillard, Rational solutions to the KdV equation from Riemann theta functions, *Jour. Of Theor. And Appl. Phys.*, V. 17, N. 3, 1-5, 2023

- [69] P. Gaillard, On Hadamard's coefficients for two dimensional Schrödinger operators, *Int. Jour. Of Theor. Phys.*, V. 62, N. 205, 1-16, 2023
- [70] P. Gaillard, Rational solutions to the Burgers' equation from particular polynomials, *WSEAS Trans. On Math*, DOI: 10.37394/23206.2023.22.94, 867-873, 2023
- [71] P. Gaillard, Rational solutions to the Gardner equation from particular polynomials, *WSEAS Equ.*, DOI: 10.37394/232021.2023.3.2, V. 3, 1-24, 2023
- [72] P. Gaillard, Rational solutions to the KPI equation as multi-lumps with a one degree of summation, *Int. J. Appl. Comput. Math*, doi.org/10.1007/s40819-024-01694-9, 1-14, 2024
- [73] P. Gaillard, Rational solutions to the sixth equation of the NLS hierarchy, *ISRG Journal Of Multi. Stud.*, V. 2, N. 6, 66-71, 1-14, 2024
- [74] C. S. Gardner, J. M. Green, M. D. Kruskall, R. M Miura, Method for solving the Korteweg-de Vries equation, *Phys. Rev. Let.*, V. 19, N. 19, 1095-1097, 1967
- [75] R. Hirota, Exact solution of the Korteweg-de Vries equation for multiple collisions of solitons, *Phys. Rev. Let.*, V. 27, N. 18, 1192-1194, 1971
- [76] A.R. Its, V.B. Matveev, Hill's operator with finitely many gaps, *Funct. Anal. and Appl.*, V. 9, 69-70, 1975
- [77] D.J .Korteweg, G. de Vries, On the change of form of long wawes adwancing in a rectangular canal, and on a new type of long stationary waves, *Phil. Mag.*, V. 39, 442-443, 1895
- [78] M. Kovalyov, On a class of solutions of KdV, *Jour. Of Diff. Equ.*, V. 213, 1-80, 2005
- [79] P. D. Lax, Periodic solutions of the KdV equation, *Comm. Pur. Applied Math.*, V. 28, 141-188, 1975
- [80] J. Liu, Exact solution of rational expansion method to the variable coefficient combined KdV equation with forced term, *Pure Appl. Math.*, V. 28, N. 5, 705-710, 2012
- [81] W.X. Ma, Y. You, Solving the KdV equation by its bilinear form wronskian solutions, *Trans. Of The A. M. S.*, V. 357, N. 5, 1753-1778, 2004
- [82] W.X. Ma, Lump solutions to the KP equation, *Physics Letters A*, 1975-1978, 2015
- [83] W.X. Ma, Linear superposition of Wronskian rational solutions to the KdV equation, *Commun. Theor. Phys.*, V. 73, N. 6, 065001-1-5, 2021

- [84] V.B. Matveev, Generalized Wronskian Formula for solutions of the KdV equation, Phys. Lett. A 166, 205-208, 1992
- [85] M.S. Osman, A.M. Wazwaz, An efficient algorithm to construct multi-soliton rational solutions of the (2 + 1)-dimensional KdV equation with variable coefficients, Appl. Math. Comput., V. 321, 282-289, 2018 Appl. Math. Comput. 321 (2018), 282-289.
- [86] N.J. Zabusky, M.D. Kruskal, Interaction of solitons in a collisionless plasma and the recurrence of initial states, Phys. Rev. Lett., V. 15, 240-243, 1965
- [87] V.E. Zakharov, L.D. Faddeev, Korteweg-de Vries equation: A completely integrable Hamiltonian system, Func. Anal. and its Appl., V. 5, N. 4, 280-287, 1971