



HAL
open science

Metabolic balance of [3H]-BADGE in rodents: a spotlight on the complex in vivo fate of the diglycidyl ether analogue of bisphenol A

Elodie N Person, Lidia de la Fuente, Isabelle Jouanin, Emilien L Jamin, Florence Blas-Y-Estrada, Sandrine Bruel, Daniel Zalko, Nicolas J Cabaton

► To cite this version:

Elodie N Person, Lidia de la Fuente, Isabelle Jouanin, Emilien L Jamin, Florence Blas-Y-Estrada, et al.. Metabolic balance of [3H]-BADGE in rodents: a spotlight on the complex in vivo fate of the diglycidyl ether analogue of bisphenol A. Gordon Research Conference on Environmental Endocrine Disruptors, Jun 2018, Les Diablerets, Vaud, Switzerland. hal-04678671

HAL Id: hal-04678671

<https://hal.science/hal-04678671>

Submitted on 27 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Introduction

Bisphenol A diglycidyl ether = BADGE (2,2-Bis[4-(glycidyloxy)phenyl]propane, CAS N°1675-54-3, MW 340.4)

- Used in the manufacture of **epoxy resins** and **polycarbonates** for **food packaging**
- Synthesized by a condensation reaction between **epichlorohydrin** and **Bisphenol A**
- Protective coatings**: car and marine industry, food packaging, adhesives, dental sealants, etc.^[1, 2]
- Reported **human daily intake**: approximately **0.16 µg/kg b.w./day**^[2, 3]




- Tolerable Daily Intake, BADGE + hydrolysis products = 0.15 mg/kg b.w.^[4]
- Migration** of BADGE + hydrolysis products does not exceed: 9 mg/kg in food, 1 mg/kg for chlorinated derivatives^[4]
- Possible **endocrine disruptor** (PPARγ antagonist)^[5, 6]

Objectives: Investigate the **fate of BADGE** in male and female Wistar rats
Explore BADGE metabolic balance and identify the different metabolic pathways involved in its **biotransformation** for **toxicological purposes**
Identify candidate biomarker metabolites, with the perspective of further human biomonitoring studies

Methodology

Chemicals:



[³H]-BADGE (radiochemical purity > 97 %, specific activity 10.9 GBq/mmol) was synthesized in-house from radio-labelled [³H]-Bisphenol A (Moravek). Unlabeled BADGE was purchased from Sigma Aldrich.

Animal experiment:

50 µg/kg [³H]-BADGE (gavage)

3 male and 3 female 8-week Wistar rats were forced an adjusted single [³H]-BADGE dose of **50 µg/kg b.w.** (1.6 MBq/kg b.w.) dissolved in corn oil. After **24h** spent in individual stainless steel metabolic cages, with water and food supplied *ad libitum*, rats were weighed and euthanized.

Metabolic Balance and Tissue Distribution:

Urine, feces, blood, liver, genital tract, glands, adipose tissue, muscle, digestive tract (and its content), brain, kidney, spleen, lung, heart, bladder, pancreas, skin and the remaining carcass were **collected for radioactivity quantification**. **Extractions** were performed on feces and digestive contents. Radioactivity was measured in liquid samples using a **Packard scintillation analyzer** (a)

Tri-Carb 2910TR, with Ultima Gold as the scintillation cocktail (Perkin Elmer). Feces were dried and ground before **solubilization** in **Soluene** (Perkin Elmer). Radioactivity in homogenized tissues, organs, blood, and carcasses was determined by complete **combustion** using an **Oxidizer 307** (b, Perkin Elmer), followed by quantification on a scintillation analyzer. At least 3 replicates were analyzed for each sample.

Metabolic profiling and identification:

High-performance-liquid-chromatography coupled to on-line radioactivity detection (Radio-HPLC). Excreted metabolites were separated on an Ultimate 3000 (c, ThermoScientific) HPLC system, with a Zorbax SB-C18 column (5 µm ; 4.6 x 250 mm, Agilent) coupled to a Flo-One A610TR detector (d, Perkin Elmer), using 2 mL/min Flow-Scint II (Perkin Elmer) as the scintillation cocktail.

Analytical conditions:

Flow rate	1 mL/min	Column temperature	25 °C	Analysis time	55 min
Phases	A: Ammonium acetate buffer 20 mM pH 3,5 / Acetonitrile (95:5 v/v) B: Ammonium acetate buffer 20 mM pH 3,5 / Acetonitrile (10:90 v/v)				
Gradient	0-4 min: 100 % A, 4-6 min: 100-75 % A, 6-16 min: 75 % A, 16-18 min: 75-70 % A, 18-25 min: 70 % A, 25-27 min: 70-30 % A, 27-38 min: 30 % A, 38-40 min: 30-0 % A, 40-50 min: 0 % A, 50-51 min: 0-100 % A, 51-55 min: 100 % A				

Preliminary results

Metabolic balance of [³H]-BADGE following a single oral dose administered to male and female Wistar rats

For each animal, radioactivity remaining in the feeding material was quantified to calculate the nominal [³H]-BADGE dose administered: **46.8 ± 2.2 µg/kg b.w.** (1.5 ± 0.07 MBq/kg b.w.).




Fig. 1: 24h metabolic balance in male (A) and female (B) Wistar rats. Results (mean, n = 3 for each group) are expressed as % of the administered radioactivity (t-test, *** p-value ≤ 0.001, Graphpad®)

- 92.1 ± 9.4 %** of the administered radioactivity recovered at 24h following BADGE administration (n = 6).
- Feces was the main route of excretion** (55.1 ± 16.8 % in males, 66.2 ± 6.7 % in females), and the **digestive tract** accounted for 27.5 ± 6.7 % of the administered radioactivity in males and 15.8 ± 4.1 % in females, respectively.
- Significant sex-related differences were observed for **urine**: 4.5 ± 0.9 % recovered in males, 11.8 ± 0.9 % in females.




Fig. 3: Residual levels of radioactivity in tissues of male and female Wistar rats. Results (mean ± SEM) are expressed in ng of BADGE equivalent/g tissue (ppb) (t-test, * p-value ≤ 0.05, Graphpad®)

Digestive tract contents




Fig. 2: 24h metabolic balance: residual radioactivity in digestive tract contents of male and female Wistar rats. Results (mean ± SEM) are expressed in µg of BADGE equivalent (t-test, * p-value ≤ 0.05, Graphpad®)

STO: stomach, DUO: duodenum, JEJ: jejunum, ILE: ileon, CAE: caecum, COL: colon

Radiochromatographic profiling of urine and feces




Fig. 4: Typical Radio-HPLC profiles for urine and feces collected over 24h

- Sex-related differences** were highlighted for **urine and fecal metabolite profiles**
- Complex radiochromatographic profiles underline the **chemical reactivity of BADGE**

Conclusions

In this **first *in vivo* 24h-study** of [³H]-BADGE carried out in male and female Wistar rats, radioactivity was found to be **excreted** mainly in **feces**, with residues found in all tissues, including reproductive organs. **Sex-related differences** were highlighted for **urine, liver, lung** as well as **duodenum, caecum and colon contents**. Qualitative differences are also suggested by first Radio-HPLC analyzes carried out for urine and feces. BADGE elimination over 24h (urine + feces) appears to be faster in female rats compared to males (77.9 ± 6.2 % in females vs. 59.6 ± 16.2 % in males). Preliminary HPLC Radio-profiling analyzes strongly suggest a **complex metabolism** of this molecule. **Further HPLC-MS/MS investigations** are being carried out, with the aim to identify unknown **BADGE metabolites**.

References

[1] Xue J. *et al.*, 2018, Environ Res. 162:35-40.
 [2] Poole A. *et al.*, 2004, Food Addit Contam. 21(9):905-19.
 [3] Wang L. *et al.*, 2015, Environ Sci Technol. 49(5):3150-7.
 [4] European Commission regulation N°1895/2005 on the restriction of use of certain epoxy derivatives in materials and articles intended to come into contact with food, 2005

[5] Wright HM. *et al.*, 2000, J Biol Chem. 275(3):1873-7.
 [6] Satoh K. *et al.*, 2004, Food Chem Toxicol. 42(6):983-93.

All the experimental procedure was carried out in accordance with the accepted standards of animal care and validated by the local ethical committee for animal experimentation N°086 (project number: APAFIS#1308-201706161630824 v3) and by the French Ministry of Higher Education, Research and Innovation (agreement: C3155513)