

A photograph of a person standing on a beach at night, holding a flashlight. The beam of light illuminates the sandy ground and the water. The sky is dark with some stars visible.

AGU23

WIDE. OPEN. SCIENCE.

#AGU23

Influence of aquatic metabolism on marsh carbon dynamics and associated atmospheric CO₂ fluxes

Mayen J.^{1,2}, Polsenaere P.², Regaudie de Gioux A.³, Deborde J.², Collin K.¹, Le Merre Y.¹, Foucault É.⁴, Ouisse V.⁴, André L.^{5,6}, Arnaud M.^{7,2}, Kostyrka P.², Lemeele J.-C.⁸, Lelong V.⁹, Vagner M.¹⁰, Souchu P.¹

¹IFREMER, Littoral, LER/MPL, France, ²IFREMER, Littoral, LER/PC, France, ³IFREMER, Dyneco, Pelagos, France, ⁴MARBEF (Montpellier Univ., CNRS, Ifremer, IRD), France, ⁵BRGM, France, ⁶STO (Orléans Univ., CNRS, BRGM), France, ⁷Institute of Ecology and Environmental Sciences Paris, Sorbonne Univ., France, ⁸LPO, Réserve Naturelle Nationale de Lilleau des Niges, France, ⁹LPO, Réserve Naturelle de Moëze-Oléron, France, ¹⁰LEMAR (CNRS, Brest Univ., IRD, Ifremer), France

AGU23

B022 - Coastal Wetland Carbon and Nitrogen Cycles: Recent Advances in Measurements, Modeling, and Syntheses – 12/16/2023

WIDE. OPEN. SCIENCE.

AGU23

MAYEN JÉRÉMY

Ph.D. student in coastal biogeochemistry
at IFREMER





WIDE. OPEN. SCIENCE.

AGU23 ➤ MARSH CO₂ PUMP AND CARBON EXPORT


- « Salt marshes are coastal wetlands that are flooded and drained by salt water brought in by the tides » (NOAA)

Processes and fluxes of carbon in salt marshes:



WIDE. OPEN. SCIENCE.

What is the contribution of metabolic processes in the water column to carbon dynamics in a tidal salt marsh?




WIDE. OPEN. SCIENCE.

AGU23 > STUDY SITE: A TIDAL SALT MARSH

Klijn et al. (2015)

Mayen et al. in prep.



01/03/2021, low tide

22/07/2021, incoming tide

27/04/2021, high tide




WIDE. OPEN. SCIENCE.

AGU23

SAMPLING STRATEGY AND FIELD TECHNIQUE

Mayen et al. in prep.




AGU23 ➤ TEMPORAL VARIATIONS OF BIOGEOCHEMICAL PARAMETERS


Mayen et al. in prep.

→ nMDS (dist. eucl.) with Chla, pCO₂, DIC, TA, POC, DOC, PON, DON, DIN, DIP, DSi

A




C



Maybe put all the figures
at the top to avoid blank
space

B



- Strong control of seasonal factor (between 24-h cycles) and tidal factor (high tide vs low tide) **is not only winter that is different?**
- Low control of diurnal factor (daytime vs night-time)


WIDE. OPEN. SCIENCE.

AGU23 ➤ TEMPORAL VARIATIONS OF WATER pCO₂

Mayen et al. in prep.

pCO₂(water) < pCO₂(air) : undersaturated in CO₂ with respect to the atmosphere (CO₂ sink)

pCO₂(water) > pCO₂(air) : oversaturated in CO₂ with respect to the atmosphere (CO₂ source)



shelf sea?

- Marsh waters (high tides) influenced by shelf: slightly oversaturated in CO₂ (except in spring)

- Channel waters (low tides) influenced by salt marshes: strongly oversaturated in CO₂ in winter (LT/Night) and strongly undersaturated in CO₂ in spring and summer (LT/Day and LT/Night)

what about fall?

AGU23


HORIZONTAL EXCHANGES IN CARBON

Mayen et al. in prep.


very interesting

$$\text{Total Carbon (TC)} = \text{POC} + \text{DOC} + \text{DIC} (\text{HCO}_3^- + \text{CO}_3^{2-} + \text{CO}_2)$$


$$\text{Total Nitrogen (TN)} = \text{PON} + \text{DON} + \text{DIN} (\text{NO}_3^- + \text{NO}_2^- + \text{NH}_4^+)$$



- Winter: NH_4^+ and DIC (HCO_3^-) increases from high to low tide (tidal pumping) \rightarrow strong mineralization of marsh organic matter
- Spring and summer: DOC and DON increases from high to low tides **WIDE. OPEN. SCIENCE.**
- strong marsh primary production (phytoplankton, macroalgae and terrestrial macrophytes)





- Autotrophy at day (except to C2-HT/Day and C4-HT/Day) and heterotrophy at night (except to C2-HT/Night)
- Higher production and respiration rates of planktonic communities in channel waters (LT) than in coastal waters (HT)
- Highest planktonic autotrophy : C2-LT/Day



AGU23 ➤ CONCLUSIONS

- Winter: microbial mineralization **IS IT FROM ORGANIC MATTER? IT IS QUITE COUNTERINTUITIVE** of organic matter in the salt marsh → strong increases of DIC (and DIP, NH_4^+) by tidal pumping
- Spring and summer: lateral exportation of CO_2 -depleted and DOC-concentrated waters (high primary production in the salt marsh)
- Significant contribution of planktonic metabolism to CO_2 undersaturation water exports (and microphytobenthos on mudflats) **very nice, often microphytobenthos are overlooked in budget**
- Horizontal exchanges in salt marshes has a major impact on water carbon dynamics
- Low contribution of planktonic metabolism on the atmospheric C uptake by the salt marsh footprint



WIDE. OPEN. SCIENCE.

12


THANK YOU




Important role of salt marshes in horizontal
and vertical exchanges of carbon

AGU23

WIDE. OPEN. SCIENCE.









WIDE. OPEN. SCIENCE.



A



B



WIDE. OPEN. SCIENCE.