



Underwater Fluxes of biogenic constituents and their Ecological Implications

Jérémie Mayen, Pierre Polsenaere, Philippe Geairon, Jean-Michel Chabirand, Jonathan Deborde, James Grizon, Jean-Christophe Lemesle, Benoit Poitevin, Christine Dupuy, Marie Vagner

► To cite this version:

Jérémie Mayen, Pierre Polsenaere, Philippe Geairon, Jean-Michel Chabirand, Jonathan Deborde, et al.. Underwater Fluxes of biogenic constituents and their Ecological Implications. ASLO, Jun 2021, Online, France. hal-04678865

HAL Id: hal-04678865

<https://hal.science/hal-04678865v1>

Submitted on 27 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spatial and temporal variations in water pCO₂ and atmospheric CO₂ exchanges over a temperate salt marsh system

*J. Mayen^a, P. Polsegaere^a, P. Geairon^a, J.-M. Chabirand^a,
J. Debord^a, J. Grizon^a, J.-C. Lemesle^b, B. Poitevin^c,
C. Dupuy^d, M. Vagner^e*

^a IFREMER, LER-PC, ^b LPO, Réserve Naturelle de Lilleau des Niges,
^c Pôle-Nature de l'Ecomusée du Marais Salant, ^d LIENSS UMR7266
CNRS-Université La Rochelle, ^e LEMAR, UMR 6539 CNRS



AGIR pour la
BIODIVERSITÉ




ifremer

MARSH CO₂ PUMP AND EXPORT


Ifremer

- « Salt marshes are coastal wetlands that are flooded and drained by salt water brought in by the tides. »
(National Oceanic and Atmospheric Administration)

(Acclimaterra, 2018)



Dyked salt marshes



(Watson, 2015)

- CO₂ sink, organic carbon burial and C export by the tide :

Tg C yr⁻¹ (10¹² g C yr⁻¹)

Intertidal marsh
(5.0 × 10⁹ m²)

NPP/CO₂
Fixation = 10.1

Estuary/bay
(8.75 × 10⁹ m²)

Inner shelf
(28.9 × 10⁹ m²)

(Cai, 2011)

Degassing = 1.64

Degassing = 0.41

Degassing = 1.88


River input
DIC = 0.6
DOC = 0.9
Marsh DIC export = 0.83
Expected marsh OC export = 5.5

Total terrestrial DIC export = 1.5, OC = 6.4
Net DIC export = 2.6–5.2
Expected OC export = 2.3–4.9

OC burial = 0.29

► OBJECTIVES

- Study the dynamics of (i) water pCO₂ and (ii) atmospheric CO₂ exchanges in relation to biotic and abiotic controlling factors along an intertidal bay – channel – salt marshes continuum at seasonal, diurnal and tidal scales



Understanding the evolution of the ecological functions of coastal wetlands (CO₂ sinks, C sequestration in sediments) in the face of global change.






FIER D'ARS COASTAL WETLAND - FRANCE




- Intertidal bay on the Ré island – France
- Complex of salt marshes and anthropised mudflats (1,200 ha): **coastal wetland**

- Bay – channel – salt marshes continuum



4 interconnected stations with different management practices and configurations



01/05/2019



Station b: intertidal system

10/02/2020



Station c: nature reserve

11/02/2020



Station d: salt-farming activity




Station b: intertidal system

BIOGEOCHEMICAL MEASUREMENTS

- *In situ* measurements of pCO_2 , $[\text{chl } \alpha]$ and biogeochemical parameters (oxygen, salinity, temperature ...)
- 24-hour cycles of *in situ* measurements at the seasonal and spatial scales → high frequency (1 minute)

- 5 multiparameter and specific probes (autonomous) in sub-surface waters:

- C-SENSE (X2): pCO_2 measurements (0-2000 ppmv + 0-4000 ppmv, $\% \text{Er} = 3\%$ of range)
- EXO2 (YSI): biogeochemical parameter measurements (T , S , TU , pH , O_2)
- Aquatic MiniPAR (PME)
- C3-fluorimeter: chlorophyll- α concentration



► pCO₂ DYNAMICS ALONG THE CONTINUUM

1. Major influence of station typologies on pCO₂

- a: Fier d'Ars intertidal bay
- b: Vieux port channel
- c: A1 salt marsh (nature reserve)
- d: Loix salt marsh (salt-farming activity)


Intertidal bay and channel (a and b)

- Strong tidal influences → High turbidity → limited phytoplankton activity
- Few aquatic vegetation


Salt marshes (c and d)

- Photosynthesis of primary producers (macroalgae/ phytoplankton/ *Ruppia maritima*)

PCA - Biplot




Source: factoextra



► SAISONAL pCO₂ DYNAMIC IN MARSHES


2. Major influence of management practices on pCO₂

- Summer 2019:**
 - **c:** photosynthesis activity of aquatic macroalgae
 - **d:** zooplankton bloom → phytoplankton cell degradation



Station c, 22/07/2019

Autumn 2019: ➤ **c:** Macroalgae degradation → Organic matter mineralisation → high pCO₂



Winter 2020: no significant difference on pCO₂


$$\text{pCO}_2(c) > \text{pCO}_2(d)$$

- a:** Fier d'Ar's intertidal bay
- b:** Vieux port channel
- c:** A1 salt marsh (nature reserve)
- d:** Loix salt marsh (salt-farming activity)

DIURNAL AND TIDAL pCO_2 DYNAMICS IN A MARSH

Ifremer

- Indigenous biological activity (primary production/respiration) at the diurnal scale
- Coastal water incomings (tidal advection): increase pCO_2 in winter, spring and summer




METABOLISM OF FIER D'ARS COASTAL WETLAND



- Estimated water-air CO_2 flux (mmol $\text{CO}_2 \text{ m}^{-2} \text{ h}^{-1}$): $\text{FCO}_2 = \alpha \ k (\text{pCO}_2\text{eau} - \text{pCO}_2\text{air})$

(Takahashi et al., 2002)



- α : Fier d'Ars intertidal bay
 b : Vieux port channel
 c : A1 salt marsh (nature reserve)
 d : Loix salt marsh (salt-farming activity)

➤ CONCLUSION AND PERSPECTIVES

- **Significant influences of pCO₂ dynamics:**
 - ✓ Station typologies within the coastal zone (Bay, channel, marsh)
 - ✓ Management practices: water turnover
- Diurnal and tidal scales: **non-temperature effects on pCO₂ dynamics**
 - ✓ Biological activity (photosynthesis/respiration)
 - ✓ Different tidal effects according to water column CO₂ saturation
- **Métabolism:**
 - ✓ Channel: CO₂ source to atmosphere
 - ✓ Salt marshes: CO₂ sink
- Study: (1) the atmospheric CO₂ fluxes by Eddy Covariance (*in situ* measurements)
(2) the salt marsh aquatic metabolism

Jérémie MAYEN, PhD student
Ifremer (French Research Institute for the Exploitation of the Sea)

E-mail : jeremy.mayen@ifremer.fr
Lab page : <https://wwz.ifremer.fr/littoral/Laboratoires-Environnement-Ressources/LER-Pertuis-Charentais>



Thank you for your attention !