



HAL
open science

Members of the gut microbiota modulate the bioavailability of the antiseizure medication lamotrigine

Zehra Esra Ilhan, Catherine Philippe, Andrea Marchetto, Philippe Gérard, Stéphane Auvin, Patricia Lepage

► To cite this version:

Zehra Esra Ilhan, Catherine Philippe, Andrea Marchetto, Philippe Gérard, Stéphane Auvin, et al.. Members of the gut microbiota modulate the bioavailability of the antiseizure medication lamotrigine. 19th International Symposium on Microbial Ecology, Aug 2024, Cape Town / Le Cap, South Africa. hal-04678898

HAL Id: hal-04678898

<https://hal.science/hal-04678898v1>

Submitted on 27 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Members of the gut microbiota modulate the bioavailability of the antiseizure medication lamotrigine



Zehra Esra Ilhan^a, Catherine Philippe^a, Andrea Marchetto^a, Philippe Gérard^a, Stéphane Auvin^{b,c,d}, Patricia Lepage^a

^aUniversité Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, Domaine de Vilvert, Jouy-en-Josas, 78350, France; ^bUniversité de Paris, INSERM NeuroDiderot, Paris, France; ^cAPHP, Robert Debré University Hospital, Pediatric Neurology Department, Paris, France; ^dInstitut Universitaire de France, (IUF), Paris, France

Background




Antiseizure medications are not effective in 1/3 of the pediatric patients.

OBJECTIVES

- Assess the impact of antiseizure medications on gut microbiome dynamics and function
- Elucidate the role of gut microbiome on bioavailability of lamotrigine

Methods




Results


1. Among the 10 antiseizure medications, lamotrigine disturbed the microbial communities the most.

Lamotrigine had an impact similar to kanamycin.


- ASMs:**
valproate, lacosamide, levetiracetam, carbamazepine, topiramate, lamotrigine, gabapentin, ethosuximide, zonisamide, phenytoin.
- Antibiotics:**
kanamycin, ciprofloxacin, azithromycin, vancomycin, chloramphenicol




3. Lamotrigine disappearance rate was inoculum specific.



4. Lamotrigine-degrading cultures were enriched in Enterococcus durans, Phocaeicola dorei, Eggerthella sp. and Faecalibacterium prausnitzii species.




Lamotrigine treatment reduced predicted abundance of oxidoreductases but increased ligases and hydrolases.




Based on predicted genes, lamotrigine treated and DMSO-control samples formed separate clusters.

2. Lamotrigine influenced the relative abundance of key genera in a dose-dependent manner.




5. Lamotrigine impacted microbiome function more than its composition: altered the production and consumption of 114 metabolites.



Production of several amino acid metabolites were reduced and catabolism of dipeptides were enriched in lamotrigine treated cultures.

6. Genera in lamotrigine-degrading cultures positively correlated with methionine and urea metabolites.



Summary

- Lamotrigine disturbed microbial community dynamics.
- Microbial communities dominated by *E. durans*, *P. dorei*, *Eggerthella* sp. and *F. prausnitzii* reduced bioavailability of lamotrigine.
- Dipeptide consumption and production of several amino acid metabolites increased in lamotrigine treated microbial communities.

