



Spatial and ontogenetic variability of omega-3 fatty acid composition in European sea bass *Dicentrarchus labrax* from two French estuaries

Mickaël Péron, Antoine Bideau, Fabienne Le Grand, David Mazurais,
Philippe Soudant, Marie Vagner

► To cite this version:

Mickaël Péron, Antoine Bideau, Fabienne Le Grand, David Mazurais, Philippe Soudant, et al.. Spatial and ontogenetic variability of omega-3 fatty acid composition in European sea bass *Dicentrarchus labrax* from two French estuaries. Colloque d'Ecophysiologie Animale, 2021, Montpellier, France. hal-04678966

HAL Id: hal-04678966
<https://hal.science/hal-04678966v1>

Submitted on 27 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spatial and ontogenetic variability of omega-3 fatty acid composition in European sea bass *Dicentrarchus labrax* from two French estuaries

Mickaël Péron, Antoine Bideau, Fabienne Le Grand, David Mazurais, Philippe Soudant, Marie Vagner

Lipids

Source & **reserve** of energy → Triglycerides → **Neutral lipids (NL)**

Dietary influence

Skalli *et al*, 2006

Lipids

Source & **reserve** of energy → Triglycerides → **Neutral lipids (NL)**

Dietary influence

Skalli *et al*, 2006

Structure & function of membranes → Phospholipids → **Polar lipids (PL)**

Lipids


Source & **reserve** of energy → Triglycerides → **Neutral lipids (NL)**

Dietary influence


Skalli *et al*, 2006

Structure & function of membranes → Phospholipids → **Polar lipids (PL)**


Fatty acids



Saturated
(*e.g* Palmitic acid)



Mono-unsaturated
(*e.g* Oleic acid)



Poly-unsaturated
(*e.g* Eicosapentaenoic acid)

Lipids

Source & **reserve** of energy → Triglycerides → **Neutral lipids (NL)**

Dietary influence

Skalli *et al*, 2006

Structure & function of membranes → Phospholipids → **Polar lipids (PL)**

Fatty acids

**Long chain n-3
Poly-unsaturated** →



Physiology

Growth

Disease preventing

Behavior

Neural developement



Essential for humans and animals

Lipids

Source & **reserve** of energy → Triglycerides → **Neutral lipids (NL)**


Dietary influence

Skalli *et al*, 2006

Structure & function of membranes → Phospholipids → **Polar lipids (PL)**

Fatty acids

EPA





20:5 n-3


DHA



22:6 n-3




EPA



DHA

- Low / no ability to synthesize n-3 fatty acids for fish

-> *Limited efficiency of enzymes (elongases & desaturases)*




- Low / no ability to synthesize n-3 fatty acids for fish

-> *Limited efficiency of enzymes (elongases & desaturases)*

- Difference between marine and freshwater systems

-> *No selective pressure on highly abundant n-3 systems*



- Low / no ability to synthesize n-3 fatty acids for fish

-> *Limited efficiency of enzymes (elongases & desaturases)*

- Difference between marine and freshwater systems

-> *No selective pressure on highly abundant n-3 systems*

What about euryhaline fish ?


European sea bass

Dicentrarchus labrax :

juvenile stages in estuaries

Habitat / life stage effect on n-3 tissue composition

Low in-natura information




What about euryhaline fish ?



Dicentrarchus labrax


- How is the n-3 FA composition modulated ?
- How does it vary in natural habitat
- Does it depend on sampling site / life stage ?



NourDem

Summer 2019 (July-August)

Three different life
stages : G₁ / G₂ / G₃




$$N = 12 - G_1$$

$$N = 10 - G_2$$

$$N = 12 - G_3$$

76 juveniles



$$N = 18 - G_1$$

$$N = 12 - G_2$$

$$N = 12 - G_3$$




Lipid extraction and separation


Reserve lipids (NL)
Membrane lipids (PL)



Liver = Highly influenced
by trophic differences

Caballero *et al*,
1999




Fatty acid profiles with
GC-FID







Percentage of main FA in reserve lipids (NL)




Percentage of main FA in membrane lipids (PL)





Seine estuary



Percentage of main FA in reserve lipids (NL)





Percentage of main FA in membrane lipids (PL)




- High variations in reserve lipids -> Dietary influence
- DHA = main FA in membranes
- EPA / DHA ratio matching litterature (3x factor)



Quantity of DHA in reserve lipids (NL)




Quantity of DHA in reserve lipids (NL)





Quantity of DHA in membrane lipids (PL)



Quantity of DHA in reserve lipids (NL)





Quantity of DHA in membrane lipids (PL)




- Lower DHA quantity in G1
- Prey lipid composition influence ?
- DHA retention in tissue : PL > NL
- Retention / Synthesis plasticity : *Influence of environment / trophic level*



Quantity of EPA in reserve lipids (NL)




Quantity of EPA in reserve lipids (NL)





Quantity of EPA in membrane lipids (PL)



Quantity of EPA in reserve lipids (NL)





Quantity of EPA in membrane lipids (PL)




- Different profiles between reserve & membrane
DHA & EPA = different roles
Tocher, 2003
- Used for eicosanoids production ?
- Used to be converted into DHA?



Percentage of main FA in reserve lipids (NL)




Percentage of main FA in reserve lipids (NL)





Percentage of main FA in membrane lipids (PL)



Percentage of main FA in reserve lipids (NL)





Percentage of main FA in membrane lipids (PL)



- EPA and DHA levels varying with age -> Different trophic dynamics ?
- EPA / DHA used as energy source ?


Quantity of DHA in reserve lipids (NL)

NS





Quantity of DHA in reserve lipids (NL)

NS




Quantity of DHA in membrane lipids (PL)

NS





Quantity of DHA in reserve lipids (NL)

NS





Quantity of DHA in membrane lipids (PL)

NS




- No trophic differences or membrane differences -> Effect of sampling sites ?

Quantity of EPA in reserve lipids (NL)




Quantity of EPA in reserve lipids (NL)




NS


Quantity of EPA in membrane lipids (PL)





NS



Quantity of EPA in reserve lipids (NL)



Quantity of EPA in membrane lipids (PL)



- No trophic differences or membrane differences -> Effect of sampling sites

What was observed *in natura* ?

Selective DHA retetion in membrane lipids in
Loire & Seine

What was observed *in natura* ?

Selective DHA retetion in membrane lipids in
Loire & Seine

Life stage effect on DHA composition in
Seine

What was observed *in natura* ?

Selective DHA retetion in membrane lipids in
Loire & Seine

Life stage effect on DHA composition in
Seine

No differences for EPA quantities in
membrane lipids

Many factors could influence the levels of EPA / DHA


- Habitat related factors :

- Trophic chain length
- Abundance of preys
- Water parameters

Many factors could influence the levels of EPA / DHA

- Habitat related factors :

- Trophic chain length
- Abundance of preys
- Water parameters




Otoliths micro-chemistry : elemental composition of water fish
lived in : salinity, metals

Many factors could influence the levels of EPA / DHA

- Habitat related factors :

- Trophic chain length
- Abundance of preys
- Water parameters



Otoliths micro-chemistry : elemental composition of water fish
lived in : salinity, metals

- Individuals related factors :

- Capacities of retention / synthesis




Transcriptomic data on n-3 PUFA metabolism genes :
elongases, desaturases

Many factors could influence the levels of EPA / DHA

- Habitat related factors :

- Trophic chain length
- Abundance of preys
- Water parameters



Otoliths micro-chemistry : elemental composition of water fish
lived in : salinity, metals

- Individuals related factors :

- Capacities of retention / synthesis
- Type of tissue



Analyzing other tissues : Muscle, brain



A wide-angle photograph of a harbor at sunset. The sky is filled with dramatic orange and yellow clouds. The sun is low on the horizon, casting a bright reflection on the calm water. On the left, there's a row of multi-story buildings, some with signs like "PROSPECTUS" and "HOTEL". In the center, a white bridge arches across the water. Several sailboats and small boats are moored along the right bank, which is lined with dark trees. The overall atmosphere is peaceful and scenic.

*Thank you for
your attention*