



HAL
open science

Chemical distance for smooth Gaussian fields in higher dimension

David Vernotte

► **To cite this version:**

David Vernotte. Chemical distance for smooth Gaussian fields in higher dimension. 2024. hal-04681135

HAL Id: hal-04681135

<https://hal.science/hal-04681135v1>

Preprint submitted on 29 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Chemical distance for smooth Gaussian fields in higher dimension.

David Verotte

August 29, 2024

Abstract

Gaussian percolation can be seen as the generalization of standard Bernoulli percolation on \mathbb{Z}^d . Instead of a random discrete configuration on a lattice, one considers a continuous Gaussian field f and studies the topological and geometric properties of the random excursion set $\mathcal{E}_\ell(f) := \{x \in \mathbb{R}^d \mid f(x) \geq -\ell\}$ where $\ell \in \mathbb{R}$ is called a level. It is known that for a wide variety of fields f , there is a phase transition at some ℓ_c . When $\ell > \ell_c$, the excursion set $\mathcal{E}_\ell(f)$ presents a unique unbounded component while if $\ell < \ell_c$ there are only bounded components in $\mathcal{E}_\ell(f)$. In the supercritical regime, $\ell > \ell_c$, we study the geometry of the unbounded cluster. Inspired by the work of Peter Antal and Agoston Pisztora for the Bernoulli model [1], we introduce the chemical distance between two points x and y as the Euclidean length of the shortest path connecting these points and staying in $\mathcal{E}_\ell(f)$. In this paper, we show that when $\ell > -\ell_c$ then with high probability, the chemical distance between two points has a behavior close to the Euclidean distance between those two points.

Contents

1	General introduction and notations	2
2	On the local uniqueness event	6
2.1	Approximation of the field	8
2.2	Deterministic argument for local uniqueness	9
2.3	Proof of Proposition 2.4	11
3	Stochastic domination argument	14
3.1	Comparison with a Bernoulli process of high parameter	14
3.2	Global structures	15
4	Local control of the chemical distance and proof of the main theorem	17
4.1	Local control of the chemical distance around a point	17
4.2	Proof of Theorem 1.8	22

1 General introduction and notations

This paper studies some geometric properties of excursion sets of some random Gaussian field in the Euclidean space of dimension $d \geq 2$. The continuous model of random Gaussian field and the percolation model associated to it have gathered a lot of interest over the past few years (see for instance [3],[15],[12],[18],[6] for an incomplete list of references). We present the model as a continuous analogue of the classical discrete percolation (Bernoulli percolation). In the discrete case, say site percolation on \mathbb{Z}^d for convenience, one consider a configuration $(\omega_i)_{i \in \mathbb{Z}^d} \in [0, 1]^{\mathbb{Z}^d}$, where each ω_i is a uniform random variable over $[0, 1]$ and all ω_i are mutually independent. Given a parameter $p \in [0, 1]$ one can consider the random set

$$\mathcal{G}_p(\omega) := \{i \in \mathbb{Z}^d \mid \omega_i \leq p\}. \quad (1)$$

The set \mathcal{G}_p is a random set and its properties are well known. In particular it is known that there is a phase transition at some $p_c(d) \in]0, 1[$, when $p < p_c(d)$ all clusters of $\mathcal{G}_p(\omega)$ are finite (almost surely) while when $p > p_c$, $\mathcal{G}_p(\omega)$ presents a unique infinite cluster. Note that the configuration ω can be seen as a random function

$$\omega : \mathbb{Z}^d \rightarrow [0, 1].$$

The continuous model replaces ω with a random continuous function

$$f : \mathbb{R}^d \rightarrow \mathbb{R},$$

and for a level $\ell \in \mathbb{R}$ the analogue of $\mathcal{G}_p(\omega)$ is

$$\mathcal{E}_\ell(f) := \{x \in \mathbb{R}^d \mid f(x) \geq -\ell\}. \quad (2)$$

In the following, we properly define the model, explaining how the random function f is defined. We then give the main theorems related to the percolation model and finally we introduce the chemical distance and state our main theorem.

In the rest of the paper, f will denote a centered, stationary, continuous, Gaussian field on \mathbb{R}^d . That is, f is a random collection $(f(x))_{x \in \mathbb{R}^d} \in \mathbb{R}^{\mathbb{R}^d}$ such that

- For all $x_1, \dots, x_n \in \mathbb{R}^d$ the vector $(f(x_1), \dots, f(x_n))$ is a random Gaussian vector that is centered (Gaussian field).
- For all $x \in \mathbb{R}^d$, $f(x + \cdot)$ and $f(\cdot)$ have the same law (stationarity).
- Almost surely $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is continuous.

An example of such a field is the Bargmann-Fock field. Consider $(a_{i_1, \dots, i_d})_{i_1, \dots, i_d \geq 0}$ a countable family of mutually independent standard Gaussian random variables. If $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ we define

$$f(x) := e^{-\frac{1}{2}\|x\|^2} \sum_{i_1, \dots, i_d \geq 0} a_{i_1, \dots, i_d} \frac{x_1^{i_1} \dots x_d^{i_d}}{\sqrt{i_1! \dots i_d!}}. \quad (3)$$

Here and in the following, $\|\cdot\|$ denotes the usual Euclidean norm. A simple computation shows that f is well defined and by dominated convergence it is almost surely continuous (in

fact \mathcal{C}^∞). Moreover f is clearly a Gaussian field that is centered and $\mathbb{E}[f(x)f(y)] = e^{-\frac{1}{2}\|x-y\|^2}$ depends only on $x - y$ implying that f is stationary. It is good to have in mind that our study applies for this specific field. However, we present another way to construct such fields. This is done via the white noise representation.

Definition 1.1. A white noise W on \mathbb{R}^d is a centered Gaussian field indexed by function of $L^2(\mathbb{R}^d)$ such that for any $\varphi_1, \varphi_2 \in L^2(\mathbb{R}^d)$ we have

$$\mathbb{E}[W(\varphi_1)W(\varphi_2)] = \int_{\mathbb{R}^d} \varphi_1(x)\varphi_2(x)dx.$$

Consider a function $q : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying the following assumption for some $\beta > d$.

Assumption 1.2. Let $\beta > d$. We say that q satisfy Assumption 1.2 for β if

- q is \mathcal{C}^{10} and $\partial^\alpha q \in L^2(\mathbb{R}^d)$ for all $\alpha \in \mathbb{N}^d$ such that $\|\alpha\|_1 \leq 10$.
- $\max(|q(x)|, \|\nabla q(x)\|) = O(\|x\|^{-\beta})$.
- q is isotropic ($q(x)$ only depends on $\|x\|$).
- $q \geq 0$.
- q is not identically equal to the zero function.

Given a function q that satisfy Assumption 1.2 for some $\beta > d$ one can define

$$f := q * W,$$

where $*$ denotes convolution. That is, for all $x \in \mathbb{R}^d$ we set $f(x) := W(q(x - \cdot))$, where $q(x - \cdot)$ is the function $y \mapsto q(x - y)$ which is in fact in $L^2(\mathbb{R}^d)$. It follows that f is a centered Gaussian field. Its covariance kernel can be computed: $\mathbb{E}[f(x)f(y)] = (q * q)(x - y)$. This shows that the field f is stationary. Moreover, by dominated convergence one can instead consider a continuous (in fact \mathcal{C}^4) modification of f which we do in the rest of the paper.

Remark. If we set $q(x) = \left(\frac{2}{\pi}\right)^{d/4} e^{-\|x\|^2}$ then $f = q * W$ obtained has the law of the Bargmann-Fock field previously introduced. Thus, note that the Bargmann-Fock field satisfies Assumption 1.2 for all $\beta > d$.

The white noise representation of the field is useful to define approximations of the field that have good properties (like finite range dependence see Definitions 2.5 and 2.6).

We now introduce the percolation model associated to the continuous setting. Given a real parameter $\ell \in \mathbb{R}$ which we call a *level*, we define the *excursion set at level ℓ* as

$$\mathcal{E}_\ell(f) := \{x \in \mathbb{R}^d \mid f(x) \geq -\ell\}. \quad (4)$$

We remark that since f is random, the set $\mathcal{E}_\ell(f)$ is also a random set. It appears that for f fixed the set $\mathcal{E}_\ell(f)$ increases as ℓ increases. Thus, the probability that $\mathcal{E}_\ell(f)$ contains an unbounded component is increasing in ℓ and we define the critical probability for percolation as

$$\ell_c := \sup \{\ell \in \mathbb{R} \mid \mathbb{P}(\mathcal{E}_\ell(f) \text{ has no unbounded connected component}) = 1\}. \quad (5)$$

It turns out that when $d \geq 2$, then for a large class of Gaussian field we have $\ell_c \in]-\infty, \infty[$, that is, there exists a non trivial phase transition. It is important to have in mind that the critical level ℓ_c is to be compared with the critical probability $p_c(d)$ in the Bernoulli setting on \mathbb{Z}^d . For our continuous model, a lot is known about the case $d = 2$. In particular the following theorem can be thought of as an analogue of the Theorem of Kesten [9] concerning the value of the critical probability of Bernoulli bond percolation on \mathbb{Z}^2 .

Theorem 1.3 ([16], [12]). *In the case $d = 2$, if q satisfies Assumption 1.2 for some $\beta > 2$ then we have $\ell_c = 0$.*

In the case $d \geq 3$ we also have the following result.

Theorem 1.4 ([6]). *In the case $d \geq 3$, if q satisfies Assumption 1.2 for some $\beta > d$ then we have $\ell_c < 0$.*

Remark. One may remark that the critical parameter ℓ_c in the case of dimension $d = 2$ is mostly independent of the law of the field we consider it will always be 0, however for $d \geq 3$, the exact value of ℓ_c is always negative showing in words that is strictly easier to percolate in the space than in the plane, but the exact value of ℓ_c may depend on the exact law of f .

Remark. Note that the statements of the two theorems 1.3 and 1.4 may hold under weaker assumptions. We refer the reader to [12] and [6] to see the exact assumptions that were made on the law of f , as for the sake of clarity we prefer to not write the details here.

When $\ell > \ell_c$ we are in the supercritical phase, when $\ell < \ell_c$ we are in the subcritical phase. This phase transition has received a lot of attention over the past few years. One result of interest is the following

Theorem 1.5 ([18]). *If $d \geq 2$ and if q satisfy Assumption 1.2 for some $\beta > d$, then if $\ell > \ell_c$, the set $\mathcal{E}_\ell(f)$ almost surely presents a unique unbounded component.*

This theorem already gives a good description of the supercritical phase. The set $\mathcal{E}_\ell(f)$ contains only one unbounded connected component and (countably) many finite ones. In this article we aim to obtain a more geometric information about the unbounded component of $\mathcal{E}_\ell(f)$. In order to do so we introduce the following notations and definitions

Definition 1.6. Let $\ell \in \mathbb{R}$ and A, B be two subsets of \mathbb{R}^d , we define the event $\left\{ A \overset{\mathcal{E}_\ell(f)}{\longleftrightarrow} B \right\}$, also denoted by $\left\{ A \overset{f \geq -\ell}{\longleftrightarrow} B \right\}$, as the event that there exists a connected component of $\mathcal{E}_\ell(f)$ that intersects both A and B . When $A = \{x\}$ and $B = \{y\}$ are two singletons we simply use the notation $\left\{ x \overset{\mathcal{E}_\ell(f)}{\longleftrightarrow} y \right\}$ instead of $\left\{ \{x\} \overset{\mathcal{E}_\ell(f)}{\longleftrightarrow} \{y\} \right\}$.

We then define the chemical distance as a generalisation of graph distance for the case of Bernoulli percolation.

Definition 1.7. Given $x, y \in \mathbb{R}^d$ and a subset $\mathcal{E} \subset \mathbb{R}^d$ we define the set $\Gamma(\mathcal{E}, x, y)$ as

$$\Gamma(\mathcal{E}, x, y) := \{ \gamma : [0, 1] \rightarrow \mathcal{E} \mid \gamma \text{ is continuous and rectifiable, } \gamma(0) = x, \gamma(1) = y \} \quad (6)$$

The set $\Gamma(\mathcal{E}, x, y)$ can be empty, for instance if x or y are not in \mathcal{E} or are not in the same connected component of \mathcal{E} . The chemical distance between x and y is denoted by $d_{\text{chem}}^\mathcal{E}(x, y)$,

it takes values in $\mathbb{R}_+ \cup \{\infty\}$ and is defined as

$$d_{\text{chem}}^{\mathcal{E}}(x, y) := \inf\{\text{length}(\gamma) \mid \gamma \in \Gamma(\mathcal{E}_\ell(f), x, y)\}, \quad (7)$$

with the convention: $\inf \emptyset = \infty$. Here, $\text{length}(\gamma)$ denotes the Euclidean length of the curve γ (γ being rectifiable).

Remark. One may suggest the case where we have two points x and y in the same connected component of $\mathcal{E}_\ell(f)$ but with no rectifiable path between them hence having $x \xleftrightarrow{\mathcal{E}_\ell(f)} y$ but $d_{\text{chem}}^{\mathcal{E}_\ell(f)}(x, y) = \infty$. We will see that we do not need to concern ourselves with such a possibility since an analysis of our arguments shows that this event has zero probability.

Our main theorem is the following

Theorem 1.8. *Assume q satisfies Assumption 1.2 with some $\beta > d$ and that $\ell > -\ell_c(q)$. Then for any $\delta > 0$, there exists a constant $\kappa' > 0$ such that*

$$\mathbb{P}\left(0 \xleftrightarrow{\mathcal{E}_\ell(f)} x \text{ and } d_{\text{chem}}^{\mathcal{E}_\ell(f)}(0, x) > \|x\| \log^{\kappa(\delta)}(\|x\|)\right) = O\left(\|x\|^{-\kappa'}\right), \quad (8)$$

where $\kappa(\delta) = (1 + \delta)(d - 1) \left(\frac{1}{2} + \frac{1}{2\beta - d}\right)$.

We make a few comments about Theorem 1.8. The first observation is that this theorem is to be compared with a result of Peter Antal and Agoston Pisztora concerning the chemical distance in the context of Bernoulli percolation.

Theorem 1.9 ([1]). *In the context of Bernoulli percolation, if $d \geq 2$ and $p > p_c(d)$, we have constants $c, \rho > 0$ (depending on d and p) such that*

$$\mathbb{P}\left(0 \xleftrightarrow{\mathcal{G}_p(\omega)} x \text{ and } d_{\text{chem}}^{\mathcal{G}_p(\omega)}(0, x) > \rho \|x\|\right) = O\left(e^{-c\|x\|}\right).$$

Where $d_{\text{chem}}^{\mathcal{G}_p(\omega)}$ denotes the graph distance induced in $\mathcal{G}_p(\omega)$.

A major difference between the two theorems is that our is not optimal in the sense that it only holds for $\ell > -\ell_c$ instead of the expected full supercritical region $\ell > \ell_c$. This is due to the lack of estimates on the probability of some local uniqueness event (see Definition 2.3 for the precise definition of this event). This event is a non-monotonous event and we only manage to prove that it has high probability in the regime $\ell > -\ell_c$. Note that in [19], Theorem 1.8 was proven for the case of dimension $d = 2$ for the full supercritical region $\ell > 0$, this is due to the fact that in dimension 2 one can use planarity arguments and work only with increasing events. However, if this local uniqueness event indeed has high probability in the whole region $\ell > \ell_c$ we expect our proof of Theorem 1.8 to hold in this region (at the very least our argument would work in the region $\ell > 0$). Another difference between the two theorems is the logarithm factor that appears in Theorem 1.8. This is mainly due to the fact that contrary to Bernoulli percolation where the status of all sites are independent, our field f has values that are heavily correlated. This is problematic when we try to develop a renormalization argument, since we need to consider boxes of a logarithmic scale instead of boxes of fixed size. Finally, a last difference is the speed of decrease in the two theorems.

We obtain a polynomial decay in Theorem 1.8 instead of an exponential decay in Theorem 1.9. The reason for this is the fact that a continuous field f can have level sets that contort a lot. While for Bernoulli percolation there is a minimal scale, this is absolutely not the case for a continuous field. Hence we need to do a local control around the two points we want to connect 0 and x to ensure that the chemical distance "generated" near those points is not too high. This is done via a quantitative implicit function theorem. However the estimates obtained are not strong enough to guarantee an exponential decay. This is the only place in the proof where we lose a super-polynomial decay.

Compared to previous work in dimension 2 [19] where planarity was heavily used, the arguments used here are much closer to the ones in [1]. The strategy of the proof can be summarized as follows. In Section 2 we prove that in the regime $\ell > -\ell_c$ a local uniqueness event (see Definition 2.3) holds with high probability for f but also for a discretize and finite range version approximation of the field f . In Section 3, we prepare a renormalization argument. The content of this section is pretty classical but was formulated to adapt to our framework. The purpose of the renormalization is to build a path of adjacent boxes in which the field f (rather its finite range approximation) satisfies the local uniqueness event. This will allow us to pick up a path in $\mathcal{E}_\ell(f)$ of controllable length, starting not too far from 0 and arriving not too far from x . Section 4 is dedicated to proving that one can connect 0 and x to this path by paying a reasonable cost (in terms of chemical distance). This part of the paper is inspired by the arguments developed in [3]. We conclude Section 4 with the proof of Theorem 1.8.

Acknowledgements : I am very grateful to my PhD advisor Damien Gayet for introducing me to this problem as well as for many insightful discussions and for his remarks on a preliminary version of this paper.

2 On the local uniqueness event

In this section we present the local uniqueness event and show that it has high probability in the region $\ell > -\ell_c$.

We begin by properly defining the local uniqueness event as the intersection of two events, one is an "existence" event while the other one is a "uniqueness" event (the later being the hard and interesting part). First we introduce some notations. For $R > 0$ we define the box $B_R \subset \mathbb{R}^d$ as

$$B_R := [-R, R]^d. \tag{9}$$

For $i \in \{1, \dots, d\}$ we define the two i -faces $F_R^{i,+}$, $F_R^{i,-}$ of B_R as

$$\begin{aligned} F_R^{i,-} &:= [-R, R]^{i-1} \times \{-R\} \times [-R, R]^{d-i} \\ F_R^{i,+} &:= [-R, R]^{i-1} \times \{+R\} \times [-R, R]^{d-i} \end{aligned}$$

Definition 2.1. For $R > 1$ and $\ell \in \mathbb{R}$, we define the existence event $\text{Exist}(R, \ell)$ as the event that: for all $i \in \{1, \dots, d\}$, there exists a connected component of $B_R \cap \mathcal{E}_\ell(f)$ that intersects both $F_R^{i,-}$ and $F_R^{i,+}$ (see Figure 1 for an illustration of this event).




Figure 1: Illustration of the event $\text{Exist}(R, \ell)$ for $d = 3$. We have three different crossings (represented in different colors), each joining two opposite faces of the box.

We also define the uniqueness event.

Definition 2.2. For $R > 1$, $\ell \in \mathbb{R}$, we can consider the set $B_R \cap \mathcal{E}_\ell(f)$ as a union of disjoint connected components $B_R \cap \mathcal{E}_\ell(f) = \bigsqcup_{i \in I} \mathcal{C}_i$. Given some $\kappa \in]0, 1[$, the uniqueness event $\text{Unique}(R, \ell, \kappa)$ is the event that all \mathcal{C}_i of diameter at least κR are connected within $B_{R(1+\kappa)} \cap \mathcal{E}_\ell(f)$.

Remark. One could define the uniqueness event as the event that at most one of the \mathcal{C}_i in the above definition has diameter greater than κR , however, in order to avoid many technicalities later on, we prefer to work with this weaker definition.

The local uniqueness event in the box B_R is simply the intersection of the existence event and the uniqueness event in this box, more formally we make the following definition.

Definition 2.3. For $R > 1$, $\ell \in \mathbb{R}$ and $\kappa \in]0, 1[$ we define the local uniqueness event $\mathcal{A}(R, \ell, \kappa)$ as

$$\mathcal{A}(R, \ell, \kappa) := \text{Exist}(R, \ell) \cap \text{Unique}(R, \ell, \kappa). \quad (10)$$

We make a few comments about this local uniqueness event. These comments are not useful for the proof of our theorem but give some context about this type of event and its importance. First, note that having a high probability of the event $\mathcal{A}(R, \ell, \kappa)$ can be seen as a quantitative statement of Theorem 1.5. Indeed the proof of Theorem 1.5 uses non quantitative ideas coming from ergodicity and the Cameron-Martin formula and does not quite describe the geometry of the unbounded cluster. Second, recall that an event is said to be *increasing* if increasing values of the field f can only favor the realisation of the event (that is E is an increasing event if for all f, g we have $(f \leq g \text{ and } f \in E) \Rightarrow g \in E$). The existence event in Definition 2.1 is clearly an increasing event, as the set $\mathcal{E}_\ell(f)$ can only grow

when f increases. However, the uniqueness event in Definition 2.2 is neither increasing nor decreasing. Thus, our event \mathcal{A}_R itself is neither increasing nor decreasing. Hence, the whole machinery designed to deal with monotonous event fails to work with the event $\mathcal{A}(R, \ell, \kappa)$ so that it is not trivial to obtain good estimates about the probability of this event. If we consider Bernoulli percolation it is proved (see for instance [14]) that the event $\mathcal{A}(R, \ell, \kappa)$ has high probability in the whole supercritical phase (it is trivial that this event can not have high probability in the subcritical phase). It is conjectured (see for instance [6]) that the same is true in the context of continuous Gaussian field. In dimension $d \geq 3$ little is known about the probability of this event. In [6] a weaker result concerning some form of a cluster "present everywhere" was proved, but did not guarantee high probability for the event $\mathcal{A}(R, \ell, \kappa)$ even for $\ell > 0$. In [4] and [5] analogue of the event $\mathcal{A}(R, \ell, \kappa)$ were proven to have high probability in the whole the region $\ell > \ell_c$ but for different percolation models. In our paper we prove that the event $\mathcal{A}(R, \ell, \kappa)$ holds with high probability as soon as $\ell > -\ell_c$ for a large class of Gaussian fields. By Theorem 1.4, this result is weaker than the expected region $\ell > \ell_c$, but it still gives some non trivial region in which the result holds.

This section is dedicated to the proof of the following statement.

Proposition 2.4. *Assume that q satisfies assumption 1.2 for some $\beta > d$. Suppose $\ell > -\ell_c$, $\kappa \in]0, 1[$, then there exist constants $c, C > 0$, such that, for all $R > 1$,*

$$\mathbb{P}(f \in \mathcal{A}(R, \ell, \kappa)) \geq 1 - CR^d e^{-cR}. \quad (11)$$

The strategy to prove Proposition 2.4 is to argue that when $\ell > -\ell_c$ then the set $\{f \leq -\ell\}$ does not percolates (all components are bounded), and use a deterministic argument that states that it is difficult to observe two huge connected components of $\{f \geq -\ell\}$ without having also a huge connected component in $\{f \leq -\ell\}$.

2.1 Approximation of the field

We introduce approximations of the field f that have good properties (finite-range-dependence, piece-wise constant). This relies on the white noise decomposition of the field f .

In the following of the paper, we fix an arbitrary function $\chi : \mathbb{R}^d \rightarrow [0, 1]$ such that

- χ is \mathcal{C}^∞ .
- $\chi(x) = 1$ for all x such that $\|x\| \leq \frac{1}{4}$.
- $\chi(x) = 0$ for all x such that $\|x\| \geq \frac{1}{2}$.

With this function fixed we can make the following definition.

Definition 2.5. For $r > 1$ we define $\chi_r(x) := \chi(x/r)$ and we define the random field f_r as

$$f_r := (q\chi_r) * W. \quad (12)$$

Since $q\chi_r$ has compact support included in the Euclidean ball of radius $r/2$ we see that the field f_r is r -dependent, meaning that two events that depends on values of the field f_r at distance at least r are independent.

Definition 2.6. For $\varepsilon > 0$, and a function $g : \mathbb{R}^d \rightarrow \mathbb{R}$ is a function we define the ε -discretization of g as the function $g^\varepsilon : \mathbb{R}^d \rightarrow \mathbb{R}$ defined by $g^\varepsilon(x) := g(y)$ where y is the unique point in $\varepsilon\mathbb{Z}^d$ such that $x \in y + [-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}]^d$.

We will often write f_r^ε which should be interpreted as $(f_r)^\varepsilon$, that is the ε -discretization of the field f_r . It is good to have in mind that the fields f_r and f_r^ε are good local approximations of the field f as r is big and ε is small. This can be seen in the following classical proposition which is an application of the Borell-TIS inequality.

Proposition 2.7 (see for instance [17, Proposition 2.1]). *Assume that q satisfies Assumption 1.2 for some $\beta > d$. There exists a constant $c > 0$ such that*

$$\forall r > 1, \quad \forall s > \frac{1}{r^{\beta - \frac{d}{2}}}, \quad \mathbb{P} \left(\sup_{x \in B_1} |f_r(x) - f(x)| \geq s \right) \leq e^{-cs^2 r^{2\beta - d}}. \quad (13)$$

$$\forall r > 1, \quad \forall \varepsilon > 0, \quad \forall s > \varepsilon, \quad \mathbb{P} \left(\sup_{x \in B_1} |f_r^\varepsilon(x) - f_r(x)| \geq s \right) \leq e^{-cs^2/\varepsilon^2}. \quad (14)$$

Where we recall that B_1 denotes $B_1 = [-1, 1]^d$.

2.2 Deterministic argument for local uniqueness

We now present a deterministic argument that guarantees the fact that some set presents the local uniqueness property.

Definition 2.8. Let $R > 1, \kappa \in]0, 1[$ and recall that $B_R = [-R, R]^d$. Let $E \subset \mathbb{R}^d$ be a subset. We say that

- E has the (R, κ) *small clusters property*, if all the connected components of $E \cap B_{R(1+\kappa)}$ have diameter less than κR .
- E has the (R, κ) *unique cluster property*, if all the connected components of $E \cap B_R$ of diameter greater than κR are connected within $E \cap B_{R(1+\kappa)}$.
- E has the R *crossing property*, if for all $i \in \{1, d\}$ there exists a connected component of $E \cap B_R$ that intersects the two faces $F_R^{i,-}$ and $F_R^{i,+}$.

In this context the diameter refer to the usual Euclidean diameter.

We argue that if a set has some kind of small cluster property then its complement in the box has some form of a unique cluster property. The proof we provide next is purely geometrical and deterministic (although we will soon apply it to random sets).

Proposition 2.9. *Let $\kappa \in]0, 1[$, $E^- \subset \mathbb{R}^d$ and $E^+ \subset \mathbb{R}^d$ be two \mathcal{C}^1 -differentiable submanifolds of dimension d with boundary (these manifolds don't need to be connected and can have countably many connected components). Suppose also that their boundary is the same $E^0 = \partial E^- = \partial E^+$ and that we can write $\mathbb{R}^d = (E^- \setminus E^0) \sqcup E^0 \sqcup (E^+ \setminus E^0)$. Then, if for some $R > 1$, E^0 only intersects ∂B_R transversely and if E^- has the (R, κ) small clusters property, then E^+ has the (R, κ) unique cluster property and the R crossing property.*

Proof. We can write $E^- \cap B_{R(1+\kappa)} = \sqcup_{i=1}^n \mathcal{C}_i$ where the \mathcal{C}_i are connected. Since E^- has the (R, κ) small clusters property, we have by definition that for all i , $\text{diam}(\mathcal{C}_i) \leq \kappa R$. For each i we can consider the topological boundary of \mathcal{C}_i , that is we can write $\partial \mathcal{C}_i = \sqcup_{j=1}^{n_i} S_j^{(i)}$ where the $S_j^{(i)}$ are connected compact hypersurfaces. Because of our hypothesis, all these hypersurfaces are subsets of $E^0 \cup \partial B_{R(1+\kappa)}$. The Brouwer separation theorem allows to state that each of these compact hypersurfaces S separates the space \mathbb{R}^d into two connected parts, the interior $\text{Int}(S)$ that is bounded and the exterior of S that is unbounded $\text{Ext}(S)$. Without restriction of generality since the \mathcal{C}_i are connected, we can assume that for all $2 \leq j \leq n_i$ we have $S_j^{(i)} \subset \text{Int}(S_1^{(i)})$. This allows us to write $\mathcal{C}_i = \text{Int}(S_1^{(i)}) \setminus \bigcup_{j=2}^{n_i} \text{Int}(S_j^{(i)})$. Denote by \mathcal{C} the set of points $x \in E^+ \cap B_R$ such that the connected component of x in $E^+ \cap B_R$ is of diameter greater than κR . Our goal is to prove that any two points of \mathcal{C} are connected within $E^+ \cap B_{R(1+\delta)}$. For this purpose, consider two points x and y in \mathcal{C} . Since x is connected to a point z within $B_{R(1+\kappa)} \cap E^+$ at distance at least κR from x , we deduce that x does not belong to any $\text{Int}(S_1^{(i)})$ (otherwise this would contradict the fact that all \mathcal{C}_i have diameter less than κR). By symmetry the same is true for y . Now, consider γ the segment that joins the two points x and y (the segment stays in B_R by convexity of B_R , however it may leave E^+). Follow the segment from x to y until we either reach y or reach a point z_1 in some \mathcal{C}_i . In the later case the previous discussion allow us to state that the point z_1 belongs to some $S_1^{(i)}$. We can then look at \tilde{z}_1 the last intersection point between γ and $S_1^{(i)}$ and replace the segment $[z_1, \tilde{z}_1]$ of the path γ by some path in $S_1^{(i)}$ that connects the two points z_1 and \tilde{z}_1 (see Figure 2 for an illustration). Since z_1 is at distance at least κR from the boundary $\partial B_{R(1+\kappa)}$ and since \mathcal{C}_i is of diameter at most κR we see that \mathcal{C}_i does not intersects $\partial B_{R(1+\kappa)}$. Thus, the outer boundary $S_1^{(i)}$ is included in E^0 . Using this observation, we see that we can indeed find a path between z_1 and \tilde{z}_1 in E^0 (and also in $B_{R(1+\kappa)}$). By repeating this procedure until we reach y we have connected the two points x and y within $E^+ \cap B_{R(1+\kappa)}$ proving that E^+ has the (R, κ) unique cluster property. Note that \mathcal{C} can be shown to be non empty simply by choosing two points in $B_R \setminus \bigcup_{i=1}^n \text{Int}(S_1^{(i)})$ at distance κR from one each other and connecting them as done above. This concludes the proof that E^+ has the (R, κ) unique cluster property. In order to show that E^+ has the R crossing property, it is enough for each direction $j \in \{1, \dots, d\}$ to find two points w_1^j, w_2^j such that $w_1^j \in F_R^{j,-} \cap \mathcal{C}$ and $w_2^j \in F_R^{j,+} \cap \mathcal{C}$ as they will be connected in $B_R \cap E^+$ via \mathcal{C} . In order to algorithmically build such the point w_i^j , one way is to draw a segment starting from one point in \mathcal{C} towards the center of the face $F_R^{j,-}$. Should the segment encounter one of \mathcal{C}_i that does not intersects $F_R^{j,-}$ we deviate the path by following the boundary of \mathcal{C}_i until we can continue on our initial trajectory by staying in E^+ . If the segment reaches some \mathcal{C}_i that intersects $F_R^{j,-}$ we deviate our path by following the boundary of \mathcal{C}_i until we reach a point in $F_R^{j,-}$. Otherwise, our segment will simply reach a point in $F_R^{j,-}$. The path we built starts from a point in \mathcal{C} and only visits points in E^+ . This guarantees that all points visited by the path are in \mathcal{C} . Moreover the path eventually visits one point w_1^j in $F_R^{j,-}$. This concludes the proof of the proposition. \square

We conclude by a remark on the definition of the unique cluster property. One may find more natural to define the following uniqueness property:

Definition 2.10. Say that $E \subset \mathbb{R}^d$ has the (R, κ) *strong unique cluster property* if among




Figure 2: Illustration of the proof of Proposition 2.9.

the connected components of $E \cap B_R$ at most one has diameter greater than κR .

We also believe that an analogue of Proposition 2.9 with the (R, δ) unique cluster property replaced by the (R, δ) strong unique cluster property should hold. However a lot of technicalities are generated due to the possible intersections between the boundary of the box B_R with the boundary of the clusters of E^- . In order to avoid dealing these difficulties we prefer to work with the definition of the (R, κ) unique cluster property which is enough for our purpose.

2.3 Proof of Proposition 2.4

We want to prove Proposition 2.4 and in fact we will prove the more general result that also concerns the approximation f_r^ε of the field f .

Proposition 2.11. *Assume that q satisfies Assumption 1.2 for some $\beta > d$. Let $\ell > -\ell_c$, $\kappa \in]0, 1[$. Then, there exist constants $c, C > 0$, such that for all $R > 1$, $r > 1$, $\varepsilon < 1$,*

$$\mathbb{P}(f \in \mathcal{A}(R, \ell, \kappa)) \geq 1 - CR^d e^{-cR}. \quad (15)$$

$$\mathbb{P}(f_r^\varepsilon \in \mathcal{A}(R, \ell, \kappa)) \geq 1 - CR^d (e^{-cR} + e^{-cr^{2\beta-d}} + e^{-c\varepsilon^{-2}}). \quad (16)$$

The strategy is to show that $\mathcal{E}_\ell(f)$ (resp $\mathcal{E}_\ell(f_r^\varepsilon)$) have their complement that have the (R, κ) small cluster property (see Definition 2.8) to deduce via Proposition 2.9 that these sets have

the (R, κ) unique cluster property. First we recall an estimate about a crossing probability for Gaussian field.

Theorem 2.12 ([16], [12] for $d = 2$, [17] for $d \geq 3$). *Assume that q satisfies Assumption 1.2 for some $\beta > d$. For all $\ell < \ell_c$, there exist constants $c, C > 0$ such that for all $R \geq 1$,*

$$\mathbb{P}\left(B_1 \xleftrightarrow{f \geq -\ell} \partial B_R\right) \leq Ce^{-cR}. \quad (17)$$

We will use this theorem in the form of the equivalent corollary (that is just a direct consequence of the equality in law of f and $-f$).

Corollary 2.13. *Assume that q satisfies Assumption 1.2 for some $\beta > d$. For $\ell > -\ell_c$, there exist constant $c, C > 0$ such that for all $R \geq 1$,*

$$\mathbb{P}\left(B_1 \xleftrightarrow{f \leq -\ell} \partial B_R\right) \leq Ce^{-cR}. \quad (18)$$

Proof. This is simply using the fact that f and $-f$ have the same law (because f is a centered Gaussian field). Hence the two sets

$$\mathcal{E}_\ell(f) = \{x \in \mathbb{R}^d \mid f(x) \geq -\ell\}, \text{ and}$$

$$\mathcal{E}_\ell(-f) = \{x \in \mathbb{R}^d \mid -f(x) \geq -\ell\}$$

have the same law. The later set can be written as $\{x \in \mathbb{R}^d \mid f(x) \leq -(-\ell)\}$. Since taking $\ell < \ell_c$ is equivalent to taking $-\ell > -\ell_c$ the corollary holds. \square

We use this estimate to show that with high probability a box $B_R = [-R, R]$ does not contain a path of $\{f \leq -\ell\}$ with high diameter.

Proposition 2.14. *Assume that q satisfies Assumption 1.2 for some $\beta > d$. For all $\ell > -\ell_c$ and $\kappa \in]0, 1[$, there exist constants constants $c, C > 0$ such that, for all $R > 1, r > 1, \varepsilon < 1$,*

$$\begin{aligned} & \mathbb{P}\left(\{x \in \mathbb{R}^d \mid f_r^\varepsilon(x) \leq -\ell\} \text{ has the } (R, \kappa) \text{ small clusters property}\right) \\ & \geq 1 - CR^d(e^{-cR} + e^{-c\varepsilon^{-2}} + e^{-cr^{2\beta-d}}) \end{aligned} \quad (19)$$

Moreover, if we replace f_r^ε by f we get the following

$$\begin{aligned} & \mathbb{P}\left(\{x \in \mathbb{R}^d \mid f(x) \leq -\ell\} \text{ has the } (R, \kappa) \text{ small clusters property}\right) \\ & \geq 1 - CR^d e^{-cR} \end{aligned} \quad (20)$$

Proof. Fix $\ell' \in \mathbb{R}$ such that $\ell > \ell' > -\ell_c$. Cover the box $B_{R(1+\kappa)} = [-(1+\kappa)R, (1+\kappa)R]^d$ with $N \leq CR^d$ boxes of size 1 (where C is a constant depending only on the dimension d) and denote $b_1, b_2, \dots, b_N \in \mathbb{R}^d$ the centers of all those boxes. On the event that $\{f \leq -\ell'\}$ does not have the (R, κ) small clusters property, we can find x and y in $B_{R(1+\kappa)}$ such that x and y are distant of at least κR and such that x and y both are in a same connected component of $\{f \leq -\ell'\}$. Let i be one index such that $x \in b_i + [-1, 1]^d$. As soon as R is

big enough (compared to $\frac{1}{\kappa}$ and d) we have $y \notin b_i + \left[-\frac{\kappa R}{2\sqrt{d}}, \frac{\kappa R}{2\sqrt{d}}\right]^d$. Hence on the event that $\{f \leq -\ell'\}$ does not have the (R, κ) small clusters property, we have

$$\exists 1 \leq i \leq N, \quad b_i + [-1, 1]^d \xleftrightarrow{f \leq -\ell'} \partial \left(b_i + \left[-\frac{\kappa R}{2\sqrt{d}}, \frac{\kappa R}{2\sqrt{d}}\right]^d \right).$$

Applying stationarity (to assume $b_i = 0$) and applying Corollary 2.13, and doing an union bound on all possible i we see that

$$\begin{aligned} & \mathbb{P}(\{f \leq -\ell'\} \text{ does not have the } (R, \kappa) \text{ small clusters property}) \\ & \leq \sum_{i=1}^N \mathbb{P} \left(b_i + [-1, 1]^d \xleftrightarrow{f \leq -\ell'} \partial \left(b_i + \left[-\frac{\kappa R}{2\sqrt{d}}, \frac{\kappa R}{2\sqrt{d}}\right]^d \right) \right) \\ & \leq C' R^d e^{-c'R}, \end{aligned} \tag{21}$$

where c' and C' are constants that only depend on ℓ , κ and d . This yields the conclusion (20) for the field f . For the field f_r^ε , we know by local comparison (see Proposition 2.7) that for r big enough and ε small enough (compared with $|\ell - \ell'|$) we have a constant $c > 0$ depending on the law of f and on $|\ell - \ell'|$ such that :

$$\mathbb{P} \left(\sup_{B_1} |f - f_r^\varepsilon| > |\ell - \ell'| \right) \leq e^{-cr^{2\beta-d}} + e^{-c\varepsilon^{-2}}. \tag{22}$$

Hence, using again an union bound we see that

$$\mathbb{P} \left(\sup_{B_{R(1+\kappa)}} |f - f_r^\varepsilon| > |\ell - \ell'| \right) \leq CR^d (e^{-cr^{2\beta-d}} + e^{-c\varepsilon^{-2}}). \tag{23}$$

Now, if $\sup_{B_{R(1+\kappa)}} |f - f_r^\varepsilon| < |\ell - \ell'|$ then for all $x \in B_{R(1+\kappa)}$ we see that

$$f_r^\varepsilon(x) \leq -\ell \Rightarrow f(x) \leq -\ell'.$$

Thus, if $\sup_{B_{R(1+\kappa)}} |f - f_r^\varepsilon| < |\ell - \ell'|$ and if $\{x \in \mathbb{R}^d \mid f(x) \leq -\ell'\}$ has the (R, κ) small cluster property then so does $\{x \in \mathbb{R}^d \mid f_r^\varepsilon(x) \leq -\ell\}$. By an union bound we conclude :

$$\begin{aligned} & \mathbb{P}(\{x \in \mathbb{R}^d \mid f_r^\varepsilon(x) \leq -\ell\} \text{ does not have the } (R, \kappa) \text{ small clusters property}) \\ & \leq \mathbb{P}(\{x \in \mathbb{R}^d \mid f(x) \leq -\ell'\} \text{ does not have the } (R, \kappa) \text{ small clusters property}) \\ & \quad + \mathbb{P} \left(\sup_{B_{R(1+\kappa)}} |f - f_r^\varepsilon| > |\ell - \ell'| \right) \\ & \leq CR^d (e^{-cr^{2\beta-d}} + e^{-c\varepsilon^{-2}}) + CR^d e^{-cR}. \end{aligned}$$

Adjusting constants we get (19). □

Finally, we are ready to prove that when $\ell > -\ell_c$ the local uniqueness event $\mathcal{A}(R, \ell, \kappa)$ has high probability.

Proof of Proposition 2.11. We will do the proof for f , the proof is straightforward to adapt to f_r^ε . Denote

$$\begin{aligned} E^+ &:= \{x \in \mathbb{R} \mid f(x) \geq -\ell\}, \\ E^- &:= \{x \in \mathbb{R}^d \mid f(x) \leq -\ell\}, \\ E^0 &:= \{x \in \mathbb{R}^d \mid f(x) = -\ell\}. \end{aligned}$$

We also fix $R > 1$. Applying Lemma A.9 of [15], we see that almost surely, the sets E^+, E^-, E^0 satisfy the hypotheses of Proposition 2.9 for B_R . Hence, whenever E^- has the (R, κ) small clusters property, then E^+ will have the (R, κ) unique cluster property and R crossing property. It only remains to apply our estimates of Proposition 2.14, to see that both events $\{f \in \text{Unique}(R, \ell, \kappa)\}$ and $\{f \in \text{Exist}(R, \ell)\}$ have probability no less than $1 - CR^d e^{-cR}$. \square

Remark. The way we proved that the existence event $\text{Exist}(R, \ell)$ has high probability may seem unsatisfactory. We argue that it is possible to also derive this result very easily for all $\ell > 0$ simply by considering the percolation model induced in two-dimensional planes and use the results in [12] to show that in the plane rectangle are crossed with high probability. Also, since the event $f \in \text{Exist}(R, \ell)$ is an increasing connection event, the machinery developed by Severo in [17] together with Theorem 3.1 in [14] might allow to prove this result for all $\ell > \ell_c$, but a lot remains to be done in this direction.

3 Stochastic domination argument

In this section we introduce several tools for a renormalization argument. The idea is that we want to build a path of adjacent boxes into a renormalized lattice where in each box, the field f_r^ε will satisfy the local uniqueness event (see Definition 2.3). In order to do so we first recall a classical stochastic domination theorem allowing us to compare a measure on $\{0, 1\}^{\mathbb{Z}^d}$ with high marginal and finite range correlation with a classical Bernoulli process of high parameter. Typically one can have in mind the law of a configuration $\omega \in \{0, 1\}^{\mathbb{Z}^d}$ where $\omega_i = 1$ if and only if the field f_r^ε satisfy the local uniqueness event in the box $\frac{r}{10}i + B_r$. Then we introduce the notion of global structure as some path in the renormalized lattice that is of controllable length and come close to two selected points, and argue that such structures appear with high probability. This last part relies on argument developed in [7] and [1].

3.1 Comparison with a Bernoulli process of high parameter

We briefly recall the notion of stochastic domination for two measures on $\{0, 1\}^{\mathbb{Z}^d}$ and then state a classical stochastic domination result.

First we make the following definition.

Definition 3.1. Consider a countable set V , and two probability measures μ_1 and μ_2 on $(\{0, 1\}^V, \mathcal{B}(\{0, 1\})^{\otimes V})$. We say that μ_1 is stochastically dominated by μ_2 and we denote

$\mu_1 \prec \mu_2$ if there exist a coupling (X^1, X^2) of law μ where X^1 has law μ_1 and X^2 has law μ_2 and where

$$\mu(\forall i \in V, X_i^1 \leq X_i^2) = 1.$$

Definition 3.2. Let $d \geq 2$ and $p \in [0, 1]$, denote π_p the law on $\{0, 1\}^{\mathbb{Z}^d}$ given by independent and identically distributed Bernoulli random variable of parameter p . We immediately have

$$p \leq p' \Rightarrow \pi_p \prec \pi_{p'}.$$

We also recall the notion of measures that are finite-range dependent.

Definition 3.3. Let $M \in \mathbb{N}$. A probability measure μ on $\{0, 1\}^{\mathbb{Z}^d}$ is said to be M -dependent, where $M \in \mathbb{N}$ is a parameter, if for all subsets A, B of \mathbb{Z}^d such that $d_\infty(A, B) \geq M$ the two σ -algebras $\mathcal{F}_A := \sigma(\omega_i, i \in A)$ and $\mathcal{F}_B := \sigma(\omega_i, i \in B)$ are μ -independent.

A classical and general tool to compare a probability measure μ with some π_p is the following theorem. Basically the idea is that a measure μ with finite-range dependence stochastically dominates a Bernoulli process $\pi_{1-\alpha}$ with small $\alpha > 0$ as soon as the marginal of μ are close enough to 1.

Theorem 3.4 ([10], [1]). *For any $M \in \mathbb{N}$, $d \geq 1$ and $p \in [0, 1]$, there exist a quantity $\alpha(p, M, d) \in [0, 1]$ such that*

$$\alpha(p, M, d) \xrightarrow[p \rightarrow 1]{} 0, \tag{24}$$

and such that for any law μ on $\{0, 1\}^{\mathbb{Z}^d}$ satisfying:

- μ is M -dependent,
- $\inf_{i \in \mathbb{Z}^d} \mu(\omega_i = 1) \geq p$,

we have

$$\pi_{1-\alpha(p,d,M)} \prec \mu. \tag{25}$$

Remark. Note that the fact that $\alpha(p, M, d)$ can be as small as we wish as soon as p get close to 1 is the core statement of the claim, otherwise we could just take $\alpha(p, M, d) \equiv 1$ since we always have $\pi_0 \prec \mu$, but of course this is not what we are interested in. Furthermore, an analysis of the proofs in [1] says that we can always choose $\alpha(p, d, M) \leq 4(1-p)^{\frac{1}{(2M+1)^d}}$

3.2 Global structures

We introduce some terminology related to percolation on discrete graph and we also define the notion of *global structure* that will be used in the proof of Theorem 1.8.

Definition 3.5. Consider $A \subset \mathbb{Z}^d$ We say that two sites $x, y \in A$ are connected (resp. \star -connected) within A if one can find a finite sequence x_0, \dots, x_n of points of A such that

- $x_0 = x, x_n = y$,
- $\forall 0 \leq i \leq n-1, \|x_{i+1} - x_i\|_1 = 1$ (resp. $\|x_{i+1} - x_i\|_\infty = 1$).

We say that A is connected (resp. \star -connected) if any two points in A are connected (resp. \star -connected) within A .

Remark. Note that one should not confound the notion of being connected (as a subset of \mathbb{Z}^d) with the notion of being connected (as a subset of \mathbb{R}^d). In the rest of this section, the above definition takes precedence.

In the following, let $\omega \in \{0, 1\}^{\mathbb{Z}^d}$ be a configuration. We now state the definition of a *global structure* around two points.

Definition 3.6. Let $x \in \mathbb{Z}^d \setminus \{0\}$ and $C, \delta > 0$ be constants. A set $G \subset \mathbb{Z}^d$ is said to be a *global structure* around 0 and x with constants C, δ if the following is satisfied:

- $\forall y \in G, \omega_y = 1,$
- $|G| \leq C \|x\|_1,$
- G is connected,
- $d(0, G) \leq \log^{1+\delta}(\|x\|_1),$ and
- $d(x, G) \leq \log^{1+\delta}(\|x\|_1).$

We also define $\mathcal{G}(x, C, \delta)$ the event that there exists a global structure around 0 and x with constants C, δ .

In this section we prove the following result

Proposition 3.7. *For all $d \geq 2$, there exists $0 < \alpha_d < 1$ such that for all $\delta > 0$ there exist constants $c, C, C_0 > 0$ so that, if $p \in [1 - \alpha_d, 1]$, we have*

$$\forall x \in \mathbb{Z}^d \setminus \{0\}, \pi_p(\mathcal{G}(x, C_0, \delta)) > 1 - Ce^{-c \log^{1+\delta}(\|x\|_1)}. \quad (26)$$

Proof. We follow the argument in [14]. Consider a deterministic path of length $N = \|x\|_1$ in \mathbb{Z}^d joining 0 and x . We denote this path x_0, x_1, \dots, x_N with $x_0 = 0, x_N = x$ and $\|x_{i+1} - x_i\|_1 = 1$ for all $0 \leq i \leq N - 1$. For each $0 \leq i \leq N$ we can consider the biggest \star -connected component of $\{j \in \mathbb{Z}^d \mid \omega_j = 0\}$ that contains i (that is, the union of all sets that simultaneously are \star -connected, contain i and only contain vertices j such that $\omega_j = 0$). Denote this \star -connected component by \mathcal{C}_i (note that if $\omega_i = 1$ then $\mathcal{C}_i = \emptyset$). For a set $\mathcal{C} \subset \mathbb{Z}^d$ we define the *boundary* of \mathcal{C} as

$$\partial\mathcal{C} := \{i \in \mathbb{Z}^d \mid i \notin \mathcal{C} \text{ and } \exists j \in \mathcal{C}, \|i - j\|_\infty = 1\}.$$

We define $\overline{\mathcal{C}}_i$ as

$$\overline{\mathcal{C}}_i := \begin{cases} \{i\} & \text{if } \mathcal{C}_i = \emptyset \\ \mathcal{C}_i \cup \partial\mathcal{C}_i & \text{otherwise} \end{cases}.$$

We also define

$$A := \bigcup_{0 \leq i \leq N} \overline{\mathcal{C}}_i.$$

Using the arguments developed in [7] (see also section 4 in [1]) it is known that there exists $\alpha_d > 0$ such that for $1 > p > 1 - \alpha_d$ we have constants $C_0, c > 0$ such that :

$$\pi_p(|A| > C_0 N) \leq \exp(-cN). \quad (27)$$

We further claim that with high probability A is contained in a $\log^{1+\delta}(\|x\|_1)$ neighborhood of the path x_0, x_1, \dots, x_N . More precisely, a classical Peierls argument shows that if $p > 1 - \frac{1}{3^d}$ we have a constant $c > 0$ such that for all $n \geq 0$

$$\pi_p(|\mathcal{C}_0| > n) \leq \exp(-cn). \quad (28)$$

Denote \mathcal{N} the event that A is contained in a $\log^{1+\delta}(\|x\|_1)$ neighborhood of the path x_0, \dots, x_N , that is \mathcal{N} is the event

$$\{\forall a \in A, \exists 0 \leq i \leq N, \|a - x_i\|_1 \leq \log^{1+\delta}(\|x\|_1)\}.$$

Thus choosing n of order $\log^{1+\delta}(\|x\|_1)$ in (28) and doing an union bound for all clusters \mathcal{C}_i for $0 \leq i \leq N$, we obtain (adjusting constants)

$$\pi_p(\mathcal{N}) \leq C e^{-c \log^{1+\delta}(\|x\|_1)}. \quad (29)$$

Finally notice that if G denotes the external boundary of A , then G is a set of open sites that is connected and that surrounds the segment x_0, \dots, x_N , hence the conclusion. \square

4 Local control of the chemical distance and proof of the main theorem

In this section we propose an argument to control the probability of having an unusual high chemical distance in a box. We will then use these estimates together with the result of the previous section to give the proof of Theorem 1.8

4.1 Local control of the chemical distance around a point

We introduce some notations that were defined in [19]. The definitions were made for the two dimensional case, but they straightforwardly adapt to a higher dimension setting.

First, we recall Definition 1.7 of the chemical distance and we make the following definition.

Definition 4.1. Let \mathcal{E} and \mathcal{C} be two subsets of \mathbb{R}^d . The *chemical diameter* of \mathcal{C} within \mathcal{E} is defined as

$$\text{diam}_{\text{chem}}^{\mathcal{E}}(\mathcal{C}) := \sup\{d_{\text{chem}}^{\mathcal{E}}(x, y) \mid x, y \in \mathcal{C}\}. \quad (30)$$

Since we will be interested in a set that may have several connected components we make the following definition.

Definition 4.2. Let $s > 1$ and $\mathcal{E} \subset \mathbb{R}^d$ be a subset. We assume that $\mathcal{E} \cap B_s$ has finitely many connected components denoted by $\mathcal{C}_1, \dots, \mathcal{C}_n$. We define

$$S(s, \mathcal{E}) := \max_{1 \leq i \leq n} \text{diam}_{\text{chem}}^{\mathcal{E}}(\mathcal{C}_i). \quad (31)$$

In the case where $\mathcal{E} \cap B_s$ has infinitely many connected components, we arbitrarily define $S(s, \mathcal{E}) = \infty$.

Our claim is the following

Proposition 4.3. *Assume that q satisfies Assumption 1.2 for some $\beta > d$. For all $\ell \in \mathbb{R}$, for all $\eta > 0$ there exist constants $C, c, \kappa, \kappa' > 0$ such that*

$$\forall s > 10, \forall x > 10, \mathbb{P}(S(s, \mathcal{E}_\ell(f)) \geq x) \leq C \frac{s^\kappa}{x^{\kappa'}} + e^{-cs^\eta} \quad (32)$$

The estimate in Proposition 4.3 is pretty weak, we believe the same result to hold with a much faster decay but we could not find a proof of such result. However, this estimate will only be used two times in two small boxes of logarithmic scale. In order to prove Proposition 4.3, we will show that the box B_s can be filled by small boxes of size ε such that, with high probability, for each such small box B , the level set $\{f = -\ell\}$ either avoids B or cuts it into only two parts. This property can be interpreted in some sense as the level set $\{f = -\ell\}$ not being too much tortuous. Our argument is inspired by the work in [3]. We state a few results from [3] that we will use in the proof of Proposition 4.3. First, note that if $A \subset \mathbb{R}^d$ is a set and $f \in \mathcal{C}^2(\mathbb{R}^d, \mathbb{R})$ we use the Euclidean nature of \mathbb{R}^d to confound the first derivative df with the gradient ∇f and the second derivative d^2f with the Hessian matrix $H(f)$. We also define $\|df\|_{\mathcal{C}^1(A)} := \sup_{x \in A} \|\nabla f(x)\| + \sup_{x \in A} \|H(f)(x)\|$.

Proposition 4.4 (Quantitative implicit function theorem). *Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a \mathcal{C}^2 function. Take $U \subset \mathbb{R}^d$ an open set and $x \in U$, and positive constants $k > \lambda > 0$. If the following is satisfied :*

- $f(x) = 0$,
- $x + [-1, 1]^d \subset U$,
- $\|\nabla f(x)\| > \lambda$,
- $\|df\|_{\mathcal{C}^1(U)} \leq k$.

Then, if $\varepsilon = \frac{\lambda^2}{4k^2d^{3/2}}$ there exists a \mathcal{C}^2 function $\phi : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$, and a direction $i \in \{1, \dots, d\}$ such that

$$\forall y \in x + [-\varepsilon, \varepsilon]^d, f(y) = 0 \Leftrightarrow y_i = \phi(y_1, \dots, \widehat{y}_i, \dots, y_d).$$

Proof. Since we have $\|\nabla f(x)\| > \lambda$ we can find a $i \in \{1, \dots, d\}$ such that

$$\left| \frac{\partial f}{\partial x_i}(x) \right| \geq \frac{\lambda}{\sqrt{d}}.$$

Fix $\delta = \frac{\lambda}{2dk}$ (note that we have $\delta \leq 1$ by hypothesis), by the mean value theorem we have for all $y \in x + [-\delta, \delta]^d$

$$\left| \frac{\partial f}{\partial x_i}(x) - \frac{\partial f}{\partial x_i}(y) \right| \leq \|x - y\| \|df\|_{C^1(U)} \leq \sqrt{d}\delta k = \frac{\lambda}{2\sqrt{d}}.$$

Dividing by $\left| \frac{\partial f}{\partial x_i}(x) \right|$ we obtain

$$\forall y \in x + [-\delta, \delta]^d, \left| 1 - \left(\frac{\partial f}{\partial x_i}(x) \right)^{-1} \frac{\partial f}{\partial x_i}(y) \right| \leq \frac{1}{2}.$$

We can apply Theorem A.2 in [3] with a $M \leq \frac{\sqrt{d}}{\lambda}$ and $C \leq k$ which directly implies our claim for $\varepsilon = \frac{\delta}{2MC} \geq \frac{\lambda^2}{4d^{3/2}k^2}$. \square

We now state some well-known estimates about the behaviour of the derivative of the random field f in a box.

Proposition 4.5. *Assume that q satisfies Assumptions 1.2 for some $\beta > d$ and let $\ell \in \mathbb{R}$. Then, there exist constants $c, C, \kappa_1, \kappa_2 > 0$ such that for all $u \geq 0$ and $s \geq 10$,*

$$\mathbb{P} \left(\|f\|_{C^2(B_s)} \geq C \log^{\frac{1}{2}}(s) + u \right) \leq \exp \left(-c \frac{u^2}{2} \right). \quad (33)$$

$$\mathbb{P}(\exists x \in B_s \mid f(x) = -\ell \text{ and } \|\nabla f(x)\| < \lambda) \leq C \lambda^{\frac{1}{\kappa_1}} s^{\kappa_2}. \quad (34)$$

Proof. Concerning (33), this is an application of the Borell-TIS inequality, see also Lemma 5.2 in [3] for a similar estimate. For the proof of (34), this is a restatement of Lemma 5.3 in [3]. \square

Since the proof of Proposition 4.3 requires covering various sets with a controllable number of small boxes of size ε we introduce the following definition.

Definition 4.6. Let $A \subset \mathbb{R}^d$ and $\varepsilon > 0$. A set $\mathcal{R} \subset A$ is called a ε -covering of A if

$$A \subset \bigcup_{x \in \mathcal{R}} B(x, \varepsilon),$$

where $B(x, \varepsilon)$ denotes the open box $B(x, \varepsilon) := x +]-\varepsilon, \varepsilon[^d$. We also define

$$N(A, \varepsilon) := \inf\{|\mathcal{R}| \mid \mathcal{R} \text{ is a } \varepsilon\text{-covering of } A\}. \quad (35)$$

Note that $N(A, \varepsilon)$ can eventually be infinite (when A is unbounded).

Lemma 4.7. *If $A \subset B \subset \mathbb{R}^d$ and $\varepsilon > 0$ then*

$$N(A, 2\varepsilon) \leq N(B, \varepsilon).$$

Proof. The statement is trivial if $N(B, \varepsilon) = \infty$, otherwise choose $\mathcal{R} = \{x_1, \dots, x_n\} \subset B$ a ε -covering of B . We have $A \subset B \subset \bigcup_{i=1}^n B(x_i, \varepsilon)$. Find $0 \leq m \leq n$ and reorder the x_i 's so that $A \subset \bigcup_{i=1}^m B(x_i, \varepsilon)$ and so that A intersects all $B(x_i, \varepsilon)$ for $1 \leq i \leq m$. For $1 \leq i \leq m$ we choose $y_i \in A \cap B(x_i, \varepsilon)$ and since $B(x_i, \varepsilon) \subset B(y_i, 2\varepsilon)$ we immediately that y_1, \dots, y_m is a 2ε -covering of A which yields the conclusion. \square

Corollary 4.8. *There exists a constant C depending only on d such that for all $A \subset B_R$ and $\varepsilon > 0$ we have*

$$N(A, \varepsilon) \leq C \left(\frac{R}{\varepsilon} \right)^d.$$

Proof. We apply Lemma 4.7 to $B = B_R$ and we note that B_R can obviously be covered with $O\left(\left(\frac{R}{\varepsilon}\right)^d\right)$ balls of the form $B(x, \frac{\varepsilon}{2})$ \square

Now we are ready to prove Proposition 4.3.

Proof of Proposition 4.3. Fix $s \geq 10$, $x \geq 10$ and $\eta > 0$. Let $\lambda > 0$ to be fixed later. Denote

$$k := s^\eta.$$

Consider the two events:

$$\begin{aligned} \mathcal{A} &:= \left\{ \|df\|_{C^1(B_s + [-1, 1]^d)} \leq k \right\}, \\ \mathcal{B} &:= \left\{ \forall y \in B_s, f(y) = -\ell \Rightarrow \|\nabla f(x)\| > \lambda \right\}. \end{aligned}$$

Applying Proposition 4.5, we see that

$$\mathbb{P}(\mathcal{A}) \geq 1 - e^{-cs^{2\eta}} \quad \text{and} \quad \mathbb{P}(\mathcal{B}) \geq 1 - C\lambda^{\frac{1}{\kappa_1}} s^{\kappa_2}. \quad (36)$$

Set $\varepsilon = \frac{\lambda^2}{4d^{3/2}k^2}$ as in Proposition 4.4. Applying Corollary 4.8 to the set $B_s \cap \{f = -\ell\}$ with $\frac{\varepsilon}{2}$, we can find a constant C_d (depending only on d) and a collection of $N \leq C_d \left(\frac{s}{\varepsilon}\right)^d$ points denoted $(x_i)_{1 \leq i \leq N}$ such that for all i , $f(x_i) = -\ell$ and such that

$$\forall y \in B_s, f(y) = -\ell \Rightarrow \exists 1 \leq i \leq N, y \in B(x_i, \frac{\varepsilon}{2}).$$

Moreover, by the triangular inequality, if $y \in B_s$ is such that $d(y, \{f = -\ell\}) \leq \frac{\varepsilon}{2}$ then there exists $1 \leq i \leq N$ such that $y \in B(x_i, \varepsilon)$.

Consider the set $F := \{y \in B_s \mid \exists z \in B_s, f(z) = -\ell \text{ and } d(y, z) \leq \frac{\varepsilon}{2}\}$ and its complement $G := B_s \setminus F$. The previous observation shows that one has :

$$F \subset \bigcup_{i=1}^N B(x_i, \varepsilon).$$

Applying Corollary 4.8, we can also cover G by less than N balls of radius $\frac{\varepsilon}{2\sqrt{d}}$ and of centers y_i chosen in G .

$$G \subset \bigcup_{i=1}^N B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right).$$

Note that by construction, for each ball $B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right)$ in the covering of G this ball does not intersect $\{f = -\ell\} \cap B_s$.

Under the event $\mathcal{A} \cap \mathcal{B}$ take any two points $z_1, z_2 \in B_s \cap \mathcal{E}_\ell(f)$. We claim the following.

Claim 1. *There exists a deterministic constant $C'_d > 0$ depending only on d , such that under one of the following hypotheses.*

- *There exists $i \in \{1, \dots, N\}$ such that z_1 and z_2 are in $\overline{B(x_i, \varepsilon)}$,*
- *There exists $i \in \{1, \dots, N\}$ such that z_1 and z_2 are in $\overline{B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right)}$ and z_1 is in $B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right)$,*

then there exists a rectifiable path in $\mathcal{E}_\ell(f)$ of length less than $C'_d\varepsilon$ that connects z_1 and z_2 .

Proof of Claim 1. Assume first that z_1 and z_2 are in some $\overline{B(x_i, \varepsilon)} = x_i + [-\varepsilon, \varepsilon]^d$. We apply Proposition 4.4 at x_i and we see that the box $\overline{B(x_i, \varepsilon)}$ is separated in two by the hypersurface $\{f = -\ell\}$. Since by hypothesis we have $z_1, z_2 \in \mathcal{E}_\ell(f)$ we deduce that the two points z_1 and z_2 are on the same side of this hypersurface. These two points can connect within $\mathcal{E}_\ell(f)$ first by traveling towards one face of the box and then travelling along this face of the box to connect. This connection was made of length less than $4\varepsilon + 2\sqrt{d}\varepsilon$.

Now assume that z_1 and z_2 are in some $\overline{B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right)}$. By a previous observation, the set $B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right) \cap B_s$ does not intersect $\{f = -\ell\}$, thus by continuity of f this set $B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right) \cap B_s$ is either included in $\{f > -\ell\}$ or in $\{f < -\ell\}$. By the assumption on z_1 the later case can be excluded. By taking the closure we see that $\overline{B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right)} \subset \mathcal{E}_\ell(f)$. Now by convexity, the result follows as we can simply join the two points z_1 and z_2 by a straight segment. \square

We can now conclude the proof of Proposition 4.3. Consider two points y and z that are connected in $B_s \cap \mathcal{E}_\ell(f)$, and denote $\gamma : [0, 1] \rightarrow B_s \cap \mathcal{E}_\ell(f)$ a continuous function such that $\gamma(0) = y$ and $\gamma(1) = z$. We want to find a path between y and z of controlled length. To do so consider the following construction. We start from $z_0 := y$, if $z_0 \in F$ we know that we have some i such that $z_0 \in B(x_i, \varepsilon)$ and we define $\mathcal{B}_0 := B(x_i, \varepsilon)$ (for such an arbitrarily chosen i). Otherwise there exists some i such that $z_0 \in B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right)$ and we define $\mathcal{B}_0 := B\left(y_i, \frac{\varepsilon}{2\sqrt{d}}\right)$ for such a i (again arbitrarily chosen). We then define $t_0 := \sup\{t > 0 \mid \gamma(t) \in \mathcal{B}_0\}$ and $z_1 = \gamma(t_0)$ and we repeat this procedure until we reach step M where $t_{M-1} = 1$. We have obtained a sequence $(z_k)_{0 \leq k \leq M}$ (with $z_0 = y$ and $z_M = z$) and boxes $(\mathcal{B}_k)_{0 \leq k \leq M-1}$. Moreover, by construction, all boxes obtained in this construction are different (this also guarantees the fact that the procedure ends). Since there are at most $2N$ boxes to choose from, we have $M \leq 2N$. Furthermore, for all k we have $z_k \in \mathcal{B}_k$ and $z_{k+1} \in \overline{\mathcal{B}_k}$. Additionally, we have $z_k, z_{k+1} \in B_s \cap \mathcal{E}_\ell(f)$ (this is due to the fact that the z_k are part of the path γ). We can thus apply Claim 1 to find a path in $\mathcal{E}_\ell(f)$ joining z_k and z_{k+1} of length less than $C'_d\varepsilon$. By gluing together these paths, we obtain a rectifiable path between y and z of length less than $2NC'_d\varepsilon$. Since the points y and z were arbitrary, on the event $\mathcal{A} \cap \mathcal{B}$ we have

$S(s, \mathcal{E}_\ell(f)) \leq 2NC'_d \varepsilon \leq 2C_d C'_d \frac{s^d}{\varepsilon^{d-1}}$. By our choice of ε we see that we have a constant C''_d depending on d such that

$$S(s, \mathcal{E}_\ell(f)) \leq C''_d s^d (s^\eta)^{2(d-1)} \lambda^{-2(d-1)}.$$

Finally, we choose

$$\lambda = \left(\frac{C''_d s^{d+2\eta(d-1)}}{x} \right)^{\frac{1}{2(d-1)}},$$

which yields that under the event $\mathcal{A} \cap \mathcal{B}$ we have $S(s, \mathcal{E}_\ell(f)) \leq x$. Using estimates (36) and adjusting constants, we get the conclusion of Proposition 4.3. \square

4.2 Proof of Theorem 1.8

We now give the proof of Theorem 1.8.

Proof of Theorem 1.8. In this proof c, C denote positive constants that may change from line to line. By isotropy, it is enough to look at the chemical distance between $0 \in \mathbb{R}^d$ and $(x, 0, \dots, 0) \in \mathbb{R}^d$ where $x > 1$ is a real number (that can be thought of as big). Thus in the following, we assume that x is fixed (and big enough for future purpose). Fix also some parameter $\delta > 0$, that is free for now. We introduce the following quantities

$$r = R = \log^{\frac{1+\delta}{2\beta-d}}(x), \quad \varepsilon = \frac{1}{\log^{\frac{1+\delta}{2}}(x)}.$$

Also, consider, ℓ and ℓ' such that :

$$\ell > \ell' > -\ell_c.$$

We will work at the level ℓ' with f_r^ε to then recover information about the field f at level ℓ .

We introduce our renormalization procedure in order to use Theorem 3.4. Define the event \mathcal{A}_0 as

$$\mathcal{A}_0 := \left\{ f_r^\varepsilon \in \mathcal{A} \left(R, \ell', \frac{1}{100} \right) \right\},$$

where we recall Definition 2.3 of the event $\mathcal{A}(R, \ell, \kappa)$. Note that we decide to work with $\kappa = \frac{1}{100}$ simply to fix a certain value for κ but the exact value is not crucial for the rest of the argument. For $i \in \mathbb{Z}^d$ we define the event \mathcal{A}_i as the translation of the event \mathcal{A}_0 of vector $\frac{R}{10}i$. We also denote by \mathbf{B}_i the box $\mathbf{B}_i = \frac{R}{10}i + B_R$ and by $\tilde{\mathbf{B}}_i$ the box $\tilde{\mathbf{B}}_i = \frac{R}{10}i + B_{R(1+\frac{1}{100})}$. With these notations, note that the event \mathcal{A}_i can simply be restated as the occurrence of the existence event and the uniqueness event in the box \mathbf{B}_i .

Consider the configuration $(\omega_i)_{i \in \mathbb{Z}^d} \in \{0, 1\}^{\mathbb{Z}^d}$ such that $\forall i \in \mathbb{Z}^d, \omega_i = \mathbf{1}_{\mathcal{A}_i}$, denote by μ the probability measure on $\{0, 1\}^{\mathbb{Z}^d}$ whose law is the law of $\omega \in \{0, 1\}^{\mathbb{Z}^d}$. By construction, using the fact that f_r^ε is r -dependent, one see that μ is 100-dependent (where 100 is clearly not optimal but will not intervene later on). We can apply Theorem 3.4 and find $\alpha(x, \delta) \in [0, 1]$ such that

$$\pi_{1-\alpha(x, \delta)} \prec \mu.$$

Furthermore, applying Proposition 2.11 and stationarity, one see that

$$\begin{aligned}
& \inf_{i \in \mathbb{Z}^d} \mu(\omega_i = 1) \\
&= \mathbb{P} \left(f_r^\varepsilon \in \mathcal{A} \left(R, \ell', \frac{1}{100} \right) \right) \\
&\geq 1 - CR^d \left(e^{-cR} + e^{-cr^{2\beta-d}} + e^{-c\varepsilon^{-2}} \right) \\
&\geq 1 - C \left(\log^{\frac{1+\delta}{2\beta-d}}(x) \right)^d \left(e^{-c \log^{\frac{1+\delta}{2\beta-d}}(x)} + e^{-c \log^{1+\delta}(x)} + e^{-c \log^{1+\delta}(x)} \right)
\end{aligned}$$

This yields

$$\inf_{i \in \mathbb{Z}^d} \mu(\omega_i = 1) \xrightarrow{x \rightarrow \infty} 1.$$

By the conclusion of Theorem 3.4, we see that $\alpha(x)$ can be chosen so that

$$\alpha(x, \delta) \xrightarrow{x \rightarrow \infty} 0.$$

In the following, we choose to only work with $x > x_0$ so that we have $1 - \alpha(x, \delta) > 1 - \alpha_d$ where α_d is given by Proposition 3.7. Let $X \in \mathbb{N}^*$ such that $x \in \frac{R}{10}X + B_R$. Recall Definition 3.6 of the event $\mathcal{G}(X, C, \delta)$. By Proposition 3.7, we can choose $C_0 > 0$ such that

$$\pi_{1-\alpha_d}(\mathcal{G}(X, C_0, \delta)) \geq 1 - Ce^{-c \log^{1+\delta}(X)}, \quad (37)$$

for some constants $c, C > 0$. Since, μ stochastically dominates $\pi_{1-\alpha_d}$ and since the event $\mathcal{G}(X, C_0, \delta)$ is an increasing event we get

$$\mu(\mathcal{G}(X, C_0, \delta)) \geq 1 - Ce^{-c \log^{1+\delta}(X)} \quad (38)$$

Under the realization of this event, denote $\mathcal{C} \subset \mathbb{Z}^d$ a global structure around 0 and X with constants C_0, δ . Since for all $i \in \mathcal{C}$ the event \mathcal{A}_i occurs, and using the overlap of the boxes, we can glue together the "big clusters" of $\{f_r^\varepsilon \geq -\ell'\} \cap \mathbf{B}_i$ of all boxes \mathbf{B}_i for $i \in \mathcal{C}$ within the union of the $\tilde{\mathbf{B}}_i$ for $i \in \mathcal{C}$ (this is due to Definition 2.2 of the uniqueness event with $\kappa = \frac{1}{100}$). In particular this implies that there exists a path γ in $\{f_r^\varepsilon \geq -\ell'\}$ that stays in the $\tilde{\mathbf{B}}_i$ for $i \in \mathcal{C}$ and such that

$$\max(d(0, \gamma), d(x, \gamma)) \leq R \log^{1+\delta}(X), \quad (39)$$

where d is the Euclidean distance in \mathbb{R}^d . Since we have $X \leq x$ and $R = \log^{\frac{1+\delta}{2\beta-d}}(x)$ we obtain that

$$\max(d(0, \gamma), d(x, \gamma)) \leq \log^{1+\delta + \frac{1+\delta}{2\beta-d}}(x). \quad (40)$$

Note that since such a path is confined in at most C_0X boxes $\tilde{\mathbf{B}}_i$ and since f_r^ε is piece-wise constant on each small cube of size ε , this path γ can be chosen of Euclidean length less than $C'X\varepsilon^{\frac{R^d}{\varepsilon^d}}$ where C' is a constant that may depend on d and C_0 . This is a very rough upper bound that we obtain by dividing each box $\tilde{\mathbf{B}}_i$ into $C' \left(\frac{R}{\varepsilon}\right)^d$ small boxes of size ε on which the field f_r^ε is constant, and then choosing γ so that if γ visits one of this small boxes it will never visit it again. With the additional observation that $X \leq \frac{x}{R}$, the above can be

restated as follows: under the event $\mathcal{G}(X, C_0, \delta)$ there exists a continuous rectifiable path γ taking values in $\{f_r^\varepsilon \geq -\ell'\}$ such that (40) is verified and such that

$$\text{length}(\gamma) \leq C' x \log^{(1+\delta)(d-1)(\frac{1}{2} + \frac{1}{2\beta-d})}(x) \quad (41)$$

Moreover, the above construction shows that γ can be chosen included in a box of the form $B_{C''x}$ where $C'' > 0$ is a constant depending on C_0, d .

Applying Proposition 2.7 we see that for x big enough we have a constant $c > 0$ such that

$$\mathbb{P}\left(\|f - f_r^\varepsilon\|_{B_{C''x}} > \ell - \ell'\right) \leq e^{-c \log^{1+\delta}(x)}. \quad (42)$$

Under the event $\mathcal{H} := \left\{\|f - f_r^\varepsilon\|_{B_{C''x}} \leq \ell - \ell'\right\}$ we observe that the path γ is also included in $\mathcal{E}_\ell(f)$ (since on this event we have $f_r^\varepsilon(z) \geq -\ell'$ implies $f(z) \geq f'(z) - (f'(z) - f(z)) \geq -\ell$ for all $z \in B_{C''x}$).

On the event that 0 and x are connected within $\mathcal{E}_\ell(f)$ we argue that with high probability we can connect 0 and x to the path γ in a neighborhood of 0 and x . In fact, consider the event \mathcal{U}_1 as

$$\mathcal{U}_1 := \left\{f \in \text{Unique}\left(2 \log^{1+\delta+\frac{1+\delta}{2\beta-d}}(x), \ell, \frac{1}{100}\right)\right\}. \quad (43)$$

and define the event \mathcal{U}_2 as the translated of the event \mathcal{U}_1 by x . (recall the definition of the uniqueness event in Definition 2.2). Applying Proposition 2.11 we see that for x big enough we have

$$\mathbb{P}(\mathcal{U}_1) = \mathbb{P}(\mathcal{U}_2) \geq 1 - C e^{-c \log^{1+\delta}(x)}. \quad (44)$$

Under the event $\{0 \xleftrightarrow{\mathcal{E}_\ell(f)} x\}$ and the event \mathcal{U}_1 we know that 0 connects to the path γ within the box centered at 0 of radius $2\left(1 + \frac{1}{100}\right) \log^{1+\delta+\frac{1+\delta}{2\beta-d}}(x)$ (a similar connection happens around x via the event \mathcal{U}_2). Now it only remains to argue that this connection can be done with a path of reasonable Euclidean length. For this, we use the estimate of Proposition 4.3. More precisely define the event \mathcal{S}_1 as

$$\mathcal{S}_1 = \left\{S\left(2\left(1 + \frac{1}{100}\right) \log^{1+\delta+\frac{1+\delta}{2\beta-d}}(x), \mathcal{E}_\ell(f)\right) \leq x \log^{(1+\delta)(d-1)(\frac{1}{2} + \frac{1}{2\beta-d})}(x)\right\}, \quad (45)$$

and define the event \mathcal{S}_2 as the translated of the event \mathcal{S}_1 by x (recall Definition 4.2 of $S(s, E)$). Applying Proposition 4.3 and using stationarity we see that we have a constant $\kappa' > 0$ such that for x big enough

$$\mathbb{P}(\mathcal{S}_1) = \mathbb{P}(\mathcal{S}_2) \geq 1 - x^{-\kappa'}. \quad (46)$$

Under the event \mathcal{S}_1 it is known that the path between 0 and γ can be chosen of Euclidean length less than $x \log^{(1+\delta)(d-1)(\frac{1}{2} + \frac{1}{2\beta-d})}(x)$ (similarly for the connection between x and γ using \mathcal{S}_2). Hence, gluing these paths together we finally obtain a path between 0 and x in $\mathcal{E}_\ell(f)$ of length less than $3x \log^{(1+\delta)(d-1)(\frac{1}{2} + \frac{1}{2\beta-d})}(x)$. Doing an union bound and adjusting constants we finally obtain that for x big enough.

$$\mathbb{P}\left(0 \xleftrightarrow{\mathcal{E}_\ell(f)} x \text{ and } d_{\text{chem}}^{\mathcal{E}_\ell(f)}(0, x) > 3x \log^{(1+\delta)(d-1)(\frac{1}{2} + \frac{1}{2\beta-d})}(x)\right) \leq x^{-\kappa'}.$$

That is precisely the conclusion of Theorem 1.8.

□

References

- [1] Peter Antal and Agoston Pisztora, *On the chemical distance for supercritical bernoulli percolation*, The Annals of Probability **24** (1996), no. 2, 1036–1048.
- [2] Jean-Marc Azaïs and Mario Wschebor, *Level sets and extrema of random processes and fields*, Found. Comput. Math. **10** (2010), no. 4, 481–484.
- [3] Vincent Beffara and Damien Gayet, *Percolation of random nodal lines*, Publications mathématiques de l’IHÉS **126** (2017), no. 1, 131–176.
- [4] Hugo Duminil-Copin, Subhajit Goswami, Pierre-François Rodriguez, and Franco Severo, *Equality of critical parameters for percolation of gaussian free field level sets*, Duke Mathematical Journal (2023).
- [5] Hugo Duminil-Copin, Subhajit Goswami, Pierre-François Rodriguez, Franco Severo, and Augusto Teixeira, *A characterization of strong percolation via disconnection*, 2023.
- [6] Hugo Duminil-Copin, Alejandro Rivera, Pierre-François Rodriguez, and Hugo Vanneuille, *Existence of an unbounded nodal hypersurface for smooth gaussian fields in dimension $d > 3$* , The Annals of Probability **51** (2023), no. 1, 228–276.
- [7] Luiz Fontes and Charles M. Newman, *First Passage Percolation for Random Colorings of \mathbb{Z}^d* , The Annals of Applied Probability **3** (1993), no. 3, 746 – 762.
- [8] Svante Janson, *Gaussian hilbert spaces*, no. 129, Cambridge university press, 1997.
- [9] Harry Kesten, *The critical probability of bond percolation on the square lattice equals $1/2$* , Communications in mathematical physics **74** (1980), no. 1, 41–59.
- [10] T. M. Liggett, R. H. Schonmann, and A. M. Stacey, *Domination by product measures*, The Annals of Probability **25** (1997), no. 1, 71 – 95.
- [11] Stephen Muirhead, Alejandro Rivera, Hugo Vanneuille, and Laurin Köhler-Schindler, *The phase transition for planar Gaussian percolation models without FKG*, The Annals of Probability **51** (2023), no. 5, 1785 – 1829.
- [12] Stephen Muirhead and Hugo Vanneuille, *The sharp phase transition for level set percolation of smooth planar Gaussian fields*, Annales de l’Institut Henri Poincaré, Probabilités et Statistiques **56** (2020), no. 2, 1358 – 1390.
- [13] Fedor Nazarov and Mikhail Sodin, *Asymptotic laws for the spatial distribution and the number of connected components of zero sets of gaussian random functions*, Zurnal matematicheskoy fiziki, analiza, geometrii **12** (2015).
- [14] Agoston Pisztora, *Surface order large deviations for ising, potts and percolation models*, Probability Theory and Related Fields **104** (1996), no. 4, 427–466.

- [15] Alejandro Rivera and Hugo Vanneuille, *Quasi-independence for nodal lines*, Annales de l'Institut Henri Poincaré, Probabilités et Statistiques **55** (2019), no. 3, 1679 – 1711.
- [16] Alejandro Rivera and Hugo Vanneuille, *The critical threshold for bargmann–fock percolation*, Annales Henri Lebesgue **3** (2020), 169–215.
- [17] Franco Severo, *Sharp phase transition for Gaussian percolation in all dimensions*, Annales Henri Lebesgue **5** (2022), 987–1008 (en).
- [18] ———, *Uniqueness of Unbounded Component for Level Sets of Smooth Gaussian Fields*, International Mathematics Research Notices **2024** (2023), no. 11, 9002–9009.
- [19] David Verotte, *Chemical distance in the supercritical phase of planar gaussian fields*, arXiv preprint arXiv:2312.14205 (2023).