



**HAL**  
open science

## Editorial: Social Interaction in Animals: Linking Experimental Approach and Social Network Analysis

Cédric Sueur, Frédéric Mery

► **To cite this version:**

Cédric Sueur, Frédéric Mery. Editorial: Social Interaction in Animals: Linking Experimental Approach and Social Network Analysis. *Frontiers in Psychology*, 2017, 8, pp.35. 10.3389/fpsyg.2017.00035 . hal-04681710

**HAL Id: hal-04681710**

**<https://hal.science/hal-04681710v1>**

Submitted on 29 Aug 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Editorial: Social Interaction in Animals: Linking Experimental Approach and Social Network Analysis

Cédric Sueur<sup>1\*</sup> and Frédéric Mery<sup>2</sup>

<sup>1</sup> Centre National de la Recherche Scientifique (CNRS), Institut Pluridisciplinaire Hubert Curien UMR 7178, Université de Strasbourg, Strasbourg, France, <sup>2</sup> Evolution, Génomes, Comportement and Ecologie, Centre National de la Recherche Scientifique (CNRS), Institut de Recherche pour le Développement (IRD), Université Paris-Sud, Université Paris-Saclay, Gif-sur-Yvette, France

**Keywords:** fitness, information, social transmission, health, disease, social structure, RSiena

## Editorial on the Research Topic

### Social Interaction in Animals: Linking Experimental Approach and Social Network Analysis

Understanding the link between individual behavior and population organization and functioning has long been central to ecology and evolutionary biology (Krause et al., 2009; Sueur et al., 2011; Kurvers et al., 2014). Behavior is a response to intrinsic and extrinsic factors including individual state, ecological factors, or social interactions. Within a group, each individual can be seen as part of a network of social interactions varying in strength, type, and dynamic. The structure of this network can deeply impact the ecology and evolution of individuals, populations, and species.

Three studies in this present issue tried to understand how group members are socially structured in non-human primates. Borgeaud et al. used a stochastic actor-oriented model (RSiena Package, Snijders, 2001) to test the dynamics of relationships of three groups of wild vervet monkeys. They found that triadic closure was significant in all three groups while degree popularity was significant in only two groups. Moreover, the dynamics of relationships according to the attributes of sex, matriline, and age differed significantly among groups.

In another way, Sosa showed that in Barbary macaques, females are more central, more active, and have a denser ego network in the social network than males; thus, they contribute in a greater way to the cohesive structure of the network. High-ranking individuals are likely to receive fewer agonistic behaviors than low-ranking individuals, and high-ranking females receive more allogrooming. Revealing the positions, the roles, and the interactional behavioral patterns of individuals can help understand the mechanisms that shape the overall structure of a social network.

Naud et al. studied another species of primates, the Mandrills. The objective of their study was to investigate how the group spatial distribution of a semi-free ranging colony of Mandrills in a food competition context relates to its social organization. Their results showed that high-ranking individuals were more observed in proximity of the feeding zone but that affiliative relationships were also associated with individual spatial distributions and explain more the individual distribution than dominance hierarchy.

These studies showed that within a group social interactions can take many forms and may significantly affect an individual's fitness (Silk et al., 2003; Formica et al., 2012; Kurvers et al., 2014). These interactions may result in complex systems at the group-level, such as in the case of collective decisions (Sueur et al., 2012). Among them, social transmission of information has been studied mostly in vertebrates (Whiten and van Schaik, 2007). Duboscq et al. reviewed the context and

## OPEN ACCESS

### Edited by:

Jeffrey R. Stevens,  
University of Nebraska–Lincoln, USA

### Reviewed by:

Daizaburo Shizuka,  
University of Nebraska–Lincoln, USA

### \*Correspondence:

Cédric Sueur  
cedric.sueur@iphc.cnrs.fr

### Specialty section:

This article was submitted to  
Comparative Psychology,  
a section of the journal  
Frontiers in Psychology

**Received:** 13 December 2016

**Accepted:** 06 January 2017

**Published:** 19 January 2017

### Citation:

Sueur C and Mery F (2017) Editorial:  
Social Interaction in Animals: Linking  
Experimental Approach and Social  
Network Analysis.  
Front. Psychol. 8:35.  
doi: 10.3389/fpsyg.2017.00035

the methodology of experiments testing social transmission of information. However, they also discussed the reasons why social transmission sometimes does not occur despite being expected to and spanned a full range of mechanisms and processes including the constraints imposed by the social networks in which animals are embedded.

In a study on zebra finches, Fernandez et al. designed a method analyzing group vocal network semi-automatically. They wanted to test the hypothesis that the social structure of the group influences the parameters of the group vocal network. Using Markov analysis and cross-correlation analyses, they showed that juveniles as well as adults were more likely to respond to individuals of their own age-class.

In insects, social learning has been unambiguously demonstrated in social Hymenoptera but this probably reflects limited research effort and recent evidence show that even non-eusocial insects such as *Drosophila*, cockroaches, and crickets can copy the behavior of others (Battesti et al., 2012, 2015; Waters and Fewell, 2012). In this way, Pasquaretta et al. also used the RSiena package to analyze the dynamic of the interaction network of the fruit fly *Drosophila melanogaster* during social learning experiments. This work showed the importance of new methodologies in social network analyses to better understand causes and effects of animal social networks properties. The study of the processes which may facilitate or prevent this transmission and the analyses of the relationship between social network structure and efficiency of social transmission became in recent years an emerging and promising field of research (Sueur, 2011; Pasquaretta et al., 2014).

For instance, a number of recent studies have used Network Based Diffusion Analysis (NBDA) to detect the role of social transmission in the spread of a novel behavior through a population (Franz and Nunn, 2009; Hoppitt et al., 2010). Whalen and Hoppitt presented in this special issue a unified framework for performing NBDA in a Bayesian setting, and demonstrated how the Watanabe Akaike Information Criteria (WAIC) can be used for model selection. They performed a large scale simulation study and found that NBDA using WAIC could recover the correct model of social transmission under a wide range of cases, including under the presence of random effects, individual level variables, and alternative models of social transmission.

On another topic, Senior et al. worked on an integrated model approach between social network analysis and nutritional

behavior. Animals have evolved complex foraging strategies to obtain a nutritionally balanced diet and associated fitness benefits. This nutritional behavior can also influence animal social interactions and affect group structures. Senior et al. demonstrated how social network analyses can be integrated into such a nutritional modeling framework. They illustrated their approach by examining the case of nutritionally mediated dominance hierarchies and demonstrated how metrics from social network analyses can be used to predict the fitness of agents in these simulations.

Health is a component of fitness also very well studied in Animal Behavioral Sciences (Abbot et al., 2011; MacIntosh et al., 2011; Rico-Uribe et al., 2016). In their study, McCowan et al. argued that nonhuman primate social systems are sufficiently complex to serve as model systems to study links between social life and health as we might observe in Humans. The influence of social contexts influencing health and fitness in non-human primates might help us to improve human health.

Finally, Golemić et al. used a layer motif approach to understand social networks of kindergarten children and concluded that this method can be applicable on a more general scale to any group of individuals where interactions and identities can be readily observed and scored.

Using different animal species, including humans, this special issue investigated and showed how the structure of a group affects social interaction, information transfer, and collective decisions; but also how individuals treat different sources of information according to their sociality and the latest methodologies used to understand these processes.

## AUTHOR CONTRIBUTIONS

All authors listed, have made substantial, direct and intellectual contribution to the work, and approved it for publication.

## FUNDING

This project is funded by an ANR programme Blanc (ANR 12 BSV7 0013 02) to FM and CS and to an ANR Programme Santé Publique (ANR-15-CE36-0005-01) to CS. CS is funded by the University of Strasbourg Institute for Advanced Study (USIAS).

## REFERENCES

- Abbot, P., Abe, J., Alcock, J., Alizon, S., Alpedrinha, J. A., Andersson, M., et al. (2011). Inclusive fitness theory and eusociality. *Nature* 471, E1–E4. doi: 10.1038/nature09831
- Battesti, M., Moreno, C., Joly, D., and Mery, F. (2012). Spread of social information and dynamics of social transmission within *Drosophila* groups. *Curr. Biol.* 22, 309–313. doi: 10.1016/j.cub.2011.12.050
- Battesti, M., Pasquaretta, C., Moreno, C., Tesse, S., Joly, D., Klensch, E., et al. (2015). Ecology of information: social transmission dynamics within groups of non-social insects. *Proc. R. Soc. Lond. B Biol. Sci.* 282:20142480. doi: 10.1098/rspb.2014.2480
- Formica, V. A., Wood, C. W., Larsen, W. B., Butterfield, R. E., Augat, M. E., Hougen, H. Y., et al. (2012). Fitness consequences of social network position in a wild population of forked fungus beetles (*Bolitotherus cornutus*). *J. Evol. Biol.* 25, 130–137. doi: 10.1111/j.1420-9101.2011.02411.x
- Franz, M., and Nunn, C. L. (2009). Network-based diffusion analysis: a new method for detecting social learning. *Proc. R. Soc. B Biol. Sci.* 276, 1829–1836. doi: 10.1098/rspb.2008.1824
- Hoppitt, W., Boogert, N. J., and Laland, K. N. (2010). Detecting social transmission in networks. *J. Theor. Biol.* 263, 544–555. doi: 10.1016/j.jtbi.2010.01.004
- Krause, J., Lusseau, D., and James, R. (2009). Animal social networks: an introduction. *Behav. Ecol. Sociobiol. (Print)*. 63, 967–973. doi: 10.1007/s00265-009-0747-0

- Kurvers, R. H. Krause, J., Croft, D. P., Wilson, A. D., and Wolf, M. (2014). The evolutionary and ecological consequences of animal social networks: emerging issues. *Trends Ecol. Evol. (Amst)*. 29, 326–335. doi: 10.1016/j.tree.2014.04.002
- MacIntosh, A. J., Alados, C. L., and Huffman, M. A. (2011). Fractal analysis of behaviour in a wild primate: behavioural complexity in health and disease. *J. R. Soc. Interface* 8, 1497–1509. doi: 10.1098/rsif.2011.0049
- Pasquaretta, C., Levé, M., Claidière, N., van de Waal, E., Whiten, A., MacIntosh, A. J., et al. (2014). Social networks in primates: smart and tolerant species have more efficient networks. *Sci. Rep.* 4:7600. doi: 10.1038/srep07600
- Rico-Uribe, L. A., Caballero, F. F., Olaya, B., Tobiasz-Adamczyk, B., Koskinen, S., Leonardi, M., et al. (2016). Loneliness, social networks, and health: a cross-sectional study in three countries. *PLoS ONE* 11:e0145264. doi: 10.1371/journal.pone.0145264
- Silk, J. B., Alberts, S. C., and Altmann, J. (2003). Social bonds of female baboons enhance infant survival. *Science* 302, 1231–1234. doi: 10.1126/science.1088580
- Snijders, T. A. B. (2001). The statistical evaluation of social network dynamics. *Soc. Methodol.* 31, 361–395. doi: 10.1111/0081-1750.00099
- Sueur, C. (2011). “Social network, information flow and decision-making efficiency: a comparison of humans and animals,” in *Social Networking and Community Behavior Modeling: Qualitative and Quantitative Measures*, ed. M. Safar (Hershey, PA: IGI Global), 164–177.
- Sueur, C., Deneubourg, J.-L., and Petit, O. (2012). From social network (centralized vs. decentralized) to collective decision-making (unshared vs. shared consensus). *PLoS ONE* 7:e32566. doi: 10.1371/journal.pone.0032566
- Sueur, C., Jacobs, A., Amblard, F., Petit, O., and King, A. J. (2011). How can social network analysis improve the study of primate behavior? *Am. J. Primatol.* 73, 703–719. doi: 10.1002/ajp.20915
- Waters, J. S., and Fewell, J. H. (2012). Information processing in social insect networks. *PLoS ONE* 7:e40337. doi: 10.1371/journal.pone.0040337
- Whiten, A., and van Schaik, C. P. (2007). The evolution of animal “cultures” and social intelligence. *Philos. Trans. R. Soc. B Biol. Sci.* 362, 603–620. doi: 10.1098/rstb.2006.1998

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer DS and handling Editor declared their shared affiliation, and the handling Editor states that the process nevertheless met the standards of a fair and objective review.

Copyright © 2017 Sueur and Mery. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.