

Remote sensing for mapping Natura 2000 habitats in the Brière marshes (France, Loire-Atlantique): setting up a long-term monitoring strategy to understand changes

Thomas Lafitte, Marc Robin

► To cite this version:

Thomas Lafitte, Marc Robin. Remote sensing for mapping Natura 2000 habitats in the Brière marshes (France, Loire-Atlantique): setting up a long-term monitoring strategy to understand changes. EGU General Assembly 2024, Apr 2024, Vienne (Autriche), Austria. hal-04689282

HAL Id: hal-04689282 https://hal.science/hal-04689282v1

Submitted on 6 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Slide 1 :

Presentation of the project

I come from France, I'm going to present you some results of my thesis.

Slide 2 :

- location : west Atlantic coast in France

-thesis work on the Regional natural park of Briere

-more precisely on the Natura 2000 sector over 17 000 hectares

-main aim of the thesis is to study the changing dynamics of plant formations on the entire sector, in order to enable prospective modelling

Slide 3 :

-three different maps were available at the beginning of the study

-field sampling efforts are very costly and time-consuming, and are hampered by difficult access to the extensive marsh areas

-not spatially exhaustive, present a simplification of habitat mosaics, making modelling impossible.

Slide 4 :

-The first step involves an hyperspectral and LiDAR aerial survey across the whole sector.

-It is completed by the acquisition of a World View 3 scene, in order to assess the contributions of the different types of data \rightarrow because hyspec aerial survey very costly and hard to realize. And to set-up the observatory

-At the same time, floristic field surveys were carried out to characterize the textural and spectral variability of the images \rightarrow transformed into ROI

-Automatic classification methods were then applied.

-differences between the variables used : only 19 for WV3 included the DHM from the LiDAR survey

-results : 13 habitats and a Out Of Bag

Slide 5 :

-in Briere, two invasive alien species are very endangering the habitats

-the trees are also invasive

Slide 6 :

-a complete mapping over the 17000 hectares of marshland has been carried out,

-according to the Eunis declination

-WV3 is not spatially exhaustive but it's a test, with no big differences

Slide 7 :

-we can follow Crassula helmsii and Ludwigia , 0.00% of confusion with other species in the matrix of confusion

-for 1950, 1977, 1999 photointerpretation \rightarrow very time consuming and boring

So in 2023 we can have the distribution and the species

Slide 8 :

-as a result : Complete mapping, according to the Eunis declination

-2024 : National LiDAR survey planned by the National Geographic Institute

-2025 : a new WV03 image + DHM from de NGI LiDAR

Questions

Why didn't you use a hyperspectral satellite ?

Satellite such as ENDMAP is 30 meters resolution, it is not enough

How did you obtain the ROI :

1-identify the texture and spectral variability of the image by applying a vegetation indice

2-go on the localisation

3-describe the dominant species

What is the cost ?

50 \$ of the kilometer square, for the hyspec hard to say because there is the hyspec +lidar but it is much more

The update of the acquisition of the image ?

To study the alien species, maybe 2 images in summer in July in the end of august

For classic communities, updated each year

Link between the research and the Park ?

The CIFRE programme is a partnership between a French company, a laboratory and a PhD student

According to climate change how will the swamp evolve ?

Is the main goal of the thesis, we will do a prospective modelisation of the changing dynamics of plant formations

Less precipitations and a longer warm period, high temperature

We will include the management practices in the model

Explication of the alien specie distribution ?

North factor : species competition + lower topographical level too much water during a longer period

South factor : salt concentration in the soil

Future of sea level rise ?

We know that in the estuary sea level will rise from 60 to 1meter until 2100 so plant communities will be very impacted, maybe the swamp will be a gulf with salt water