



HAL
open science

On Sparsity and Sub-Gaussianity in the Johnson-Lindenstrauss Lemma

Aurélien Garivier, Emmanuel Pilliat

► **To cite this version:**

Aurélien Garivier, Emmanuel Pilliat. On Sparsity and Sub-Gaussianity in the Johnson-Lindenstrauss Lemma. 2024. hal-04690731v1

HAL Id: hal-04690731

<https://hal.science/hal-04690731v1>

Preprint submitted on 6 Sep 2024 (v1), last revised 23 Sep 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License

ON SPARSITY AND SUB-GAUSSIANITY IN THE JOHNSON-LINDENSTRAUSS LEMMA ^{*,**}

AURÉLIEN GARIVIER AND EMMANUEL PILLIAT¹

Abstract. We provide a simple proof of the Johnson-Lindenstrauss lemma for sub-Gaussian variables. We extend the analysis to identify how sparse projections can be, and what the cost of sparsity is on the target dimension. The Johnson-Lindenstrauss lemma is the theoretical core of the dimensionality reduction methods based on random projections. While its original formulation involves matrices with Gaussian entries, the computational cost of random projections can be drastically reduced by the use of simpler variables, especially if they vanish with a high probability. In this paper, we propose a simple and elementary analysis of random projections under classical assumptions that emphasizes the key role of sub-Gaussianity. Furthermore, we show how to extend it to sparse projections, emphasizing the limits induced by the sparsity of the data itself.

Résumé. Nous présentons ici une preuve simple du lemme de Johnson-Lindenstrauss pour les variables sous-Gaussiennes, qui permet d'identifier à quel point les matrices de projections peuvent être creuses et avec quelles conséquences pour la dimension cible. Le lemme de Johnson-Lindenstrauss est au cœur des méthodes de réduction de dimension par projections aléatoires. Son énoncé initial impliquait des matrices de variables Gaussiennes, mais il a ensuite été montré que des variables plus simples, pouvant être nulles avec une probabilité importante, présentaient les mêmes garanties théoriques tout en réduisant drastiquement le coût de calcul. Nous proposons dans cet article une analyse simple et élémentaire des projections aléatoires qui met en lumière le rôle clé de la sous-Gaussianité. En outre, nous montrons comment étendre cette analyse aux matrices creuses, en mettant au jour les limites induites par des données elles-même parcimonieuses.

2020 Mathematics Subject Classification. 62, 60.

August 30th 2024.

INTRODUCTION

The celebrated Johnson-Lindenstrauss lemma [9] ensures the existence low-distortion embeddings of points from high-dimensional into low-dimensional Euclidean space. If $x_1, \dots, x_n \in \mathbb{R}^p$, where p is a (large) integer, and if $\epsilon > 0$ is a tolerance parameter, then there exists a matrix $A \in \mathcal{M}_{d,p}(\mathbb{R})$ such that

$$\forall 1 \leq i, j \leq n, \quad (1 - \epsilon)\|Ax_i - Ax_j\|^2 \leq \|x_i - x_j\|^2 \leq (1 + \epsilon)\|Ax_i - Ax_j\|^2 \quad (1)$$

Keywords and phrases: Johnson-Lindenstrauss, Random Projections, Sparsity

* *Chaire SeqALO (ANR-20-CHIA-0020-01)*

** *PEPR IA project FOUNDRY (ANR-23-PEIA-0003)*

¹ Univ. Lyon ENS de Lyon UMPA UMR 5669 / LIP UMR 5668 46 allée d'Italie F-69364 Lyon cedex 07

as soon as

$$d \geq \frac{8 \log(n)}{\epsilon^2 - \epsilon^3}. \quad (2)$$

The classical proof of this result is an elegant illustration of the Probabilistic Method [2]: when drawing the entries of A at random from independent Gaussian distributions, Property (1) is satisfied with positive probability when the output space is large enough. It results from a simple deviation bound for the chi-square distribution, and hence builds on the specificity of the Gaussian distribution. This proof is not only mathematically remarkable, but it also gives mathematical foundations for *random projections*, a simple and computationally efficient dimensionality reduction technique in unsupervised machine learning (see e.g. [3, 6, 8, 14, 15] and references therein).

In 2001, [1] however showed that random projections can easily be extended to non-Gaussian matrices. In particular, Rademacher, or $\{-1, 0, 1\}$ -valued entries can just as well be chosen, leading to even simpler algorithms suitable for database applications. The proof provided in this article relies on moment bounds and is somewhat specific to those two distributions. It is generally considered [11] that "a uniform distribution is easier to generate than normals, but the analysis is more difficult". Even faster methods for sparse data or streams were then devised [5, 10] using random hashing constructions and more involved moment bounds. Very recently and concurrently to our work, [12] has proposed a unified analysis of sparse Johnson-Lindenstrauss methods based on the Hanson-Wright inequality, while [7] tries to identify the optimal rate of sparsity in the data as a function of the dimension d , the number of points n and the tolerance parameter ϵ .

The main contribution of this paper is twofold. First, the pedagogical purpose of this note is to highlight that *sub-Gaussianity* is indeed an elementary property of random matrix entries that suffices to ensure the success of random projections. Contrary to [12], our analysis is entirely elementary, and exploits sub-Gaussianity in an original way. A connection to the Hanson-Wright inequality is proposed at the end of the paper. To begin, we give here a simple proof that any 1-sub-Gaussian law with variance 1 offers the same guarantees as the Gaussian law. Our analysis explains simply why $\{-1, 0, 1\}$ -valued variables with a proportion up to $2/3$ of coefficients equal to 0 are a safe choice, but also makes it possible to design many variants, and to go further in the understanding of much sparser random projections. Interestingly, our treatments of the lower- and the upper bound of (1) are not totally symmetric. While the upper deviations of sub-Gaussian variables can be handled by Chernoff's bound just as those of the Gaussian law, the lower deviations can obviously be much smaller (after all, constant variables are sub-Gaussian) and hence require a different argument. The second purpose of this paper is to build on this analysis to clearly emphasize the conditions on the data under which much sparser projection matrices can be considered. The take-home message is that the distances are preserved if and only if the projection matrix entries are non-zero with a probability larger than the inverse of the number s of significant coefficients in the data.

The paper is organized as follows. Section 1 treats the deviations of an average of squared sub-Gaussian variables. The obtained bound is applied in Section 2 to derive the classical Johnson-Lindenstrauss lemma for sub-Gaussian random matrices. We discuss in Section 3 a few examples of choices of the distribution P for random projections. We explain in Section 4 why much sparser choices are possible. We show in Section 5 that the quasi-isometry property still holds for matrices as sparse as can be, at the price of poly-logarithmic terms in the target dimension. The optimality of this result is discussed in Section 6, together with a connection to the Hanson-Richt inequality.

1. CHERNOFF'S METHOD FOR SQUARED SUB-GAUSSIAN VARIABLES

Let X be a random variable assumed to be 1-sub-Gaussian, which means that $\forall \lambda \in \mathbb{R}, \mathbb{E}[e^{\lambda X}] \leq e^{\lambda^2/2}$. This implies in particular that $\mathbb{E}[X] = 0$ and that $\text{Var}[X] \leq 1$. We derive in this section a deviation bound for the empirical mean of independent copies of X^2 . Chernoff's method requires to bound the exponential moments $\mathbb{E}[e^{\ell X^2}]$ of X^2 with $\ell > 0$ for the right deviations and with $\ell < 0$ for the left deviations.

1.1. The upper bound

We start with the right deviations, for which we will see right away that a reduction to the Gaussian case is possible without further assumption. Following [16] (Theorem 2.6), and remarking that for all $x \in \mathbb{R}$, and $\ell > 0$,

$$e^{\ell x^2} = \int_{-\infty}^{\infty} e^{\lambda x} \frac{e^{-\frac{\lambda^2}{4\ell}}}{2\sqrt{\pi\ell}} d\lambda,$$

if X is 1-sub-Gaussian we obtain by Fubini's theorem that for every $\ell \in (0, 1/2)$

$$\begin{aligned} \mathbb{E} \left[e^{\ell X^2} \right] &= \mathbb{E} \left[\int_{-\infty}^{\infty} e^{\lambda X} \frac{e^{-\frac{\lambda^2}{4\ell}}}{2\sqrt{\pi\ell}} d\lambda \right] = \int_{-\infty}^{\infty} \mathbb{E} [e^{\lambda X}] \frac{e^{-\frac{\lambda^2}{4\ell}}}{2\sqrt{\pi\ell}} d\lambda \\ &\leq \int_{-\infty}^{\infty} e^{\frac{\lambda^2}{2}} \frac{e^{-\frac{\lambda^2}{4\ell}}}{2\sqrt{\pi\ell}} d\lambda = \int_{-\infty}^{\infty} e^{-\frac{\lambda^2(1-2\ell)}{4\ell}} \frac{d\lambda}{2\sqrt{\pi\ell}} = \frac{1}{\sqrt{1-2\ell}}, \end{aligned}$$

which holds with equality if and only if $X \sim \mathcal{N}(0, 1)$. Equivalently: observe that if $G \sim \mathcal{N}(0, 1)$, Fubini's theorem implies that

$$\begin{aligned} \mathbb{E}_X \left[e^{\ell X^2} \right] &= \mathbb{E}_X \left[\mathbb{E}_G \left[e^{\sqrt{2\ell} X G} \right] \right] = \mathbb{E}_G \left[\mathbb{E}_X \left[e^{\sqrt{2\ell} G X} \right] \right] \\ &\leq \mathbb{E}_G \left[e^{\ell G^2} \right] = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{\ell u^2} e^{-\frac{u^2}{2}} du = \frac{1}{\sqrt{1-2\ell}} \end{aligned} \quad (3)$$

with equality if and only if $X \sim \mathcal{N}(0, 1)$.

Hence all sub-Gaussian variables have exponential moments bounded by those of a Gaussian law, which permits the right-deviations to be handled the usual way. If Z_1, \dots, Z_d are independent random variables with the same distribution as X^2 , then for every positive ϵ , Markov's inequality implies that

$$\mathbb{P} \left(\frac{Z_1 + \dots + Z_d}{d} \geq 1 + \epsilon \right) = \mathbb{P} \left(e^{\ell(Z_1 + \dots + Z_d)} \geq e^{d\ell(1+\epsilon)} \right) \leq \frac{\mathbb{E}[e^{\ell Z_1}]^d}{e^{d\ell(1+\epsilon)}} = e^{-d(\ell(1+\epsilon) - \ln \mathbb{E}[e^{\ell Z_1}])}.$$

The concave function $\ell \mapsto \ell(1+\epsilon) - \ln \mathbb{E}[e^{\ell X}] = \ell(1+\epsilon) + \frac{1}{2} \log(1-2\ell)$ is maximized at ℓ^* such that $1+\epsilon = \frac{1}{1-2\ell^*}$, that is at $\ell^* = \frac{1}{2} \left(1 - \frac{1}{1+\epsilon} \right) = \frac{\epsilon}{2(1+\epsilon)}$. Hence, $\mathbb{P}(Z_1 + \dots + Z_d \geq (1+\epsilon)d) \leq e^{-dI(\epsilon)}$ with

$$I(\epsilon) = \ell^*(1+\epsilon) - \ln \mathbb{E}[e^{\ell^* X}] = \frac{\epsilon - \log(1+\epsilon)}{2}.$$

This expression can be slightly simplified in many different ways. Let us illustrate the very useful "Pollard trick": taking $g(\epsilon) = \epsilon - \log(1+\epsilon)$, since $g(0) = g'(0) = 0$ and since $g''(\epsilon) = 1/(1+\epsilon)^2$ is convex, by Jensen's inequality

$$\frac{\epsilon - \log(1+\epsilon)}{\epsilon^2/2} = \int_0^1 g''(s\epsilon) 2(1-s) ds \geq g'' \left(\epsilon \int_0^1 s 2(1-s) ds \right) = g'' \left(\frac{\epsilon}{3} \right),$$

and hence $I(\epsilon) = \frac{\epsilon - \log(1+\epsilon)}{2} \geq \frac{\epsilon^2}{4 \left(1 + \frac{\epsilon}{3}\right)^2} \geq \frac{\epsilon^2 - \epsilon^3}{4}$. In summary,

$$\mathbb{P} \left(\frac{Z_1 + \dots + Z_d}{d} \geq 1 + \epsilon \right) \leq e^{-d \left(\frac{\epsilon^2 - \epsilon^3}{4} \right)}. \quad (4)$$

1.2. The lower bound

There is no hope to prove that $\mathbb{E}\left[e^{-\ell X^2}\right] \leq \frac{1}{\sqrt{1+2\ell}}$ for any $\ell > 0$ for all 1-sub-Gaussian distributions, since it is for example not the case if $X = 0$ almost surely. In the context of the Johnson-Lindenstrauss lemma, it is very natural to assume that the entries of the random matrix have variance 1, so that at least $\mathbb{E}[\|Ax_i - Ax_j\|^2] = \|x_i - x_j\|^2$. Under this assumption, it is maybe possible to bound the negative exponential moments bounded by those of the standard Gaussian. and to conclude (as in the Gaussian case) by remarking that $I(-\epsilon) \geq I(\epsilon)$, i.e. that the left-deviations of the Chi-square are lighter than the right deviations. But we do unfortunately not have a proof for that.

Instead, we remark that if $\text{Var}[X] = 1$, the sub-Gaussianity inequality $\mathbb{E}[e^{\lambda X}] \leq e^{\lambda^2/2}$ implies by Taylor expansion around $\lambda = 0$ that $\mathbb{E}[X^4] \leq 3$. Using that $e^{-u} \leq 1 - u + \frac{u^2}{2}$, we obtain that $\mathbb{E}\left[e^{-\ell X^2}\right] \leq 1 - \ell + \frac{3\ell^2}{2}$ and hence

$$\mathbb{P}\left(\frac{Z_1 + \dots + Z_d}{d} \leq 1 - \epsilon\right) \leq e^{-d(\ell(-1+\epsilon) - \ln(1 - \ell + \frac{3\ell^2}{2}))}.$$

Since $-\ln(1 - u) \geq u + u^2/2$,

$$\ell(-1 + \epsilon) - \ln\left(1 - \ell + \frac{3\ell^2}{2}\right) \geq \ell\epsilon - \frac{3\ell^2}{2} + \frac{(\ell - 3\ell^2/2)^2}{2} \geq \ell\epsilon - \ell^2 - \frac{3\ell^3}{2} = \frac{\epsilon^2}{4} - \frac{3\epsilon^3}{16}$$

for $\ell = \epsilon/2$. It follows that

$$\mathbb{P}\left(\frac{Z_1 + \dots + Z_d}{d} \leq 1 - \epsilon\right) \leq e^{-d\left(\frac{\epsilon^2}{4} - \frac{3\epsilon^3}{16}\right)} \leq e^{-d\left(\frac{\epsilon^2 - \epsilon^3}{4}\right)}. \quad (5)$$

2. APPLICATION TO THE JOHNSON-LINDENSTRAUSS LEMMA

In the sequel, we assume that $A_{i,j} = T_{i,j}/\sqrt{d}$, $1 \leq i \leq d, 1 \leq j \leq p$, where the $(T_{i,j})$ centered, standard independent variables of a 1-sub-Gaussian distribution P :

$$\mathbb{E}[Z_{i,j}] = 0, \quad \text{Var}[T_{i,j}] = 1, \quad \mathbb{E}\left[e^{\lambda T_{i,j}}\right] \leq e^{\frac{\lambda^2}{2}}.$$

For a vector $y \in \mathbb{R}^p$, define $Y = Ay$ and for all $i \in \{1, \dots, d\}$

$$Z_i = \frac{\sqrt{d} Y_i}{\|y\|} = \sum_{j=1}^p \frac{y_j}{\|y\|} T_{i,j}.$$

Then, as for all $\lambda \in \mathbb{R}$

$$\mathbb{E}\left[e^{\lambda Z_i}\right] = \prod_{j=1}^p \mathbb{E}\left[e^{\lambda \frac{y_j}{\|y\|} T_{i,j}}\right] \leq \prod_{j=1}^p e^{\frac{y_j^2 \lambda^2}{2\|y\|^2}} = e^{\frac{\lambda^2}{2}},$$

Z_i is 1-sub-Gaussian. Since

$$\frac{\|Ay\|^2}{\|y\|^2} = \frac{1}{d} \sum_{i=1}^d \left(\frac{\sqrt{d} Y_i}{\|y\|}\right)^2 = \frac{1}{d} \sum_{i=1}^d Z_i^2,$$

Equations (4) and (5) yield:

$$\mathbb{P}\left(\frac{\|Ay\|^2}{\|y\|^2} \notin [1 - \epsilon, 1 + \epsilon]\right) = \mathbb{P}\left(\frac{1}{d} \sum_{i=1}^d Z_i^2 > 1 + \epsilon\right) + \mathbb{P}\left(\frac{1}{d} \sum_{i=1}^d Z_i^2 < 1 - \epsilon\right) \leq 2e^{-d\left(\frac{\epsilon^2 - \epsilon^3}{4}\right)} \leq \frac{2}{n^2}$$

as soon as $d \geq \frac{8 \log(n)}{\epsilon^2 - \epsilon^3}$. By the union bound,

$$\mathbb{P} \left(\bigcup_{1 \leq i < j \leq n} \left\{ \|A(x_i - x_j)\|^2 \notin [(1 - \epsilon)\|x_i - x_j\|^2, (1 + \epsilon)\|x_i - x_j\|^2] \right\} \right) \leq \frac{n(n-1)}{n^2} < 1,$$

hence giving the desired conclusion.

Observe that the constant 8 in Condition (2) is the best that can be obtained from this proof. The dependency in $1/\epsilon^2$ also appears to be necessary, but the second-order term ϵ^3 is slightly improvable. In the Gaussian case, the proof above allows to use

$$d = \frac{4 \log(n)}{\epsilon - \log(1 + \epsilon)} \leq \frac{8 \log(n)}{\epsilon^2} \left(1 + \frac{\epsilon}{3}\right)^2,$$

as we saw in Section 1.1. For sub-Gaussian variables, the simple expression (2) covers at the same time left- and right-deviations. Also note that choosing $d \geq \frac{4 \log(n^2/\delta)}{\epsilon^2 - \epsilon^3}$ permits Property (1) to hold with probability at least $1 - \delta$.

3. WHAT DISTRIBUTION SHOULD WE USE IN RANDOM PROJECTIONS?

We have thus seen that any 1-sub-Gaussian distribution of variance 1 presents just the same guarantees as the standard Gaussian for random projections. This is for example the case of $P = \frac{\delta_{-1} + \delta_1}{2}$, or of $P = \mathcal{U}([-\sqrt{3}, \sqrt{3}])$, which are very fast and easy laws to sample. To see it, observe that their exponential moment functions are $\cosh(\lambda)$ and $\sinh(\sqrt{3}\lambda)/(\sqrt{3}\lambda)$, which are both easily seen to be upper-bounded by $e^{\lambda^2/2}$.

One may wonder, after [1], how *sparse* a random projection matrix can be (sparse matrices require fewer computations). A sparse choice can be written $X = \epsilon U$, where $\epsilon \sim \mathcal{B}(q)$ and U is a centered random variable. The requirement $\text{Var}[X] = 1$ implies $\mathbb{E}[U^2] = 1/q$. 1-sub-Gaussianity then requires $\mathbb{E}[X^4] = q\mathbb{E}[U^4] \leq 3$, and since $\mathbb{E}[U^4] \geq \mathbb{E}[U^2]^2 = 1/q^2$ this implies that $q \geq 1/3$. Moreover, the choice $q = 1/3$ is possible only if $\mathbb{E}[U^4] \geq \mathbb{E}[U^2]^2$, that is if $U^2 = 1/q$ almost surely. The choice

$$P = \frac{q}{2} \delta_{-\frac{1}{\sqrt{q}}} + (1 - q) \delta_0 + \frac{q}{2} \delta_{\frac{1}{\sqrt{q}}} \quad (6)$$

with $q = 1/3$ is indeed the suggestion of Achlioptas, and it is 1-sub-Gaussian. The justification of this choice in [1] is pretty involved, while we here only need to check that for all $\lambda \in \mathbb{R}$,

$$\mathbb{E}[e^{\lambda X}] = 1 - p + p \cosh\left(\frac{\lambda}{\sqrt{p}}\right) = 1 + \sum_{k=1}^{\infty} \frac{\lambda^{2k}}{p^{k-1}(2k)!} \leq e^{\lambda^2/2} = 1 + \sum_{k=1}^{\infty} \frac{\lambda^{2k}}{2^k k!}$$

whenever $q \geq 1/3$. A sufficient condition for the inequality is that for all $k \geq 1$,

$$\frac{1}{q^{k-1}(2k)!} \leq \frac{1}{2^k k!} \iff q^{k-1} \geq \frac{2^k k!}{(2k)!}. \quad (7)$$

For $k = 1$ this is always true, for $k = 2$ it requires that $q \geq \frac{4 \times 2}{24} = \frac{1}{3}$. A simple induction shows that if $q \geq 1/3$, the condition is also satisfied for all $k \geq 3$. Reciprocally, if $q < 1/3$ then $\mathbb{E}[e^{\lambda X}] - e^{\lambda^2/2} \sim_{\lambda \rightarrow 0} -c\lambda^4$ for a positive constant c , and P is not 1-sub-Gaussian. This shows that Achlioptas' suggestion is the only "optimal" choice in terms of sparsity. Nevertheless, many other choices are possible, such as for example $P = \frac{1}{12} \delta_{-2} + \frac{1}{6} \delta_{-1} + \frac{1}{2} \delta_0 + \frac{1}{6} \delta_1 + \frac{1}{12} \delta_2$.




FIGURE 1. Admissible value ϵ , in function of the sparsity parameter q of the random projection entries. Each data point $x_i \in \mathbb{R}^{10000}$ has independent Gaussian entries. The projection matrix A has independent entries with distribution $\frac{q}{2}\delta_{-\frac{1}{\sqrt{dq}}} + (1-q)\delta_0 + \frac{q}{2}\delta_{\frac{1}{\sqrt{dq}}}$. The target dimension is $d = 500$. The blue line shows $\min_{i,j} \frac{\|A(x_i-x_j)\|^2}{\|x_i-x_j\|^2}$, while the red line shows $\max_{i,j} \frac{\|A(x_i-x_j)\|^2}{\|x_i-x_j\|^2}$. The value $q = 1/3$ seems to play no special role, much sparser matrices seem to respect pairwise distances just as well.

4. TOWARDS SPARSER MATRICES

In the previous section, we showed that the minimal probability q for the non-zeros values of a suitable 1-sub-Gaussian distribution P is $1/3$. In fact, this result was proven in [11] with somehow more complicated moment arguments. It allows to take a target dimension $d \geq 8 \log(n)/(\epsilon^2 - \epsilon^3)$ – see (2) – to get a ϵ -quasi-isometry with nonzero probability, whatever the data x . The previous analysis remains however quite conservative in that the sub-Gaussianity of Z_i is deduced from the sub-Gaussianity of *each* of its summands. We may expect to gain a lot of sparsity by using the fact that a sum can be a lot more concentrated than each of its components. Figure 1 suggests that, at least under certain conditions on the data, much sparser matrices may be considered.

Let indeed $U \in \mathbb{R}^{d \times p}$ be a matrix of iid 1-sub-Gaussian entries with variance 1, and $\zeta \in \mathbb{R}^{d \times p}$ be a matrix of iid Bernoulli variables of parameter q independent from U that is used to *mask* a proportion $1 - q$ of the coefficients. We assume that for all i, k ,

$$A_{ik} = \frac{1}{\sqrt{dq}} \zeta_{ik} U_{ik}, \quad (8)$$

and write as before $Y = Ay$ and $Z_i = \frac{\sqrt{d}Y_i}{\|y\|_2}$.

Following the previous analysis, we need to bound $\mathbb{E}[\exp(\lambda Z_i^2)]$, and we know how to do it from $\mathbb{E}[\exp(\lambda Z_i)]$ when Z_i is sub-Gaussian thanks to the argument of Inequality (3). Since $Z_i = \sum_{k=1}^p y_k \frac{\zeta_{ik} U_{ik}}{\|y\|_2 \sqrt{q}}$ is a sum of many small contributions, for any fixed λ we can bound $\ln \mathbb{E}[\exp(\lambda Z_i)]$ using only the local behaviour of $\psi(\lambda) := \ln \mathbb{E}[\exp(\lambda U_{i,k})]$ around 0, which is of order $\lambda^2/2$ even when ψ is not upper-bounded by that quantity. But using Inequality (3) would require a *uniformly* control of ψ , which we cannot provide. We are hence obliged to take another path, by conditioning on the mask variables (ζ_{ik}) and focusing on the "typical" behaviour.

Namely, let for each $i \in \{1, \dots, d\}$ and for $0 \leq \epsilon \leq 1$ let

$$G_i = \left\{ \left(1 - \frac{\epsilon}{3}\right) \|y\|_2^2 \leq \sum_{k=1}^p \frac{y_k^2}{q} \zeta_{ik} \leq \left(1 + \frac{\epsilon}{3}\right) \|y\|_2^2 \right\}.$$

By Bernstein's inequality applied on the $\left[0, \frac{\|y\|_\infty^2}{q\|y\|_2^2}\right]$ -valued independent variables $\left(\frac{y_i^2}{q\|y\|_2^2}\zeta_{ik}\right)_{1 \leq k \leq p}$, which have variance $\frac{y_i^4}{q^2\|y\|_2^4} q(1-q) \leq \frac{y_i^4}{q\|y\|_2^2}$,

$$\mathbb{P}(\bar{G}_i) \leq 2 \exp\left(-\frac{\epsilon^2/18}{\sum_{k=1}^p \frac{y_i^4}{q\|y\|_2^4} + \frac{\|y\|_\infty^2 \epsilon}{9q\|y\|_2^2}}\right) \leq 2 \exp\left(-\frac{q\epsilon^2\|y\|_2^4}{18\|y\|_4^4 + 2\|y\|_\infty^2\|y\|_2^2}\right),$$

which is smaller than $1/(2d)$ as soon as

$$q \geq \frac{18\|y\|_4^4 + 2\|y\|_\infty^2\|y\|_2^2}{\epsilon^2\|y\|_2^4} \log(2d). \quad (9)$$

On the event G_i , the behaviour of Y_i is as expected: conditioning on $(\zeta_{i,k})_k$,

$$\begin{aligned} \mathbb{E}[\exp(\lambda Z_i) \mathbb{1}_{G_i}] &= \prod_{k=1}^p \mathbb{E}\left[\exp\left(\lambda y_k \frac{\sqrt{d}\zeta_{ik}U_{ik}}{\|y\|_2\sqrt{dq}}\right) \mathbb{1}_{G_i}\right] \\ &= \prod_{k=1}^p \mathbb{E}\left[\mathbb{E}\left[\exp\left(\lambda \frac{y_k\zeta_{ik}U_{ik}}{\|y\|_2\sqrt{q}}\right) \mathbb{1}_{G_i} \middle| \zeta_{i,1}, \dots, \zeta_{i,k}\right]\right] \\ &\leq \prod_{k=1}^p \mathbb{E}\left[\exp\left(\frac{\lambda^2 y_k^2 \zeta_{ik}}{2\|y\|_2^2 q}\right) \mathbb{1}_{G_i} \middle| \zeta_{i,1}, \dots, \zeta_{i,k}\right] \\ &= \mathbb{E}\left[\exp\left(\frac{\lambda^2}{2} \sum_{k=1}^p \frac{y_k^2}{\|y\|_2^2 q} \zeta_{ik}\right) \mathbb{1}_{G_i} \middle| \zeta_{i,1}, \dots, \zeta_{i,k}\right] \\ &\leq \exp\left(\frac{\lambda^2}{2} \left(1 + \frac{\epsilon}{3}\right)\right) \end{aligned}$$

so that $Z_i/\sqrt{1+\epsilon/3}$ is 1-sub-Gaussian and by Equation (3)

$$\mathbb{E}\left[\exp\left(\frac{\ell Z_i^2}{1+\frac{\epsilon}{3}}\right) \mathbb{1}_{G_i}\right] \leq \frac{1}{\sqrt{1-2\ell}}.$$

Left deviations may be treated similarly. Hence, on the event $G = \bigcap_{i=1}^d G_i$, the behaviour of $\|Y\|$ is just as before: for all $\epsilon \leq 1$, by Equations (4) and (5)

$$\begin{aligned} \mathbb{P}\left(G \cap \left\{\frac{\|Ay\|^2}{\|y\|^2} \notin [1-\epsilon, 1+\epsilon]\right\}\right) &\leq \mathbb{P}\left(G \cap \left\{\frac{1}{d} \sum_{i=1}^d \frac{Z_i^2}{1+\frac{\epsilon}{3}} > \frac{1+\epsilon}{1+\frac{\epsilon}{3}}\right\}\right) + \mathbb{P}\left(G \cap \left\{\frac{1}{d} \sum_{i=1}^d \frac{Z_i^2}{1-\frac{\epsilon}{3}} < \frac{1-\epsilon}{1-\frac{\epsilon}{3}}\right\}\right) \\ &\leq \mathbb{P}\left(G \cap \left\{\frac{1}{d} \sum_{i=1}^d \frac{Z_i^2}{1+\frac{\epsilon}{3}} > 1+\frac{\epsilon}{3}\right\}\right) + \mathbb{P}\left(G \cap \left\{\frac{1}{d} \sum_{i=1}^d \frac{Z_i^2}{1-\frac{\epsilon}{3}} < 1-\frac{\epsilon}{3}\right\}\right) \\ &\leq 2e^{-d\left(\frac{\epsilon^2-\epsilon^3}{36}\right)}. \end{aligned}$$

Consequently,

$$\begin{aligned} & \mathbb{P} \left(\bigcup_{1 \leq i < j \leq n} \left\{ \|A(x_i - x_j)\|^2 \notin [(1 - \epsilon)\|x_i - x_j\|^2, (1 + \epsilon)\|x_i - x_j\|^2] \right\} \right) \\ & \leq P(\bar{G}) + \sum_{1 \leq i < j \leq n} \mathbb{P} \left(G \cap \left\{ \frac{\|A(x_i - x_j)\|^2}{\|x_i - x_j\|^2} \notin [1 - \epsilon, 1 + \epsilon] \right\} \right) < \frac{1}{2} + n^2 e^{-d \left(\frac{\epsilon^2 - \epsilon^3}{36} \right)} \leq 1 \end{aligned}$$

as soon as q satisfies Eq.(9) and $d \geq \frac{36 \log(2n^2)}{\epsilon^2 - \epsilon^3}$. The next section contains a discussion on sparsity conditions ensuring Condition (9). For now, it may just be observed that if all the non-zero coefficients of y are of the same order of magnitude $1/\sqrt{p}$, then q is allowed to be of order $\log(d)/(\epsilon^2 p)$. The cost in terms of target dimension is only a multiplicative constant (that is not optimized in the previous reasoning). In the sequel, we investigate the minimal order of magnitude for the value of q ensuring the quasi-isometry property (1) with a dimension d of order $1/\epsilon^2$, up to a polylogarithmic factor.

5. VERY SPARSE MATRICES

As in the previous section, we consider a matrix $U \in \mathbb{R}^{d \times p}$ with iid 1-sub-Gaussian entries of variance 1, a matrix $\zeta \in \mathbb{R}^{d \times p}$ iid of Bernoulli distributions of parameter q and independent of U , and we consider all i, k the random projection matrix A defined for $1 \leq i \leq d$ and $1 \leq k \leq p$ by

$$A_{ik} = \frac{1}{\sqrt{dq}} \zeta_{ik} U_{ik}. \quad (10)$$

The 1-sub-Gaussian choice U_{ik} is $\frac{1}{6}\delta_{-\sqrt{3}} + \frac{2}{3}\delta_0 + \frac{1}{6}\delta_{\sqrt{3}}$ maximizes the average sparsity of A , which has on average a proportion $q/3$ of non-zero coefficients. We apply the matrix A to n points x_1, \dots, x_n in a high-dimensional space \mathbb{R}^p , and we look for the minimal conditions under which the quasi-isometry property (1) still holds with high probability. We define

$$d_0 := d_0(n, \delta, \epsilon) = \frac{12}{\epsilon^2} \log(3n/\delta) \left(1 + \sqrt{4 \log(nd/\delta)} + 2 \log(nd/\delta) \right)^2. \quad (11)$$

Theorem 1 states that d_0 is a sufficient projection dimension as soon as q catches the sparsity pattern of x :

Theorem 1. *Let x_1, \dots, x_n be arbitrary vector in \mathbb{R}^p and let $A \in \mathbb{R}^{d \times p}$ be a random matrix with independent entries $A_{ik} = \frac{1}{\sqrt{dq}} \zeta_{ik} U_{ik}$ with*

$$q \geq \max_{i \neq j} \frac{\|x_i - x_j\|_\infty^2}{\|x_i - x_j\|_2^2}. \quad (12)$$

Then for any $\delta \in (0, 1)$ and any $d \geq d_0(n, \delta, \epsilon)$, the ϵ -quasi-isometry property (1) holds with probability at least $1 - \delta$.

Hence, we can take q as small as $\max_{i \neq j} \frac{\|x_i - x_j\|_\infty^2}{\|x_i - x_j\|_2^2}$ while keeping the original guarantee of Johnson Lindenstrauss (1) with nonzero probability under the same condition (2) up to a polylogarithmic factor: we require $d \geq d_0(n, 1, \epsilon)$ instead of $d \geq 8 \log(n)/(\epsilon^2 - \epsilon^3)$.

Condition (12), can be understood as a "not-too-high-sparsity" condition on the differences $x_i - x_j$, which we formalize as follows. For any constant $\kappa \in (0, 1)$ and integer any $s \in \{1, \dots, p\}$, we say that a vector v is (κ, κ', s) -full if $\|v\|_\infty \leq \sqrt{\kappa'/s}$ and if it has at least s coordinates whose absolute value are at least equal to $\sqrt{\kappa/s}$, that is

$$\|v\|_\infty \leq \sqrt{\kappa'/s} \quad \text{and} \quad |\{k : |v_k| \geq \sqrt{\kappa/s}\}| \geq s. \quad (13)$$

This implies in particular that $\|v\|^2 \geq \kappa \geq \frac{\kappa}{\kappa'} s \|v\|_\infty^2$. Hence, if a set of vectors $\{x_1, \dots, x_n\}$ is such that all the differences $x_i - x_j$ are (κ, κ', s) -full for $i \neq j$, then a sufficient condition implying (12) is

$$q \geq \left(\frac{\kappa'}{\kappa}\right) \frac{1}{s}. \quad (14)$$

In other words, we can take a matrix A which has only a proportion $q \gtrsim 1/s$ of nonzero coefficients. This condition is for instance very weak in the dense case where the differences $x_i - x_j$ are (κ, κ', p) -full for all $i \neq j$, since it only requires A to have a proportion nonzero coefficients of order $q \gtrsim 1/p$. In that case, all the coefficients of each difference $x_i - x_j$ are uniformly spread over the p dimensions, in the sense that up to constants κ, κ' , $|x_{ik} - x_{jk}| \asymp 1/\sqrt{p}$ for any k .

Condition (12) becomes however much stronger when there exists a difference $x_i - x_j$ which is s -sparse for a small s , that is $|\{k : x_{ik} \neq x_{jk}\}| \leq s$. Indeed, In such a sparse case, $\|x_i - x_j\|_\infty^2 / \|x_i - x_j\|_2^2 \geq 1/s$ implying that the condition $q \geq 1/s$ is necessary to satisfy (12).

Proof of Theorem 1. Let \odot be the Hadamard product, so that $A = \frac{1}{\sqrt{dq}} \zeta \odot U$. We assume that y is unit vector of \mathbb{R}^p representing one of the unit vector $\frac{x_i - x_j}{\|x_i - x_j\|}$, and we write as before $Y = Ay$. The coefficients of $w_i = \frac{1}{\sqrt{dq}} \zeta_i \odot y$ are equal to $w_{ik} = \frac{1}{\sqrt{dq}} \zeta_{ik} y_k$, and

$$Y_i^2 = \frac{1}{dq} \sum_{k', k} \zeta_{ik} \zeta_{ik'} U_{ik} U_{ik'} y_k y_{k'} = (U_i^T w_i)^2.$$

The upper bound

$Y_i / \|w_i\|$ is 1-sub-Gaussian conditionally to ζ . Hence, if G is a standard Gaussian random variable, it holds conditionally to ζ that for any ℓ in $[0, 1/(2 \max \|w_i\|^2))$,

$$\mathbb{E}_U \left[e^{\ell Y_i^2} \right] = \mathbb{E}_G \left[\mathbb{E}_U \left[e^{\sqrt{2\ell} Y_i G} \right] \right] \leq \mathbb{E}_G \left[e^{\ell \|w_i\|^2 G^2} \right] = \frac{1}{\sqrt{1 - 2\ell \|w_i\|^2}} \leq \exp \left(\ell \|w_i\|^2 + \frac{\ell^2 \|w_i\|^4}{1 - 2\ell \|w_i\|^2} \right),$$

where the last inequality comes from the fact that $-\frac{1}{2} \log(1 - 2\ell \|w_i\|^2) - \ell \|w_i\|^2 = \int_0^\ell \frac{2s \|w_i\|^4}{1 - 2s \|w_i\|^2} ds \leq \frac{\ell^2 \|w_i\|^4}{1 - 2\ell \|w_i\|^2}$. Hence, conditionally to ζ , we have that

$$\mathbb{P}_\zeta (\|Ay\|^2 \geq 1 + \epsilon) \leq \mathbb{E}_\zeta \left[\exp \left(\ell \sum_{i=1}^d \|w_i\|^2 + \frac{d\ell^2 \max_i \|w_i\|^4}{1 - 2\ell \max_i \|w_i\|^2} - \ell(1 + \epsilon) \right) \right]. \quad (15)$$

Let us now integrate according to ζ . The w_{ik} 's are iid random variables distributed according to $\frac{y_k}{\sqrt{dq}} \mathcal{B}(q)$. Moreover, $\text{Var}(w_{ik}^2) \leq \frac{y_k^4}{d^2 q^2} \mathbb{E}[\zeta_{ik}^4] \leq \frac{1}{d^2 q} y_k^4$. Bernstein's inequality together with a union bound over the d possible indices $i = 1, \dots, d$ gives that with probability at least $1 - \delta/(3n^2)$,

$$\max_i \|w_i\|^2 \leq \frac{1}{d} + \sqrt{2 \frac{1}{d^2 q} \|y\|_4^4 \log(3n^2 d / \delta)} + \frac{1}{dq} \|y\|_\infty^2 \log(3n^2 d / \delta).$$

The assumption (12) implies that $q \geq \|y\|_\infty^2$. Since $\|y\|^2 = 1$, $\|y\|_4^4 \leq \|y\|_\infty^2$ and the following event G holds with probability at least $\delta/(3n^2)$:

$$G = \left\{ \max_i \|w_i\|^2 \leq \frac{1}{d} \left(1 + \sqrt{4 \log(3nd/\delta)} + 2 \log(3nd/\delta) \right) \right\} = \left\{ \forall i, \|w_i\|^2 \leq \frac{\Psi}{d} \right\}. \quad (16)$$

where for simplicity we write $\Psi = 1 + \sqrt{4 \log(3nd/\delta)} + 2 \log(3nd/\delta)$. Using the inequality $e^u \leq 1 + u + (e-2)u^2$ for any $u \in [0, 1]$, we have that for any $\ell \in [0, \frac{dq}{\|y\|_\infty^2})$,

$$\begin{aligned} \mathbb{E} \left[\exp \left(\ell \sum_{i=1}^d \|w_i\|^2 \right) \right] &= \prod_{i,k} \left(q \exp\left(\frac{\ell}{dq} y_k^2\right) + 1 - q \right) \\ &\leq \prod_{i,k} \exp \left(q \left(\exp\left(\frac{\ell}{dq} y_k^2\right) - 1 \right) \right) \\ &\leq \exp \left(\ell + (e-2) \frac{\ell^2}{dq} \|y\|_4^4 \right) \\ &\leq \exp \left(\ell + (e-2) \frac{\ell^2}{d} \right). \end{aligned}$$

Let us now integrate the conditional probability $\mathbb{P}_\zeta(\|Ay\|^2 \geq 1 + \epsilon)$ over ζ . For any $\ell \in [0, d/(4\Psi))$, we have

$$\begin{aligned} \mathbb{P}(\|Ay\|^2 \geq 1 + \epsilon) &\leq \mathbb{E} \left[\exp \left(\ell \sum_{i=1}^d \|w_i\|^2 + \frac{d\ell^2 \max_i \|w_i\|^4}{1 - 2\ell \max_i \|w_i\|^2} - \ell(1 + \epsilon) \right) \mathbf{1}_G \right] + \frac{\delta}{3n^2} \\ &\leq \mathbb{E} \left[\exp \left(\ell \sum_{i=1}^d \|w_i\|^2 + 2\ell^2 \frac{A^2}{d} - \ell(1 + \epsilon) \right) \right] + \frac{\delta}{3n^2} \\ &\leq \exp \left((e-2) \frac{\ell^2}{d} + 2\ell^2 \frac{\Psi^2}{d} - \ell\epsilon \right) + \frac{\delta}{3n^2} \end{aligned}$$

The second inequality comes from Equation 15, which holds true under the event G defined in (16). The third inequality comes from the fact that $\ell \leq d/(4\Psi^2) \leq d \leq dq/\|y\|_\infty^2$ and the above upper bound on $\mathbb{E}[\exp(\ell \sum_{i=1}^d \|w_i\|^2)]$.

Choosing $\ell = d\epsilon/(2(e-2) + 4\Psi^2) \leq d/(4\Psi^2)$, we get

$$\mathbb{P}(\|Ay\|^2 \geq 1 + \epsilon) \leq \exp \left(-\frac{d\epsilon^2}{2(e-2) + 4\Psi^2} \right) + \frac{\delta}{3n^2}.$$

Hence, if $d \geq d_0 = 12 \log(3n/\delta) \Psi^2 / \epsilon^2$, we obtain that

$$\mathbb{P}(\|Ay\|^2 \geq 1 + \epsilon) \leq \exp \left(-2 \log \frac{3n}{\delta} \right) + \frac{\delta}{3n^2} \leq \frac{2\delta}{3n^2}.$$

A union bound all the $n(n-1)/2 \leq n^2$ pairs gives that

$$\mathbb{P} \left(\bigcup_{1 \leq i < j \leq n} \left\{ \|A(x_i - x_j)\|^2 \geq (1 + \epsilon) \|x_i - x_j\|^2 \right\} \right) \leq 2\delta/3. \quad (17)$$

The lower bound

For the lower bound, we use the same arguments as in section 1.2. We still have that $\mathbb{E}[(Ay)_i^2] = 1/d$ for any $i \in \{1, \dots, d\}$, but we since the variables $(Ay)_i$ are not sub-Gaussians, do not have the bound $\mathbb{E}[(Ay)_i^4] \leq 3$. Instead, we bound the fourth moment as follows:

$$\begin{aligned}\mathbb{E}[(Ay)_i^4] &\leq \frac{3}{d^2 q^2} \sum_{k \neq k'} q^2 y_k^2 y_{k'}^2 + \frac{1}{d^2 q^2} \sum_{k=1}^d q y_k^4 \mathbb{E}[U_{ik}^4] \\ &\leq \frac{3}{d^2} + \frac{3\|y\|_\infty^2}{d^2 q} \leq \frac{6}{d^2}.\end{aligned}$$

Hence,

$$\mathbb{P}(\|Ay\|^2 \leq 1 - \epsilon) \leq \exp\left(d \ln\left(1 - \frac{\ell}{d} + 3\frac{\ell^2}{d^2}\right) + \ell(1 - \epsilon)\right) \leq \exp\left(3\frac{\ell^2}{d} - \ell\epsilon\right).$$

Choosing $\ell = \epsilon/6$, we obtain that

$$\mathbb{P}(\|Ay\|^2 \leq 1 - \epsilon) \leq \exp\left(-\frac{d\epsilon^2}{12}\right).$$

If $d \geq d_0 \geq 24 \log(3n/\delta)$, then we obtain

$$\mathbb{P}(\|Ay\|^2 \leq 1 - \epsilon) \leq \delta/(3n^2).$$

Hence, from a union bound over the at most n^2 possible pairs x_i, x_j , we obtain that

$$\mathbb{P}\left(\bigcup_{1 \leq i < j \leq n} \left\{\|A(x_i - x_j)\|^2 \leq (1 - \epsilon)\|x_i - x_j\|^2\right\}\right) \leq \delta/3. \quad (18)$$

We conclude from the upper bound (17) and the lower bound (18) that if $q \geq \max_{i \neq j} \frac{\|x_i - x_j\|_\infty^2}{\|x_i - x_j\|_2^2}$ and if $d \geq d_0(n, \delta, \epsilon)$, the ϵ -quasi isometry property (1) holds with probability at least $1 - \delta$, that is

$$\mathbb{P}\left(\bigcup_{1 \leq i < j \leq n} \left\{\|A(x_i - x_j)\|^2 \notin [(1 - \epsilon)\|x_i - x_j\|^2, (1 + \epsilon)\|x_i - x_j\|^2]\right\}\right) \leq 2\delta/3 + \delta/3 \leq \delta.$$

□

6. OPTIMALITY OF CONDITION (12) AND FILTERING EFFECT

It turns out that the condition $q \gtrsim 1/s$ is optimal in some sense if we impose the dimension d to be of order $1/\epsilon^2$ up to a polylogarithm. Experimentally, Figures 2 and 3 dually confirm that the random projection quality remains roughly the same as long as the proportion of non-zero coefficients is clearly above the minimum between $1/s$ and $1/3$. Indeed, a consequence of Theorem 2 is that if $d \asymp 1/\epsilon^2$ up to a polylog and if $q \ll 1/s$, then $\|Ay\|^2 \notin [1 - \epsilon, 1 + \epsilon]$ with positive probability, for any $(1, 1, s)$ -full vector y – see (13) for the definition of (κ, κ', s) full vectors. A first idea toward the proof of this optimality result is the following. Let y be $(1, 1, s)$ -full vector, or, equivalently, $\|y\| = 1$ and $y_i \in \{-1/\sqrt{s}, 1/\sqrt{s}, 0\}$. If $A \in \mathbb{R}^{d \times p}$ is any random matrix whose coefficients are independent and such that for all (i, k) , $\mathbb{P}(A_{ik} \neq 0) \leq q$, then,

$$\mathbb{P}(Ay = 0) \geq \mathbb{P}(\forall (i, k) \in d \times S, A_{ik} = 0) = (1 - q)^{ds} = e^{ds \ln(1-q)} \geq e^{-\frac{dsq}{1-q}}.$$

Hence, if $q \leq 1/(2ds)$, then $\mathbb{P}(Ay = 0) > 1/e$. In other words, there is no hope to satisfy the quasi-isometry property (1) with high probability if $q \leq 1/(2ds)$. This argument miss however the regim where $\epsilon^2/s \lesssim q \lesssim 1/s$ if $d \asymp 1/\epsilon^2$. The following theorem provides a general optimality result for all $q < 1/(240s)$, and hence fills the gap between ϵ^2/s and $1/s$ when $d \asymp 1/\epsilon^2$ up to a polylogarithmic factor.




FIGURE 2. Admissible value ϵ in function of the sparsity parameter q of the random projection entries (logarithmic scale), for different values of the sparsity s of the data. Each data point $x_i \in \mathbb{R}^{10000}$ has exactly s non-zero components, which are independent Gaussian entries. The coefficients of the projection matrix A are independent and have distribution $\frac{q}{2}\delta_{-\frac{1}{\sqrt{dq}}} + (1 - q)\delta_0 + \frac{q}{2}\delta_{\frac{1}{\sqrt{dq}}}$. The target dimension is $d = 500$. The blue line shows $\min_{i,j} \frac{\|A(x_i - x_j)\|^2}{\|x_i - x_j\|^2}$, while the red line shows $\max_{i,j} \frac{\|A(x_i - x_j)\|^2}{\|x_i - x_j\|^2}$. Observe that the scales of the ordinates are different between the plots.

It can be observed that quasi-isometry is ensured whenever $q \times s$ is sufficiently large.

Theorem 2. Assume that $A \in \mathbb{R}^{d \times p}$ has iid coefficients distributed according to distribution P with parameter q – see (6). Let $y \in \mathbb{R}^p$ be a unit vector with coordinates in $\{-1/\sqrt{s}, 1/\sqrt{s}, 0\}$. If $dq\epsilon^2 \leq 1/2$, $qs < 1/240$, then

$$P(\|Ay\|^2 \in [1 - \epsilon, 1 + \epsilon]) \leq 1 - e^{-5000}.$$

In other words, if $d \asymp 1/\epsilon^2$ up to a polylog, then Theorem 2 only requires that $q \lesssim 1/s$ up to a polylog. We take a probability $1 - e^{-5000}$ that is very close to 1 is the theorem to match the two regimes where $dqs \gtrsim 1$ and $dqs \lesssim 1$. In the proof of Theorem 2, we also show that in the sub-case where $dqs \geq 1/2048$, the probability of success $P(\|Ay\|^2 \in [1 - \epsilon, 1 + \epsilon])$ is smaller than $1/2$.

Proof of Theorem 2. Let y be a unit vector of \mathbb{R}^p . If $dqs \leq 1/2048$, then we have that

$$P(\|Ay\|^2 \notin [1 - \epsilon, 1 + \epsilon]) \geq \mathbb{P}(Ay = 0) \geq e^{-ds \frac{q}{1-q}} \geq e^{-5000},$$

which proves the result in that case.




FIGURE 3. Admissible value ϵ , in function on the sparsity parameter s of the data (logarithmic scale), for different values of the sparsity q of the projection matrix. Each data point of the $n = 100$ data point $x_i \in \mathbb{R}^{10000}$ has independent coefficients that are non-zero with probability s/p ; the non-zero coefficients are independent and uniformly distributed on $\{-1, +1\}$. The projection matrix A has independent entries with distribution $\frac{q}{2}\delta_{-\frac{1}{\sqrt{dq}}} + (1-q)\delta_0 + \frac{q}{2}\delta_{\frac{1}{\sqrt{dq}}}$. The target dimension is $d = 500$. The blue line shows $\min_{i,j} \frac{\|A(x_i - x_j)\|^2}{\|x_i - x_j\|^2}$, while the red line shows $\max_{i,j} \frac{\|A(x_i - x_j)\|^2}{\|x_i - x_j\|^2}$.

We observe that the values $q \geq 1/3$ ensure the quasi-isometry property whatever the data. For smaller values of q , the number s of non-zero coefficients needs to be larger than $1/q$.

In what follows, we assume that $dqs \geq 1/2048$. Chebychev's inequality implies that

$$\begin{aligned} \mathbb{P}(\|Ay\|^2 \in [1 - \epsilon, 1 + \epsilon]) &= \mathbb{P}((\|Ay\|^2 - 1)^2 \leq \epsilon^2) \\ &\leq \frac{\text{Var}[(\|Ay\|^2 - 1)^2]}{\mathbb{E}[(\|Ay\|^2 - 1)^2] - \epsilon^2}. \end{aligned}$$

Subsequently, we give a lower bound of $\mathbb{E}[(\|Ay\|^2 - 1)^2]$ and an upper bound of $\text{Var}(\|Ay\|^2 - 1)^2$. For simplicity, let us write X for a random variable following the distribution of one of the coefficients of A . X can be written $\frac{1}{\sqrt{dq}}\zeta \cdot U$ where (ζ, U) are independent random variables respectively distributed according to a Bernoulli variable of parameter q , and to a Rademacher variable. This implies in particular that for any $k \geq 1$, $\mathbb{E}[X^{2k+1}] = 0$ and $\mathbb{E}[X^{2k}] = \frac{1}{d^k q^{k-1}}$.

Lower bound of $\mathbb{E}[(\|Ay\|^2 - 1)^2]$.

$$\begin{aligned}
\mathbb{E} [(\|Ay\|^2 - 1)^2] &= \mathbb{E} [\|Ay\|^4] - 1 \\
&= \mathbb{E} \left[\left(\sum_{i=1}^d \sum_{k=1}^p \sum_{l=1}^p A_{ik} A_{il} y_k y_l \right)^2 \right] - 1 \\
&= \sum_{i_1, i_2, k_1, k_2, l_1, l_2} \mathbb{E} \left[\prod_{u \in \{1, 2\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] - 1,
\end{aligned}$$

where the final sum is over all $(i_1, i_2) \in [d]^2$ and all $(k_1, k_2, l_1, l_2) \in [p]^4$. Let us fix i_1, i_2 such that $i_1 = i_2$. Since $\mathbb{E}[A_{ik}] = \mathbb{E}[A_{ik}^3] = 0$ for any i, k , either $k_1 = k_2 = l_1 = l_2$ or there is exactly two pairs of equal indices among (k_1, k_2, l_1, l_2) . Since there are exactly 3 possible ways of matching 2 pairs among the four indices, we have that

$$\sum_{k_1, k_2, l_1, l_2} \mathbb{E} \left[\prod_{u \in \{1, 2\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] - 1 = \mathbb{E}[X^4] \|y\|_4^4 + 3\mathbb{E}[X^2]^2 (\|y\|_2^4 - \|y\|_4^4) = \frac{1}{d^2 q s} + \frac{3}{d^2} \left(1 - \frac{1}{s}\right).$$

If $i_1 \neq i_2$, then we necessarily have that $k_1 = l_1$ and $k_2 = l_2$ for nonzero contributions. Hence, in that case,

$$\sum_{k_1, k_2, l_1, l_2} \mathbb{E} \left[\prod_{u \in \{1, 2\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] - 1 = \mathbb{E}[X^2]^2 \|y\|_2^4 = \frac{1}{d^2}.$$

Combining the two cases, we obtain that

$$\mathbb{E} [(\|Ay\|^2 - 1)^2] = \frac{d}{d^2 q s} + \frac{3d}{d^2} \left(1 - \frac{1}{s}\right) + \frac{d(d-1)}{d^2} - 1 \geq \frac{1}{d q s}. \quad (19)$$

Upper bound of $\text{Var}(\|Ay\|^2 - 1)^2$.

$$\begin{aligned}
\text{Var} [(\|Ay\|^2 - 1)^2] &\leq \mathbb{E} [(\|Ay\|^2 - 1)^4] = \mathbb{E}[\|Ay\|^8] - 4\mathbb{E}[\|Ay\|^6] + 6\mathbb{E}[\|Ay\|^4] - 4\mathbb{E}[\|Ay\|^2] + 1 \\
&\leq \mathbb{E}[\|Ay\|^8] - 4\mathbb{E}[\|Ay\|^6] + \frac{6}{d q s} + 3 + \frac{18}{d}.
\end{aligned}$$

The inequality comes from the above computation of $\mathbb{E}[\|Ay\|^4]$. In what follows, we first upperbound $\mathbb{E}[\|Ay\|^8]$ and then we lowerbound $\mathbb{E}[\|Ay\|^6]$. The idea of lowerbounding $\mathbb{E}[\|Ay\|^6]$ is to cancel out the terms of constant order or of order $1/(dqs)$.

From the same computation as for $\mathbb{E}[\|Ay\|^4]$, we have that

$$\mathbb{E}[\|Ay\|^8] = \sum_{(i_u), (k_u), (l_u)} \mathbb{E} \left[\prod_{u \in \{1, 2, 3, 4\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right], \quad (20)$$

where the sum is over all $((i_u)_{u=1, \dots, 4}, (k_u)_{u=1, \dots, 4}, (l_u)_{u=1, \dots, 4}) \in [d]^4 \times [p]^8$. Let us consider the following sets for the indices (i_u) :

- (1) $(i_u) \in I_1$ if the i_u 's are pairwise distinct. in that case, $|I_1| = d(d-1)(d-2)(d-3) \leq d^4$
- (2) $(i_u) \in I_2$ if there are exactly two equal indices among the i_u 's. In other words, (i_u) is a permutation of (i, i, i', i'') where i, i', i'' are pairwise distinct. Here, $|I_2| = 6d(d-1)(d-2) \leq 6d^3$

- (3) $(i_u) \in I_3$ if there are exactly three equal indices among the i_u 's, i.e (i_u) is a permutation of (i, i, i, i') where $i \neq i'$. Here, $|I_3| = 4d(d-1) \leq 4d^2$
- (4) $(i_u) \in I_4$ if all the i_u 's are equal. Here, $|I_4| = d$
- (5) $(i_u) \in I_5$ if there are exactly two pairs of equal indices among the i_u 's. Here, $|I_5| = 3d(d-1) \leq 3d^2$

The sets (I_v) are disjoint, and the reader can check that the sum of their sizes is equal to d^4 . Let us fix $(i_u) \in [d]^4$, and consider the five following cases, each corresponding to one of the sets (I_v) .

- (1) If $(i_u) \in I_1$, then the expectation of the product over $u \in \{1, 2, 3, 4\}$ is nonzero only if $k_u = l_u$ for all $u \in \{1, 2, 3, 4\}$. Hence,

$$\sum_{(k_u), (l_u)} \mathbb{E} \left[\prod_{u \in \{1, 2, 3, 4\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] = \frac{1}{d^4} \|y\|_2^8 = \frac{1}{d^4}.$$

- (2) If $(i_u) \in I_2$, we assume that without loss of generality that (i_1, i_2, i_3) are pairwise distinct and that $i_3 = i_4$. In that case, we have a nonzero contribution only if $k_1 = l_1$, $k_2 = l_2$ and if either $(k_3 = l_3 = k_4 = l_4)$ or there are two matching pairs among the indices (k_3, l_3, k_4, l_4) (3 possible matching). Hence, using the fact that $\|y\|_2 = 1$ and $\|y\|_4^4 = 1/s$:

$$\sum_{(k_u), (l_u)} \mathbb{E} \left[\prod_{u \in \{1, 2, 3, 4\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] \leq \frac{1}{d^4 q} \|y\|_2^4 \|y\|_4^4 + \frac{3}{d^4} \|y\|_2^8 = \frac{1}{d^4 q s} + \frac{3}{d^4}.$$

- (3) If $(i_u) \in I_3$, we assume that i_1, i_2 are distinct and that $i_2 = i_3 = i_4$. In that case, we have a nonzero contribution if $k_1 = l_1$ and if either $(k_2 = l_2 = k_3 = l_3 = k_4 = l_4)$ or if there are 3 matching pairs among $(k_1, l_2, k_3, l_3, k_4, l_4)$ ($5 \cdot 3 = 15$ possible matchings). Hence, using also that $q s \leq 1$,

$$\sum_{(k_u), (l_u)} \mathbb{E} \left[\prod_{u \in \{1, 2, 3, 4\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] \leq \frac{1}{d^4 q^2} \|y\|_2^2 \|y\|_6^6 + \frac{15}{d^4} \|y\|_2^8 \leq \frac{16}{d^4 q^2 s^2}.$$

- (4) If $(i_u) \in I_4$, then there is a nonzero contribution in one of the three following cases. Either the k_u 's and l_u 's are all equal, or there are 2 groups among the k_u 's and l_u 's, each made of 4 indices that are all equal ($\frac{1}{2} \binom{8}{4} = 35$ possibilities), or there are 4 matching pairs ($7 \cdot 5 \cdot 3 = 105$ possible matching). Hence,

$$\sum_{(k_u), (l_u)} \mathbb{E} \left[\prod_{u \in \{1, 2, 3, 4\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] \leq \frac{1}{d^4 q^3} \|y\|_8^8 + \frac{35}{d^4 q^2} \|y\|_4^8 + \frac{105}{d^4} \|y\|_2^8 \leq \frac{141}{d^4 q^3 s^3}.$$

- (5) If $(i_u) \in I_5$, assume without loss of generality that $i_1 = i_2$, $i_3 = i_4$ and $i_2 \neq i_3$. Then there are two possibilities for each pairs (i_1, i_2) and (i_3, i_4) . Either $k_1 = l_1 = k_2 = l_2$ (resp. $k_3 = l_3 = k_4 = l_4$) or there are three pairs of equal indices among k_1, l_1, k_2, l_2 (resp. k_3, l_3, k_4, l_4). This gives

$$\sum_{(k_u), (l_u)} \mathbb{E} \left[\prod_{u \in \{1, 2, 3, 4\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] \leq \left(\frac{1}{d^2 q} \|y\|_4^4 + \frac{3}{d^2} \|y\|_2^4 \right)^2 \leq \frac{16}{d^4 q^2 s^2}.$$

Decomposing the equation (20) into these five above cases and using the assumption $d q s \geq 1$, we obtain that

$$\mathbb{E}[\|Ay\|^8] \leq 1 + \frac{6}{d q s} + \frac{18}{d} + \frac{4 * 16}{d^2 q^2 s^2} + \frac{141}{d^3 q^3 s^3} + \frac{3 * 16}{d^2 q^2 s^2} \leq 1 + \frac{6}{d q s} + \frac{18}{d} + \frac{253}{d^2 q^2 s^2},$$

which implies that

$$\begin{aligned} \mathbb{V}\text{ar} [(\|Ay\|^2 - 1)^2] &\leq \mathbb{E}[\|Ay\|^8] - 4\mathbb{E}[\|Ay\|^6] + \frac{6}{dqs} + 3 + \frac{12}{d} \\ &\leq 4 + \frac{12}{dqs} - 4\mathbb{E}[\|Ay\|^6] + \frac{253}{d^2q^2s^2} + \frac{30}{d}. \end{aligned}$$

We now show that the term $4 + \frac{12}{dqs}$ is smaller than $4\mathbb{E}[\|Ay\|^6]$. Doing the same reasoning as above, we can write

$$\mathbb{E}[\|Ay\|^6] = \sum_{(j_u), (k_u), (l_u)} \mathbb{E} \left[\prod_{u \in \{1,2,3\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right], \quad (21)$$

where the sum is over all $(j_u), (k_u), (l_u)$ in $[d]^3 \times [p]^6$. The product is always non-negative, and we consider the sets

$$J_1 = \{(j, j, j) : j \in [d]\} \text{ and } J_2 = \{(j_1, j_2, j_3) : \text{two of the } j_u \text{ are equal and distinct from the other one}\}.$$

We have that $|J_1| = d^3$ and $|J_2| = 3d(d-1)$, so that

$$\begin{aligned} \mathbb{E}[\|Ay\|^6] &\geq \sum_{(j_u) \in J_1, (k_u), (l_u)} \mathbb{E} \left[\prod_{u \in \{1,2,3\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] + \sum_{(j_u) \in J_2, (k_u), (l_u)} \mathbb{E} \left[\prod_{u \in \{1,2,3\}} A_{i_u k_u} A_{i_u l_u} y_{k_u} y_{l_u} \right] \\ &= \|y\|_2^6 + 3d(d-1) \frac{1}{d^3q} \|y\|_4^4 \|y\|_2^2 \\ &= 1 + \frac{3}{dqs} - \frac{3}{d^2qs} \geq 1 + \frac{3}{dqs} - \frac{3}{d^2q^2s^2} \end{aligned}$$

To conclude, we obtain

$$\begin{aligned} \mathbb{V}\text{ar} [(\|Ay\|^2 - 1)^2] &\leq 4 + \frac{12}{dqs} - 4\mathbb{E}[\|Ay\|^6] + \frac{253}{d^2q^2s^2} + \frac{30}{d} \\ &\leq \frac{256}{d^2q^2s^2} + \frac{30}{d}. \end{aligned}$$

Combining this latter upper bound with (19), we conclude that

$$P(\|Ay\|^2 \in [1 - \epsilon, 1 + \epsilon]) \leq \frac{\frac{256}{d^2q^2s^2} + \frac{30}{d}}{\frac{1}{dqs} - \epsilon^2} \leq \frac{\frac{256}{dqs} + 30qs}{1 - dqs\epsilon^2} \leq 1/2,$$

where we used in the last inequality the assumption that $dqs\epsilon^2 \leq 1/2$, $qs \leq 1/240$ and $dqs \geq 2048$. This concludes the proof of Theorem 2. \square

6.1. Concentration of non-negative quadratic forms

The upper bound given in section 5 is in fact strongly connected to the Hanson-Wright inequality for sub-Gaussian random variables – see e.g. [13], and [12] for an application to the Johnson-Lindenstrauss lemma. While this inequality is known with precise constants for Gaussian chaos of order 2 – see Example 2.12 of [4], the constants are unclear in the literature when the random variables are only assumed to be sub-Gaussians. We conclude this paper by giving a precise statement of the Hanson Wright inequality for sub-Gaussian vectors and when the quadratic form is assumed to be non-negative.

We say that a vector $X \in \mathbb{R}^n$ is sub-Gaussian if for any $u \in \mathbb{R}^d$,

$$\mathbb{E}[e^{u^T X}] \leq e^{\frac{1}{2}\|u\|_2^2}. \quad (22)$$

In particular, if Z_1, \dots, Z_d are independent real random variable and 1-sub-Gaussian, that is $\mathbb{E}[e^{\lambda Z_i}] \leq e^{\lambda^2/2}$, then for any orthogonal matrix P , PZ is a sub-Gaussian vector. In contrast to [13], we do not require in the following Theorem the coordinates of X to be independent.

Theorem 3. *Let S be any $d \times d$ symmetric matrix with non-negative eigenvalues, and X be a sub-Gaussian vector in the sense of (22). Then, for any $\ell \in [0, 1/(2\|S\|_{op})]$,*

$$\mathbb{E}_X[e^{\ell X^T S X}] \leq \exp(\ell \text{Tr}(S) + \frac{\ell^2 \|S\|_F^2}{1-2\ell\|S\|_{op}}).$$

As a consequence of Theorem 3 and following the same computations as in Theorem 10 of [4], it holds that with probability at least $1 - \delta$,

$$X^T S X \leq \sqrt{4\|S\|_F^2 \log(1/\delta)} + 2\|S\|_{op} \log(1/\delta),$$

for any $\delta \in (0, 1)$.

Proof of Theorem 3. Let us write $S = P \text{Diag}(\mu_1, \dots, \mu_n) P^T$, where $\mu_1 \geq \dots \geq \mu_n \geq 0$ and P is an orthogonal matrix. Let also Y be the sub-Gaussian vector equal to PX , and $\ell > 0$. From a Fubini argument,

$$\mathbb{E}_X[e^{\ell X^T S X}] = \mathbb{E}_X\left[\sum_{i=1}^n \ell \mu_i Y_i^2\right] = \mathbb{E}_G[\mathbb{E}_X[e^{\sum_{i=1}^n \sqrt{2\ell\mu_i} Y_i G_i}]] \leq \mathbb{E}_G[e^{\sum_{i=1}^n \frac{1}{2} \ell \mu_i G_i^2}],$$

where G_1, \dots, G_n stand for independent standard and centered Gaussian random variables. Then,

$$\mathbb{E}_G[e^{\sum_{i=1}^n \frac{1}{2} \ell \mu_i G_i^2}] = \prod_{i=1}^n \frac{1}{\sqrt{1-2\ell\mu_i}} = e^{\ell \text{Tr}(S)} \prod_{i=1}^n \exp(-\frac{1}{2} \log(1-2\ell\mu_i) - \ell\mu_i) \leq \exp(\ell \text{Tr}(S) + \sum_{i=1}^n \frac{\ell^2 \mu_i^2}{1-2s\mu_i}),$$

where the first inequality comes from the inequality $-\frac{1}{2} \log(1-2\ell\mu_i) - \ell\mu_i = \int_0^\ell \frac{2s\mu_i}{1-2s\mu_i} ds \leq \frac{\ell^2 \mu_i^2}{1-2\ell\mu_i}$. We conclude the proof by remarking that $\sum_{i=1}^n \frac{\ell^2 \mu_i^2}{1-2s\mu_i} \leq \frac{\ell^2 \|S\|_F^2}{1-2s\|S\|_{op}}$. \square

REFERENCES

- [1] Dimitris Achlioptas. Database-friendly random projections. In *Proceedings of the Twentieth ACM SIGMOD-SIGACT-SIGART Symposium on Principles of Database Systems*, PODS '01, page 274–281, New York, NY, USA, 2001. Association for Computing Machinery.
- [2] Noga Alon and Joel H. Spencer. *The Probabilistic Method*. Wiley, New York, second edition, 2004.
- [3] Ella Bingham and Heikki Mannila. Random projection in dimensionality reduction: applications to image and text data. In *Proceedings of the Seventh ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, KDD '01, page 245–250, New York, NY, USA, 2001. Association for Computing Machinery.
- [4] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities*. Oxford University Press, Oxford, 2013. A nonasymptotic theory of independence, With a foreword by Michel Ledoux.
- [5] Moses Charikar, Kevin Chen, and Martin Farach-Colton. Finding frequent items in data streams. *Theoretical Computer Science*, 312(1):3–15, 2004. Automata, Languages and Programming.
- [6] Kenneth L. Clarkson and David P. Woodruff. Numerical linear algebra in the streaming model. In *Proceedings of the Forty-First Annual ACM Symposium on Theory of Computing*, STOC '09, page 205–214, New York, NY, USA, 2009. Association for Computing Machinery.

- [7] Mikael Møller Høgsgaard, Lior Kamma, Kasper Green Larsen, Jelani Nelson, and Chris Schwiegelshohn. Sparse dimensionality reduction revisited. In Ruslan Salakhutdinov, Zico Kolter, Katherine Heller, Adrian Weller, Nuria Oliver, Jonathan Scarlett, and Felix Berkenkamp, editors, *Proceedings of the 41st International Conference on Machine Learning*, volume 235 of *Proceedings of Machine Learning Research*, pages 18454–18469. PMLR, 21–27 Jul 2024.
- [8] P. Indyk. Algorithmic applications of low-distortion geometric embeddings. In *Proceedings 42nd IEEE Symposium on Foundations of Computer Science*, pages 10–33, 2001.
- [9] William Johnson and Joram Lindenstrauss. Extensions of lipschitz maps into a hilbert space. *Contemporary Mathematics*, 26:189–206, 01 1984.
- [10] Daniel M. Kane and Jelani Nelson. Sparser Johnson-Lindenstrauss Transforms. *J. ACM*, 61(1), jan 2014.
- [11] Ping Li, Trevor J. Hastie, and Kenneth W. Church. Very sparse random projections. In *Proceedings of the 12th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, KDD '06, page 287–296, New York, NY, USA, 2006. Association for Computing Machinery.
- [12] Yingru Li. Simple, unified analysis of Johnson-Lindenstrauss with applications. 2024.
- [13] Mark Rudelson and Roman Vershynin. Hanson-wright inequality and sub-gaussian concentration. 2013.
- [14] Tamas Sarlos. Improved approximation algorithms for large matrices via random projections. In *2006 47th Annual IEEE Symposium on Foundations of Computer Science (FOCS'06)*, pages 143–152, 2006.
- [15] Santosh Vempala. The random projection method. 01 2004.
- [16] Martin J. Wainwright. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2019.