



HAL
open science

Mask side-effects are related to gender in long-term CPAP: results from the InterfaceVent real-life study

Celia Vidal, Fanny Bertelli, Jean-Pierre Mallet, Raphael Gilson,
Jean-Christian Borel, Frédéric Gagnadoux, Arnaud Bourdin, Nicolas Molinari,
Dany Jaffuel

► To cite this version:

Celia Vidal, Fanny Bertelli, Jean-Pierre Mallet, Raphael Gilson, Jean-Christian Borel, et al.. Mask side-effects are related to gender in long-term CPAP: results from the InterfaceVent real-life study. *Respiratory Research*, 2024, 25 (1), pp.331. 10.1186/s12931-024-02965-1 . hal-04694849

HAL Id: hal-04694849

<https://hal.science/hal-04694849v1>

Submitted on 12 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Mask side-effects are related to gender in long-term CPAP: results from the InterfaceVent real-life study

Celia Vidal ^{1 2}, Fanny Bertelli ², Jean-Pierre Mallet ^{2 3 4}, Raphael Gilson ², Jean-Christian Borel ⁵, Frédéric Gagnadoux ⁶, Arnaud Bourdin ^{2 3 4}, Nicolas Molinari ^{1 2}, Dany Jaffuel ^{7 8 9}

¹ IDESP, INSERM, PreMEDical INRIA, CHU Montpellier, Montpellier University, Montpellier, France.

² Groupe Adène, Montpellier, France.

³ Department of Respiratory Diseases, Arnaud de Villeneuve Hospital, CHRU Montpellier, Montpellier University Hospital, 371, Avenue Doyen Giraud, 34295, Montpellier Cedex 5, France.

⁴ PhyMedExp (INSERM U 1046, CNRS UMR9214), Montpellier University, Montpellier, France.

⁵ Université Grenoble-Alpes, CHUGA Et INSERM U1300, Grenoble, France.

⁶ Departement of Respiratory and Sleep Medicine, University Hospital of Angers, Angers, France.

⁷ Groupe Adène, Montpellier, France. dany.jaffuel@wanadoo.fr.

⁸ Department of Respiratory Diseases, Arnaud de Villeneuve Hospital, CHRU Montpellier, Montpellier University Hospital, 371, Avenue Doyen Giraud, 34295, Montpellier Cedex 5, France. dany.jaffuel@wanadoo.fr.

⁹ PhyMedExp (INSERM U 1046, CNRS UMR9214), Montpellier University, Montpellier, France. dany.jaffuel@wanadoo.fr.

Keywords:

Leaks; Side-effects; Sleep apnea; Women.

Abstract

Background:

Over the past three decades, our understanding of sleep apnea in women has advanced, revealing disparities in pathophysiology, diagnosis, and treatment compared to men. However, no real-life study to date has explored the relationship between mask-related side effects (MRSEs) and gender in the context of long-term CPAP.

Methods:

The InterfaceVent-CPAP study is a prospective real-life cross-sectional study conducted in an apneic adult cohort undergoing at least 3 months of CPAP with unrestricted mask-access (34 different masks, no gender specific mask series). MRSE were assessed by the patient using visual analog scales (VAS). CPAP-non-adherence was defined as a mean CPAP-usage of less than 4 h per day. The primary objective of this ancillary study was to investigate the impact of gender on the prevalence of MRSEs reported by the patient. Secondary analyses assessed the impact of MRSEs on CPAP-usage and CPAP-non-adherence depending on the gender.

Results:

A total of 1484 patients treated for a median duration of 4.4 years (IQ₂₅₋₇₅: 2.0-9.7) were included in the cohort, with women accounting for 27.8%. The prevalence of patient-reported mask injury, defined as a VAS score ≥ 5 ($p = 0.021$), was higher in women than in men (9.6% versus 5.3%). For nasal pillow masks, the median MRSE VAS score for dry mouth was higher in women ($p = 0.039$). For oronasal masks, the median MRSE VAS score for runny nose was higher in men ($p = 0.039$). Multivariable regression analyses revealed that, for both women and men, dry mouth was independently and negatively associated with CPAP-usage, and positively associated with CPAP-non-adherence.

Conclusion:

In real-life patients treated with long-term CPAP, there are gender differences in patient reported MRSEs. In the context of personalized medicine, these results suggest that the design of future masks should consider these gender differences if masks specifically for women are developed. However, only dry mouth, a side effect not related to mask design, impacts CPAP-usage and non-adherence.

Trial registration:

INTERFACEVENT IS REGISTERED WITH CLINICALTRIALS.GOV
(NCT03013283).FIRST REGISTRATION DATE IS 2016-12-23.

Background

The prevalence of sleep apnea syndrome (SAS) in adults over the age of 35 ranges from 5.9% to 79.2%, depending on the clinical symptoms and apnea/hypopnea scoring criteria used [1, 2]. In 2024, Continuous Positive Airway Pressure (CPAP) remains the cornerstone of SAS treatment, despite major advances in alternative therapies [3–5]. CPAP-adherence is associated with improved quality of life (QoL) [6, 7], and reduced incident or recurrent major adverse cardiovascular events [8, 9].

Over the past three decades, our understanding of SAS in women has grown, highlighting the existence of disparities between women and men in pathophysiology, diagnosis and treatment [10–12]. The apnea–hypopnea index (AHI) is lower in women than in men, and QoL is worse in women [11, 13]. Some studies also suggest that CPAP-adherence is poorer in women [12].

Although our knowledge about the specificities of SAS in women is increasing, real gaps in the research persist, as highlighted by a recent editorial advocating for targeted research on gender disparities in SAS [14]. While several manufacturers have developed women’s and men’s versions of certain series of their masks, no external study has validated the concept of a gender-specific mask. To date, there is no long-term study reporting gender differences in MRSEs, nor studies reporting the impact of gender-related MRSEs on CPAP-adherence. Therefore, the primary objective of the present study was to investigate the impact of gender on the prevalence of patient-reported MRSEs. Secondary analyses assessed the impact of MRSEs on CPAP-usage and CPAP-nonadherence (defined as a mean CPAP-usage of less than 4 h per day) depending on gender.

Methods

Study design and study population

This study presents an ancillary analysis of the InterfaceVent study, which was exhaustively described in a prior publication [15]. Briefly, the InterfaceVent study (ClinicalTrials.gov: NCT03013283) was a prospective, real-life cross-sectional study conducted from February 7, 2017 to April 1, 2019 in adults undergoing at least 3 months of CPAP or non-invasive ventilation. We herein report results for SAS-patients treated exclusively by CPAP. SAS was defined according to the French Social Security (FSS) system criteria: (1) Apnea Hypopnea Index (AHI) $\geq 30/h$ (or AHI $\geq 15/h$ and more than 10/h respiratory-effort-related arousals), and (2) associated with sleepiness and at least three of the following symptoms: snoring, headaches, hypertension, reduced vigilance, libido disorders, nocturia. These criteria must be met in order for patients to be reimbursement by the FSS. The Apard ADENE group, a non-profit home care provider, provided care to patients following an initial prescription by one of the 336 device-prescribing physicians in the Occitanie region of France. Patient inclusion occurred during one of the routine home visits, conducted by one of the 32 Apard technicians, that are required to receive reimbursement for CPAP treatment by the FSS-single payer system. No CPAP-adherence threshold was required for reimbursement, and patients with poor compliance were not systematically excluded (for exclusion criteria, see [15]). Patients

had unlimited access to 34 masks (see [15]). No specific women's or men's versions of the mask series were available at the time of study.

Collected data

Side-effect visual analogue scales (VAS; see below), the Epworth-Sleepiness-Scale and the EQ-5D-3L questionnaire were administered by a technician employed by the home care provider during a scheduled visit, as previously described [15].

An 11-point VAS (0 = no reported side-effect to 10 = very uncomfortable side-effect) was used to assess the following MRSEs: dry mouth, partner disturbance due to leaks, patient-reported leaks, noisy mask, heavy mask, painful mask, mask injury, painful harness, harness injury, redness of the eyes, itchy eyes, dry nose, stuffy nose, and runny nose. Importantly, the technician did not help patients fill out the questionnaires and the VAS.

Statistical analyses

Continuous data were expressed as medians with their associated quartile ranges due to non-Gaussian distributions. Qualitative parameters were expressed as numbers and percentages. Gender effect was evaluated using Wilcoxon–Mann–Whitney test for quantitative data, and Chi-square or Fisher tests for qualitative effects. Side-effects and mask model, according to gender and mask type, were compared using Chi-square or Fisher tests. For these last comparisons, corrected p-value with False Discovery Rate correction were performed. To visualize correlations between MRSEs for a given gender and mask type, a principal component analysis was performed for women and men on nasal (NM), oronasal (ONM) and nasal pillow masks (NPM). Univariate and Multivariable logistic and linear regression analyses were used to study, by gender, associations between CPAP-usage and CPAP-non-adherence (defined as a mean CPAP usage of less than 4 h per day) versus explanatory variables (demographic data, Epworth-Sleepiness-Scale (ESS) score, EQ-5D-3L-questionnaires, device/mask data and MRSEs). All statistical analyses were performed with R (V.4.3.1).

Results

Population baseline characteristics according to mask type and gender are summarized in Table 1. A total of 1484 patients (27.8% women) were included in the analysis. Patients received significantly different mask types according to gender ($p = 0.002$). Specifically, 58.6% of women received NM versus 52.5% of men, 19.6% of women received NPM versus 16.4% of men, and 21.8% of women received oronasal masks versus 31.1% of men. The median BMI was higher in women than in men (32.5 kg/m² vs. 30.5 kg/m², $p < 0.001$), and more than half of patients were obese (62.2% for women vs. 53.9% for men, $p = 0.007$). Initial AHI ≥ 30 /h occurred less often in women (81.2% vs. 88.1%, $p = 0.001$). Women were more likely to live alone compared to men (46.2% vs. 20.7%, $p < 0.001$), and experienced anxiety/depression more frequently (51.5% vs. 34.8%, $p < 0.001$). Furthermore, median

EQ-5D-3L health VAS (0–100 score) was lower in women than in men (60.6 vs. 70.2, $p < 0.001$). Active workers were 19.8% and 20.9% for women and men respectively ($p = 0.628$), active smokers were 11.9% and 11.8% for women and men respectively ($p = 0.952$). Median global leaks and global large leaks were lower in women than in men for NM ($p < 0.001$ and $p = 0.039$). Mean pressure was lower in women for NM ($p = 0.004$). The median CPAP-usage was lower in women than in men for both NM and NPM ($p = 0.005$ and $p < 0.001$). Women's lower CPAP-usage is also reflected in their higher level of nonadherence (10.9% vs. 6.5%, $p = 0.005$).

CPAP-usage and non-adherence according to mask type and gender are depicted in Fig. 1. For women, median CPAP-usage was higher in NM than in NPM (6.5 h/day vs. 6.2 h/day, $p = 0.036$), reflecting greater nonadherence in NPM compared with NM (17.3% vs. 7.4%, $p = 0.031$). For men, median CPAP-usage was higher in NM than in ONM (7.0 h/day vs. 6.5 h/day, $p = 0.038$), reflecting greater non-adherence in ONM compared with NM (9.3% vs. 4.6%, $p = 0.017$).

Prevalence of mask related side-effects reported by the patient depending on gender and mask type

Gender differences in terms of specific VAS scores according to mask type are depicted in Fig. 2. When we compared genders, we found that median MRSE VAS scores for partner disturbing leaks were lower in women ($p < 0.001$) for NM, were lower for women for runny nose ($p = 0.039$) for ONM, and were higher in women for dry mouth ($p = 0.039$) for NPM.

MRSE frequencies (VAS score ≥ 1 and VAS score ≥ 5) according to gender were analyzed (see Additional file 1). Partner disturbing leaks were lower in women than in men ($p < 0.001$ for VAS score ≥ 1 and $p = 0.004$ for VAS score ≥ 5); the prevalence of mask injury was higher in women than in men (9.6% and 5.3%, respectively) for a VAS score ≥ 5 ($p = 0.021$).

Mask differences in terms of specific VAS scores according to gender were analyzed (see Additional file 2). When we compared mask types, we found that, for women, a median MRSE VAS score for runny nose was higher in NPM than in ONM ($p = 0.045$), and median MRSE VAS score for patient reported leaks was higher in ONM than in NM ($p = 0.012$). For men, median MRSE VAS scores for dry mouth, patient reported leaks, partner disturbing leaks, itchy eyes and red eyes were higher in ONM than in NM ($p < 0.001$, $p < 0.001$, $p = 0.042$, $p = 0.034$ and $p = 0.013$, respectively). Median MRSE VAS scores for dry mouth, itchy eyes and red eyes were higher in ONM than in NM ($p < 0.001$, $p < 0.001$ and $p < 0.001$, respectively). Median MRSE VAS score for itchy eyes was higher in NM than in NPM ($p = 0.025$).

Principal Component Analysis (PCA) suggested that there were comparable groups of side effects according to gender for NM and ONM (Additional file 3). For NPM, the groups were more dispersed, with different positioning of leak items by gender.

Mask series and gender

There was no difference in the distribution of mask series according to gender (see Additional file 4). However, there were significant differences in the distribution of mask series for each type of mask by gender (see Additional file 5). For NPM, there was a significant difference

in Swift Fx ® (51.1%) and Nuance pro ® (31.2%) for men but no significant difference for women. For both women and men, Mirage Fx ® and Swift Fx ® were the most popular mask series for NM and NPM, respectively. For ONM, Simplus ® was the most popular for women and Quattro ® for men.

Mask related side-effects, CPAP-usage and CPAP-non-adherence according to gender

Tables 2 and 3 summarize univariate and multivariable linear and logistic regression analyses evaluating the impact of explanatory variables on CPAP-usage and CPAP-non-adherence according to gender. For women, in the model explaining CPAP-usage, the latter was independently associated with higher BMI, higher mean pressure, non-active smokers, higher VAS score for partner disturbing leaks and lower VAS score for dry mouth. In the model explaining CPAP-non-adherence, the latter was independently associated with only lower VAS score for partner disturbing leaks.

Table 1 Population characteristics according to mask type and gender

	Whole population (N= 1484)		Nasal mask (N= 804)		Nasal pillow mask (N= 257)		Oronasal mask (N= 423)					
	Women, N= 413	Men, N= 1071	Women, N= 242	Men, N= 562	Women, N= 81	Men, N= 176	Women, N= 90	Men, N= 333				
			P value						P value			
Demographics												
Age (years)	67.0 [61.0; 74.0]	67.0 [60.0; 74.0]	0.818 ¹	67.0 [61.0; 74.0]	0.533 ¹	65.0 [58.0; 72.0]	66.0 [60.0; 73.3]	0.333 ¹	67.0 [59.3; 72.8]	68.0 [60.0; 74.0]	0.641 ¹	
BMI (kg/m ²)	32.5 [27.7; 36.9]	30.5 [27.5; 34.0]	<0.001 ¹	32.5 [27.9; 37.2]	30.2 [27.3; 33.8]	<0.001 ¹	30.3 [26.4; 36.0]	30.2 [27.3; 33.7]	0.850 ⁰	33.6 [28.7; 36.8]	31.0 [28.1; 34.5]	0.014 ¹
Diagnosis of AHI	36.0 [30.0; 49.0]	40.2 [32.0; 38.0]	<0.001 ¹	36.0 [30.0; 50.0]	40.4 [32.0; 57.2]	0.005 ¹	36.0 [30.0; 46.0]	40.0 [31.0; 57.5]	0.020 ⁰	36.0 [31.0; 49.0]	41.0 [31.0; 59.0]	0.097 ¹
Presence of partner (events/h)	218 (53.8)	832 (79.3)	<0.001 ²	124 (52.3)	441 (79.3)	<0.001 ²	45 (57.0)	140 (81.4)	<0.001 ²	49 (55.1)	251 (78.2)	<0.001 ²
Epworth scale												
ESS (0-24 score)	5.0 [3.0; 8.0]	5.0 [3.0; 9.0]	0.347 ¹	4.0 [2.0; 8.0]	5.0 [3.0; 8.8]	0.043 ¹	7.0 [3.0; 10.0]	5.0 [3.0; 9.0]	0.149 ¹	6.0 [3.0; 9.0]	6.0 [3.0; 9.0]	0.803 ¹
RES (%)	63 (15.3)	177 (16.5)	0.551 ¹	32 (13.2)	89 (15.8)	0.342 ²	14 (17.3)	32 (18.2)	0.862 ²	17 (18.9)	56 (16.8)	0.644 ²
Device												
CPAP-usage (h/day)	6.5 [5.0; 7.5]	6.8 [5.7; 7.8]	<0.001 ¹	6.5 [5.3; 7.8]	7.0 [5.8; 7.9]	0.005 ¹	6.2 [4.8; 7.0]	6.8 [5.8; 7.8]	<0.001 ¹	6.5 [5.0; 7.4]	6.5 [5.5; 7.8]	0.098 ¹
Non-adherence (%)	45 (10.9)	70 (6.5)	0.005 ²	18 (7.4)	26 (4.6)	0.108 ²	14 (17.3)	13 (7.4)	0.016 ²	13 (14.4)	31 (9.3)	0.157 ²
Current AHI ^{low} (events/h)	1.6 [0.8; 3.0]	2.1 [1.0; 4.2]	<0.001 ¹	1.4 [0.7; 2.8]	1.8 [1.0; 4.0]	<0.001 ¹	1.6 [0.9; 3.0]	1.6 [0.8; 3.0]	0.924 ¹	1.9 [1.1; 3.1]	2.9 [1.5; 5.5]	<0.001 ¹
Treatment duration (years)	3.6 [1.5; 8.4]	5.0 [2.3; 10.2]	<0.001 ¹	3.6 [1.4; 8.3]	5.1 [2.0; 10.9]	0.010 ¹	3.2 [1.4; 6.7]	6.5 [3.3; 10.6]	<0.001 ¹	4.0 [1.7; 9.2]	4.2 [2.3; 9.2]	0.626 ¹
Mean pressure (cmH ₂ O)	8.0 [6.4; 9.6]	8.3 [6.8; 10.0]	<0.001 ¹	7.6 [6.0; 9.5]	8.0 [6.6; 9.8]	0.004 ¹	7.3 [6.2; 8.8]	7.7 [6.5; 9.4]	0.208 ¹	9.1 [7.9; 10.6]	9.1 [7.6; 10.8]	0.926 ¹
90th/95th pressure (cmH ₂ O)	10.0 [8.0; 11.6]	10.0 [8.4; 11.8]	0.033 ¹	9.8 [7.9; 11.3]	10.0 [8.0; 11.5]	0.061 ¹	9.5 [7.9; 10.8]	9.6 [8.0; 11.0]	0.414 ¹	11.0 [9.6; 12.0]	10.9 [9.3; 12.0]	0.520 ¹
Fixed pressure (cmH ₂ O)	41 (10.0)	150 (14.0)	0.037 ²	24 (10.0)	80 (14.2)	0.098 ²	8 (9.9)	25 (14.2)	0.335 ²	9 (10.0)	45 (13.5)	0.375 ²
Comfort mode (%)	65 (15.7)	169 (15.8)	0.984 ²	38 (15.7)	92 (16.4)	0.814 ²	13 (16.1)	29 (16.5)	0.931 ²	14 (15.6)	48 (14.4)	0.786 ²
Heated humidifier (%)	255 (61.7)	616 (58.5)	0.247 ²	143 (59.1)	291 (51.8)	0.056 ²	57 (64.2)	107 (60.8)	0.602 ²	60 (66.7)	228 (68.5)	0.745 ²

Table 1 (continued)

	Whole population (N= 1484)			Nasal mask (N= 804)			Nasal pillow mask (N= 257)			Oronasal mask (N= 423)		
	Women, N= 413	Men, N= 1071	P value	Women, N= 242	Men, N= 562	p-value	Women, N= 81	Men, N= 176	p-value	Women, N= 90	Men, N= 333	p-value
Heated breathing tube (%)	27 (6.5)	32 (3.0)	0.002²	12 (5.0)	14 (2.5)	0.070 ²	8 (9.9)	5 (2.8)	0.028³	7 (7.8)	13 (3.9)	0.157 ³
Mask availability since 2013 (%)	170 (41.2)	440 (41.2)	0.979 ²	66 (27.3)	171 (30.5)	0.353 ²	42 (51.9)	61 (34.9)	0.010²	62 (68.9)	208 (62.7)	0.274 ²
Unintentional leaks (l/min)	2.4 [0.0; 7.2]	2.5 [0.0; 7.5]	0.601 ¹	2.5 [0.0; 8.0]	2.5 [0.0; 8.4]	0.681 ¹	1.2 [0.0; 4.5]	2.4 [0.0; 7.3]	0.271 ¹	0.0 [0.0; 3.6]	1.2 [0.0; 7.3]	0.258 ¹
Unintentional large leaks (%)	0.1 [0.0; 0.8]	0.1 [0.0; 0.9]	0.828 ¹	0.1 [0.0; 0.7]	0.0 [0.0; 0.6]	0.530 ¹	0.0 [0.0; 0.8]	0.1 [0.0; 0.6]	0.994 ¹	0.1 [0.0; 1.0]	0.3 [0.0; 3.8]	0.202 ¹
Global leaks (l/min)	28.0 [22.0; 34.0]	36.0 [30.0; 45.5]	<0.001¹	27.0 [21.0; 33.8]	33.0 [30.0; 43.0]	<0.001¹	31.0 [25.8; 33.5]	36.0 [28.0; 43.0]	0.261 ¹	28.0 [23.0; 39.0]	37.5 [33.1; 48.2]	0.171 ¹
Global large leaks (%)	0.4 [0.0; 2.6]	1.0 [0.2; 6.4]	0.015¹	0.3 [0.0; 2.2]	1.0 [0.1; 5.0]	0.039¹	0.8 [0.2; 4.7]	0.9 [0.4; 5.4]	0.769 ¹	1.6 [0.1; 2.7]	5.0 [0.2; 8.8]	0.530 ¹

Leaks were obtained using CPAP built-in software [15]. Data are reported as medians and quartiles or numbers and percentages of total as appropriate. ¹Wilcoxon-Mann-Whitney test; ²Chi-square test; ³Fisher exact test. Bolds variables are statistically significant at the 5% threshold

AH: Apnea-Hypopnea Index; AHI₁₀₋₃₀: AHI reported by device; BMI: Body Mass Index; CPAP: continuous positive airway pressure; ESS: epworth sleepiness scale; N= number of patients responding, Non-adherence: CPAP-usage under 4 h per day; RES: residual excessive sleepiness (ESS score > 10); WAS: visual analogue scale




Fig. 1 CPAP-usage (a) and CPAP-non-adherence (b) according to mask type and gender




Fig. 2 Visual analogue scales for mask related side-effect scores according to gender and mask type

For men, in the model explaining CPAP-usage, the latter was independently associated with higher age, BMI, treatment duration, p90/95th pressure, lower ESS score, availability of the mask before 2013, higher VAS score for harness injury, and lower VAS score for dry nose. In the model explaining CPAP-non-adherence, the latter was independently associated with lower age and BMI, living after 2013, and higher VAS score for dry mouth.

Table 2 Univariate analyses by gender

Variable	CPAP-usage (h/day)				CPAP-non-adherence (<4 h/day)							
	Women, N = 413		Men, N = 1071		Women, N = 413		Men, N = 1071					
	Estimate	95% CI	p-value	Estimate	95% CI	p-value	OR	95% CI	p-value			
Demographics												
Age (years)	0.01	- 0.01, 0.03	0.205	0.02	0.01, 0.03	<0.001	1.00	0.97, 1.03	0.953	0.97	0.95, 0.99	0.003
BMI (kg/m ²)	0.05	0.02, 0.08	<0.001	0.05	0.03, 0.07	<0.001	0.95	0.90, 1.00	0.075	0.95	0.90, 1.00	0.043
Diagnostic AHI (events/h)	0.01	0.00, 0.02	0.089	0.01	0.00, 0.02	0.001	1.00	0.98, 1.02	0.680	0.99	0.98, 1.01	0.264
Active smokers	-0.54	- 1.11, 0.02	0.059	0.14	- 0.19, 0.47	0.403	1.56	0.65, 3.74	0.320	1.63	0.85, 3.13	0.147
Mustache	NA	NA	NA	-0.05	- 0.30, 0.20	0.706	NA	NA	NA	1.00	0.57, 1.76	0.992
Beard	NA	NA	NA	-0.20	- 0.48, 0.08	0.161	NA	NA	NA	0.86	0.44, 1.68	0.661
Active workers	-0.53	- 0.98, - 0.07	0.025	-0.47	- 0.73, - 0.21	<0.001	1.46	0.70, 3.05	0.311	1.71	0.99, 2.96	0.055
Presence of partner	0.28	- 0.08, 0.63	0.128	0.11	- 0.15, 0.38	0.395	0.72	0.38, 1.36	0.310	0.45	0.27, 0.76	0.003
Epworth scale												
ESS (0-24 score)	-0.09	- 0.13, - 0.05	<0.001	-0.05	- 0.07, - 0.02	<0.001	1.09	1.02, 1.16	0.007	1.04	0.99, 1.10	0.144
RES	-0.84	- 1.32, - 0.35	<0.001	-0.27	- 0.56, 0.01	0.058	2.26	1.10, 4.66	0.027	1.29	0.70, 2.36	0.419
Device												
Current AHI _{low} (events/h)	0.04	- 0.04, 0.11	0.355	-0.02	- 0.04, 0.01	0.259	0.96	0.83, 1.12	0.621	1.07	1.02, 1.12	0.002
Treatment duration (years)	0.05	0.02, 0.09	0.002	0.06	0.04, 0.08	<0.001	0.96	0.89, 1.03	0.218	0.91	0.86, 0.96	<0.001
Mean pressure (cmH ₂ O)	0.15	0.07, 0.23	<0.001	0.10	0.05, 0.15	<0.001	0.95	0.82, 1.10	0.496	0.93	0.83, 1.05	0.231
90th/95th pressure (cmH ₂ O)	0.12	0.04, 0.20	0.003	0.08	0.03, 0.13	<0.001	0.96	0.84, 1.11	0.588	0.94	0.84, 1.05	0.298
Fixed pressure	0.56	- 0.03, 1.16	0.063	0.33	0.03, 0.64	0.031	0.62	0.18, 2.09	0.440	0.45	0.18, 1.15	0.095
Comfort mode	-0.34	- 0.83, 0.14	0.166	-0.18	- 0.47, 0.11	0.227	2.16	1.05, 4.45	0.036	1.11	0.58, 2.12	0.746
Heated humidifier	-0.42	- 0.78, - 0.06	0.024	-0.23	- 0.45, - 0.02	0.034	1.27	0.66, 2.45	0.472	1.49	0.89, 2.49	0.129
Heated breathing tube	-0.52	- 1.24, 0.19	0.152	-0.17	- 0.78, 0.45	0.600	1.02	0.30, 3.55	0.970	2.77	1.03, 7.44	0.043
Mask												
Interface	-	-	0.029	-	-	0.200	-	-	0.026	-	-	0.022
Nasal	-	-	-	-	-	-	-	-	-	-	-	-
Nasal pillow	-0.61	- 1.07, - 0.15	0.010	-0.04	- 0.34, 0.26	0.806	2.60	1.23, 5.50	0.012	1.64	0.83, 3.27	0.157
Oronasal	-0.30	- 0.75, 0.14	0.180	-0.22	- 0.45, 0.02	0.077	2.10	0.98, 4.49	0.055	2.12	1.23, 3.63	0.007
Mask availability since 2013	-0.44	- 0.80, - 0.08	0.016	-0.34	- 0.55, - 0.12	0.002	1.57	0.84, 2.92	0.153	2.93	1.76, 4.88	<0.001
Device reported leaks (0-100 score)	0.00	- 0.01, 0.01	0.685	0.00	- 0.01, 0.01	0.863	1.01	0.99, 1.03	0.374	1.00	0.99, 1.02	0.610
Side-effects (0-10 VAS score)												
Patient reported leaks	-0.02	- 0.08, 0.04	0.458	0.01	- 0.03, 0.05	0.611	1.00	0.91, 1.11	0.942	1.01	0.93, 1.09	0.902
Partner disturbing leaks	0.05	- 0.01, 0.11	0.079	0.02	- 0.01, 0.05	0.262	0.80	0.68, 0.94	0.007	0.93	0.86, 1.00	0.067

Table 2 (continued)

Variable	CPAP-usage (h/day)				CPAP-non-adherence (< 4 h/day)							
	Women, N= 413		Men, N= 1071		Women, N= 413		Men, N= 1071					
	Estimate	95% CI	p-value	Estimate	95% CI	p-value	OR	95% CI	p-value			
Noisy mask	-0.03	-0.10, 0.04	0.350	0.00	-0.05, 0.04	0.817	1.06	0.95, 1.19	0.291	0.97	0.88, 1.07	0.513
Heavy mask	-0.08	-0.19, 0.02	0.102	-0.04	-0.11, 0.02	0.194	1.00	0.83, 1.19	0.966	0.92	0.77, 1.09	0.310
Mask pain	-0.07	-0.16, 0.02	0.119	-0.01	-0.07, 0.05	0.678	1.08	0.94, 1.24	0.285	1.06	0.94, 1.20	0.311
Mask injury	-0.01	-0.09, 0.07	0.772	0.05	-0.01, 0.12	0.107	1.03	0.90, 1.17	0.702	0.98	0.84, 1.14	0.783
Harness pain	-0.03	-0.13, 0.07	0.522	-0.01	-0.08, 0.06	0.882	1.05	0.90, 1.22	0.560	1.11	0.97, 1.26	0.119
Harness injury	0.01	-0.12, 0.14	0.887	0.09	0.01, 0.17	0.020	1.05	0.86, 1.28	0.658	0.88	0.69, 1.12	0.293
Red eyes	0.04	-0.03, 0.11	0.235	0.03	-0.01, 0.07	0.159	0.96	0.85, 1.09	0.554	1.04	0.94, 1.14	0.450
Itchy eyes	0.03	-0.03, 0.10	0.300	0.04	0.00, 0.09	0.051	1.01	0.89, 1.13	0.929	0.98	0.88, 1.10	0.773
Dry nose	-0.05	-0.10, 0.00	0.069	-0.06	-0.09, -0.03	<0.001	1.00	0.90, 1.10	0.928	1.04	0.96, 1.12	0.308
Stuffed nose	-0.06	-0.12, 0.00	0.034	-0.02	-0.06, 0.02	0.397	1.02	0.92, 1.13	0.668	0.99	0.91, 1.09	0.904
Runny nose	0.01	-0.06, 0.08	0.719	0.02	-0.02, 0.06	0.298	0.99	0.87, 1.12	0.883	0.97	0.88, 1.07	0.586
Dry mouth	-0.05	-0.10, 0.00	0.043	-0.03	-0.06, 0.00	0.090	1.04	0.95, 1.12	0.409	1.06	0.99, 1.13	0.092
Number* [0/14]	-0.03	-0.08, 0.01	0.150	-0.01	-0.03, 0.02	0.500	1.02	0.94, 1.10	0.667	1.01	0.96, 1.07	0.632

Bolds variables are statistically significant at the 5% threshold. Italics variables are included in multivariable analyses; *Number of side effects; dichotomous data created when the VAS scale for the side effect was above or equal to 1. This variable was not included in the multivariable analyses because of its collinearity with MRSE-variables

AHI: Apnea-hypopnea Index; AHI_{low}: AHI reported by device; BMI: body mass index; CI: Exact Confidence Interval; CPAP: continuous positive airway pressure; ESS: Epworth Sleepiness Scale; MA: not applicable; OR: Odds Ratio; VAS: Visual Analogue Scale

Table 3 Multivariable regressions for CPAP-usage and CPAP-non-adherence as variables-of-interest, summary of significant explanatory-variables including side-effects

Variable	CPAP-usage (h/day)				CPAP-non-adherence (<4 h/day)					
	Women, N= 413		Men, N= 1071		Women, N= 413		Men, N= 1071			
	Estimate	95% CI	p-value	Estimate	95% CI	p-value	OR	95% CI	p-value	
Demographics										
Age (years)				0.02	0.01, 0.03	<0.001		0.96	0.93, 0.98	<0.001
BMI (kg/m ²)	0.04	0.01, 0.07	0.016	0.04	0.02, 0.07	<0.001		0.93	0.88, 0.99	0.015
Active smokers	-0.89	-1.50, -0.27	0.005					0.43	0.24, 0.77	0.005
Presence of partner										
Epworth scale										
ESS (0-24 score)				-0.04	-0.07, -0.02	0.001				
Device										
Current AHI _{low} (events/h)				0.04	0.02, 0.06	<0.001		1.08	1.03, 1.14	0.001
Treatment duration (years)										
Mean pressure (cmH ₂ O)	0.16	0.06, 0.25	0.002							
90th/95th pressure (cmH ₂ O)				0.05	0.00, 0.11	0.034				
Mask										
Mask availability since 2013				-0.30	-0.53, -0.07	0.009		2.72	1.56, 4.74	<0.001
Side-effects										
Partner disturbing leaks (0-10 VAS score)	0.07	0.01, 0.13	0.026				0.80	0.68, 0.94	0.007	
Hairness Injury (0-10 VAS score)				0.13	0.04, 0.21	0.003				
Dry nose (0-10 VAS score)				-0.06	-0.09, -0.02	0.004				
Dry mouth (0-10 VAS score)	-0.10	-0.15, -0.04	<0.001					1.08	1.00, 1.16	0.050

Note that for models, explanatory-variables with a p-value <0.15 at the univariate level were fed into multivariable analyses using stepwise selection. A backward elimination was then applied and only explanatory-variables with a p-value <0.05 at the multivariable level remain in the definitive models. Bold variables are statistically significant at the 5% threshold

AHI: Apnea-Hypopnea Index; AHI_{low}: AHI reported by device; BMI: body mass index; CI: exact confidence interval; CPAP: continuous positive airway pressure; ESS: epworth sleepiness scale; OR: odds ratio; VAS: visual analogue scale.

Discussion

To our knowledge, this study is the first to report, in a large cohort of patients treated with long-term CPAP, the gender-specific prevalence of several MRSEs and their impact on CPAP-usage and non-adherence. The main results reported here suggest that: 1) there are disparities in MRSEs according to gender and mask type; 2) different MRSEs are independently associated with CPAP-usage and non-adherence according to gender.

External validity of our study

In a 2021 narrative review, Bouloukaki et al. estimated the men-women ratio for sleep apnea syndrome to be 1.5:1 (i.e., 40% women) [13]. In our study, 27.8% of the patients are women. This is comparable to the prevalence of 32.3% recently published in a study by Prigent et al., which involved 25,846 patients in France receiving identical care [16]. However, our prevalence of women is higher than that of the ISAACC cohort (16.5%) and lower than the prevalence in the PLSC (39% women) and HynoLauss (53% women) cohorts [17]. Importantly, the characteristics of our population of women are comparable to those reported in the literature, with a lower quality of life (including more symptoms of depression), more obesity, less severe initial SAS, and less CPAP-adherence than men [11–14, 18].

Mask related side-effects depending on gender

The study identified only three MRSEs (dry mouth, runny nose and mask injury) that differed between men and women, with only mask injury potentially justifying the design of a gender-specific mask.

Dry mouth is a MRSE known to be associated with a decrease in CPAP-adherence (Bachour and Maasilta, 2004; Rotty et al., 2021). For NPMs, the median MRSE VAS score for dry mouth was significantly higher in women, and the use of heated breathing tubes was significantly elevated in this group. Similar trends were observed for NMs and ONMs, but did not reach significance. This observation is of crucial importance considering that future NPM designs cannot directly mitigate this side effect, and interventions by technicians or patients, such as the use of heated humidifiers and heated breathing tubes, should be considered to limit dry mouth. Indeed, a recent meta-analysis has reported the efficacy of heated humidifiers in addressing dry mouth (Hu et al., 2023), and the use of heated humidifiers is recommended with a moderate quality of evidence (Patil et al., 2019a; Patil et al., 2019b).

For ONMs, the median MRSE VAS score for runny nose was significantly higher in men. In a previous study, runny nose was associated with residual excessive sleepiness (RES, defined as an Epworth-Sleepiness-Scale score of ≥ 11), but not CPAP-adherence, in univariate analysis [15]. To mitigate this side effect, the use of heated humidifiers and/or topical steroids are proposed, but the quality of evidence is low [4, 5].

The prevalence of patient-reported mask injury was higher in women than in men. In a previous study, we found that mask injury was associated with RES, but not CPAP-adherence, in univariate analysis [15]. If masks specifically dedicated to women are developed in the future, they should take these gender specificities into account.

In accordance with the findings of previous studies [19, 20], our study observed that patient-reported leaks were the most prevalent MRSE, but no gender differences in prevalence were found.

Factors influencing CPAP-adherence depending on gender

In line with previous long term cohort studies, we reported that: i) for both women and men, BMI was independently and positively associated with CPAP adherence [21]; ii) for men, the presence of a partner was positively and independently associated with CPAP adherence [21]; iii) for men, treatment duration and age were also independently and positively associated with CPAP-usage [18]. Partner disturbing leaks, for women, and harness injury, for men, were positively associated with CPAP-usage and CPAP-adherence. We observe these results as an association and not as a cause, since prolonged CPAP-usage increases the risk of mask-related injuries in patients and potentially causes discomfort for their partners, particularly in cases where sleep disturbances arise from noisy air leaks or skin irritation due to leak exposure.

Study limitations

The long-term design of our study serves as both a strength and a limitation. Patients may have been treated with various mask series and mask types prior to inclusion. Therefore, we cannot ascertain whether the prevalence of different masks or MRSEs was influenced by different mask sequences. Furthermore, it is important to keep in mind that since our patients were treated with long-term CPAP our observations are not validated for patients treated with short-term CPAP.

Patients were enrolled in the study from February 7, 2017, to April 1, 2019. This is also a strength and a limitation. It is a strength because the women in the cohort received the same treatment as the men, as none of the women used masks specifically designed for women, such as the “for her” series by ResMed. However, it is a limitation because only 17% of the patients were treated with NPM, despite the increasing use of this type of mask [22, 23]. Furthermore, recent minimal contact masks, which are now available, were not used in our study, representing another limitation.

The lack of data on comorbidities, as other diseases and medications may impact the probability of MRSE. Additionally, in men, the presence of a beard may be a factor affecting MRSEs.

Conclusion

In patients undergoing long-term CPAP therapy, gender differences in MRSEs have been observed. The study identified three MRSEs that differed between men and women, with only one potentially justifying the creation of a gender-specific mask. Women were more affected by dry mouth with NPMs, and men by runny nose with ONMs; however, these side effects cannot be directly addressed in future mask designs. On the other hand, the higher incidence of mask injuries reported by women could guide the development of masks specifically designed for them. In the context of personalized medicine, our results suggest that future

mask designs should take these gender differences into account when developing masks specifically for women. Nonetheless, our study highlights that only dry mouth, a side effect not related to mask design, impacts CPAP-usage and non-adherence.

Abbreviations

AHI Apnea–Hypopnea-Index
BMI Body Mass Index
CPAP Continuous Positive Airway Pressure
ESS Epworth Sleepiness Scale
FSS French Social Security
MRSEs Mask related side-effects
NM Nasal mask
NPM Nasal pillow mask
ONM Oronasal mask
PCA Principal component analysis
QoL Quality of life
RES Residual excessive sleepiness
SAS Sleep apnea syndrome
VAS Visual analogue scales

Acknowledgements

The authors would like to thank the Apard team for their help in carrying out this study (administrative Team: Pierre Coulot, Bernard Alsina, Valerie Bachelier, Christophe Jeanjean, Philippe Lansard, Joel Nogue, Magali Partyka; technician team: Matthieu Alberti, Julien Bauchu, Yannick Baudelot, Gregory Bel, Julien Bernard, Julien Bourrel, Frederic Bousquet, David Crespy, Olivier Cubero, Eric Deghal, Fabien Deville, Sebastien Faure, Laure Ferraz, Olivier Gaubert, Georges Guichard, Franck Issert, Renaud Lopez, Mounia Maachou, Clement Maurin, David Minguez, Fabien Moubeche, Christophe Pinotti, Liva Ranaivo, Lazhar Saighi, Frederic Sola, Frederic Tallavignes, Olivier Tramier, Jean-Michel Tribe, Jean-Marc Uriol, Romain Vernet). The authors would like to thank Dr Sarah Skinner for editing the article.

Author contributions

DJ accessed all data and takes responsibility for the integrity and accuracy of the analysis. All authors contributed to and approved the final submitted manuscript. CV: data collection, analysis, and manuscript preparation; FB: data collection; JPM: study design analysis and manuscript preparation; RG: data collection and analysis; JCB: manuscript preparation; FG: manuscript preparation; AB: study design, analysis and manuscript preparation; NM: study design, data collection, analysis, and manuscript preparation; DJ: study design, analysis, and manuscript preparation.

Funding

No funding.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

InterfaceVent is registered with ClinicalTrials.gov (NCT03013283). The protocol complied with the Declaration of Helsinki and was reviewed and approved by an independent ethics committee (Comite de Protection des Personnes Sud Mediterranee 1; reference number RO-2016/50). All participants had given their written informed consent.

Consent for publication

Not applicable.

Competing interests

CV declares working for Adene home healthcare provider company. JPM declares grants/funds: Adene, Novartis, Chiesy, GSK, DPC-ORL; personal fees from Pulmon X. RG declares working for Adene home healthcare provider company. JCB declares consultant fees from AGIR a dom, a French Homecare provider. FG declares receipt of personal fees from AIR LIQUIDE SANTE, INSPIRE, BIOPROJET, RESMED and SEFAM; payment for presentations from BIOPROJET, CIDELEC, INSPIRE, RESMED, and SEFAM; non-financial support from ASTEN SANTE. AB declares grants/funds: Boehringer Ingelheim; personal fees: AstraZeneca, Boehringer Ingelheim, Chiesi, GlaxoSmithKline, Novartis, Sanofi-Regeneron; clinical trial investigator: Acceleron, Actelion, Galapagos, Merck Sharpe & Dohme, Nuvaira, Pulmonx, United Therapeutic, Celltrion, Vertex. NM declares grants/funds: GlaxoSmithKline; personal fees: Sanofi-Regeneron. DJ reports personal fees from Lowenstein, Jazz, Bioprojet, Adene, Bastide, LVL, GSK, Astra, ALK, Bohringer Ingelheim, Sanofi, Philips Healthcare, and Resmed, personal fees and nonfinancial support from Sefam and Nomics, grants and personal fees from Novartis, outside the submitted work.

References

1. Benjafield AV, Ayas NT, Eastwood PR, Heinzer R, Ip MSM, Morrell MJ, et al. Estimation of the global prevalence and burden of obstructive sleep apnoea: a literature-based analysis. *Lancet Respir Med*. 2019;7:687–98. 10.1016/S2213-2600(19)30198-5 - [DOI](#) - [PMC](#) - [PubMed](#)
2. Heinzer R, Marti-Soler H, Haba-Rubio J. Prevalence of sleep apnoea syndrome in the middle to old age general population. *Lancet Respir Med*. 2016;4:e5-6. 10.1016/S2213-2600(16)00006-0 - [DOI](#) - [PubMed](#)
3. Randerath W, Verbraecken J, de Raaff CAL, Hedner J, Herkenrath S, Hohenhorst W, et al. European Respiratory Society guideline on non-CPAP therapies for obstructive

- sleep apnoea. *Eur Respir Rev.* 2021;30: 210200. 10.1183/16000617.0200-2021 - [DOI](#) - [PMC](#) - [PubMed](#)
4. Patil SP, Ayappa IA, Caples SM, Kimoff RJ, Patel SR, Harrod CG. Treatment of adult obstructive sleep apnea with positive airway pressure: an american academy of sleep medicine clinical practice guideline. *J Clin Sleep Med.* 2019;15:335–43. 10.5664/jcsm.7640 - [DOI](#) - [PMC](#) - [PubMed](#)
 5. Patil SP, Ayappa IA, Caples SM, Kimoff RJ, Patel SR, Harrod CG. Treatment of adult obstructive sleep apnea with positive airway pressure: an american academy of sleep medicine systematic review, meta-analysis, and GRADE assessment. *J Clin Sleep Med.* 2019;15:301–34. 10.5664/jcsm.7638 - [DOI](#) - [PMC](#) - [PubMed](#)
 6. Yu J, Zhou Z, McEvoy RD, Anderson CS, Rodgers A, Perkovic V, et al. Association of positive airway pressure with cardiovascular events and death in adults with sleep apnea: a systematic review and meta-analysis. *JAMA.* 2017;318:156–66. 10.1001/jama.2017.7967 - [DOI](#) - [PMC](#) - [PubMed](#)
 7. Weaver TE, Maislin G, Dinges DF, Bloxham T, George CFP, Greenberg H, et al. Relationship between hours of CPAP use and achieving normal levels of sleepiness and daily functioning. *Sleep.* 2007;30:711–9. 10.1093/sleep/30.6.711 - [DOI](#) - [PMC](#) - [PubMed](#)
 8. Gervès-Pinquier C, Bailly S, Goupil F, Pigeanne T, Launois S, Leclair-Visonneau L, et al. Positive airway pressure adherence, mortality, and cardiovascular events in patients with sleep apnea. *Am J Respir Crit Care Med.* 2022;206:1393–404. 10.1164/rccm.202202-0366OC - [DOI](#) - [PubMed](#)
 9. Azarbarzin A, Zinchuk A, Wellman A, Labarca G, Vena D, Gell L, et al. Cardiovascular benefit of continuous positive airway pressure in adults with coronary artery disease and obstructive sleep apnea without excessive sleepiness. *Am J Respir Crit Care Med.* 2022;206:767–74. 10.1164/rccm.202111-2608OC - [DOI](#) - [PMC](#) - [PubMed](#)
 10. Bublitz M, Adra N, Hijazi L, Shaib F, Attarian H, Bourjeily G. A narrative review of sex and gender differences in sleep disordered breathing: gaps and opportunities. *Life.* 2022;12:2003. 10.3390/life12122003 - [DOI](#) - [PMC](#) - [PubMed](#)
 11. Martínez-García MÁ, Sierra-Párraga JM, Garcia-Ortega AA. Obstructive sleep apnea in women: we can do more and better. *Sleep Med Rev.* 2022;64: 101645. 10.1016/j.smrv.2022.101645 - [DOI](#) - [PubMed](#)
 12. Patel SR, Bakker JP, Stitt CJ, Aloia MS, Nouraiie SM. Age and sex disparities in adherence to CPAP. *Chest.* 2021;159:382–9. 10.1016/j.chest.2020.07.017 - [DOI](#) - [PMC](#) - [PubMed](#)
 13. Bouloukaki I, Tsiligianni I, Schiza S. Evaluation of obstructive sleep apnea in female patients in primary care: time for improvement? *Med Princ Pract.* 2021;30:508–14. 10.1159/000518932 - [DOI](#) - [PMC](#) - [PubMed](#)
 14. Martínez-García MÁ, Labarca G. Obstructive sleep apnea in women: scientific evidence is urgently needed. *J Clin Sleep Med.* 2022;18:1–2. 10.5664/jcsm.9684 - [DOI](#) - [PMC](#) - [PubMed](#)
 15. Rotty M-C, Suehs CM, Mallet J-P, Martinez C, Borel J-C, Rabec C, et al. Mask side-effects in long-term CPAP-patients impact adherence and sleepiness: the InterfaceVent real-life study. *Respir Res.* 2021;22:17. 10.1186/s12931-021-01618-x - [DOI](#) - [PMC](#) - [PubMed](#)
 16. Prigent A, Blanloeil C, Jaffuel D, Serandour AL, Barlet F, Gagnadoux F. Seasonal changes in positive airway pressure adherence. *Front Med.* 2024;11:1302431.10.3389/fmed.2024.1302431 - [DOI](#) - [PMC](#) - [PubMed](#)

17. Solelhac G, Sánchez-de-la-Torre M, Blanchard M, Berger M, Hirotsu C, Imler T, et al. Pulse wave amplitude drops index: a biomarker of cardiovascular risk in obstructive sleep apnea. *Am J Respir Crit Care Med*. 2023;207:1620–32. 10.1164/rccm.202206-12230C - [DOI](#) - [PMC](#) - [PubMed](#)
18. Prigent A, Blanloeil C, Serandour A-L, Barlet F, Gagnadoux F, Jaffuel D. A biphasic effect of age on CPAP adherence: a cross-sectional study of 26,343 patients. *Respir Res*. 2023;24:234. 10.1186/s12931-023-02543-x - [DOI](#) - [PMC](#) - [PubMed](#)
19. Bachour A, Vitikainen P, Virkkula P, Maasilta P. CPAP interface: satisfaction and side effects. *Sleep Breath*. 2013;17:667–72. 10.1007/s11325-012-0740-0 - [DOI](#) - [PubMed](#)
20. Borel JC, Tamisier R, Dias-Domingos S, Sapene M, Martin F, Stach B, et al. Type of mask may impact on continuous positive airway pressure adherence in apneic patients. *PLoS ONE*. 2013;8: e64382. 10.1371/journal.pone.0064382 - [DOI](#) - [PMC](#) - [PubMed](#)
21. Gagnadoux F, Le Vaillant M, Goupil F, Pigeanne T, Chollet S, Masson P, et al. Influence of marital status and employment status on long-term adherence with continuous positive airway pressure in sleep apnea patients. *PLoS ONE*. 2011;6: e22503. 10.1371/journal.pone.0022503 - [DOI](#) - [PMC](#) - [PubMed](#)
22. Laaka A, Hollmén M, Bachour A. Evaluation of CPAP mask performance during 3 years of mask usage: time for reconsideration of renewal policies? *BMJ Open Respir Res*. 2021;8: e001104. 10.1136/bmjresp-2021-001104 - [DOI](#) - [PMC](#) - [PubMed](#)
23. Zonato AI, Rosa CFA, Oliveira L, Bittencourt L. Comparing CPAP masks during initial titration for obstructive sleep apnea syndrome: 1 year experience. *Braz J Otorhinolaryngol*. 2022;88(Suppl 5):S63–8. 10.1016/j.bjorl.2021.10.007 - [DOI](#) - [PMC](#) - [PubMed](#)