

ST38 E. coli evolution in response to the natural conjugative plasmids carrying the carbapenemase gene blaOXA 48

Pengdbamba Dieudonné Zongo, Nicolas Cabanel, Philippe Glaser, Isabelle Rosinski-Chupin

▶ To cite this version:

Pengdbamba Dieudonné Zongo, Nicolas Cabanel, Philippe Glaser, Isabelle Rosinski-Chupin. ST38 E. coli evolution in response to the natural conjugative plasmids carrying the carbapenemase gene blaOXA 48. 17e Congrès national de la société française de microbiologie, Société Française de microbiologie - SFM, Oct 2022, Montpellier, France. hal-04695579

HAL Id: hal-04695579 https://hal.science/hal-04695579v1

Submitted on 12 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ST38 *E. coli* evolution in response to the natural conjugative plasmids carrying the carbapenemase gene *bla*_{0XA-48}

Pengdbamba Dieudonné ZONGO (pengdbamba-dieudonne.zongo@pasteur.fr)^{1,2}, Nicolas Cabanel¹, Philippe Glaser¹, Isabelle ROSINSKI-CHUPIN¹

¹EERA, Institut Pasteur, Paris, France ² Sorbonne Université, Paris, France

Introduction and objectives

Carbapenems are last resort ß-lactams used to treat multidrug resistant Gram-negative bacteria. However, the increased incidence of carbapenem resistance is threatening their therapeutic efficacy. The emergence of *E. coli* clones expressing carbapenemases and their dissemination in the community are particularly worrisome. Sequence Type (ST) 38 is the most prevalent ST among carbapenemase-producing *E. coli* isolated in France and in other European countries. By phylogenetic analysis, we showed the world-wide dissemination of at least three OXA-48 producing ST38 lineages with chromosomal integrations of the bla_{OXA-48} gene, contrasting with the general mode of bla_{OXA-48} dissemination in *Enterobacterales* through conjugative transfer of IncL pOXA-48 plasmids. We are aiming to characterize the benefit of chromosomal integrations and the involved mechanisms.

Materials and methods

We introduced three different variants of pOXA-48 by conjugation into different naïve ST38 *E. coli* strains and assessed the fitness cost by monitoring growth in rich and poor media. We assessed the plasmid stability by ten days passages without selective pressure. We also performed a 28-day experimental evolution of the different transconjugants using meropenem as selective pressure to look for mutations that may lower potential fitness cost and to select *bla*_{OXA-48} chromosomal integration. Evolved strains were analyzed by whole genome sequencing.

Results, discussion and conclusion

The three pOXA-48 variants were unstable in three different ST38 *E. coli* genetic backgrounds. Particularly, growth curve analyses revealed a strong fitness decrease in one of these backgrounds with an up to 26% increase in generation time whatever the plasmid variant. *bla*_{OXA-48} integrations into chromosome were observed during experimental evolution. They were dependent on both the genetic background of the transconjugant and on the pOXA-48 type and led to a fitness improvement. Alternatively, an increase in both the fitness of evolved transconjugants and in plasmid stability was reached in transconjugants inactivated or mutated in an operon of unknown function. We hypothesize that proteins encoded by this operon might interact with pOXA-48 plasmids. Altogether our results provide new information on the emergence of *E. coli* lineages expressing OXA-48 carbapenemases.

Key words: OXA-48, E. coli ST38, carbapenem resistance, plasmid stability, evolution