

Photophysical, Cyclic Voltammetry, Electron Paramagnetic Resonance, X-ray Molecular Structure, DFT Calculations and Molecular Docking Study of a new Mn(III) Metalloporphyrin

Hayfa Mkacher ^a, Bouzid Gassoumi ^b, Nour Ehouda Dardouri ^a, Soumaya Nasri ^{a,c}, Frédérique Loiseau ^d, Florian Molton ^d, Thierry Roisnel ^e Ilona Turowska-Tyrk^f, Houcine Ghala ^b, Samir Acherar ^g and Habib Nasri ^{*,a}

^a: University of Monastir. Laboratory of Physical Chemistry of Materials. Faculty of Science of Monastir. Avenue de l'Environment. 5019 Monastir. Tunisia.

^b: Laboratory of Advanced Materials and Interfaces (LIMA). University of Monastir. Faculty of Science of Monastir. Avenue of Environment. 5000 Monastir. Tunisia.

^c: Department of Chemistry, College of Science, Majmaah University, Majmaah 11952, Saudi Arabia.

^d: Department of Molecular Chemistry. CNRS UMR 5250. Université Grenoble Alpes. F-38000 Grenoble. France.

^e: Institute of Chemical Sciences of Rennes, UMR 6226, University of Rennes 1, Beaulieu Campus, Rennes 35042, France.

^f: Faculty of Chemistry, University of Warsaw, Pasteura 1, 02-093 Warsaw, Poland.

^g: Université de Lorraine. CNRS. LCPM. F-54000 Nancy. France.

Contents

1. IR spectroscopy.....	2
2. X-ray Molecular structure.....	4
3. Hirshfeld surface analysis.....	7
4. EPR spectroscopy.....	8

1. IR spectroscopy




Figure S1. Neat IR spectrum of H_2TClPP .




Figure S2. Neat IR spectrum of $[\text{Mn}^{\text{III}}(\text{TCIPP})(\text{OAc})]$.




Figure S3. Neat IR spectrum of $[\text{Mn}^{\text{III}}(\text{TCIPP})\text{SO}_3\text{CF}_3]$.




Figure S4. Neat IR spectrum of Complex I.

2. . X-ray Molecular structure




Figure S5. Top view of the $[\text{Mn}^{\text{III}}(\text{TClPP})(\text{DMAP})_2]^+$ ion complex showing the dihedral angle φ between the DMAP plane axial ligands and the closest N(pyrrole)-Mn-N(DMAP).

Table S1. Inter- and intra-molecular interactions for **I**.

<i>Hydrogen bond</i>			
D ^a -H...A ^b	Symmetry	D...A [Å]	D-H...A [°]
C7-H7...O3	1-x, 1-y, 1-z	3.150(4)	126
C29-H29...O1	x, -1+y, z	3.334(4)	143
C48-H48...O2	x, -1+y, z	3.335(5)	162
C53-H53...O3	-1+x, -1+y, z	3.416(4)	160
C64-H64...O1	1-x, 1-y, 1-z	2.959(9)	148

<i>Interactions C-H…Cg</i>			
C-H…Cg (π)	Symmetry code	C…Cg (π)	C–H…Cg
C23-H23…Cg1	-x, 1-y, 1-z	3.884(4)	164
C51-H51C…Cg10	1+x, y, z	3.528(4)	132
C58-H58B…Cg9	-1+x, y, z	3.405(4)	133
C61-H61A…Cg14	1+x, y, z	3.749(12)	161

<i>Inter- and Intra-molecular interactions C-X…Cg</i>				
C–X…Cg (π)	Symmetry code	C…Cg	X…Cg	C–H…Cg
C30-Cl2…Cg11	x,-1+y,z	4.949(3)	3.4337(15)	143.39(12)
C42-Cl4…Cg13	x, 1+y, z	4.855(3)	3.4195(15)	137.59(12)
C64-Cl6…Cg14	1-x, 1-y, 1-z	4.710(9)	3.598(3)	118.10(3)
C59-F1…Cg3	1+x, 1+y, z	4.356(5)	3.300(4)	136.30(3)
C59-F2…Cg11	1-x, 1-y, 1-z	4.442(4)	3.297(4)	144.10(3)
C65-Cl9…Cg14	1-x, 1-y, 1-z	4.920(2)	3.811(16)	119.90(9)

^a: D = donor, ^b: A = acceptor, X =heteroatom, Cg =centroid.

Cg1: is the centroid of the N1/C1-C4 pyrrole five-membered ring.

Cg3: is the centroid of the N3/C11-C14 pyrrole five-membered ring.

Cg9: is the centroid of the N5/C45-C49 phenyl six-membered ring.

Cg10: N7/C52-C56 phenyl six-membered ring.

Cg11: C21-C26 a phenyl ix-membered ring.

Cg13: C33-C38 a phenyl six-membered ring.

Cg14: C39-C44 a phenyl six-membered ring.




Figure S6. Hydrogen bonds interactions in complex I according to the direction [100].




Figure S7. Inter-and intra-molecular interactions C–H...Cg in the crystal lattice of I view along the [010] direction.




Figure S8. Inter-and intramolecular interactions C–X…Cg in the crystal structure of **I** view along the [100] direction.

3. Hirshfeld surface analysis




Figure S9. Hirshfeld surface mapped over d_{norm} and decomposed fingerprint plots for the dominant interactions for **I**

4. EPR spectroscopy




Figure S10. Experimental EPR spectra for complex I in parallel (a) and perpendicular (b) modes at 5 K.