

Identification of a Streptomyces necrotic elicitor involved in antifungal activity, plant defense stimulation and bacteria fitness in the rhizosphere

Clément Nicolle, Damien Gayrard, Aurélien Amiel, Alba Noel, Sylvie Lautru,

Thomas Rey, Bernard Dumas

▶ To cite this version:

Clément Nicolle, Damien Gayrard, Aurélien Amiel, Alba Noel, Sylvie Lautru, et al.. Identification of a Streptomyces necrotic elicitor involved in antifungal activity, plant defense stimulation and bacteria fitness in the rhizosphere. ICPP 2023 - 12th International Congress of Plant Pathology, Aug 2023, Lyon, France. hal-04701417

HAL Id: hal-04701417 https://hal.science/hal-04701417v1

Submitted on 18 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Identification of a Streptomyces specialized metabolite involved

in antifungal activity, plant defense stimulation and bacteria fitness in the rhizosphere

Clément NICOLLE¹, Damien GAYRARD², Marion HORTALA¹, Aurélien AMIEL², Maya Long³, Alba NOEL³, Sylvie LAUTRU³, Thomas REY², Bernard DUMAS¹

¹Laboratory of Research in Plant Science, University of Toulouse, CNRS, UPS, Toulouse INP, Auzeville Tolosane, France

² De Sangosse, Pont-Du-Casse, France

³ Institute for Integrative Biology of the Cell, CEA, CNRS, University of Paris-Sud, Orsay, France

|--|

Identifying new strains from rhizosphere and understanding their biological activities offers new Arabidopsis thaliana. approaches to plant protection.

activate hypersensitive response (HR) in

Streptomyces sp. AgN23 was isolated from grapevine rhizosphere and screened for its abilities to produce

Gaining deeper understanding of the interaction between AgN23 and Arabidopsis thaliana is necessary to assess its potential as a **biocontrol** agent.

Findings

Identification of a candidate specialized metabolite in AgN23 culture extracts

 $(\mathbf{2})$ Implementation of a reverse genetic approach

Evaluation of the **mode of action** of the metabolit (4)

(3) genes, metabolomic rerouting)

Importancce of the compounds biosynthesis for the strain fitness

reroutes plant specialized metabolome

Validation of the role of the candidate compound

Mock

AgN23 WT



 $(\mathbf{2})$

Jin plant defense pathways activation (HR, defense AgN23 knock-out mutants for galbonolides are impaired in plant

pSC004-22



Figure 1: Left, Photograph of Arabidopsis thaliana rhizoplane colonized by AgN23 10 days after inoculation. **Right**, Confocal fluorescence image of AgN23-GFP around Arabidopsis root.



Figure 3: Arabidopsis leaf area measurement following growth within non- or inoculated potting soil.



Figure 2: Observation of Arabidopsis thaliana dwarf root phenotype 10 days after inoculation with AgN23.



Figure 4: AgN23 genome copy number in *Arabidopsis* rhizosphere 8 weeks after soil inoculation.

defense activation

pSC004-10



Figure 13: In planta expression level of *pr1*, *pad3*, and *pad4* genes responsible for







Figure 14: Peak area of camalexin in Arabidopsis 10 days after inoculation with WT or KO strains

Figure 5: AgN23 modulates Arabidopsis root metabolome by inducing biosynthesis of indolic compounds such as camalexin.

⁷AgN23 produces galbonolides, antifungals inhibiting ceramide synthesis in Arabidopsis thaliana



GalB GalC GalD GalA GalE AJZ84270 QTI87461 QTI87463 QTI87462 AJZ86980

Figure 8: Galbonolides biosynthetic gene cluster with putative AgN23 genomic loci (NCBI assembly GCA_001598115.2). Knock-out gene is shown salicylic acid-dependent systemic acquired resistance, synthesis of camalexin and positive regulation of SA accumulation, respectively

4 ⁹Galbonolides biosynthesis modulates AgN23 fitness in the rhizosphere of Arabidopsis thaliana

pSC004-22

pSC004-10

Figure 15: Left, Photographs of Arabidopsis 6 weeks within WT or KO inoculated potting soil. **Right,** AgN23 genome copy number in Arabidopsis rhizosphere 6 weeks after inoculation.

Conclusion

Figure 10: Left, Conversion of NBD-C6-ceramide to NBD-C6-IPC by Arabidopsis thaliana inositol phosphorylceramide synthase (AtIPCS2). **Right**, AtIPCS2 activity following treatments with extracts from culture supernatant of AgN23 WT or KO mutants

References

Ahkami, et al. (2017). Rhizosphere engineering: Enhancing sustainable plant ecosystem productivity. Rhizosphere 3: 233–243. Denny & Smith. (2004). Rafts and sphingolipid biosynthesis in the kinetoplastid parasitic protozoa. Molecular Microbiology 53: 725–733. Gayrard, D., Nicolle, C., et al. (2023). Genome Sequence of the Streptomyces Strain AgN23 Revealed Expansion and Acquisition of Gene Repertoires Potentially Involved in Biocontrol Activity and Rhizosphere Colonization. PhytoFrontiersTM: PHYTOFR-11-22-0131-R.

Mandala, et al. (1998). Rustmicin, a Potent Antifungal Agent, Inhibits Sphingolipid Synthesis at Inositol Phosphoceramide Synthase. Journal of Biological Chemistry 273: 14942–14949. Vergnes, et al. (2020). Phyllosphere colonisation by a soil Streptomyces sp. promotes plant defense responses against fungal infection.: 43.

