



HAL
open science

Glaucoma and Computer Use: Do Contrast and Color Enhancements Improve Visual Comfort in Patients?

Clémentine Garric, Jean-François Rouland, Quentin Lenoble

► **To cite this version:**

Clémentine Garric, Jean-François Rouland, Quentin Lenoble. Glaucoma and Computer Use: Do Contrast and Color Enhancements Improve Visual Comfort in Patients?. *Ophthalmology Glaucoma*, 2021, 4 (5), pp.531-540. 10.1016/j.ogla.2021.01.006 . hal-04702663

HAL Id: hal-04702663

<https://hal.science/hal-04702663v1>

Submitted on 13 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Title page**

2 **TITLE**

3 *Glaucoma and computer use: do contrast and color enhancements improve visual comfort in*
4 *patients?*

5 **RUNNING HEAD**

6 Glaucoma and Computer Use

7 **AUTHORS**

8 Clémentine Garric ^{1,2}, Jean-François Rouland ^{1,2,3} & Quentin Lenoble ^{1,2}

9 ¹ Univ. Lille, Inserm, CHU Lille, U1172 - LiNCog (JPARC) - Lille Neurosciences & Cognition, F-59000 Lille, France

10 ² Univ. Lille, CNRS, CHU Lille, UMR 9193, SCALab, Sciences Cognitives et, Sciences Affectives, F-59000 Lille,
11 France.

12 ³ Ophthalmology Department, Claude Huriez Hospital, University of Lille, Lille, France

13 **CORRESPONDING AUTHOR**

14 Quentin Lenoble, Faculté de Médecine de Lille,
15 Pôle Recherche, Siem et. Ouest, 1 Place de Verdun, Lille 59000, France
16 (email: quentin.lenoble@univ-lille.fr)

17 **FINANCIAL SUPPORT**

18 PhD grant: Centre Hospitalier Universitaire de Lille & Région de Lille

19 **CONFLICT OF INTEREST**

20 No conflicting relationship exists for any author.

21 **KEYWORDS**

22 Low vision, glaucoma, computer use, quality of life, eye-tracking, image enhancement

23 **ABBREVIATIONS**

24 CGI: Computer Graphical Interface

25 CLV: Central Loss of Vision

26 CS: Contrast Sensitivity

27 HFA: Humphrey Field Analyzer

28 I.T.: Information Technology

29 NEI VFQ-25: National Eye Institute Visual Functioning Questionnaire

30 PCA: Principal Component Analysis

31 PLV: Peripheral Loss of Vision

32 POAG: Primary Open-Angle Glaucoma

33

34 Abstract

35 **Purpose:** To estimate the impact of glaucoma on computer use and assess specific adaptations of the
36 graphical interface to this form of visual impairment.

37 **Design:** Prospective experimental cohort study.

38 **Participants:** Forty-nine participants were recruited: 16 patients with primary open-angle glaucoma
39 (POAG, mean±SD: 62.7±5.6 years old), 17 age-matched participants (mean±SD: 59.1±8.3 years old)
40 and 16 young controls (mean±SD: 23.3±2.1 years old).

41 **Methods:** An ophthalmologic examination prior to the study evaluated the level of visual loss (MD),
42 visual acuity (LogMar) and contrast sensitivity (CS) of the POAG patients. Each participant underwent
43 the following measurements: an Information Technology (I.T.) experience questionnaire, a
44 preference task monitored by eye-tracking, and a feed-back session. The experimental task was
45 based on ecological computer scenes following three enhancement levels (low, medium and high),
46 determined by gradual modulation of contrast, luminance and color. Participants were asked to
47 select the most readable and comfortable stimulus between four images displayed on the screen: the
48 original computer scene and three enhanced versions.

49 **Main outcome measures:** Clinical, oculomotor and subjective data were computed together in a
50 multivariate model by using a principal component analysis (PCA).

51 **Results:** The PCA revealed three principal components accounting for 72% of the total variance of the
52 data and showed a greater need for enhanced computer scenes in glaucoma patients, an equal
53 preference for low and medium enhancement within the three groups, and significantly longer
54 oculomotor behavior in the patients group. Subjective reports of difficulty to use I.T. due to vision
55 were correlated with visual impairment and high enhancement preference. Contrast sensitivity (CS)
56 was critical to explain the main variations of the data. A reduced CS had a significant effect on the
57 preference for enhanced computer scenes ($r = -0.431$; $p < 0.002$) and a less effective exploration
58 velocity ($r = 0.428$; $p < 0.002$).

59 **Conclusions:** Glaucoma alters the global exploration of computer scenes. High enhancement of the
60 graphical interface could improve visual comfort during computer use. Subjective patients' reports
61 underline the importance of including I.T. questions in visual-related quality-of-life questionnaires.

62 Introduction

63 Glaucoma is the progressive degeneration of the optic nerve fiber, resulting in loss from peripheral to
64 central vision that can critically lead to blindness. The pathology usually affects adults over 40 years
65 old, and its prevalence is increasing considerably with the aging of the world population.¹
66 Appropriate treatments tend to limit the progression in visual field loss but damage is irreversible
67 and affects everyday life functioning.^{2,3} Previous studies have shown that the supposedly spared
68 central vision in glaucoma presents certain impairments, such as a decrease in contrast sensitivity⁴⁻⁶
69 and alteration of luminance adaptation.⁷ Furthermore, in tasks involving higher levels of visual
70 cognition capacities, some authors have demonstrated that glaucoma affects complex visual
71 processing, such as object categorization or face recognition in ways that are not predictable by their
72 extent of visual field loss.⁸⁻¹⁰ Ganglion cell damage in glaucoma may also affect oculomotor behavior.
73 For instance, by measuring eye movements towards static and moving targets in central vision,
74 Lamirel et al. (2014) demonstrated that, compared to age-matched controls, saccades in patients
75 presented delayed latency and decreased precision.¹¹ Altogether, these studies bring new
76 understanding of the disease by testing the limits of patients' visual capacities at threshold stimuli on
77 a computer screen (low contrast, decreasing size, filtered spatial frequency, repetitive target
78 movements, ...). However, while most of the experimental paradigms are computer-dependent, few
79 investigations have focused on the impact of glaucoma on Information Technology (I.T.) use with
80 ecological measures. Recently, Asfaw et al. assessed eye movements during free exploration of
81 natural scenes on computer¹² and found altered saccadic movements and oculomotor compensation
82 strategies in patients with glaucoma. However, participants were tested monocularly and these
83 results were found by comparing worse and better eye recordings. To our knowledge, no study has
84 yet investigated the binocular visual performance of patients with glaucoma on ecological computer
85 tasks compared to normal aging subjects. We hypothesized that impairment due to glaucoma,
86 previously measured at threshold on experimental paradigms, could have an impact on common

87 computer use. The specific knowledge of the limits of visual functions could lead to new tools to
88 enhance the I.T. experience of patients.

89 A translational area of research between vision science, ophthalmology and technology aims at
90 developing high-tech aids for the rehabilitation of the visually impaired. Moshtael et al. (2015)
91 reviewed the digital image processes used to maximize the residual vision in low-vision patients.¹³ To
92 name a few, contrast, contour enhancement and edge detection techniques were used on images or
93 videos to create high-tech aids and assess visual performance improvement in low-vision patients
94 (i.e. object recognition, reading). In this review, only one study among the twelve selected
95 considered peripheral loss of vision (PLV) and the peripheral deficit condition was an artificially
96 simulated tunnel vision.¹⁴ The main clinical target for these aids remains patients with central vision
97 loss, i.e. age-related macular degeneration. Considering the previously mentioned central vision
98 impairment of patients with glaucoma, such technological rehabilitation could benefit PLV patients.
99 One may hypothesize that modulating specific parameters of the computer graphical interface (CGI),
100 such as increasing contrast along with adjusting luminance intensity, could create a more suitable
101 digital environment for glaucomatous patients. Furthermore, results from the central loss of vision
102 (CLV) aid literature could also provide relevant tracks to follow. Watson et al. (2012) hypothesized
103 that by developing stimuli enhanced according to the visual properties of the residual peripheral
104 vision in patients with CLV, they could improve their capacity to read or recognize facial
105 expressions.¹⁵ Jittered stimuli (with low contrast and low spatial frequency) improved the visual
106 abilities of CLV patients with advanced symptoms, but not in the group with moderate vision loss. In
107 line with these findings, we postulated that progressive enhancement of stimuli according to the
108 visual properties of central vision (i.e. a progressive contrast and color heightening in computer
109 scenes) could be adapted for various glaucoma stages and could optimize the readability of
110 computer scenes for PLV patients.

111 Further evidence of the lack of information on the I.T. experience of glaucomatous patients, or low-
112 vision patients in general, can be found in visually related quality of life (QoL) questionnaires such as
113 the 25-item *National Eye Institute Visual Functioning Questionnaire* (NEIVFQ-25¹⁶). While these
114 questionnaires are commonly used to measure everyday life difficulties of low-vision patients,^{17,18}
115 patient-reported I.T. experience is still not included.

116 In this context, the aim of the present study was to fill the gap between the lack of information
117 concerning the possible impact of glaucoma on I.T. use (subjective experience and oculomotor
118 behavior) and graphical interface adaptation on computer screens for PLV patients. To do so, we
119 built gradual enhanced versions of ecological computer scenes and tested three groups of
120 participants (young controls, age-matched controls and glaucoma patients) on a preference task,
121 monitored by an eye-tracker. We aimed at exploring the I.T. experience and its enhancement in
122 glaucomatous and normal aging populations, considering the four critical dimensions of the study:
123 subjective I.T. experience, visual assessment, enhancement preference measures and oculomotor
124 behavior.

125 **Methods**

126 **Participants**

127 Forty-nine participants were recruited for this study: 16 patients with primary open-angle glaucoma
128 (POAG, mean±SD: 62.7±5.6 years old), 17 age-matched participants (mean±SD: 59.1±8.3 years old)
129 and 16 young controls (mean±SD: 23.3±2.1 years old). All participants provided written informed
130 consent before the experiment. They were required to have a corrected binocular visual acuity of at
131 least 0.2 logarithm of the minimum angle of resolutions units (log MAR), no neurological history and
132 no ophthalmological condition other than glaucoma, except for uncomplicated cataract surgery.
133 According to the results on the Humphrey Field Analyzer (HFA; Carl Zeiss Meditec, Dublin, CA), three
134 possible stages of the pathology were defined on the basis of the worst eye Mean Deviation (MD)
135 score as: early (from 0.00 to -6.00 dB), moderate (going from -6.01 to -12.00 dB) and severe

136 glaucoma (lower than -12.00 dB). Clinical data is detailed in **Table 1**. Each participant over 60 years
137 old (controls and glaucoma patients) was cognitively assessed with the Mini Mental State
138 Examination (MMSE¹⁹) and had to reach a score above 26/30 to be included. Binocular contrast
139 sensitivity was assessed with a Pelli-Robson chart and scored in log CS. These clinical evaluations
140 were followed by the experimental procedure: an I.T. experience questionnaire, a forced-choice
141 experiment (the preference task) monitored by eye-tracking, and a feedback session. This protocol
142 was approved by our institutional ethics committee (VAO 2019-346-S71) and adhered to the
143 Declaration of Helsinki.

144 *[Insert Table 1 - Demographic and clinical data of participants with POAG]*

145 **I.T experience questionnaire**

146 The I.T. experience questionnaire was created for this study to measure the global experience of the
147 participants towards information technologies (original French version available in **Supplemental**
148 **Material**). The aim was to collect information efficiently about the type of device used, the frequency
149 of use and the estimated user level (five items), as well as an estimation of potential difficulties
150 encountered due to vision (four items). Based on the NEIVFQ-25 format, which is widely used to
151 quantify quality of life in low-vision patients, the last four items were focused on everyday life I.T.
152 common tasks (i.e. reading, searching for icons on a computer screen, reaching a target with a
153 mouse) and the answers were given on a scale from (1) “No difficulty at all” to (5) “Use stopped
154 because of visual difficulties”. An average difficulty score was computed from these four items.

155 Demographic details by groups are shown in **Table 2**.

156 *[Insert Table 2- Contrast sensitivity and I.T. experience demographic data]*

157 **Apparatus**

158 Participants were seated in a dimly lit room with controlled artificial lighting. The distance from the
159 screen (15.6-inch screen; resolution 1920x1080 pixels) was set at 57 cm with a chin rest. The
160 computer was a Dell Precision M4800 equipped with an Intel Core I7 processor and 16 GB RAM. The
161 stimuli were displayed synchronously with eye-tracking recording thanks to the *Experiment Center*
162 software and a RED-m eye-tracker model (250 Hz) by *SensoMotoric Instruments (SMI)*. Participants
163 were tested binocularly, wearing their usual correction.

164 **Stimuli**

165 To create ecological computer scenes as stimuli, we gathered 30 screenshots of web pages of
166 everyday life use of the internet (weather website, recipes, maps, ...). According to the intensity of
167 contrast, color and luminance from original computer scenes, three experimental levels of image
168 enhancement were created using the *Gnu Image Manipulation Program (GIMP, 2.10.18)* free and
169 open source. Pictures were handled in RGB mode (Red-Green-Blue) by the program. Pixel properties
170 on RGB values were converted by different levels of percentage on contrast, balanced luminance,
171 and color saturation for each picture. The three levels of image enhancement were defined as
172 follows: (1) *Low enhancement* was processed by increasing contrast (mean±SD: 58% ± 25) and
173 balancing luminance (mean±SD: -33% ± 28). Visually, *low enhancement* pictures depict a slight
174 increase in contrast. (2) *Medium enhancement* corresponded to a *low enhancement* processing
175 merged with color increment (mean±SD: 96% ± 26). Pictures with *medium enhancement* appeared to
176 the participants as more contrasted and more colored than the original picture. (3) *High*
177 *enhancement* processing applied high saturation of the original version without counterbalancing
178 luminance. Descriptively, the high enhancement was is radically different from the original and
179 accentuated contrast, luminance, and color perceptions in a way that contours and lines were more
180 detectable. In the end, the image database included 120 stimuli (922 × 518 pixels), namely four
181 versions of each of the 30 screenshots (**Fig. 1**): *original* (i.e. unenhanced), *low*, *medium* and *high*

182 *version*. To rigorously validate the stimuli set, the physical properties of the 120 stimuli were
183 analyzed, compared among the three enhancement levels, and adjusted using GIMP.

184 *[Insert Figure 1: Experimental design of one trial of the preference task]*

185 **Procedure**

186 The preference task was used to measure the most readable and comfortable image enhancement of
187 ecological computer scenes. Participants were tested in two blocks of 30 trials, after calibration of
188 the eye tracker and two training trials. The trial structure began with a fixation phase ranging from
189 400 ms to 600 ms, with a red fixation cross at the center of the screen. The fixation phase was
190 followed by the simultaneous display of four versions of one computer scene (**Fig. 1**) and a cursor at
191 the center (on the red fixation cross). Participants were instructed to carefully observe the four
192 stimuli and to manually select the most readable and comfortable display, thanks to the cursor on
193 the screen. There was no time restriction and the next trial started after the selection. The 30 trials of
194 each block corresponded to the 30 randomly ordered screenshots. After the end of the first block
195 and a short break (duration decided by the participant), the same stimuli were shown in a different
196 randomized order during the second block. Each version (*original, low, medium and high*
197 *enhancement*) was equally represented over the 60 trials on the four portions of the screen (i.e.
198 upper left, upper right, lower left, lower right) and not displayed in the same position between the
199 first and the second block. At the end of the two blocks, open-ended feedback questions were
200 answered. The answers were recorded to let the participants describe their preference and their
201 choices during the experiment. Critical key words and preference types were analyzed for each of the
202 49 participants, then gathered into different categories to classify their answers on a Subjective
203 Enhancement Preference Scale, ranging from (1) *None* to (5) *Highly augmented pictures*.

204 **Data Analysis**

205 The data was analyzed in two ways. First, single variable analyses were designed to assess how the
206 main variables of the experimental task were related to vision loss (comparison between healthy
207 participants and patients, along with correlation between clinical measures and performances).
208 Secondly, the Principal Component Analysis revealed correlations between objective (clinical data,
209 oculomotor data) and subjective (preference task, I.T. questionnaire) records. The analyses were
210 conducted with Statistical Package for the Social Sciences (SPSS, version 25).

211 *Single variable analyses*

212 Non-parametric tests (Kruskal-Wallis) were used to assess differences between groups for
213 demographical data (**Table 2**) and individual variables. A Bonferroni correction was conducted to test
214 groups pairwise. When variables showed a normal distribution but heterogenous population
215 variances, a pairwise comparison was performed with the Welch t-test. Correlations between clinical
216 data and behavioral data were assessed by Spearman's rank correlations. Statistical significance of
217 the correlation coefficient (r) was set at $P < 0.05$.

218 *Principal Component Analysis*

219 Principal component analysis (PCA) is a multivariate statistical method generally applied to reduce
220 high-dimensional data sets into groups of interrelated variables called principal components (PC).
221 PCA starts with a covariance matrix computation between every variable. Based on this matrix, the
222 calculation of eigenvalues and eigenvectors helps sort principal components by relevance: the first
223 principal components have the highest eigenvalue, meaning that they retain the optimized rate of
224 variation present in all original variables and eliminate redundant information. Each PC with an
225 eigenvalue below 1 provides less information than an individual variable variation (Kaiser criterion).
226 The number of PC retained usually accounts for at least 50% of the total variance of the original data
227 set. Eigenvectors are converted to factor loadings that characterize the weight and nature of the
228 correlation of an individual variable within a PC.

229 We performed a PCA on our data set to statistically identify the link between *visual assessment*, *I.T.*
230 *experience*, *preference measures* and *oculomotor behaviors* defined below. Two variables were
231 included in the visual assessment: *contrast sensitivity* (log CS) and the *clinical visual deficit* on a 4-
232 level scale: 1-None, 2-Corrected vision, 3-Monocular Glaucoma, 4-Binocular Glaucoma. The *I.T.*
233 experience was represented by the *I.T. difficulty score due to vision* (**Table 2**).

234 We evaluated four preference measures:

- 235 - *Behavioral preference* as the percentage of original, low, medium and high enhancement
236 manually answered on 60 trials ($P_{original}$, P_{low} , P_{medium} , P_{high});
- 237 - *Visual-oriented preference* as the mean dwell time per enhancement version (sum of fixation
238 durations and saccades inside an Area Of Interest (AOI)) standardized by the total mean
239 dwell time per AOI of a participant;
- 240 - *Global enhancement preference* as the odds of selecting an enhanced version over the
241 original²⁰: $\frac{P_{enhanced}}{(1-P_{enhanced})} = \frac{(P_{low} + P_{medium} + P_{high})}{P_{original}}$, greater than 1 when enhanced versions
242 were preferred over the original, equal to 1 when there was no preference and lower than 1
243 when the original version was mainly chosen;
- 244 - *Subjective enhancement preference* as the rank on the subjective enhancement preference
245 scale calculated from the feedback analysis.

246

247 Oculomotor behaviors were declined in four variables:

- 248 - *Visual attention delay* as the global mean dwell time per AOI;
- 249 - *Exploration time* as the global mean delay between stimuli display and preference decision;
- 250 - *Exploration velocity* as the global mean scanpath efficiency in pixels per second;
- 251 - *Total revisits* as the sum of fixations towards an AOI already targeted during a trial.

252

253 Altogether, 17 variables were submitted to PCA using SPSS. Principal components were retained only
254 when the eigenvalue was greater than 1. Independent variables were considered as relevant if the
255 absolute value of factor loading exceeded 0.5. The factor loading indicates whether the variable
256 within the PC is correlated positively or negatively. Each PC was identified and labeled through an
257 overview of the various classifications of variables (*visual assessment, I.T. experience, oculomotor*
258 *behavior and preference measures*) and the nature of their interactions. Individual factor scores from
259 each PC were extracted for each participant and used as behavioral covariates in the group
260 comparison analysis. For each PC, a one-way analysis of variance (ANOVA) was applied to calculate
261 differences in factor scores across the three groups. The Bonferroni adjustment for type 1 errors was
262 used for post-hoc multiple comparisons.

263 Results

264 This study included 49 participants (n=16 young controls, n=17 age-matched controls and n=16
265 glaucoma patients). The global demographic characteristics and I.T. use measures are displayed in
266 **Table 2**. Age difference was statistically validated. Contrast sensitivity in glaucoma patients was
267 significantly lower than in the two control groups ($P < 0.001$ Kruskal-Wallis test and $P < 0.001$
268 Bonferroni pairwise comparison). Among the participants, no significant difference was found
269 between the three groups regarding the frequency of computer use. Main group effects explained by
270 significant differences between young controls and glaucoma patients were found in the percentage
271 of tablet users and I.T. difficulty score due to vision ($P < 0.05$ Kruskal-Wallis test and $P < 0.05$
272 Bonferroni pairwise comparison).

273 *Single variable analyses*

274 **Did the different groups have specific enhancement preferences?**

275 The behavioral preferences ($P_{original}$, P_{low} , P_{medium} , P_{high} manually selected) were averaged by group
276 (**Figure 2A**): a main group effect was found for $P_{original}$ and P_{high} (respectively Kruskal-Wallis; $P =$
277 0.008 and $P = 0.001$). On average, young controls selected more original and fewer high-

278 enhancement versions than age-matched controls (Welch t-test; P -values < 0.05) and glaucoma
279 patients (respectively, Welch t-test P -value = .002 and .001). No behavioral preference differences
280 were found for low- and medium-enhancement versions ($P > 0.05$; Kruskal-Wallis). No significant
281 differences were found for enhancement preference between the various glaucoma stages (**Figure**
282 **2B**, $P > 0.05$; Kruskal-Wallis).

283 *[Insert Figure 2: Group comparison of enhancement preference]*

284 **Determining the effect of visual impairments on enhancement preference and oculomotor behavior**

285 Significant correlations were found across all participants between contrast sensitivity and global
286 enhancement preference (**Figure 3A**, Spearman's $\rho = -0.43$; $P = .002$), as well as between contrast
287 sensitivity and exploration velocity (**Figure 3B**, Spearman's $\rho = 0.43$; $P = .002$). However, no
288 significant correlations were found within the patient group between Mean Deviation (MD) and the
289 main variables, global enhancement (**Figure 3C**, Spearman's $\rho = -.32$; $P = .12$) and exploration
290 velocity (**Figure 3D**, Spearman's $\rho = 0.05$; $P = .42$).

291 *[Insert Figure 3: Behavioral and oculomotor data correlated to visual impairment]*

292 *Principal Component Analysis*

293 **Is there a link between the overall experimental data (objective/subjective) and clinical measures?**

294 The PCA revealed three principal components accounting for 72% of the total variance of the data.
295 PC1 involved nine variables and accounted for 31% of the total variance, PC2 involved four variables
296 and accounted for 22.7% of the total variance and PC3 involved four variables and accounted for
297 18.3% of the total variance. Principal components are detailed in **Fig. 4A** and compared by group in
298 **Fig.4B**.

299 *[Insert Figure 4: Principal Component Analysis]*

300 The first principal component (PC1) contained six preference measures in favor of enhancement
301 preference: positive factor loading for global and subjective enhancement preference, percentage
302 chosen and visual-oriented preference of high enhancement version, correlated with negative factor
303 loading of percentage chosen and visual-oriented preference of original versions. PC1 also
304 represented visual impairment with a negative factor loading of contrast sensitivity and exploration
305 velocity, and a positive factor loading on the I.T. difficulty due to vision score. PC1 was labeled
306 “*Graphical Interface Enhancement Needed*” and showed a main group effect ($F = 12.16, P < 0.001$). As
307 shown in **Fig. 4B**, this component was significantly lower in the young control group compared to the
308 age-matched group ($P = 0.02$) and the glaucoma patients ($P < 0.001$). Glaucoma patients presented a
309 higher mean factor score than age-matched controls (statistical tendency; $P = 0.11$ Bonferroni post-
310 hoc correction, $P = 0.06$ on a t-test comparison). The second principal component (PC2) only
311 contained preference measures with positive correlations between the percentages of *low* and
312 *medium enhancement* versions and the visual-oriented attention towards them. PC2 was labeled
313 “*Average Stimuli Preference*” and no main group effect was found ($F = 1.14, P = 0.34$). Results are
314 illustrated in **Fig. 4B**. The third principal component (PC3) included four visual measures: *clinical*
315 *visual deficit* score, *total* number of *revisits* on a computer screen version, *visual attention delay* and
316 *exploration time*, which were positively correlated. PC3 was labeled “*Longer oculomotor behavior*”
317 and demonstrated a main group effect ($F = 13.40, P < 0.001$). As illustrated in **Fig. 4B**, the mean factor
318 score of this component was significantly higher in glaucoma patients than in aged-matched controls
319 ($P = 0.002$) and young controls ($P < 0.001$).

320 Discussion

321 Information Technology is increasingly present in everyday life activities and requires both low-level
322 and high-level visual functions. Gathering clinical, behavioral, oculomotor and subjective data, this
323 exploratory study was designed to provide an overall view of the impact of glaucoma on I.T.
324 experience and the possible enhancement of visual comfort during computer use. This multifactorial

325 experiment evaluated the computer graphical interface enhancement preferences of patients with
326 glaucoma, age-matched controls and young controls, along with oculomotor behavior monitored by
327 eye-tracking. Using a principal component analysis, our statistical approach was innovative and
328 demonstrated correlations between subjective and objective enhancement preferences and specific
329 visual impairment.

330 *I.T. experience*

331 In this study, the socio-demographical data collected during the I.T. experience questionnaire
332 demonstrated that glaucoma patients, age-matched controls and young controls presented the same
333 rate of computer and smartphone use, despite the generation gap. Interestingly, the main difference
334 between the devices appeared to be the use of tablets, with a higher percentage of users within the
335 glaucoma group (62% of tablet users reported, compared to 29% among age-matched controls and
336 13% among young controls, **Table 2**). This suggests that while computers and smartphones could be
337 sufficient for controls, an in-between device with a larger screen could be more suitable for low-
338 vision patients. Moreover, the difficulty score, computed from five items based on NEIVFQ and
339 adapted to our I.T. experience questionnaire, was significantly different between the groups, and
340 glaucoma patients presented the highest mean score. In a highly digitized society, these results
341 emphasize the importance of including I.T.-related questions when assessing the quality of life of
342 glaucoma patients, even in the elderly.

343 *Enhancement Preference*

344 Individual variable analysis highlighted that while young controls mostly selected original computer
345 scenes, age-matched controls and glaucoma patients preferred enhanced stimuli. High enhancement
346 versions were mostly avoided by young controls and preferred by older participants. The absence of
347 significant difference between glaucoma patients and age-matched patients revealed that CGI
348 enhancement could generally benefit elderly people, with or without visual impairment. This is in line
349 with previous work demonstrating that normal aging involves a natural decline in visual functions²¹

350 and more specifically in low-level vision.^{22,23} Therefore, boosting physical parameters of computer
351 scenes, such as contrast and color, could also create a more suitable digital environment for elderly
352 people. Regarding the exploration of the impact of glaucoma stages, no differences in enhancement
353 preferences were found between early, moderate and severe glaucoma. However, in severe
354 glaucoma, the tendency to prefer gradually higher enhancement underlines a possible link between
355 the degree of vision loss and the necessity for increased CGIs. In line with this hypothesis, our data
356 demonstrate that reduced contrast sensitivity is correlated with a stronger preference for enhanced
357 computer scenes. Further investigations with larger cohorts of various glaucoma stages could
358 statistically validate the correlation between the degree of enhancement needed and the severity of
359 visual impairment.

360 *Principal component analysis and oculomotor behavior*

361 In the present study, PCA was a robust tool to establish the global links between objective,
362 subjective, experimental and clinical data. The first principal component revealed a strong
363 correlation between visual impairment and enhancement preference (objective and subjective), in
364 favor of high enhancement. A group comparison highlighted this for the “*Graphical Interface*
365 *Enhancement Needed*” component: young controls were not concerned, age-matched controls were
366 neutral and glaucoma patients largely fulfilled this profile. This result shows that on the basis of
367 everyday life reports, subjective feedback and experimental measures, glaucoma patients presented
368 a greater need for CGI enhancement than the visually healthy groups. PC2, labeled “*Average Stimuli*
369 *Preference*” demonstrated that low and medium enhancement preference was equally shared
370 between the groups. Targeting a central vision-biased enhancement in the construction of stimuli,
371 the results of PC1 and PC2 indicate that low and medium enhancement could suit all kinds of
372 populations, whereas high enhancement seems to be better suited to low-vision patients.

373 The third principal component demonstrated that glaucoma patients presented significantly longer
374 oculomotor behavior to explore ecological computer scenes. PC3, labeled “*Longer oculomotor*

375 *behavior*”, gathered a longer exploration time and visual attention delay by stimuli, as well as a
376 higher number of revisits: stimuli that had already been scanned needed to be explored again. Our
377 results suggest that these compensated oculomotor behaviors are more frequent in glaucoma
378 patients than in visually healthy groups, and that they are correlated with higher clinical visual deficit.
379 To assess whether the longer performances could be linked to a specific deficit, we computed the
380 correlation between participants’ exploration velocity and contrast sensitivity, and found that
381 reduced contrast sensitivity was associated with a decreasing exploration velocity. Previous studies
382 have shown that glaucoma patients compensate their peripheral visual field loss with various eye-
383 movement strategies, which have been measured during the exploration of a traffic scene,²⁴ natural
384 TV watching²⁵ or monocularly during natural scene exploration within central vision on a computer
385 screen.¹² Various types of compensation have already been identified: physical change in eye
386 movement (i.e. angular difference between two saccades), higher rate of oculomotor behavior
387 compared to controls or longer delays of exploration to perform a task.²⁶ Our findings indicate that
388 glaucoma patients may use these oculomotor strategies binocularly when viewing ecological
389 computer scenes.

390 This study provides new understanding regarding the I.T. experience of patients with glaucoma and
391 that of the normal-aging population. However, some limitations must be acknowledged. First, we
392 measured computer experience on a 15.6-inch laptop, which could be small for participants who are
393 used to a desktop computer. Indeed, since patients seemed to prefer tablets to a smartphone, we
394 assume that a bigger screen and high resolution are important to improve their experience.
395 Nowadays, high quality monitors are more common, quite affordable and could already constitute an
396 option to enhance residual vision in glaucomatous users. Second, by measuring oculomotor behavior
397 on a preference task, the various time-dependent data observed could be confounded between
398 visual performance and time related to decision-making. Therefore, we explored multiple reliable
399 assessments by adding oculomotor measures standardized by the distance scanned on the screen
400 (i.e. exploration velocity [px/s]) or by considering variables without a time dimension (i.e. total

401 amount of revisits on a stimulus). Third, while the comparison between the different glaucoma
402 stages is an important focus to better understand the development of specific visual enhancement
403 (adapted to the degree of vision loss), the present data set was limited by the sample size of the
404 subgroup (early, moderate, severe) of glaucomatous patients. Finally, enhancement preference may
405 be quite different from the effective enhancement of CGI adapted to patients with glaucoma. Further
406 investigations are needed to establish an objective link between enhancement and the possible
407 improvement of patients on specific computer tasks involving their visuo-spatial capacities.

408 Overall, we conclude from the subjective reports of glaucomatous patients that I.T. experience and
409 the difficulty to use digital devices due to vision issues are significant measures to consider in visual-
410 related quality of life questionnaires. Graphical enhancement generally made ecological computer
411 scenes more suitable for glaucoma patients and age-matched controls, while young controls showed
412 a distinct preference for the original versions. However, by using principal component analysis to
413 evaluate everyday life reports, subjective feedback and experimental data together, patients with
414 glaucoma demonstrated a greater need for computer graphical interface enhancement than the
415 visually healthy groups. Moreover, contrast sensitivity, which was significantly reduced in glaucoma
416 patients, was a central measure to understand enhancement preference variations and differences in
417 oculomotor behavior. Glaucoma seems to affect the global exploration of a computer screen and
418 involves a strategic change in eye movements with longer oculomotor behavior. Finally, our findings
419 provide better understanding of the impact of glaucomatous visual loss on everyday activities, such
420 as computer tasks, and give important cues for developing adapted computer graphical interfaces for
421 such visual impairment.

422 References

- 423 1. Tham Y-C, Li X, Wong TY, Quigley HA, Aung T, Cheng C-Y. Global Prevalence of Glaucoma and
424 Projections of Glaucoma Burden through 2040. *Ophthalmology*. 2014;121(11):2081-2090.
425 doi:10.1016/j.ophtha.2014.05.013
- 426 2. Sippel K, Kasneci E, Aehling K, et al. Binocular glaucomatous visual field loss and its impact on
427 visual exploration - A supermarket study. *PLoS One*. 2014. doi:10.1371/journal.pone.0106089
- 428 3. Ramulu P. Glaucoma and disability: which tasks are affected, and at what stage of disease?
429 *Curr Opin Ophthalmol*. 2009;20(2):92-98. doi:10.1097/ICU.0b013e32832401a9
- 430 4. Lek JJ, Vingrys AJ, McKendrick AM. Rapid Contrast Adaptation in Glaucoma and in Aging.
431 *Investig Ophthalmology Vis Sci*. 2014;55(5):3171. doi:10.1167/iovs.13-13229
- 432 5. Bierings RAJM, Overkempe T, van Berkel CM, Kuiper M, Jansonius NM. Spatial contrast
433 sensitivity from star- to sunlight in healthy subjects and patients with glaucoma. *Vision Res*.
434 2019;158(March 2018):31-39. doi:10.1016/j.visres.2019.01.011
- 435 6. Ichhpujani P, Thakur S, Spaeth GL. Contrast Sensitivity and Glaucoma. *J Glaucoma*.
436 2020;29(1):71-75. doi:10.1097/IJG.0000000000001379
- 437 7. Enoch J, Jones L, Taylor DJ, et al. How do different lighting conditions affect the vision and
438 quality of life of people with glaucoma? A systematic review. *Eye*. 2020;34(1):138-154.
439 doi:10.1038/s41433-019-0679-5
- 440 8. Lenoble Q, Lek JJ, McKendrick AM. Visual object categorisation in people with glaucoma. *Br J*
441 *Ophthalmol*. 2016;100(11):1585-1590. doi:10.1136/bjophthalmol-2015-308289
- 442 9. Schafer A, Rouland JF, Peyrin C, Szaffarczyk S, Boucart M. Glaucoma Affects Viewing Distance
443 for Recognition of Sex and Facial Expression. *Investig Ophthalmology Vis Sci*. 2018;59(12):4921.
444 doi:10.1167/iovs.18-24875

- 445 10. Roux-Sibilon A, Rutgé F, Aptel F, et al. Scene and human face recognition in the central vision
446 of patients with glaucoma. Williams MA, ed. *PLoS One*. 2018;13(2).
447 doi:10.1371/journal.pone.0193465
- 448 11. Lamirel C, Milea D, Cochereau I, Duong M-H, Lorenceau J. Impaired Saccadic Eye Movement in
449 Primary Open-angle Glaucoma. *J Glaucoma*. 2014;23(1):23-32.
450 doi:10.1097/IJG.0b013e31825c10dc
- 451 12. Asfaw DS, Jones PR, Mönter VM, Smith ND, Crabb DP. Does Glaucoma Alter Eye Movements
452 When Viewing Images of Natural Scenes? A Between-Eye Study. *Investig Ophthalmology Vis
453 Sci*. 2018;59(8):3189. doi:10.1167/iovs.18-23779
- 454 13. Moshtael H, Aslam T, Underwood I, Dhillon B. High Tech Aids Low Vision: A Review of Image
455 Processing for the Visually Impaired. *Transl Vis Sci Technol*. 2015;4(4):6. doi:10.1167/tvst.4.4.6
- 456 14. Al-Atabany WI, Tong T, Degenaar PA. Improved content aware scene retargeting for retinitis
457 pigmentosa patients. *Biomed Eng Online*. 2010;9(1):52. doi:10.1186/1475-925X-9-52
- 458 15. Watson LM, Strang NC, Scobie F, Love GD, Seidel D, Manahilov V. Image Jitter Enhances Visual
459 Performance when Spatial Resolution Is Impaired. *Investig Ophthalmology Vis Sci*.
460 2012;53(10):6004. doi:10.1167/iovs.11-9157
- 461 16. Mangione CM, Lee PP, Gutierrez PR, et al. Development of the 25-item National Eye Institute
462 Visual Function Questionnaire. *Arch Ophthalmol (Chicago, Ill 1960)*. 2001;119(7):1050-1058.
- 463 17. Abe RY, Diniz-Filho A, Costa VP, Gracitelli CPB, Baig S, Medeiros FA. The Impact of Location of
464 Progressive Visual Field Loss on Longitudinal Changes in Quality of Life of Patients with
465 Glaucoma. *Ophthalmology*. 2016;123(3):552-557. doi:10.1016/j.ophtha.2015.10.046
- 466 18. Riva I, Legramandi L, Rulli E, et al. Vision-related quality of life and symptom perception
467 change over time in newly-diagnosed primary open angle glaucoma patients. *Sci Rep*.
468 2019;9(1):6735. doi:10.1038/s41598-019-43203-9

- 469 19. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state" A practical method for grading the
470 cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12(3):189-198.
471 doi:10.1016/0022-3956(75)90026-6
- 472 20. Kwon M, Ramachandra C, Satgunam P, Mel BW, Peli E, Tjan BS. Contour Enhancement
473 Benefits Older Adults with Simulated Central Field Loss. *Optom Vis Sci.* 2012;89(9):1374-1384.
474 doi:10.1097/OPX.0b013e3182678e52
- 475 21. Owsley C. Aging and vision. *Vision Res.* 2011;51(13):1610-1622.
476 doi:10.1016/j.visres.2010.10.020
- 477 22. Lenoble Q, Bordaberry P, Rougier M-B, Boucart M, Delord S. Influence of Visual Deficits on
478 Object Categorization in Normal Aging. *Exp Aging Res.* 2013;39(2):145-161.
479 doi:10.1080/0361073X.2013.761910
- 480 23. McKendrick AM, Sampson GP, Walland MJ, Badcock DR. Contrast Sensitivity Changes Due to
481 Glaucoma and Normal Aging: Low-Spatial-Frequency Losses in Both Magnocellular and
482 Parvocellular Pathways. *Investig Ophthalmology Vis Sci.* 2007;48(5):2115. doi:10.1167/iovs.06-
483 1208
- 484 24. Crabb DP, Smith ND, Rauscher FG, et al. Exploring Eye Movements in Patients with Glaucoma
485 When Viewing a Driving Scene. He S, ed. *PLoS One.* 2010;5(3):e9710.
486 doi:10.1371/journal.pone.0009710
- 487 25. Crabb DP, Smith ND, Zhu H. What's on TV? Detecting age-related neurodegenerative eye
488 disease using eye movement scanpaths. *Front Aging Neurosci.* 2014;6:312.
489 doi:10.3389/fnagi.2014.00312
- 490 26. Lenoble Q, Rouland J-F. Context Association in Glaucoma Patients Using a Touch Screen. *J*
491 *Glaucoma.* 2019;28(8):737-743. doi:10.1097/IJG.0000000000001297

492

493 Figure Legends

494 **Figure 1: Experimental design of one trial of the preference task**

495 [upper left] – original computer scene, [upper right] – high enhancement, [lower right] – low enhancement and [lower left] –
496 medium enhancement.

497 **Figure 2: Group comparison of enhancement preference**

498 [A] for young controls (blue), age-matched controls (orange) and glaucoma patients (gray); [B] for dissociated glaucoma
499 stage: early (light grey), moderate (dark grey) and severe (black dotted line). ** P-value < 0.01, * P < 0.05

500 **Figure 3: Behavioral and oculomotor data correlated to visual impairment**


501 Scatterplots with fitted regression lines illustrating the relationship between: [A] **contrast sensitivity and the global**
502 **enhancement preference** (1: no preference, 1 < original version preference, > 1: enhanced version preference, > 10: only
503 few original versions selected); [B] **contrast sensitivity and exploration velocity** (scanpath in pixels per sec); [C] **MD and**
504 **Global enhancement preference**; [D] **MD and Exploration velocity**.

505 The Spearman correlation coefficient (r) is plotted on the two graphs (top-left): ** P-value < 0.01


506 **Figure 4: Principal Components Analysis**

507 [A] **Principal components compositions resulting from PCA**. Variables gathered within a principal component retain the
508 optimized rate of variation present in the overall data set. One column represents one principal component. Factor loading
509 scores illustrate the nature of the relation between the variables within a principal component: either negative (factor
510 loading < - 0.5; from light to dark blue) or positive (factor loading > 0.5; from light to dark red) correlations. [B] **Group**
511 **comparison of principal component mean factor score**: mean comparison of PC1, PC2 and PC3 factor scores. Positive or
512 negative factor scores mean that a group is respectively congruent or incongruent with the principal component. The three
513 groups are represented from left to right in each component: young controls (blue), age-matched controls (orange) and
514 glaucoma patients (grey). * P-value < 0.05 ** P < 0.01 *** P < 0.001

Original




High




+


Medium




Low








A. Group Comparison (n=49)







B. Glaucoma Stage Comparison (n=16)







| Classification of Variables | | PC1 | PC2 | PC3 |
|--|--|---|--|--|
| | | “Graphical Interface Enhancement Needed” | “Average Stimuli Preference” | “Longer oculomotor behavior” |
| Visual Assessment  | | Contrast sensitivity | | Clinical visual deficit |
| Oculomotor Behaviour  | | Exploration Velocity | | Total revisits Visual attention delay Exploration time |
| Preference Measures  | | Global enhancement preference Subjective enhancement preference $P_{high\ enhancement}$ Visual-oriented preference [High] $P_{original}$ Visual-oriented preference [Original] | $P_{low\ enhancement}$ Visual-oriented preference [Low] $P_{medium\ enhancement}$ Visual-oriented preference [Medium] | |
| I.T. Experience  | | | | Difficulty score due to vision |





Table 1 - Demographic and clinical data of participants with POAG

| <i>Participants with POAG</i> | <i>Gender</i> | <i>Age (years)</i> | <i>Binocular Visual Acuity (log MAR)</i> | <i>Worst Eye HFA [30 -2] MD (dB)</i> | <i>Extent [MO: Monocular BO: Binocular]</i> | <i>Glaucoma stage</i> | <i>Contrast sensitivity (log CS)</i> |
|-------------------------------|---------------|--------------------|--|--------------------------------------|---|-----------------------|--------------------------------------|
| G1 | M | 64 | ≤ 0.2 | -9.79 | MO | Moderate | 1.65 |
| G2 | F | 67 | ≤ 0.2 | -10.74 | BO | Moderate | 1.65 |
| G3 | F | 57 | 0 | -5.74 | BO | Early | 1.95 |
| G4 | F | 68 | - 0.1 | -24.98 | MO | Severe | 1.65 |
| G5 | M | 58 | ≤ 0.2 | -28.18 | MO | Severe | 1.35 |
| G6 | M | 59 | 0 | -6.13 | MO | Moderate | 1.80 |
| G7 | F | 63 | 0 | -2.03 | BO | Early | 1.65 |
| G8 | F | 63 | 0.1 | -9.23 | MO | Moderate | 1.65 |
| G9 | M | 62 | 0.1 | -6.42 | MO | Moderate | 1.65 |
| G10 | F | 69 | 0.2 | -7.13 | BO | Moderate | 1.50 |
| G11 | F | 73 | 0.1 | -4.95 | BO | Early | 1.80 |
| G12 | F | 66 | 0.1 | -15.92 | MO | Severe | 1.35 |
| G13 | M | 65 | ≤ 0.2 | -18.80 | BO | Severe | 1.50 |
| G14 | M | 50 | 0.1 | -4.63 | MO | Early | 1.80 |
| G15 | M | 61 | ≤ 0.2 | -13.12 | BO | Severe | 1.95 |
| G16 | F | 58 | 0.1 | -4.56 | BO | Early | 1.65 |

F, female; M, male.

Table 1- Contrast sensitivity and I.T. experience demographic data

| CHARACTERISTICS | YOUNG (N=16) | AGE- MATCHED (N=17) | GLAUCOMA PATIENTS (N=16) |
|---|----------------------------|---------------------------|--------------------------------|
| GENDER (F/M) | 8/8 | 9/8 | 9/7 |
| AGE (*) MEAN ± SD (RANGE) | 23.3*** ± 2.1 (20 – 28) | 59.1 ± 8.3 (46 – 73) | 62.7 ± 5.6 (50 – 73) |
| CONTRAST SENSITIVITY (*) MEAN ± SD | 1.93 ± 0.05 | 1.91 ± 0.1 | 1.66*** ± 0.2 |
| % OF COMPUTER USERS | 100 % | 94 % | 94 % |
| SMARTPHONE USERS | 100 % | 100 % | 100 % |
| TABLET USERS (*) | 13 % | 29 % | 62 % |
| COMPUTER USE FREQUENCY MEAN RATE ± SD (1: ONCE A DAY. 2: ONCE A WEEK. 3: ONCE A MONTH. 4 : ALMOST NEVER. 5: USE STOPPED BECAUSE OF VISUAL DIFFICULTIES) | 1.06 ± 0.25 | 1.06 ± 0.25 | 1.33 ± 0.81 |
| ESTIMATED USER LEVEL MEAN RATE ± SD (1: EXPERT. 2: FAMILIAR. 3: NORMAL. 4: BEGINNER. 5: NO SKILLS) | 1.9 ± 0.44 | 2.25 ± 0.57 | 2.5 ± 0.89 |
| I.T. DIFFICULTY SCORE DUE TO VISION (*) MEAN RATE ± SD (1:NOT AT ALL. 2: A LITTLE. 3: MODERATELY. 4: A LOT. 5: USE STOPPED BECAUSE OF VISUAL DIFFICULTIES) | 1.05 ± 0.11 | 1.16 ± 0.21 | 1.35 ± 0.50 |

(*) P-value < .05 group effect explained at least by a significant difference between young controls group and glaucoma patients; *** P < .001 differ significantly from the two other groups