



HAL
open science

Ethanolamine: A novel anti-aging agent

Patrick Rockenfeller, Didac Carmona-Gutierrez, Federico Pietrocola, Guido Kroemer, Frank Madeo

► **To cite this version:**

Patrick Rockenfeller, Didac Carmona-Gutierrez, Federico Pietrocola, Guido Kroemer, Frank Madeo. Ethanolamine: A novel anti-aging agent. *Molecular & Cellular Oncology*, 2016, 3 (1), pp.e1019023. 10.1080/23723556.2015.1019023 . hal-04702786

HAL Id: hal-04702786

<https://hal.science/hal-04702786v1>

Submitted on 19 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Ethanolamine: A novel anti-aging agent

Patrick Rockenfeller^{1,*}, Didac Carmona-Gutierrez¹, Federico Pietrocola^{2,3,4}, Guido Kroemer^{2,3,4,5,6,7}, and Frank Madeo^{1,8,*}

¹Institute of Molecular Biosciences; NAWI Graz, University of Graz; Graz, Austria; ²Equipe 11 labellisée par la Ligue contre le Cancer; Centre de Recherche des Cordeliers; Paris, France; ³Cell Biology and Metabolomics Platforms; Gustave Roussy Comprehensive Cancer Center; Villejuif, France; ⁴INSERM; U1138; Paris, France; ⁵Université Paris Descartes; Sorbonne Paris Cité; Paris, France; ⁶Université Pierre et Marie Curie, Paris, France; ⁷Pôle de Biologie; Hôpital Européen Georges Pompidou; AP-HP; Paris, France; ⁸BioTechMed-Graz; Graz, Austria

Keywords: aging, autophagy, ethanolamine, phosphatidylethanolamine, phosphatidylserine decarboxylase

Ethanolamine (Etn) is a naturally occurring aminoalcohol necessary for synthesis of the phospholipid phosphatidylethanolamine (PE), a major component of biological membranes. We recently reported that Etn treatment increases cellular PE levels, thereby inducing cytoprotective autophagy and protecting against aging across species.

Phosphatidylethanolamine (PE) is a central intermediate of lipid metabolism and a major component of biological membranes.^{1,2} Within cellular membranes, PE not only serves as a structural phospholipid but also regulates the tethering of proteins and fusion processes.³ PE orchestrates the interaction with membrane-associated proteins, both by directly ingrain proteins into membranes and as a precursor for glycosylphosphatidylinositol (GPI) anchors.^{3,4} Importantly, PE is also directly involved in the process of macroautophagy (hereafter termed autophagy), a lysosome-dependent cellular recycling mechanism⁵ that protects cells against lethal stress and extends longevity in model organisms.⁶ During autophagy, double-membraned structures that are highly abundant in PE engulf superfluous, supernumerary, or dysfunctional macromolecules or organelles contained in the cytoplasm, forming vesicles (autophagosomes). These autophagosomes then fuse with lysosomes to generate autophagolysosomes in which the luminal cargo is degraded. Importantly, PE serves as a lipid anchor for a protein essential for several steps of the autophagic process that is

called autophagy related protein 8 (Atg8) in yeast or microtubule-associated protein 1A/1B-light chain 3 (LC3) in mammalian cells.⁷ The importance of LC3 (or Atg8) in the autophagic process is reflected by the fact that many assays designed to measure autophagy rely on quantitation of the conjugation of LC3 to PE. This lipidation step increases the electrophoretic mobility of LC3 (which can be measured by immunoblot analysis) and causes its redistribution from a diffuse cytosolic pattern to so-called autophagic puncta (which can be visualized by immunofluorescence or by tagging LC3 with green fluorescent protein).

Given the widespread functions of PE as a precursor of several biosynthetic pathways, there is high demand for this metabolite (Fig. 1). A common PE pool feeds into all major cellular PE-consuming pathways, thus resulting in competition for PE between pathways.⁴ As we have recently shown, this limitation can be overcome by genetic or pharmacological interventions. Overexpression of yeast phosphatidylserine decarboxylase 1 (Psd1), which catalyzes the step from phosphatidylserine to PE,

increased the levels of intracellular PE. Similarly, external administration of ethanolamine (Etn), a precursor of PE (Fig. 1) can increase the abundance of intracellular PE.⁸ Supporting a crucial regulatory role for PE in autophagy, we observed that both external supply of PE and an increase in its internal generation similarly increased autophagic flux.⁸ Stimulation of autophagy by PE could efficiently protect cells from necrotic or apoptotic cell death.^{6,9} This is an important finding because unwarranted cell death may contribute to several age-related diseases including stroke, atherosclerosis, and cardiovascular disease, or neurodegenerative disorders such as Alzheimer's, Parkinson's, or Huntington's disease. Importantly, pharmacological Etn treatment extended the lifespan of yeast and fruit flies, as well as cultured mammalian cells,⁸ underlining the potential of Etn as a potent autophagy and longevity drug.

Given its properties as a detergent and emulsifier, Etn is widely used in cosmetic, lifestyle, and household products such as hair colors, bleaches, washing powders, disinfectants, kitchen

© Patrick Rockenfeller, Didac Carmona-Gutierrez, Federico Pietrocola, Guido Kroemer, and Frank Madeo

*Correspondence to: Patrick Rockenfeller; Email: patrick.rockenfeller@uni-graz.at, Frank Madeo; Email: frank.madeo@uni-graz.at

Submitted: 02/05/2015; Revised: 02/06/2015; Accepted: 02/09/2015

<http://dx.doi.org/10.1080/23723556.2015.1019023>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The moral rights of the named author(s) have been asserted.




Figure 1. The 'PE hourglass'. Simplified schematic of the conversion of ethanolamine (Etn) into phosphatidylethanolamine (PE) and its downstream products. Alternative pathways for PE production were omitted for clarity. PE synthesis is rate limiting because newly synthesized PE feeds into a common cellular PE pool, which is used for diverse cellular processes including glycosylphosphatidylinositol (GPI) anchor, membrane, and phosphatidylcholine (PC) synthesis as well as the generation of autophagic membranes. ADP, adenosine diphosphate; ATP, adenosine triphosphate; CMP, cytosine monophosphate; Etn, ethanolamine; CTP, cytosine triphosphate; DAG, diacylglycerol; P, phosphate; PE, phosphatidylethanolamine; PP, pyrophosphate.

cleaners, shampoos, and soaps. Thus, its toxicologic profile as an external agent

that may irritate the skin and mucosa is well studied. However, it remains to be

determined whether Etn can be used internally within the scope of increasing organismal PE levels, thereby stimulating autophagy and possibly conferring health benefits.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Funding

This work has been supported by Fonds zur Förderung der Wissenschaftlichen Forschung in Österreich (FWF), projects SFB Lipotox F3005-B12, P23490-B12 and P24381-B20 to F.M. The authors gratefully acknowledge support from NAWI Graz and Bio-TechMed. GK is supported by the Ligue contre le Cancer (équipe labélisée); Agence Nationale de la Recherche sur le cancer (ARC); Cancéropôle Ile-de-France; Institut National du Cancer (INCa); Fondation Bettencourt-Schueler; Fondation de France; Fondation pour la Recherche Médicale (FRM); the European Commission (ArtForce); the European Research Council (ERC); the LabEx Immuno-Oncology; the SIRIC Stratified Oncology Cell DNA Repair and Tumor Immune Elimination (SOCRATE); the SIRIC Cancer Research and Personalized Medicine (CARPEM); and the Paris Alliance of Cancer Research Institutes (PACRI).

References

- Bürgermeister M, Birner-Grünberger R, Nebauer R, Daum G. Contribution of different pathways to the supply of phosphatidylethanolamine and phosphatidylcholine to mitochondrial membranes of the yeast *Saccharomyces cerevisiae*. *Biochim Biophys Acta* 2004; 1686:161-8; PMID:15522832; <http://dx.doi.org/10.1016/j.bbali.2004.09.007>
- Vance JE. Thematic Review Series: Glycerolipids. Phosphatidylserine and phosphatidylethanolamine in mammalian cells: two metabolically related aminophospholipids. *J Lipid Res* 2008; 49:1377-87; PMID:18204094; <http://dx.doi.org/10.1194/jlr.R700020-JLR200>
- Van den Brink-van der Laan E, Antoinette Killian J, de Kruijff B. Nonbilayer lipids affect peripheral and integral membrane proteins via changes in the lateral pressure profile. *Biochim Biophys Acta BBA - Biomembr* 2004; 1666:275-88; PMID:15519321; <http://dx.doi.org/10.1016/j.bbame.2004.06.010>
- Wilson-Zbinden C, Santos AX da S dos, Stoffel-Studer I, van der Vaart A, Hofmann K, Reggiori F, Riezman H, Kraft C, Peter M. Autophagy Competes for a Common Phosphatidylethanolamine Pool with Major Cellular PE-Consuming Pathways in *Saccharomyces cerevisiae*. *Genetics* 2014; genetics.114.169797
- Feng Y, He D, Yao Z, Klionsky DJ. The machinery of macroautophagy. *Cell Res* 2014; 24:24-41; PMID:24366339; <http://dx.doi.org/10.1038/cr.2013.168>
- Madeo F, Tavernarakis N, Kroemer G. Can autophagy promote longevity? *Nat Cell Biol* 2010; 12:842-6; PMID:20811357; <http://dx.doi.org/10.1038/ncb0910-842>
- Ichimura Y, Kirisako T, Takao T, Satomi Y, Shimoniishi Y, Ishihara N, Mizushima N, Tanida I, Kominami E, Ohsumi M, et al. A ubiquitin-like system mediates protein lipidation. *Nature* 2000; 408:488-92; PMID:11100732; <http://dx.doi.org/10.1038/35044114>
- Rockenfeller P, Koska M, Pietroccola F, Minois N, Knittelfelder O, Sica V, Franz J, Carmona-Gutierrez D, Kroemer G, Madeo F. Phosphatidylethanolamine positively regulates autophagy and longevity. *Cell Death Differ* [Internet] 2015; 22:499-508; [cited 2015 Jan 21]; Available from: <http://www.nature.com/cdd/journal/vaop/ncurrent/full/cdd2014219a.html>
- Eisenberg T, Knauer H, Schauer A, Büttner S, Ruckstuhl C, Carmona-Gutierrez D, Ring J, Schroeder S, Magnes C, Antonacci L, et al. Induction of autophagy by spermidine promotes longevity. *Nat Cell Biol* 2009; 11:1305-14; PMID:19801973; <http://dx.doi.org/10.1038/ncb1975>