



## Quantum effects in sonochemistry

Sergey I. Nikitenko, Sara El Hakim, Tony Chave

### ► To cite this version:

| Sergey I. Nikitenko, Sara El Hakim, Tony Chave. Quantum effects in sonochemistry. ESS18 - 18th Meeting of the European Society of Sonochemistry, May 2024, Louvain, Belgium. hal-04705994

HAL Id: hal-04705994

<https://hal.science/hal-04705994v1>

Submitted on 23 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Magnetic Nitinol: New Efficient Photothermal Catalyst

*Sergey Nikitenko, Sara El Hakim, Tony Chave*

ICSM, Marcoule, France



[serguei.nikitenko@cea.fr](mailto:serguei.nikitenko@cea.fr)

[www.icsm.fr](http://www.icsm.fr)




August 10-12, 2023, UCLA

# Ascending photothermal catalysis



- *Introducing heat into photocatalytic processes can significantly improve the photoconversion efficiency*
  
- *Self-heating of the photocatalytic systems can reduce energy consumption required for the photothermal processes*



❖ *Preparation of cost-effective photothermal catalysts from earth-abundant elements is of prime importance for heterogeneous catalysis driven by concentrated solar light*


# Nitinol as a photothermal catalyst?




Nitinol, NiTi, is a famous shape-memory alloy widely used in medicine, aerospace industry, instrumentation etc.

But what about the catalysis with NiTi?

NiTi NPs



 AMERICAN  
ELEMENTS  
THE ADVANCED MATERIALS MANUFACTURER®



Particle size 20-100 nm,  
≈ 2-4 nm passivation layer:  $[\text{TiO}_2] > [\text{NiO}]$  (XPS data)



Nitinol in hot water


# Preparation of core-shell-satellite NiTi@TiO<sub>2</sub>/Ni nanoparticles



- ✓ Sonohydrothermal (SHT) treatment of pristine NiTi NPs in pure water (200°C, autogenic pressure P= 14 bar, US: 20 kHz,  $P_{ac}=17$  W, t = 3 h)



Rietveld refinement of the PXRD data

Sample	mol % ( $\pm 10\%$ )				
	NiTi cubic	NiTiI monoclinic	Ni <sub>3</sub> Ti	Ni	TiO <sub>2</sub>
NiTi	91	5	4	—	—
NiTi@TiO <sub>2</sub> /Ni	31	—	<1	35	34



The SHT treatment leads to the appearance of metallic Ni and anatase TiO<sub>2</sub>, indicating non-congruent oxidation of nitinol in hydrothermal water


# HRTEM/EDX images of NiTi@TiO<sub>2</sub>/Ni NPs



Ultra-micro tomographic image/EDX mapping of NiTi@TiO<sub>2</sub>/Ni NPs, Ni (blue) and Ti (green)

Ni NPs ( $\approx 20$  nm) are attached onto TiO<sub>2</sub> shell forming core-shell-satellite structure


# Magnetic properties



*Magnetization curves for NiTi and NiTi@TiO<sub>2</sub>/Ni NPs measured at room temperature*

- Pristine NiTi NPs show very low magnetization
- NiTi@TiO<sub>2</sub>/Ni NPs are ferromagnetic ( $M_s = 9 \text{ emu}\cdot\text{g}^{-1}$ )

# Optical spectra



*UV/vis/NIR absorption spectra of NiTi and NiTi@TiO<sub>2</sub>/Ni NPs (100 ppm) in water ( $l = 1 \text{ cm}$ )*

- ✓ NiTi: UV-NIR broad continuum of interband/intraband transitions
- ✓ NiTi@TiO<sub>2</sub>/Ni: peaks at 270 and 310 nm are ascribed to Ni SPR and TiO<sub>2</sub> bandgap

# Photothermal and photocatalytic experiments

- ✓ **Solutions:** 100 ppm of NiTi or NiTi@TiO<sub>2</sub>/Ni NPs dispersed in 1M Glycerol

## Experimental conditions:

- Glass-made continuos Ar-flow thermostated cell  
 $T = 30\text{-}100^\circ\text{C}$
- $V = 50 \text{ mL } (l = 3 \text{ cm})$
- Xe lamp:  
**400-1100 nm: 8.9 W**  
**300-420 nm: 0.6 W**
- Online mass spectrometric analysis of the outlet gases




# Light-to-heat conversion

100 ppm of NPs

Xe lamp:

400-1100 nm: 8.9 W

300-420 nm: 0.6 W




- ✓ NiTi and NiTi@TiO<sub>2</sub>/Ni NPs show strong and very similar heat generation capacity under light exposure indicating that self-heating is determined by the metallic core

# Photocatalysis under self-heating




100 ppm of catalyst, 1M Glyc, Xe lamp, steady-state temperature 75 °C



- Upon light exposure the concentration of  $H_2$ ,  $CH_4$ , and  $CO_2$  gradually increases with temperature rise reaching a steady state value simultaneously with that of temperature
- Heating at 75 °C under dark conditions does not lead to gaseous products formation indicating the photonic origin of the process
- CO is not formed**

# Photocatalysis under self-heating



- Global photocatalytic activity:  
 $\text{NiTi@TiO}_2/\text{Ni} >> \text{NiTi} \approx \text{TiO}_2$
- $\text{NiTi@TiO}_2/\text{Ni}$ :  $[\text{H}_2] >> [\text{CH}_4] > [\text{CO}_2]$
- $\text{NiTi}$  :  $[\text{CH}_4] > [\text{H}_2] \approx [\text{CO}_2]$
- $\text{TiO}_2$  :  $[\text{H}_2] > [\text{CO}_2] > [\text{CH}_4]$


Product	$E_a$ , kJ mol⁻¹
H <sub>2</sub>	19±2
CO <sub>2</sub>	16±2
CH <sub>4</sub>	59±6

At 30-90 °C kinetics follows Arrhenius law

$$E_a(\text{CH}_4) > E_a(\text{H}_2) \approx E_a(\text{CO}_2)$$

- First step: formation of H<sub>2</sub> and CO<sub>2</sub>
- Second step: CO<sub>2</sub> methanation

# H/D KIE in CH<sub>4</sub> formation




Equimolar H<sub>2</sub>O/D<sub>2</sub>O mixture:



- 5 isotopic species CH<sub>x</sub>D<sub>y</sub>, with domination of CH<sub>2</sub>D<sub>2</sub>
- H/D ratio in released methane is close to HDO in agreement with light-driven CO<sub>2</sub> methanation mechanism

# Suggested overall mechanism



# Concluding remarks



- ✓ “Magnetic nitinol” exhibits unique set of properties: strong capability of heat generation under light irradiation, good magnetization, remarkable thermally assisted photocatalytic activity, and high stability
- ✓ Noncongruent surface oxidation of nanoalloys paves the way to the preparation of new generation of catalysts with advanced photothermal properties

# Acknowledgements

*Xavier Le Goff – ICSM*

*Valérie Flaud – University of Montpellier*

*Corine Reibel – University of Montpellier*





# Thank you!