



HAL
open science

Ionizing radiation induces cross-linking of two noncovalently bound collagen mimetic peptide triple helices in the absence of a molecular environment

Marwa Abdelmouleh, Muhamed Amin, Mathieu Lalande, Thomas Schlathölter, Jean-Christophe Pouilly

► To cite this version:

Marwa Abdelmouleh, Muhamed Amin, Mathieu Lalande, Thomas Schlathölter, Jean-Christophe Pouilly. Ionizing radiation induces cross-linking of two noncovalently bound collagen mimetic peptide triple helices in the absence of a molecular environment. *Physical Chemistry Chemical Physics*, 2023, *Molecular Dynamics in the Gas Phase*, 25 (42), pp.29249-29256. 10.1039/D3CP03264G . hal-04707780

HAL Id: hal-04707780

<https://hal.science/hal-04707780v1>

Submitted on 24 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ARTICLE

Ionizing radiation induces cross-linking of two noncovalently-bound collagen mimetic peptide triple helices in the absence of molecular environment

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Marwa Abdelmouleh,^a Muhamed Amin,^{b,c} Mathieu Lalande,^a Thomas Schlathöler^{b,d} and Jean-Christophe Pouilly^{*a}

Cross-linking is a fundamental molecular process that is highly important for many applications, in particular to tune the properties of collagen-based biomaterials. Chemical reagents, the action of enzymes or physical factors such as heat or radiation can facilitate collagen cross-linking. Ionizing radiation has the advantage of being fast, efficient and free from potentially toxic reagents. Collagen cross-linking by ionizing radiation is thought to occur via a water-mediated pathway. In the past, synthesized collagen mimetic peptides have proven to be of great value for understanding the influence of the amino acid sequence on the stability of tertiary (fibrous) as well as secondary (triple helical) structures of collagen. Cross-linking of synthetic collagen mimetic peptides is often used for modifying the properties of biomaterials. In this work, for the first time we apply radiation-induced cross-linking to synthetic collagen mimetic peptides, and present an experimental and theoretical study of peptide hexamers consisting of two noncovalently-bound triple helices in the absence of molecular environment, i.e. in the gas phase. Our results show that X-ray photoabsorption of the hydroxylated hexamer leads to ionization and cross-linking of the two triple helices: thus, we find evidence that cross-linking can be achieved by ionizing radiation, without the presence of any reagent or water. We propose a cross-linking mechanism involving the creation of two radicals on hydroxyproline side-chains and their recombination, ultimately leading to covalent bond formation between the triple helices.

Introduction

The creation of bonds is at the heart of chemistry. The phenomenon of cross-linkage is due to covalent bond formation between atoms in a molecular system. These new bonds can lead to a modification of material properties. Cross-linking is therefore a key process in many fundamental domains such as polymer chemistry, material science, biochemistry, etc. This is particularly true for collagen-based biomaterials that are currently widely used for medical applications such as prostheses, wound healing, cartilage renewal and more.¹ Collagen is the most abundant protein in mammals, and the main constituent of the extracellular matrix of connective tissues such as skin, nails, cartilage, bones, etc. The advantages of collagen for designing biomaterials are thus

biocompatibility, biodegradability, availability and versatility. Furthermore, collagen is naturally cross-linked in tissues, which is crucial for their mechanical properties. Artificial cross-linking of collagen proteins for biomaterials can be achieved by chemical reagents such as glutaraldehyde, genipin or chitosan, by physical factors such as heat or UV light, or in biological processes, involving for instance the microbial transglutaminase enzyme.² Ionizing radiation also has the ability to cross-link collagen, as demonstrated by Hu *et al.*³ for irradiation of collagen thin films by X-rays, gamma-rays and ions, and by Zhang *et al.*⁴ who irradiated collagen hydrogels with gamma-rays. This technique has not become widespread yet, despite being fast, efficient and exempt from potentially toxic reagents. The mechanism proposed for radiation-induced cross-linking is indirect: radiation-induced water radiolysis is followed by hydroxyl radical attack and H abstraction from either glycine or tyrosine side-chains. The two radical sites then recombine to create a covalent bond. The same mechanism accounts for dityrosine formation by UV or gamma photon irradiation.⁵ It is important to note that in the latter case, the first step is dehydrogenation of the tyrosine hydroxyl groups, and the covalent bond is formed between two carbon atoms of the phenyl rings. Dityrosine cross-linking in collagen has recently been employed for engineering of a biomaterial for cell encapsulation.⁶ Interestingly, peptide (and also lipid) cross linking does not necessary have to proceed via water

^a CIMAP UMR 6252 CEA/CNRS/ENSICAEN/Université de Caen Normandie, Bd Becquerel 14070 Caen, France.

^b University College Groningen, University of Groningen, Groningen, The Netherlands.

^c Laboratory of Computational Biology, National Heart, Lung and Blood, Institute, National Institutes of Health, Bethesda, Maryland 20892, USA.

^d Zernike Institute for Advanced Materials, University of Groningen, Groningen, The Netherlands.

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

hydrolysis, as shown recently in experimental UV absorption studies in the gas phase.^{7,8}

Another interest of cross-linking is the control and stabilization of the triple helical geometrical structure of collagen, in particular using collagen mimetic peptides.⁹ The latter are much shorter than native collagen, thus allowing for custom peptide synthesis of a controlled amino acid sequence and crystallization to investigate the influence of mutations on the structure and stability of the triple helix.¹⁰ Assemblies of triple helices of collagen mimetic peptides are also valuable for studying collagen fibrils that constitute collagen fibers, and last but not least, they are interesting building blocks for the design of biomaterials.^{11,12} However, to our knowledge no previous studies have addressed cross-linking of collagen mimetic peptides by ionizing radiation.

In a previous work, we have experimentally shown that the triple helix structure of assemblies of $(PPG)_{10}$ and $(POG)_{10}$ (P, O and G being proline, hydroxyproline and glycine, respectively) collagen mimetic peptides can survive in the absence of a molecular environment (in the gas phase), and that the stability of this triple helix is an intrinsic property.¹³ In particular, the gas phase structure of the $[(PPG)_{10}]_6+9H^{9+}$ hexamer was found to be a noncovalent assembly of two triple helices, bound by dipole-dipole interaction but without H bonds, as in a crystal (see Fig. 1). Moreover, we have thoroughly studied the ionization and fragmentation of $(PPG)_{10}$ and $(POG)_{10}$ peptide monomers and trimers by VUV and X-ray photons as well as by carbon ion beams. We found that fragmentation proceeds in the electronic ground state of the ionized species, after internal conversion of the initial electronic excitation into vibrational modes.^{14,15} Usually, in this process, the noncovalent bonds are cleaved before the covalent ones. Consistently, the ionization of noncovalent complexes by X-ray photoabsorption leads to extensive cleavage of the complexes.^{15–17} Up to now, the irradiation of higher-order assemblies of collagen mimetic peptides has not been investigated in the gas phase.

Here, we report on the X-ray irradiation of $[(PPG)_{10}]_6+9H^{9+}$ as well as $[(POG)_{10}]_6+9H^{9+}$ hexamers in the gas phase. We employ mass spectrometry to analyze the cationic products of single photon absorption, in order to find evidence for photo-induced cross-linking of the two triple helices.




Figure 1. Crystal structure of the $(PPG)_{10}$ collagen mimetic peptide hexamer (PDB 1K6F). Water molecules are not displayed. Heavy atoms of a given peptide are depicted with space filling and in the same color.

Experimental section

All experiments were performed using a home-made tandem mass spectrometer that has been described in detail elsewhere.¹⁸ Briefly, protonated molecular systems are put in the gas phase by an electrospray ion source (ESI), focused and guided by RF electric fields through a funnel and an octopole. The molecular ion of interest is then selected by its mass over charge ratio (m/z) in a Quadrupole Mass Spectrometer (QMS), before being accumulated in a Paul trap and irradiated by the photon beam. All trapped cations are then extracted and analyzed in a Time-Of-Flight (TOF) mass spectrometer before reaching a detector consisting of one microchannel plate (MCP), a scintillator and a photomultiplier (Bipolar TOF, Photonis). This allowed a post-acceleration voltage of -9.0 kV to be applied to the front side of the MCP, and thus increases detection efficiency compared to usual MCP detectors. For instance, around $m/z = 2500$, this efficiency is more than doubled as compared to a post-acceleration voltage of -5.0 kV. X-ray photon irradiation was carried out at the BESSY II synchrotron (U49/2-PGM1 beamline, HZB, Berlin, Germany), and a mechanical shutter was used to control the irradiation time in the 100 – 500 ms range.

The $(PPG)_{10}$ and $(POG)_{10}$ collagen mimetic peptides were purchased from Peptides International (> 95% purity) and used without further purification. Powders were dissolved in equal proportions of water and methanol, with 1% of formic acid in volume. The final concentration was around 50 $\mu\text{mol/L}$. For each molecular system studied, we recorded a reference mass spectrum without X-ray exposure (beam off), and two with the beam (beam on). We then subtracted the “beam off” spectrum from the average of the two “beam on” spectra. The result is a negative peak for the parent ion, and all positive

peaks are product cations of the interaction involving a change in m/z . The resulting spectrum was smoothed by adjacent averaging over 20 points, and calibrated in m/z . Since the flux is about 10^{13} photons/s and the target density is low (10^7 - 10^8 molecules/cm³), photoabsorption events are independent. Therefore, a probability p for absorbing one photon gives the probability p^2 for two photons. Since we tuned the irradiation time to induce less than 10% depletion of the precursor ion, it implies $p^2 + p < 0.1$ (neglecting the absorption of more than two photons) and thus $p < 0.09$. At least 90% of the events that contribute to the observed mass spectra are thus due to single X-ray absorption events. We also irradiated the trap residual gas without the molecular ion beam (by turning off the electrospray needle voltage) to exclude the contribution of peaks due to residual gas contaminations.

Computational Methods

All calculations were performed using Gaussian09.¹⁹ Geometry optimization was performed in the electronic ground singlet state of the neutral molecular species studied, with the default algorithm. Two levels of density functional theory were used: the B3LYP functional with the 6-31g(d) basis set, and the wB97XD method that includes empirical dispersion as well as long-range corrections. The absence of negative frequencies was checked for molecular species with optimized geometry, to make sure that the structures were minima of the potential energy surface.

Results and discussion

Non-hydroxylated hexamer

In Fig. 2, we show the mass spectra of the cationic products from single photon absorption by the $[(\text{PPG})_{10}_6+9\text{H}]^{9+}$ hexamer. The most intense peak at $m/z = 1266$ can be attributed to the doubly-charged peptide monomer cation (M^{2+}) and/or to the sextuply-charged trimer (T^{6+}), that cannot be distinguished with the employed mass spectrometer. For the same reason, the weaker peak at $m/z = 1519$ can be assigned to non-dissociative ionization (NDI) of the precursor ion into $[(\text{PPG})_{10}_6+9\text{H}]^{10+}$, but also to the quintuply-charged trimer cation (T^{5+}). We have previously shown that collision-induced dissociation of the $[(\text{PPG})_{10}_6+9\text{H}]^{9+}$ hexamer precursor ion leads to formation of the T^{5+} and T^{4+} trimers, consistent with the expected statistical distribution of protons in the precursor ion composed of two triple helical peptide trimers.¹³ As a consequence, single photoionization of one trimer in the precursor T^{5+}/T^{4+} hexamer gives either T^{5+}/T^{5+} or T^{6+}/T^{4+} configurations, both with similar probabilities. Furthermore, inter-helix proton transfer can turn T^{6+}/T^{4+} into T^{5+}/T^{5+} . Therefore, further dissociation into two trimers will mostly form the T^{5+} trimer ($m/z = 1519$). From Fig. 2 it is clear though that the corresponding peak is relatively weak, indicating a generally low yield of trimer photoproducts. It is therefore likely that the peak at $m/z = 1266$ is mostly due to M^{2+} .





Figure 2. Mass spectra of single-photon absorption products of the collagen mimetic peptide hexamer $[(\text{PPG})_{10}_6+9\text{H}]^{9+}$ (average $m/z = 1688$). NDI, M and T stand for non-dissociative single ionization, monomer and trimer, respectively.

The two trimers constituting the $[(\text{PPG})_{10}_6+9\text{H}]^{9+}$ hexamer are bound via weak non-covalent dipole-dipole interactions, however the total binding energy is about 2 eV.¹³ The relatively high binding energy reflects the high permanent dipole moment of the triple helix of about 65 D. The dipole is well aligned with the helix axis, which maximizes the binding energy between the two triple helices in the hexamer shown in Fig. 1. Creation of a vacancy in a carbon 1s orbital of biomolecular species after photoabsorption around 290 eV with subsequent Auger decay is known to deposit an average of about 20 eV of vibrational energy into the system,²⁰ more than enough to detach both trimers from each other. The dominance of M^{2+} is evidence for a high probability of further dissociation of the triple helices into monomers after photoabsorption. We have already observed this process for photoabsorption of $[(\text{PPG})_{10}_3+7\text{H}]^{7+}$ triple helices in the same photon energy range.¹⁴ Scheme 1 illustrates this dissociation after single and double ionization of the $[(\text{PPG})_{10}_6+9\text{H}]^{9+}$ precursor. We will show later that double ionization indeed occurs in this photon energy range for a very similar system. Moreover, previous studies have proven that soft X-ray single photon absorption induces non-dissociative multiple ionization of isolated protonated proteins.²¹ Scheme 1 also indicates the statistical distribution of charges among monomers after dissociation: only M^+ and M^{2+} are expected, the former more than twice lower in abundance. This much lower abundance of M^+ compared to M^{2+} partly explains the fact that we do not observe the singly-charged monomer M^+ peak at $m/z = 2531$. Another reason is the decrease of the detection efficiency with m/z by about a factor of 2 at 2531 compared to 1266. The absence of the M^+ peak could also indicate that this monomer does not stay intact after photoionization and fragments into lower m/z ions, in particular for photon energies of 288 eV and above (see the next sub-section). A weak peak at $m/z = 844.5$ in the spectra from Fig. 2 can be assigned to the M^{3+} monomer trication. Its observation is not straightforward from scheme 1, thus M^{3+} has to come from single or double ionization and

dissociation of the precursor hexamer into monomers before charge equilibrium.



Scheme 1. Dissociation of collagen peptide hexamers after X-ray photoabsorption, single or double ionization and vibrational energy redistribution. The charges of hexamers, trimers and monomers expected from a statistical distribution are indicated.

In addition to cleavage of the noncovalent bonds between the two triple helices as well as between peptides within triple helices, we also observed breakage of covalent bonds within the monomers. The latter process gives rise to the peaks in the low m/z range of the mass spectra of Fig. 2, assigned to backbone fragments that we have already observed after photoabsorption of $(\text{PPG})_{10}$ monomers and trimers in the VUV and soft X-ray ranges.¹⁴ These fragments imply that after photoabsorption, monomers can have sufficient vibrational energy to dissociate, even after loss of all noncovalent interactions. However, the relative abundance of these fragments is lower than in the case of trimers, at a given photon energy.¹⁵ This is due to the size of the precursor ion: a hexamer is twice as large as a trimer, the deposited vibrational energy is thus distributed over twice the degrees of freedom.²⁰ Interestingly, the relative abundance of non-dissociative single ionization (leading to $[\text{((PPG)}_{10})_6+9\text{H}]^{10+}$ of $m/z = 1519$) is lower for the hexamer than for the $[\text{((PPG)}_{10})_3+7\text{H}]^{7+}$ trimer ionized after photoabsorption at the same photon energy. This can be explained by the lower binding energy of $[\text{((PPG)}_{10})_6+9\text{H}]^{9+}$ as compared to $[\text{((PPG)}_{10})_3+7\text{H}]^{7+}$. The two triple helices in $[\text{((PPG)}_{10})_6+9\text{H}]^{9+}$ are presumably bound by dipole-dipole interactions while the three peptides in $[\text{((PPG)}_{10})_3+7\text{H}]^{7+}$ are linked by strong H bonds.¹³ All these findings are in line with photoabsorption followed by internal conversion to the electronic ground state, followed by IVR and eventually fragmentation, as previously found for other protonated molecular systems, in particular collagen peptide trimers.^{14,15}

Hydroxylated hexamer

As a next step, we have investigated the influence of proline hydroxylation on photoabsorption-induced processes. To this end, ten proline residues within the $(\text{PPG})_{10}$ collagen mimetic peptide were replaced by hydroxyprolines. In Fig. 3, we show

the mass spectra of the $[\text{((POG)}_{10})_6+9\text{H}]^{9+}$ hexamer after photoabsorption in the 100 – 531 eV photon energy range. In the following, we will assume that in the gas-phase, the geometry of $[\text{((POG)}_{10})_6+9\text{H}]^{9+}$ is similar to the crystalline form (see Fig. 4), as we did for the case of $[\text{((PPG)}_{10})_6+9\text{H}]^{9+}$. For all photon energies between 100 and 288 eV, up to $m/z = 1400$ the mass spectrum is very similar to the one observed for $[\text{((PPG)}_{10})_6+9\text{H}]^{9+}$ (see Fig. 2), featuring peaks assigned to the same peptidic backbone fragments and peptide monomers with similar relative intensities. In sharp contrast, the peak at $m/z = 1615.5$ corresponding to NDI and/or the T^{5+} trimer has a much higher relative intensity for $[\text{((POG)}_{10})_6+9\text{H}]^{9+}$. This peak even dominates at 100 and 150 eV. The peak at $m/z = 1469$, observed for photon energies up to 300 eV, can unambiguously be attributed to non-dissociative double ionization (NDDI) of the precursor ion, forming $[\text{((POG)}_{10})_6+9\text{H}]^{11+}$. The equivalent peak at $m/z = 1381$ is absent from the spectra of $[\text{((PPG)}_{10})_6+9\text{H}]^{9+}$. To get more information, it is useful to look at the evolution of the relative intensity of the peaks at $m/z = 1615.5$ and 1469 from 288 to 300 eV, around the carbon K-edge energy. Previous studies on the protonated cytochrome C protein have shown that photoabsorption at 288 or 300 eV triggers C 1s excitation or ionization, followed by Auger decay, *i.e.* a total of one or two electrons are removed, respectively (see Scheme 2).²¹ Cytochrome C is a 12 kDa protein and for proteins of this size, backbone fragmentation is usually not observed after soft X-ray single photon absorption. Due to the large number of vibrational degrees of freedom, the vibrational energy of the photoionized system is insufficient for backbone fragmentation to occur.²⁰ As a consequence, for cytochrome C NDI dominates over NDDI at 288 eV, whereas the opposite is observed at 300 eV. In Fig. 3, a similar transition from NDI to NDDI is observed for $[\text{((POG)}_{10})_6+9\text{H}]^{9+}$ if we assume the peak at $m/z = 1615.5$ to be dominated by NDI, rather than by T^{5+} . This assumption is supported by the fact the peak almost vanishes at 300 eV, even though the vibrational excitation energy should be very similar for both 1s excitation (288 eV) and 1s ionization (300 eV).²⁰ Negligible T^{5+} at 300 eV thus implies negligible T^{5+} at 288 eV. Therefore, we find that non-dissociative single and double ionization is a major channel after photoionization of $[\text{((POG)}_{10})_6+9\text{H}]^{9+}$ up to 288 eV photon energy, and their relative probabilities behave like a covalent system of similar size around the carbon K-edge (see Fig. 3), which is not expected for a noncovalent complex: we will come back to this key point later.





Figure 3. Top: mass spectra of single-photon absorption products of the collagen mimetic peptide hexamer $[(\text{POG})_{10}_6+9\text{H}]^{9+}$ (average $m/z = 1795$). Bottom: relative yield of these products (area under the corresponding peaks normalized to the total yield of product ions) as a function of photon energy. NDI = non-dissociative single ionization, NDDI = non-dissociative double ionization, M^{2+} and M^{3+} are peptide monomers, and "peptidic fragments" is the total fragmentation yield of monomers.

At 300 eV, it is also interesting to notice that the M^{3+} peak around $m/z = 898$ is much more intense than at 288 eV for $[(\text{POG})_{10}_6+9\text{H}]^{9+}$ (see Fig. 3). This confirms the hypothesis that we invoked in the previous paragraph to explain M^{3+} formation from the $[(\text{PPG})_{10}_6+9\text{H}]^{9+}$ hexamer: double ionization of the precursor ion. Besides, the NDDI peak is absent from the $[(\text{PPG})_{10}_6+9\text{H}]^{9+}$ spectra, presumably because this hexamer breaks upon double ionization, due to its lower binding energy. When photon energy increases from 100 to 531 eV, the peptide backbone fragments become much more abundant and the peaks assigned to NDI, NDDI, trimers and monomers decrease in intensity (see Fig. 3), indicating that the amount of vibrational energy in the system progressively increases with photon energy. Such an observation has already been made for other protonated systems, in particular the $[(\text{PPG})_{10}_3+7\text{H}]^{7+}$ triple helix.¹⁵ The internal energy of the system after photoabsorption increases when electrons of deeper orbitals, for instance the 1s atomic orbitals of carbon at

285 – 300 eV, nitrogen around 400 eV and oxygen at 530 – 540 eV become accessible. Thus, our results are consistent with this increasing electronic energy being converted into vibrational energy in the electronic ground state of the system, provoking more fragmentation as photon energy rises. The present data confirm the general nature of this mechanism for single X-ray photoabsorption of molecular systems in the gas phase.

Observing NDI and NDDI as major processes after soft X-ray photoionization of a noncovalently-bound system such as $[(\text{POG})_{10}_6+9\text{H}]^{9+}$ is highly unusual, since weak intermolecular bonds are usually broken by the vibrational energy deposited by photoabsorption followed by internal conversion into the ground state (see the previous discussion about photoabsorption by $[(\text{PPG})_{10}_6+9\text{H}]^{9+}$). It shows that hydroxyprolines strengthen the peptide hexamer enough to prevent separation of the two triple helices upon soft X-ray photoabsorption. In the following, we explore the possible causes for this behavior.



Scheme 2. Electronic processes occurring within a molecular cation M^{n+} after X-ray photoabsorption around atomic K-edges of heavy atoms. Green arrows illustrate resonant transitions leading to excitation or deexcitation, and red arrows depict ejection to the continuum.

Discussion

A possible cause for observing non-dissociative single and double ionization of $[(\text{POG})_{10}_6+9\text{H}]^{9+}$ and not $[(\text{PPG})_{10}_6+9\text{H}]^{9+}$ following X-ray photoabsorption would be a much higher number of degrees of freedom and thus heat capacity for the hydroxylated hexamer, quenching fragmentation. However, this is not the case: hydroxyprolines bring only 3% more degrees of freedom. The only alternative explanation is a large increase of the dissociation energy of the hexamer upon hydroxylation. Previously, we have shown by collision-induced dissociation of collagen mimetic peptide triple helices that the collision energy to induce 50%

dissociation of $[(\text{POG})_{10}]_3+7\text{H}]^{7+}$ is 14% higher than that of $[(\text{PPG})_{10}]_3+7\text{H}]^{7+}$.¹³ We have attributed this difference to intra-helix stereoelectronic effects induced by the hydroxyl group of hydroxyproline. Such effects are expected to play a role in the $[(\text{POG})_{10}]_6+9\text{H}]^{9+}$ hexamer as well (since we assume that it contains two triple helices) but only regarding dissociation into *monomers*, not regarding separation of the two triple helices.¹³ Thus, stereoelectronic effects cannot explain the high abundance of NDI and NDDI for $[(\text{POG})_{10}]_6+9\text{H}]^{9+}$. Let us look at the crystal structure of $(\text{GPO})_9$ available in the Protein Data Bank (*cf.* Fig. 4):²² one notices that the two triple helices do not share any direct H bond; only water-mediated H-bonds exist between some hydroxyproline OH groups that are close to each other. In the gas phase, these water molecules are not present, which means that the corresponding H bonds do not exist. Even if several new direct OH---OH H bonds might form between hydroxyl groups after desolvation, these bonds would only slightly increase the binding energy of the hexamer since typical binding energies for such H bonds are about 0.14 eV (the binding energy of the methanol dimer held together by such a bond²³).




Figure 4. Crystal structure of the $(\text{GPO})_9$ collagen mimetic peptide hexamer (PDB 3B0S). Peptide backbones are depicted as lines, and only carbon, oxygen (in red) and nitrogen atoms (in blue) of hydroxyproline side-chains are shown. Water molecules are not displayed on the top figure, and on the bottom, only oxygen atoms of the water molecules within 5 Å around green highlighted hydroxyprolines are shown. H bonds are indicated by dashes.

We propose that the process accounting for the large increase of the stability of the hydroxylated $(\text{POG})_{10}$ hexamer upon X-ray photoionization is the creation of a covalent bond between the two triple helices (cross-linking) as a result of photoionization. Since no such process is observed for the non-hydroxylated $(\text{PPG})_{10}$ hexamer, the hydroxyl group of hydroxyproline must play a key role in the cross-linking. By drawing a parallel to the mechanism of dityrosine formation, we can assume that the first step is the same, namely H abstraction and/or proton transfer from the hydroxyl groups of two hydroxyproline side-chains to create two radicals. If the latter are close enough, they might recombine to form an O-O bond: in Fig. 4, we can see that the closest hydroxyproline

side-chain OH groups are only 5 Å apart. Moreover, they belong to different triple helices; this O-O bond would thus eventually cross-link the two triple helices. Alternatively, after H abstraction and/or proton transfer, the radicals might migrate to the carbon atoms linked to the oxygens of the two hydroxyprolines, to form two stable tertiary radicals. Recombination would create a covalent C-C bond, which has a typical bond dissociation energy (BDE) of 3.6-3.9 eV in alkanes,²⁴ while peroxides are less stable (BDE \approx 1.7 eV for the O-O bond of di-tert-butyl peroxide²⁵). This cross-linking would thus in any case significantly increase the binding energy of the hexamer and account for the presence of intense NDI and NDDI in the mass spectra of $[(\text{POG})_{10}]_6+9\text{H}]^{9+}$. This mechanism is supported by our previous studies on protonated collagen peptides containing hydroxyproline: we found that hydroxyproline side-chains undergo a radical reaction upon photoionization in the same photon energy range than the one explored here.^{14,26} It should also be noted that a similar mechanism has been proposed for the cross-linking of protonated triacylglycerol lipid dimers upon UV photoabsorption by their noncovalent complexes with diiodoaniline in the gas phase.⁸ This mechanism starts with abstraction of two I atoms from diiodoaniline, the formation of this diradical being followed by H abstraction from a carbon atom of each of the triacylglycerols, which finally induces their cross-linking *via* recombination of the two radicals. Our experimental results indicate that using ionizing photons, radical molecular ions of noncovalent complexes are created, can undergo recombination and cross-linking without the need of an external reagent such as diiodoaniline. In the next subsection, we show that quantum-chemical calculations support this conclusion.

Results of theoretical calculations

To investigate the cross-linking process between the two triple helices of the hydroxylated peptide hexamer in the gas phase, we performed quantum-chemical calculations at the density functional theory (DFT) level. All water molecules in the crystal structure of the $(\text{GPO})_9$ collagen mimetic peptide hexamer (PDB 3B0S) were removed. Since taking the whole hexamer is computationally unaffordable, we had to choose smaller systems. As a first step, only one peptide from each helix was kept, and the residues far away from the closest hydroxyproline side chains were removed: this is called system A. In addition, the H atoms of both hydroxyl groups were removed to create two radicals, and system A was subject to geometry optimization using several levels of DFT (see the "Computational methods" section). To probe the effect of the other peptides of the hexamer, system B was created by including the residues of the other peptides that share H bonds with system A: since system B is much larger than system A (244 vs. 98 atoms), only the wb97xd level was used. In all cases, during geometry optimization, an O-O covalent bond is formed by radical recombination. The same calculations were repeated for the two radicals located at the carbon atoms linked to the hydroxyproline oxygens: a C-C

covalent bond is now formed. The resulting structures for system A can be visualized in Fig. 5. Much more thorough theoretical investigations would be needed to firmly establish the mechanism responsible for ionization-induced cross-linking between the hexamers, but the present results are the first supporting the existence of such a process in collagen mimetic peptides.




Figure 5. Optimized structures of system A for radicals initially located at oxygens (left) and carbons (right) of the hydroxyproline side chains. The O-O and C-C bonds formed between the hydroxyprolines are encircled. Carbons are depicted in light grey, oxygens in red, nitrogens in blue, hydrogens in white.

Conclusions

X-ray photoionization of dimers of collagen mimetic peptide triple helices in the gas phase is in most cases followed by fragmentation due to excess vibrational energy. However, we also observe pronounced non-dissociative single and double ionization for the peptides containing hydroxyprolines. We attribute this behavior to the creation of a covalent bond between the two triple helices, after formation of two radicals and their recombination. Quantum-chemical calculations support this picture. However, the exact mechanism is still to be identified by future calculations, but also experiments such as multi-stage collision-induced dissociation tandem high-resolution mass spectrometry of the product ions of single or double ionization. Our findings show that cross-linking of molecular systems can be achieved by ionizing radiation, without the presence of any reagent or water, but also extend the possibilities of designing biomaterials by exposure of collagen mimetic peptides to ionizing radiation. It would be valuable to change the nature of the ionizing radiation, and for instance irradiate triple helix assemblies with electron or ion beams: we expect more multiple ejections of valence electrons, and thus potentially more covalent bonds to be formed. Emission of several valence electrons from multiply-deprotonated DNA oligonucleotides also occurs after X-ray photoabsorption or carbon ion beam irradiation,²⁷ another perspective is thus to seek for cross-linking of oligonucleotide dimers upon ionizing radiation.

Author Contributions

Marwa Abdelmouleh: Investigation

Muhamed Amin: Investigation, Writing – review and editing

Mathieu Lalande: Investigation, Writing – review and editing

Thomas Schlathöler: Conceptualization, Funding Acquisition, Project administration, Writing – review and editing

Jean-Christophe Pouilly: Investigation, Formal analysis, Funding Acquisition, Project administration, Supervision, Writing – original draft

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank HZB for the allocation of synchrotron radiation beamtime and local contacts for their support during experiments. The French “Conseil Régional de Normandie” is acknowledged for a PhD funding (#15P01339). This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 730872.

References

- 1 X. Yu, C. Tang, S. Xiong, Q. Yuan, Z. Gu, Z. Li and Y. Hu, Modification of Collagen for Biomedical Applications: A Review of Physical and Chemical Methods, *Current Organic Chemistry*, 2016, **20**, 1797–1812.
- 2 K. Adamiak and A. Sionkowska, Current methods of collagen cross-linking: Review, *INTERNATIONAL JOURNAL OF BIOLOGICAL MACROMOLECULES*, 2020, **161**, 550–560.
- 3 X. Hu, W. K. Raja, B. An, O. Tokareva, P. Cebe and D. L. Kaplan, Stability of Silk and Collagen Protein Materials in Space, *Scientific Reports*, 2013, **3**, 3428.
- 4 X. Zhang, L. Xu, X. Huang, S. Wei and M. Zhai, Structural study and preliminary biological evaluation on the collagen hydrogel crosslinked by gamma-irradiation, *JOURNAL OF BIOMEDICAL MATERIALS RESEARCH PART A*, 2012, **100A**, 2960–2969.
- 5 C. Giulivi, N. J. Traaseth and K. J. A. Davies, Tyrosine oxidation products: analysis and biological relevance, *Amino Acids*, 2003, **25**, 227–232.
- 6 B. Partlow, M. Applegate, F. Omenetto and D. Kaplan, Dityrosine Cross-Linking in Designing Biomaterials, *ACS BIOMATERIALS SCIENCE & ENGINEERING*, 2016, **2**, 2108–2121.
- 7 H. Nguyen, P. Andrikopoulos, L. Rulisek, C. Shaffer and F. Turecek, Photodissociative Cross-Linking of Non-covalent Peptide-Peptide Ion Complexes in the Gas Phase, *JOURNAL OF THE AMERICAN SOCIETY FOR MASS SPECTROMETRY*, 2018, **29**, 1706–1720.
- 8 S. Nie, H. T. Pham, S. J. Blanksby and G. E. Reid, Photoinduced Intermolecular Cross-Linking of Gas Phase Triacylglycerol Lipid Ions, *Eur J Mass Spectrom (Chichester)*, 2015, **21**, 287–296.
- 9 S. A. H. Hulgán and J. D. Hartgerink, Recent Advances in Collagen Mimetic Peptide Structure and Design, *Biomacromolecules*, 2022, **23**, 1475–1489.

- 10 B. Brodsky, G. Thiagarajan, B. Madhan and K. Kar, Triple-helical peptides: An approach to collagen conformation, stability, and self-association, *Biopolymers*, 2008, **89**, 345–353.
- 11 S. M. Yu, Y. Li and D. Kim, Collagen mimetic peptides: progress towards functional applications, *Soft Matter*, 2011, **7**, 7927–7938.
- 12 Y. Li and S. M. Yu, Targeting and mimicking collagens via triple helical peptide assembly, *Current Opinion in Chemical Biology*, 2013, **17**, 968–975.
- 13 M. Lalande, C. Comby-Zerbino, M. Bouakil, P. Dugourd, F. Chirot and J.-C. Pouilly, Isolated Collagen Mimetic Peptide Assemblies Have Stable Triple-Helix Structures, *Chemistry-a European Journal*, 2018, **24**, 13728–13733.
- 14 L. Schwob, M. Lalande, J. Rangama, D. Egorov, R. Hoekstra, R. Pandey, S. Eden, T. Schlathölter, V. Vizcaino and J.-C. Pouilly, Single-photon absorption of isolated collagen mimetic peptides and triple-helix models in the VUV-X energy range, *Physical Chemistry Chemical Physics*, 2017, **19**, 18321–18329.
- 15 M. Lalande, M. Abdelmouleh, M. Ryszka, V. Vizcaino, J. Rangama, A. Mery, F. Durantel, T. Schlathölter and J.-C. Pouilly, Irradiation of isolated collagen mimetic peptides by x rays and carbon ions at the Bragg-peak energy, *Physical Review A*, 2018, **98**, 062701.
- 16 M. Abdelmouleh, M. Lalande, J. El Feghaly, V. Vizcaino, A. Rebelo, S. Eden, T. Schlathölter and J.-C. Pouilly, Mass Spectral Signatures of Complex Post-Translational Modifications in Proteins: A Proof-of-Principle Based on X-ray Irradiated Vancomycin, *J. Am. Soc. Mass Spectrom.*, 2020, **31**, 1738–1743.
- 17 M. Abdelmouleh, A. Rodriguez, J. Leroux, P. Christodoulou, B. Bernay, T. Schlathölter and J. Pouilly, X-ray photoabsorption-induced processes within protonated rifamycin sodium salts in the gas phase, *EUROPEAN PHYSICAL JOURNAL D*, , DOI:10.1140/epjd/s10053-021-00092-w.
- 18 S. Bari, R. Hoekstra and T. Schlathölter, Peptide fragmentation by keV ion-induced dissociation, *Physical Chemistry Chemical Physics*, 2010, **12**, 3376–3383.
- 19 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2016., 2016.
- 20 D. Egorov, L. Schwob, M. Lalande, R. Hoekstra and T. Schlathölter, Near edge X-ray absorption mass spectrometry of gas phase proteins: the influence of protein size, *Physical Chemistry Chemical Physics*, 2016, **18**, 26213–26223.
- 21 A. R. Milosavljevic, F. Canon, C. Nicolas, C. Miron, L. Nahon and A. Giuliani, Gas-Phase Protein Inner-Shell Spectroscopy by Coupling an Ion Trap with a Soft X-ray Beamline, *Journal of Physical Chemistry Letters*, 2012, **3**, 1191–1196.
- 22 K. Okuyama, K. Miyama, K. Mizuno and H. P. Bächinger, Crystal structure of (Gly-Pro-Hyp)₉: Implications for the collagen molecular model, *Biopolymers*, 2012, **97**, 607–616.
- 23 F. C. Hagemeister, C. J. Gruenloh and T. S. Zwier, Density Functional Theory Calculations of the Structures, Binding Energies, and Infrared Spectra of Methanol Clusters, *J. Phys. Chem. A*, 1998, **102**, 82–94.
- 24 S. J. Blanksby and G. B. Ellison, Bond Dissociation Energies of Organic Molecules, *Acc. Chem. Res.*, 2003, **36**, 255–263.
- 25 R. D. Bach and H. B. Schlegel, Bond Dissociation Energy of Peroxides Revisited, *J. Phys. Chem. A*, 2020, **124**, 4742–4751.
- 26 L. Schwob, M. Lalande, D. Egorov, J. Rangama, R. Hoekstra, V. Vizcaino, T. Schlathölter and J.-C. Pouilly, Radical-driven processes within a peptidic sequence of type I collagen upon single-photon ionisation in the gas phase, *Physical Chemistry Chemical Physics*, 2017, **19**, 22895–22904.
- 27 W. Li, O. Kavatsyuk, W. Douma, X. Wang, R. Hoekstra, D. Mayer, M. S. Robinson, M. Gühr, M. Lalande, M. Abdelmouleh, M. Ryszka, J. C. Pouilly and T. Schlathölter, Multiple valence electron detachment following Auger decay of inner-shell vacancies in gas-phase DNA, *Chem. Sci.*, , DOI:10.1039/D1SC02885E.