



**HAL**  
open science

# Risk Management in Agriculture and Crop Insurance: Implementation of a Method for Estimating Reference Yields in Organic Field Crops in France

O. Mghirbi, Youssef Lebrini, David Grandgirard, Alicia Ayerdi Gotor, Laetitia Fagot

## ► To cite this version:

O. Mghirbi, Youssef Lebrini, David Grandgirard, Alicia Ayerdi Gotor, Laetitia Fagot. Risk Management in Agriculture and Crop Insurance: Implementation of a Method for Estimating Reference Yields in Organic Field Crops in France. Institut Agro Rennes-Angers et INRAe. 18th Congress of the European Society for Agronomy, Aug 2024, Rennes, France. 2024. hal-04709709

**HAL Id: hal-04709709**

**<https://hal.science/hal-04709709v1>**

Submitted on 25 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives 4.0 International License

# RISK MANAGEMENT IN AGRICULTURE AND CROP INSURANCE: IMPLEMENTATION OF A METHOD FOR ESTIMATING REFERENCE YIELDS IN ORGANIC FIELD CROPS IN FRANCE

## STUDY CONTEXT

Agriculture in France is subject to numerous constraints, notably increasing meteorological risks in recent years due to climate change. Faced with natural disasters and pressure from bio-aggressors, farmers in both conventional and organic production are confronted with a multitude of risk factors leading to variability in their agricultural yields (Arora 2019; Malhi et al., 2021). Consequently, risk management is a key issue for the sustainability of agricultural activities. Crop insurance is considered an essential tool for safeguarding against these various risks and securing farmers' income (Folus et al. 2020; Frascarelli et al., 2021; Koenig et al., 2022).

## OBJECTIVE

Compared to conventional agriculture, there are not enough reference yields for organic farming in most agricultural regions of France. This lack of reference data makes it difficult for insurers to set up crop insurance contracts appropriate to the agricultural, environmental, and climatic contexts of farmers.

The objective of this study is to develop a method for estimating reference yields in large-scale crops (winter soft wheat, maize, spring barley) in organic farming, using statistical models, in order to determine the conditions and parameters to be considered for crop insurance pricing and the contractualization of Groupama Paris Val de Loire (GPVL) members.

## STUDY AREA

Groupama PVL : 4 regions, 10 departments, 84 agricultural small regions  
Area : 50 956 km<sup>2</sup>




Figure 1. Land use of the GPVL zone

## DETERMINANTS OF YIELD

To determine the reference yield, the first step is to study the determinants of yield in large-scale crops (winter wheat, spring barley, maize grain) to identify the factors that impact yield (Poniso et al. 2015; Ben Zekri et al., 2019). Then, statistical regression analysis was used to explain the variability of yields among members based on the factors determining yield. These factors primarily include climate, soil properties, pressure from bio-aggressors, crop rotation, technical itinerary of the crop, as well as the farmer's expertise and landscape factors.




Figure 2. Crop yield determinants and yield prediction approach

## CROP YIELD ESTIMATION

The analysis of yield variability based on yield determinants has enabled the development of a method for estimating reference yields in organic farming using regression and prediction models such as Generalized Linear Regression (GLM), Partial Least Squares (PLS), Ridge regression, and Gradient Boosting. The PLS model is the most effective and suitable for yield estimation, as determined through comparison using metrics and criteria such as R<sup>2</sup> and RMSE (Table 1). The yield estimation model was applied to predict the yields of insured members of Groupama Paris Val de Loire based on their location and the spatial intrinsic characteristics of fields.




Figure 3. The approach to estimating wheat, corn and barley yield in organic and conventional farming

Table 1. Results of R<sup>2</sup> and RMSE of the different models tested in conventional (A) and organic (B) wheat, maize and barley

Model	Winter soft wheat		Maize		Spring barley	
	R <sup>2</sup>	RMSE	R <sup>2</sup>	RMSE	R <sup>2</sup>	RMSE
Lasso regression	0,695	1,108	0,555	2,107	0,504	1,282
LM	0,706	1,083	0,552	2,088	0,547	1,179
GLM	0,707	1,083	0,557	2,088	0,560	1,179
PLS	0,710	1,083	0,557	2,127	0,560	1,212
Stepwise backward	0,699	1,103	0,553	2,119	0,501	1,288
Ridge regression	0,697	1,107	0,555	2,090	0,505	1,296
Elastic net regression	0,697	1,105	0,546	2,130	0,494	1,295

Model	Winter soft wheat		Maize		Spring barley	
	R <sup>2</sup>	RMSE	R <sup>2</sup>	RMSE	R <sup>2</sup>	RMSE
Lasso regression	0,210	2,009	0,961	11,355	0,463	1,461
LM	0,390	1,640	0,572	1,580	0,398	1,279
GLM	0,501	1,640	0,735	1,580	0,588	1,279
PLS	0,509	1,651	0,735	2,439	0,588	1,644
Stepwise backward	0,410	1,906	0,949	11,936	0,597	1,505
Ridge regression	0,307	1,735	0,975	15,196	0,584	1,721
Elastic net regression	0,338	1,840	0,942	14,226	0,466	1,477

## RESULTS & DISCUSSION




Figure 4. Spatial distribution of estimated yields (quintals/ha) at ASR scale for wheat, maize and barley in conventional farming (1) and the yield ratio organic/conventional (O/C) farming (2)

The estimated yield in organic farming at ASR scale, for winter soft wheat ranges from 3.3 t/ha to 7.8 t/ha, spring barley ranges from 1.0 t/ha to 5.1 t/ha, while maize grain varies from 5.1 t/ha to 11.3 t/ha (Figure 4). The yield ratio between organic and conventional farming varies between 53% and 105% for winter soft wheat, from 53% to 108% for grain corn and from 19% to 84% for spring barley. This ratio, which means the yield decline rate, shows a remarkable gap between organic and conventional yields which varies depending on the small agricultural regions. The climate, particularly rainfall, represents the main factor which explains this yield variation.

## CONCLUSION

Each factor has its limitations and leads to a lack of precision regarding results. Several other factors have not been studied and may impact yields, including biodiversity index, sowing date, bio-aggressors management, nitrogen management, soil cover, varieties used, and rotation duration. The statistical modelling yield approach will facilitate a better assessment of factors affecting yield and more precise crop insurance pricing depending on reference yields in organic and conventional farming, thereby helping to support farmers in the face of increasing uncertainty resulting from current climatic, agronomic, and environmental challenges.

## REFERENCES

- Arora, N.K. (2019) Impact of climate change on agriculture production and its sustainable solutions. Environmental Sustainability 2, 95–96.
- Ben Zekri, Y., Barkaoui, K., Marrou, H., Mekki, I., Belhouache, H., & Wery, J. (2019). On farm analysis of the effect of the preceding crop on N uptake and grain yield of durum wheat (Triticum durum Desf.) in Mediterranean conditions. Archives of Agronomy and Soil Science, 65(5), 594–611.
- Folus, D., Casal Ribeiro, P., Lepoivre, B., & Roumigué, A. (2020). L'assurance et la protection financière de l'agriculture. Annales des Mines - Réalités industrielles, 2020, 30-38.
- Frascarelli A., Del Sarto S., Mastandrea, G. (2021) A New Tool for Covering Risk in Agriculture: The Revenue Insurance Policy. Risks, 9, 131.
- Koenig, R., Brunette, M., Delacote, P. & Tevenart, C. (2022). Assurance récolte en France : spécificité du régime et déterminants potentiels. Économie rurale, 380, 7-25.
- Malhi G.S., Kaur M., Koushik P. (2021) Impact of Climate Change on Agriculture and its Mitigation Strategies: A Review. Sustainability, 13, 1318.
- Poniso, L.C., Al-Ghailani, L.K., Meese, K.C., Polomine, J., de Valpine, P., et Kremen, C., 2015. Diversification practices reduce organic to conventional yield gap. Proceedings of the Royal Society B: Biological Sciences, volume 282, n° 1799, p. 20141396.