



HAL
open science

The seventh equation of the NLS hierarchy and multi-rogue waves

Pierre Gaillard

► **To cite this version:**

Pierre Gaillard. The seventh equation of the NLS hierarchy and multi-rogue waves. 2024. hal-04709718

HAL Id: hal-04709718

<https://hal.science/hal-04709718v1>

Preprint submitted on 25 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The seventh equation of the NLS hierarchy and multi-rogue waves

Pierre Gaillard,

Université de Bourgogne Franche Comté,
Institut de mathématiques de Bourgogne,
9 avenue Alain Savary BP 47870
21078 Dijon Cedex, France :
E-mail : Pierre.Gaillard@u-bourgogne.fr

Abstract

The following study is part of a research program of rational solutions of the hierarchy of the nonlinear Schrödinger equation. Here, we are interested in the equation of order 7 and we highlight particular solutions providing the first orders of rogue waves not yet found.

Key Words : NLS hierarchy, quasi-rational solutions.

PACS numbers :

33Q55, 37K10, 47.10A-, 47.35.Fg, 47.54.Bd

1 Introduction

Quasi-rational solutions to the seventh equation of the NLS hierarchy are constructed. We give explicit expressions of these solutions for the first orders. They depend on multi-parameters and so patterns of these solutions in the (x, t) plane according the different values of the parameters are studied. We consider the seventh equation of the NLS hierarchy of order 7 (*NLS7*) which can be written as

$$\begin{aligned} & iu_t + u_{8x} + 16|u|^2|u_{6x} + 2u^2\bar{u}_{6x} + 56\bar{u}u_xu_{5x} \\ & + 40u\bar{u}_xu_{5x} + 12uu_x\bar{u}_{5x} + 98|u|^4u_{4x} + 168|u_x|^2u_{4x} \\ & + 112\bar{u}u_{2x}u_{4x} + 72u\bar{u}_{2x}u_{4x} + 28u^2|u|^2\bar{u}_{4x} + 42u_x^2\bar{u}_{4x} \\ & + 44uu_{2x}\bar{u}_{4x} + 68uu_x\bar{u}_{3x} + 476|u|^2\bar{u}u_xu_{3x} + 252u_x\bar{u}_{2x}u_{3x} \\ & + 308u|u|^2\bar{u}_xu_{3x} + 308\bar{u}_xu_{2x}u_{3x} + 70\bar{u}u_{3x}^2 + 196u_xu_{2x}\bar{u}_{3x} \\ & + 168u|u|^2u_x\bar{u}_{3x} + 56u^3\bar{u}_x\bar{u}_{3x} + 280|u|^6u_{2x} + 1456|u|^2|u_x|^2u_{2x} \\ & + 490\bar{u}^2u_x^2u_{2x} + 238u^2\bar{u}_x^2u_{2x} + 588|u|^2u_x^2\bar{u}_{2x} + 336u^2|u_x|^2\bar{u}_{2x} \\ & + 140|u|^4u^2\bar{u}_{2x} + 42u^3(\bar{u}_{2x})^2 + 392|u|^2u|u_{2x}|^2 + 322|u|^2\bar{u}u_{2x}^2 \\ & + 182u_{2x}^2\bar{u}_{2x} + 560|u|^4\bar{u}u_x^2 + 560|u|^4u|u_x|^2 + 420\bar{u}u_x^2|u_x|^2 \\ & + 140u^3|u|^2\bar{u}_x^2 + 378|u_x|^4u + 70|u|^8u \end{aligned} \tag{1}$$

with as usual the subscripts meaning partial derivatives and \bar{u} the complex conjugate of u .

Different classical equations are included in the NLS hierarchy; the first one is the NLS equation [1, 3, 4, 5, 12, 7, 30, 15]; the second one is the mKdV equation [76, 77, 78, 79, 80]; the third one is the LPD equation [81, 84, 85, 86, 87].

Many works has been done for these first three equations of the NLS hierarchy. For example, we can quote the following works, for the NLS equation [2], the mKdV equation [77], the LPD equation [81, 82, 83]. However, very few studies have been carried out for the following orders of this hierarchy. Here we explicitly construct solutions of the order equation seven of this hierarchy.

We construct quasi rational solutions for the first orders. The related patterns of the modulus of these solutions in the plane of coordinates $(x; t)$ are studied.

2 Quasi rational solutions of order 1 to the NLS7 equation

Theorem 2.1 *The function $v(x, t)$ defined by*

$$v(x, t) = -\frac{(3 - 4x^2 - 313600t^2 + 2240it)e^{70it}}{1 + 4x^2 + 313600t^2} \quad (2)$$

is a solution to the (NLS7) equation (1).




Figure 1. Solution of order 1 to (NLS7).

We get a smooth solution of the equation (1).

3 Quasi rational solutions of order 2 of the NLS7 equation depending on 2 real parameters

Theorem 3.1 *The function $v(x, t)$ defined by*

$$v(x, t) = \frac{n(x, t)}{d(x, t)} \quad (3)$$

with

$$\begin{aligned} n(x, t) = & -(-64x^6 + 2304b_1x^5 - 768a_1^2x^4 + 144x^4 + 107520itx^4 - 215040a_1tx^4 + \\ & 768ia_1x^4 - 15052800t^2x^4 - 34560b_1^2x^4 - 18432ia_1x^3b_1 - 4992b_1x^3 + 276480b_1^3x^3 + \\ & 18432b_1a_1^2x^3 + 361267200b_1t^2x^3 + 5160960b_1a_1tx^3 - 2580480itx^3b_1 + 2580480ia_1^2tx^2 - \\ & 1244160b_1^4x^2 + 2903040a_1tx^2 - 3072a_1^4x^2 + 361267200ia_1t^2x^2 - 33718272000a_1t^3x^2 + \\ & 23224320itb_1^2x^2 - 165888b_1^2a_1^2x^2 + 293529600t^2x^2 - 1180139520000t^4x^2 + \\ & 5760a_1^2x^2 + 16859136000it^3x^2 - 1152ia_1x^2 + 58752b_1^2x^2 + 6144ia_1^3x^2 - 361267200a_1^2t^2x^2 + \\ & 165888ia_1b_1^2x^2 - 1720320a_1^3tx^2 - 3251404800b_1^2t^2x^2 - 46448640b_1^2a_1tx^2 - \\ & 806400itx^2 + 180x^2 - 202309632000it^3xb_1 - 3161088000b_1t^2x + 36864b_1a_1^4x - \\ & 29675520b_1a_1tx + 663552b_1^3a_1^2x - 50688b_1a_1^2x - 5616b_1x + 20643840b_1a_1^3tx - \\ & 663552ia_1xb_1^3 - 290304b_1^3x + 13005619200b_1^3t^2x - 4608ia_1xb_1 + 4335206400b_1a_1^2t^2x + \\ & 14161674240000b_1t^4x + 7096320itb_1x - 4335206400ia_1t^2xb_1 + 185794560b_1^3a_1tx - \\ & 92897280itxb_1^3 - 73728ia_1^3xb_1 - 30965760ia_1^2txb_1 + 2985984b_1^5x + 404619264000b_1a_1t^3x - \\ & 262080it + 23602790400000ia_1t^4 + 54792192000it^3 - 1475174400000t^4 + 3225600ia_1^2t + \\ & 13005619200ia_1t^2b_1^2 + 337182720000ia_1^2t^3 + 606928896000it^3b_1^2 + 73543680b_1^2a_1t - \\ & 30840979456000000t^6 + 139345920itb_1^4 + 8601600ia_1^4t + 660878131200000it^5 + \\ & 69120ia_1b_1^2 - 13547520itb_1^2 - 13005619200b_1^2a_1^2t^2 - 1213857792000b_1^2a_1t^3 - \\ & 278691840b_1^4a_1t + 995328ia_1b_1^4 + 221184ia_1^3b_1^2 - 45 + 812851200ia_1t^2 - 98784000t^2 + \\ & 92897280ia_1^2tb_1^2 + 3010560a_1^3t + 270950400a_1^2t^2 - 8429568000a_1t^3 - 61931520b_1^2a_1^3t + \\ & 96768b_1^2a_1^2 + 8399462400b_1^2t^2 + 1872a_1^2 + 18000b_1^2 + 846720a_1t - 720ia_1 + \\ & 1536ia_1^3 + 12288ia_1^5 + 2408448000ia_1^3t^2 - 42485022720000b_1^2t^4 - 995328b_1^4a_1^2 - \\ & 19508428800b_1^4t^2 - 110592b_1^2a_1^4 - 23602790400000a_1^2t^4 - 3440640a_1^5t - 1204224000a_1^4t^2 - \\ & 224788480000a_1^3t^3 - 1321756262400000a_1t^5 - 2985984b_1^6 - 4096a_1^6 + 518400b_1^4 + \\ & 8448a_1^4)e^{2i(a_1+35t)} \end{aligned}$$

and

$$\begin{aligned} d(x, t) = & 64x^6 - 2304b_1x^5 + 48x^4 + 15052800t^2x^4 + 215040a_1tx^4 + 768a_1^2x^4 + \\ & 34560b_1^2x^4 - 5160960b_1a_1tx^3 - 361267200b_1t^2x^3 - 276480b_1^3x^3 + 384b_1x^3 - \\ & 18432b_1a_1^2x^3 + 1180139520000t^4x^2 + 361267200a_1^2t^2x^2 + 46448640b_1^2a_1tx^2 - \\ & 203212800t^2x^2 + 1720320a_1^3tx^2 - 1152a_1^2x^2 - 1612800a_1tx^2 + 1244160b_1^4x^2 + \\ & 3072a_1^4x^2 + 165888b_1^2a_1^2x^2 + 3251404800b_1^2t^2x^2 - 17280b_1^2x^2 + 108x^2 + \\ & 33718272000a_1t^3x^2 - 20643840b_1a_1^3tx - 185794560b_1^3a_1tx - 4335206400b_1a_1^2t^2x - \\ & 404619264000b_1a_1t^3x - 2985984b_1^5x - 13005619200b_1^3t^2x + 2077286400b_1t^2x + \\ & 124416b_1^3x - 14161674240000b_1t^4x - 663552b_1^3a_1^2x - 4608b_1a_1^2x - 2448b_1x + \\ & 14192640b_1a_1tx - 36864b_1a_1^4x + 9 + 177020928000a_1t^3 + 19508428800b_1^4t^2 + \\ & 347155200t^2 - 5148057600b_1^2t^2 + 7375872000000t^4 + 30840979456000000t^6 + \\ & 69120b_1^2a_1^2 + 1411200a_1t + 110592b_1^2a_1^4 + 995328b_1^4a_1^2 - 27095040b_1^2a_1t + \\ & 23602790400000a_1^2t^4 + 3440640a_1^5t + 1204224000a_1^4t^2 + 224788480000a_1^3t^3 + \\ & 1321756262400000a_1t^5 + 2985984b_1^6 + 4096a_1^6 - 269568b_1^4 + 6912a_1^4 + 42485022720000b_1^2t^4 + \\ & 5591040a_1^3t + 1535385600a_1^2t^2 + 61931520b_1^2a_1^3t + 13005619200b_1^2a_1^2t^2 + \\ & 1213857792000b_1^2a_1t^3 + 278691840b_1^4a_1t + 1584a_1^2 + 20016b_1^2 \end{aligned}$$

is a solution to the (NLS7) equation (1).




Figure 2. Solution of order 2 to the equation (1); to the left $a_1 = 0, b_1 = 0$; in the center $a_1 = 1, b_1 =$; to the right $a_1 = 10, b_1 = 0$.




Figure 3. Solution of order 2 to the equation (1); to the left $a_1 = 0, b_1 = 1$; in the center $a_1 = 0, b_1 = 4$; to the right $a_1 = 0, b_1 = 10$.

When one or both parameters increase, three peaks appear. When only one of the parameters increases, the three peaks appear but with different orientations.

4 Quasi rational solutions of order 3 of the NLS7 equation

The solution of order 3, depending on 4 real parameters being too long, we only present in the appendix. Here we give the solution of order 3 without parameters.

Theorem 4.1 *The function $v(x, t)$ defined by*

$$v(x, t) = \frac{n(x, t)}{d(x, t)} \quad (4)$$

with

$$\begin{aligned}
n(x, t) = & -(-4096 x^{12} + 13762560 itx^{10} - 1926758400 t^2x^{10} + 18432 x^{10} - 377644646400000 t^4x^8 + \\
& 93929472000 t^2x^8 - 258048000 itx^8t + 57600 x^8 + 5394923520000 it^3x^8 - 167242629120000 itx^6t^3 + \\
& 2077045552000000 t^4x^6 - 1369202688000 t^2x^6 + 172800 x^6 + 1109606400 itx^6 + \\
& 845924007936000000 it^5x^6 - 3947645370368000000 t^6x^6 - 17129961160704000000 itx^4t^5 + \\
& 716083200 itx^4 + 875169792000 t^2x^4 - 952878366720000 itx^4t^3 + 66320442222182400000000 it^7x^4 + \\
& 1509974354165760000000 t^6x^4 - 226800 x^4 + 59903882035200000 t^4x^4 - 232121547776384000000000 t^8x^4 - \\
& 653632074547200000 t^4x^2 + 31336408949981184000000000 t^8x^2 + 267893559263232000000 it^5x^2 - \\
& 7585324185600 t^2x^2 - 7279331738306740224000000000 t^{10}x^2 - 20246979648946176000000 t^6x^2 - \\
& 9948066333273600000000 itx^2t^7 + 2599761335109550080000000000 it^9x^2 + 3826686689280000 it^3x^2 + \\
& 3969907200 itx^2 - 113400 x^2 + 407642577345177452544000000000 it^{11} + 433641600 it - \\
& 5496461881344000 it^3 + 2484529566748508160000000 it^7 - 26443775418368000000 it^5 + \\
& 5546744179200 t^2 + 167910924623649177600000000 t^8 + 26647553684872888320000000000 it^9 + \\
& 14175 - 30937159887803645952000000000 t^{10} - 95116601380541405593600000000000 t^{12} - \\
& 704818560983040000 t^4 + 55055219338248192000000 t^6)e^{70 it}
\end{aligned}$$

and

$$\begin{aligned}
d(x, t) = & 4096 x^{12} + 1926758400 t^2x^{10} + 6144 x^{10} + 34560 x^8 - 65028096000 t^2x^8 + \\
& 377644646400000 t^4x^8 + 877879296000 t^2x^6 - 1321756262400000 t^4x^6 + 149760 x^6 + \\
& 39476453703680000000 t^6x^6 + 232121547776384000000000 t^8x^4 - 156203266867200000 t^4x^4 - \\
& 680968826388480000000 t^6x^4 - 3248695296000 t^2x^4 + 54000 x^4 + 48600 x^2 \\
& + 21490487940612096000000 t^6x^2 + 1886547433881600000 t^4x^2 + 2236763289600 t^2x^2 + \\
& 7279331738306740224000000000 t^{10}x^2 + 10445469649993728000000000 t^8x^2 - \\
& 13357721658064896000000 t^6 + 1110098090091777884160000000000 t^{10} \\
& + 281244270326081126400000000 t^8 + 866898922659840000 t^4 \\
& + 95116601380541405593600000000000 t^{12} + 2025 + 3320852774400 t^2
\end{aligned}$$

is a solution to the (NLS) equation (1).

In the following, we give the patterns of the modules of the solutions according to different values of the parameters.




Figure 4. Solution of order 3 to (1); to the left $a_1 = 0, b_1 = 0, a_2 = 0, b_2 = 0$;
in the center $a_1 = 1, b_1 = 0, a_2 = 0, b_2 = 0$; to the right $a_1 = 10, b_1 = 0,$
 $a_2 = 0, b_2 = 0$.




Figure 5. Solution of order 3 to (1); to the left $a_1 = 0, b_1 = 0, 1, a_2 = 0, b_2 = 0$; in the center $a_1 = 0, b_1 = 5, a_2 = 0, b_2 = 0$; to the right $a_1 = 0, b_1 = 5, a_2 = 0, b_2 = 0$.




Figure 6. Solution of order 3 to (1); to the left $a_1 = 0, b_1 = 0, a_2 = 0, 5, b_2 = 0$; in the center $a_1 = 0, b_1 = 0, a_2 = 1, b_2 = 0$; to the right $a_1 = 0, b_1 = 5, a_2 = 2, b_2 = 0$.




Figure 7. Solution of order 3 to (1); to the left $a_1 = 0, b_1 = 0, a_2 = 0, b_2 = 0, 5$; in the center $a_1 = 0, b_1 = 0, a_2 = 0, b_2 = 2$; to the right $a_1 = 0, b_1 = 5, a_2 = 0, b_2 = 3$.

We remark the similarity with these solutions and those relative to other equations belonging to this NLS hierarchy. For example, we recover the same types of patterns like in the NLS equation [88], the mKdV equation [89], or the Lakshmanan Porsezian Daniel equation [90]. We get the structure of triangles with peaks which appear in function of the different values of the parameters.

5 Conclusion

Quasi-rational solutions to the (*NLS7*) equation have been constructed for the first orders. These N -order solutions appear as the quotient of a polynomial of degree $N(N + 1)$ in x and t for the numerator by a polynomial of degree $N(N + 1)$ in x and t for the denominator.

The solutions of order 2 depend on two real parameters, and the structure of triangles with three peaks is observed for their modules.

The solutions of order 3 depend on four real parameters. In the plane (x, t) of the coordinates, the representation of the modules of the solutions reveals the formation of triangles containing 6 peaks.

It will be relevant to study other solutions of this equations and study the patterns of their modulus.

References

- [1] V. E. Zakharov, Stability of periodic waves of finite amplitude on a surface of a deep fluid, *J. Appl. Tech. Phys*, V. 9, 86-94, (1968)
- [2] N. Akhmediev, V. Eleonski, N. Kulagin, Generation of periodic trains of picosecond pulses in an optical fiber : exact solutions, *Sov. Phys. J.E.T.P.*, V. 62, 894-899, (1985)
- [3] N. Akhmediev, A. Ankiewicz, J.M. Soto-Crespo, Rogue waves and rational solutions of nonlinear Schrödinger equation, *Phys. Rev. E*, V. 80, 026601-1-9, 2009
- [4] N. Akhmediev, A. Ankiewicz, First-order exact solutions of the nonlinear Schrödinger equation in the normal-dispersion regime, *Phys. Rev. A*, V. 47, N. 4, 3213-3221, 2009
- [5] A. Ankiewicz, D. J. Kedziora, N. Akhmediev, Rogue wave triplets, *Phys Lett. A*, V. 375, 2782-2785, 2011
- [6] P. Dubard, P. Gaillard, C. Klein, V. B. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Spe. Top.*, V. 185, 247-258, 2010

- [7] V. Eleonskii, I. Krichever, N. Kulagin, Rational multisoliton solutions of nonlinear Schrödinger equation, *Dokl. Math. Phy.*, V. 287, 606-610, 1986
- [8] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, *Max-Planck-Institut für Mathematik, MPI 02-31*, V. 161, 2002
- [9] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, *Lett. Math. Phys.*, V. 68, 77-90, 2004
- [10] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, *Lett. Math. Phys.*, V. 89, 1-12, 2009
- [11] P. Gaillard, V.B. Matveev, Wronskian and Casorati determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *J. Phys A : Math. Theor.*, V. 42, 404409-1-16, 2009
- [12] P. Dubard, P. Gaillard, C. Klein, V. B. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Spe. Top.*, V. 185, 247-258, 2010
- [13] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, *J. Phys. A : Meth. Theor.*, V. 44, 435204-1-15, 2011
- [14] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, *J. Math. Sciences : Adv. Appl.*, V. 13, N. 2, 71-153, 2012
- [15] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *J. Math. Phys.*, V. 54, 013504-1-32, 2013
- [16] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. 4, N. 4, 246-266, 2013
- [17] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. 2013, ID 645752, 1-10, 2013
- [18] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. 7, N. 45, 1-6, 2013
- [19] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. 88, 042903-1-9, 2013
- [20] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. 2013, 1-111, 2013
- [21] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. 54, 073519-1-22, 2013

- [22] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, Commun. Theor. Phys., V. 61, 365-369, 2014
- [23] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, Phys. Scripta, V. 89, 015004-1-7, 2014
- [24] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, J. Of Phys. : Conf. Ser., V. 482, 012016-1-7, 2014
- [25] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, Int. J. Mod. Phys. C, V. 26, N. 2, 1550016-1-14, 2014
- [26] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, J. Math. Phys., V. 5, 093506-1-12, 2014
- [27] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, J. Phys. : Conf. Ser., V. 574, 012031-1-5, 2015
- [28] P. Gaillard, Tenth Peregrine breather solution of the NLS, Ann. Phys., V. 355, 293-298, 2015
- [29] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation Phys. Lett. A, V. 379, 1309-1313, 2015
- [30] P. Gaillard, Other $2N-2$ parameters solutions to the NLS equation and $2N+1$ highest amplitude of the modulus of the N -th order AP breather, J. Phys. A: Math. Theor., V. 48, 145203-1-23, 2015
- [31] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, Adv. Res., V. 4, N. 5, 346-364, 2015
- [32] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, Jour. Phys. : Conf. Ser., V. 633, 012106-1-6, 2016
- [33] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, Journal of Theoretical and Applied Physics, V. 10,1-7, 2016
- [34] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, Commun. Theor. Phys, V. 65, N. 2, 136-144, 2016
- [35] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, Annals Of Physics, V. 367, 1-5, 2016

- [36] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, *Adv. Res.*, V. 10, 83-89, 2016
- [37] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, *Theor. And Math. Phys.*, V. 189, N. 1, 1440-1449, 2016
- [38] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, *Jour. of Math. Phys.*, V. 57, 063505-1-13, doi: 10.1063/1.4953383, 2016
- [39] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. 21, N. 3, 130-139, 2017
- [40] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on $2N - 2$ parameters, *Int. Jour. of Appl. Sci. And Math.*, V. 4, N. 3, 60-70, 2017
- [41] P. Gaillard, Families of rational solutions of order 5 to the KPI equation depending on 8 parameters, *New Hor. in Math. Phys.*, V. 1, N. 1, 26-31, 2017
- [42] P. Gaillard, 6-th order rational solutions to the KPI Equation depending on 10 parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. 21, N. 2, 92-98, 2017
- [43] P. Gaillard, N -Order rational solutions to the Johnson equation depending on $2N - 2$ parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. 4, N. 9, 19-37, 2017
- [44] P. Gaillard, Families of rational solutions to the KPI equation of order 7 depending on 12 parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. 4, N. 11, 24-30, 2017
- [45] P. Gaillard, Rational solutions to the Johnson equation and rogue waves, *Int. Jour. of Inn. In Sci. and Math.*, V. 6, N. 1, 14-19, 2018
- [46] P. Gaillard, Multiparametric families of solutions of the KPI equation, the structure of their rational representations and multi-rogue waves, *Theo. And Mat. Phys.*, V. 196, N. 2, 1174-1199, 2018
- [47] P. Gaillard, The Johnson equation, Fredholm and wronskian representations of solutions and the case of order three, *Adv. In Math. Phys.*, V. 2018, 1-18, 2018
- [48] P. Gaillard, Families of solutions of order 5 to the Johnson equation depending on 8 parameters, *NHIMP*, V. 2, N. 4, 53-61, 2018
- [49] P. Gaillard, Multiparametric families of solutions to the Johnson equation, *J. Phys. : Conf. Series*, V. 1141, 012102-1-10, 2018

- [50] P. Gaillard, Rational solutions to the Boussinesq equation, *Fund. Jour. Of Math. And Appl.*, V. 2, 109-112, 2019
- [51] P. Gaillard, Differential relations for the solutions to the NLS equation and their different representations, *Comm. In Adv. Math. Sci.*, V. 2, N. 4, 1-4, 2019
- [52] P. Gaillard, Multi-parametric families of solutions of order N to the Boussinesq and KP equations and the degenerate rational case, *UJMA*, V. 3, N. 2, 44-52, 2020
- [53] P. Gaillard, The mKdV equation and multi-parameters rational solutions, *Wave Motion*, V. 100, 102667-1-9, 2021
- [54] P. Gaillard, Degenerate Riemann theta functions, Fredholm and wronskian representations of the solutions to the KdV equation and the degenerate rational case, *Jour. Of Geom. And Phys.*, V. 161, 104059-1-12, 2021
- [55] P. Gaillard, $2N$ parameter solutions to the Burgers equation, *Jour. Of Appl. Nonlin. Dyn.*, V. 1, N. 1, 1-6, 2021
- [56] P. Gaillard, Multi-Parametric rational Solutions to the KdV equation, *Asi. Jour. Of Res. and Rev. In Phys.*, V. 3, N. 3, 14-21, 2021
- [57] P. Gaillard, Multiparametric solutions to the Gardner equation and the degenerate rational case, *Jour. Of Appl. Ana. And Appl.*, V. 11, N.4, 2102-2113, 2021
- [58] P. Gaillard, Other families of rational solutions to the KPI equation, *Asi. Res. Jour Of Math.*, V. 17, N. 6, 27-34, 2021
- [59] P. Gaillard, Rational solutions to the mKdV equation associated to particular polynomials, *Wave Motion*, V. 107, 102824-1-11, 2021
- [60] P. Gaillard, Multiparametric rational solutions of order N to the KPI equation and the explicit case of order 3, *Arch. Of cur. Res. Int.*, V. 21, N . 6, 58-71, 2021
- [61] P. Gaillard, Rational solutions to the KPI equation from particular polynomials, *Wave Motion*, V. 108, 102828-1-9, 2022
- [62] P. Gaillard, Rogue Waves of the Lakshmanan Porsezian Daniel Equation Depending on Multi-parameters, *As. Jour. Of Adv. Res. And Rep.*, V 16, N. 3, 32-40, 2022
- [63] P. Gaillard, Quasi-rational and rational solutions to the defocusing NLS equation, *Jour. Of Math. Sci. And Mod. Res. And Rep.*, V 5, N. 1, 24-34, 2022
- [64] P. Gaillard, Rational solutions to the KdV equation in terms of particular polynomials, *Jour. Of Nonlin. Mod. and Ana.*, V. 4, N. 4, 615-626, 2022

- [65] P. Gaillard, Rational solutions to the Painlevé II equation from particular polynomials, *Int. J. Of Appl. Nonlin. Sci.*, V. 3, N. 3, 189-196, 2022
- [66] P. Gaillard, Multi-lump solutions to the KPI equation with a zero degree of derivation, *Jour. Of Theor. And Appl. Phys.*, V. 16, N. 4, 1-8, 2022
- [67] P. Gaillard, Families of solutions to the KPI equation given by an extended Darboux transformation, *Part. Diff. Equ. And Appl.*, V. 3, N. 80, 1-14, 2022
- [68] P. Gaillard, Rogue waves of the Hirota equation in terms of quasi-rational solutions depending on multi-parameters, *WSEAS Trans. On Math.*, DOI: 10.37394/23206.2023.22.24, V. 22, 190-203, 2023
- [69] P. Gaillard, Different representations of the solutions to the cylindrical nonlinear Schrödinger equation, *Dis. Nonlin. And Compl.*, V. 12, N. 3, 539-553, 2023
- [70] P. Gaillard, Rational solutions to the KdV equation from Riemann theta functions, *Jour. Of Theor. And Appl. Phys.*, V. 17, N. 3, 1-5, 2023
- [71] P. Gaillard, On Hadamard's coefficients for two dimensional Schrödinger operators, *Int. Jour. Of Theor. Phys.*, V. 62, N. 205, 1-16, 2023
- [72] P. Gaillard, Rational solutions to the Burgers' equation from particular polynomials, *WSEAS Trans. On Math*, DOI: 10.37394/23206.2023.22.94, 867-873, 2023
- [73] P. Gaillard, Rational solutions to the Gardner equation from particular polynomials, *WSEAS Equ.*, DOI: 10.37394/232021.2023.3.2, V. 3, 1-24, 2023
- [74] P. Gaillard, Rational solutions to the KPI equation as multi-lumps with a one degree of summation, *Int. J. Appl. Comput. Math*, doi.org/10.1007/s40819-024-01694-9, 1-14, 2024
- [75] P. Gaillard, Rational solutions to the sixth equation of the NLS hierarchy, *ISRG Journal Of Multi. Stud.*, V. 2, N. 6, 1-14, 2024
- [76] P. Gaillard, Wronskian representation of the solutions to the KdV Equation, *Jour. Of App. Phys. Sci. Int.*, V. 16, N. 2, 1-11, 2024
- [77] S. Tanaka, Modified Korteweg. de Vries equation and scattering theory, *Proc. Japan Acad.*, V. 48, 466-469, 1972
- [78] M. Wadati, The exact solution of the modified Korteweg-de Vries equation, *J. Phys. Soc. Jpn.*, V. 32, 1681-1681, 1972
- [79] Y. Ono, Algebraic soliton of the modified Korteweg-de Vries equation, *Jour. Phys. Soc. Japan*, V. 41, N. 5, 1817-1818, 1976

- [80] A. Chowdury, A. Ankiewicz, N. Akhmediev, Periodic and rational solutions of modified Korteweg-de Vries equation, *Eur. Phys. J. D*, V. 70, N. 104, 1-7, 2016
- [81] A. Chowdury, A. Ankiewicz, N. Akhmediev, Periodic and rational solutions of mKdV equation, *Eur. Phys. J. D*, V. 70, N. 104, 1-7, 2016
- [82] M. Lakshmanan, K. Porsezian, M. Daniel, Effect of discreteness on the continuum limit of the Heisenberg spin chain, *Phys. Lett. A*, V. 133, N. 9, 483488, 1998
- [83] M. Lakshmanan, M. Daniel, K. Porsezian, On the integrability aspects of the one-dimensional classical continuum isotropic biquadratic Heisenberg spin chain, *J. Math. Phys.*, V. 33, N. 5, 1807-1816, 1992
- [84] M. Daniel, K. Porsezian, M. Lakshmanan, On the integrable models of the higher order water wave equation, *Phys. Lett. A*, V. 174, N. 3, 237-240, 1993
- [85] G. Akram, M. Sadaf, M. Dawood, D. Baleanu, Optical solitons for Lakshmanan Porsezian Daniel equation with Kerr law non-linearity using improved tan expansion technique, *Res. In Phys.*, V. 29, 104758-1-13, 2021
- [86] A.A. Al Qarni et al., Optical solitons for Lakshmanan Porsezian Daniel model by Riccati equation approach, *Optik*, V. 182, 922-929, 2019
- [87] R.T. Alqahtani, M.M. Babatin, A. Biswas, Bright optical solitons for Lakshmanan Porsezian Daniel model by semi-inverse variational principle, *Optik*, V. 154, 109-114, 2018
- [88] S. Arshed et al., Optical solitons in birefringent fibers for Lakshmanan Porsezian Daniel model using $\exp(-i\phi)$ -expansion method, *Optik*, V. 172, 651-656, 2018

Appendix

Solution of order 3 to the (NLS5) equation depending on 4 real parameters :
The function $v(x, t)$ defined by

$$v(x, t) = \left(1 - 24 \frac{n(x, t)}{d(x, t)} \right) e^{i(2 a_1 - 6 a_2 + 70 t)} \quad (5)$$

with

$$\begin{aligned} n(x, t) = & (675 + 69363302400 t^2 + 88473600 b_2^2 + (2x - 12 b_1 + 60 b_2)^{10} + 27000 (8 b_1 - \\ & 80 b_2)^2 + 91800 (16 a_2 - 1120 t)^2 + 2190 (4 a_1 - 24 a_2 + 560 t)^6 + 495 (4 a_1 - 24 a_2 + \\ & 560 t)^8 + 11 (4 a_1 - 24 a_2 + 560 t)^{10} - 720 (4 a_1 - 24 a_2 + 560 t)^7 (16 a_2 - 1120 t) - \\ & 3600 (4 a_1 - 24 a_2 + 560 t)^3 (48 a_2 - 15904 t) - 720 (4 a_1 - 24 a_2 + 560 t)^5 (272 a_2 - \\ & 25312 t) - 154828800 (16 a_2 - 1120 t) t + i(15422400 t + 64800 (16 a_2 - 1120 t)^3 - \\ & 870 (4 a_1 - 24 a_2 + 560 t)^7 + 25 (4 a_1 - 24 a_2 + 560 t)^9 + (4 a_1 - 24 a_2 + 560 t)^{11} - \\ & 151200 a_2 - 5529600 (8 b_1 - 80 b_2) (16 a_2 - 1120 t) b_2 - 77414400 (16 a_2 - 1120 t)^2 t - \end{aligned}$$

$$\begin{aligned}
& 90(4a_1 - 24a_2 + 560t)^8(16a_2 - 1120t) - 120(4a_1 - 24a_2 + 560t)^6(80a_2 - \\
& 11872t) + 900(4a_1 - 24a_2 + 560t)^4(464a_2 - 23520t) + (-450(4a_1 - 24a_2 + \\
& 560t)^3 - 210(4a_1 - 24a_2 + 560t)^5 + 10(4a_1 - 24a_2 + 560t)^7 + 300(4a_1 - 24a_2 + \\
& 560t)^4(16a_2 - 1120t) + 450(4a_1 - 24a_2 + 560t)(-3 + 12(8b_1 - 80b_2)^2 - 4(16a_2 - \\
& 1120t)^2) - 14400a_2 + 2620800t + 1800(4a_1 - 24a_2 + 560t)^2(16a_2 - 2016t))(2x - \\
& 12b_1 + 60b_2)^4 + (-480(4a_1 - 24a_2 + 560t)^5(8b_1 - 80b_2) + 14400(4a_1 - 24a_2 + \\
& 560t)^2(8b_1 - 80b_2)(16a_2 - 1120t) + 7200(4a_1 - 24a_2 + 560t)(8b_1 - 48b_2) - \\
& 2400(4a_1 - 24a_2 + 560t)^3(16b_1 - 128b_2) - 14400(8b_1 - 80b_2)(16a_2 - 1120t) - \\
& 460800(16a_2 - 1120t)b_2 + 12902400(8b_1 - 80b_2)t)(2x - 12b_1 + 60b_2)^3 - 21600(8b_1 - \\
& 80b_2)^2(16a_2 - 1120t) + (1710(4a_1 - 24a_2 + 560t)^5 - 60(4a_1 - 24a_2 + 560t)^7 + \\
& 5(4a_1 - 24a_2 + 560t)^9 - 900(4a_1 - 24a_2 + 560t)^3(7 + 4(8b_1 - 80b_2)^2 - 12(16a_2 - \\
& 1120t)^2) + 675(4a_1 - 24a_2 + 560t)(7 + 16(8b_1 - 80b_2)^2 + 16(16a_2 - 1120t)^2) - \\
& 345600a_2 + 38707200t - 21600(8b_1 - 80b_2)^2(16a_2 - 1120t) - 21600(16a_2 - \\
& 1120t)^3 + 9676800(4a_1 - 24a_2 + 560t)^2t - 1800(4a_1 - 24a_2 + 560t)^4(64a_2 - \\
& 1792t))(2x - 12b_1 + 60b_2)^2 + (-240(4a_1 - 24a_2 + 560t)^7(8b_1 - 80b_2) - 7200(4a_1 - \\
& 24a_2 + 560t)^4(8b_1 - 80b_2)(16a_2 - 1120t) + 10800(4a_1 - 24a_2 + 560t)(24b_1 - \\
& 400b_2 + 4(8b_1 - 80b_2)^3 + 4(8b_1 - 80b_2)(16a_2 - 1120t)^2) + 3600(4a_1 - 24a_2 + \\
& 560t)^3(24b_1 - 176b_2) + 720(4a_1 - 24a_2 + 560t)^5(56b_1 - 400b_2) + 21600(8b_1 - \\
& 80b_2)(16a_2 - 1120t) + 1382400(16a_2 - 1120t)b_2 - 38707200(8b_1 - 80b_2)t - \\
& 43200(4a_1 - 24a_2 + 560t)^2((8b_1 - 80b_2)(16a_2 - 1120t) + 32(16a_2 - 1120t)b_2 - \\
& 896(8b_1 - 80b_2)t))(2x - 12b_1 + 60b_2) + 90(4a_1 - 24a_2 + 560t)^5(-107 + 28(8b_1 - \\
& 80b_2)^2 + 12(16a_2 - 1120t)^2) + 5400(4a_1 - 24a_2 + 560t)^2(176a_2 - 22176t + \\
& 4(8b_1 - 80b_2)^2(16a_2 - 1120t) + 4(16a_2 - 1120t)^3) - 225(4a_1 - 24a_2 + 560t)^3(11 + \\
& 80(8b_1 - 80b_2)^2 + 80(16a_2 - 1120t)^2 + 4096(8b_1 - 80b_2)b_2 + 114688(16a_2 - \\
& 1120t)t) - 675(4a_1 - 24a_2 + 560t)(-7 + 56(8b_1 - 80b_2)^2 + 88(16a_2 - 1120t)^2 - \\
& 4096(8b_1 - 80b_2)b_2 - 131072b_2^2 - 102760448t^2) + 77414400(8b_1 - 80b_2)^2t + \\
& (4a_1 - 24a_2 + 560t)(2x - 12b_1 + 60b_2)^{10} + (-60a_1 + 840a_2 - 42000t + 5(4a_1 - \\
& 24a_2 + 560t)^3)(2x - 12b_1 + 60b_2)^8 + (-600a_1 - 240a_2 + 722400t - 140(4a_1 - \\
& 24a_2 + 560t)^3 + 10(4a_1 - 24a_2 + 560t)^5 + 240(4a_1 - 24a_2 + 560t)^2(16a_2 - \\
& 1120t))(2x - 12b_1 + 60b_2)^6 + (-240(4a_1 - 24a_2 + 560t)^3(8b_1 - 80b_2) - 1440(8b_1 - \\
& 80b_2)(16a_2 - 1120t) + 720(4a_1 - 24a_2 + 560t)(8b_1 - 176b_2))(2x - 12b_1 + \\
& 60b_2)^5) + 15(1 + (4a_1 - 24a_2 + 560t)^2)(2x - 12b_1 + 60b_2)^8 + (210 - 60(4a_1 - \\
& 24a_2 + 560t)^2 + 50(4a_1 - 24a_2 + 560t)^4 + 480(4a_1 - 24a_2 + 560t)(16a_2 - \\
& 1120t))(2x - 12b_1 + 60b_2)^6 + (-720(4a_1 - 24a_2 + 560t)^2(8b_1 - 80b_2) - 5760b_1 - \\
& 11520b_2)(2x - 12b_1 + 60b_2)^5 + (450(4a_1 - 24a_2 + 560t)^2 - 150(4a_1 - 24a_2 + \\
& 560t)^4 + 70(4a_1 - 24a_2 + 560t)^6 + 1200(4a_1 - 24a_2 + 560t)^3(16a_2 - 1120t) - \\
& 450 + 5400(8b_1 - 80b_2)^2 - 1800(16a_2 - 1120t)^2 + 3600(4a_1 - 24a_2 + 560t)(16a_2 - \\
& 2016t))(2x - 12b_1 + 60b_2)^4 + (-2400(4a_1 - 24a_2 + 560t)^4(8b_1 - 80b_2) + \\
& 28800(4a_1 - 24a_2 + 560t)(8b_1 - 80b_2)(16a_2 - 1120t) + 57600b_1 - 806400b_2 - \\
& 7200(4a_1 - 24a_2 + 560t)^2(16b_1 - 128b_2))(2x - 12b_1 + 60b_2)^3 + (6750(4a_1 - \\
& 24a_2 + 560t)^4 + 420(4a_1 - 24a_2 + 560t)^6 + 45(4a_1 - 24a_2 + 560t)^8 - 2700(4a_1 - \\
& 24a_2 + 560t)^2(5 + 4(8b_1 - 80b_2)^2 - 12(16a_2 - 1120t)^2) - 675 - 10800(8b_1 - \\
& 80b_2)^2 - 10800(16a_2 - 1120t)^2 + 21600(4a_1 - 24a_2 + 560t)(32a_2 - 3136t) - \\
& 7200(4a_1 - 24a_2 + 560t)^3(32a_2 + 448t))(2x - 12b_1 + 60b_2)^2 + (-1680(4a_1 - \\
& 24a_2 + 560t)^6(8b_1 - 80b_2) - 28800(4a_1 - 24a_2 + 560t)^3(8b_1 - 80b_2)(16a_2 - \\
& 1120t) - 10800(4a_1 - 24a_2 + 560t)^2(8b_1 - 272b_2) + 86400b_1 - 1209600b_2 +
\end{aligned}$$

$$\begin{aligned}
& 43200(8b_1 - 80b_2)^3 + 43200(8b_1 - 80b_2)(16a_2 - 1120t)^2 + 3600(4a_1 - 24a_2 + 560t)^4(8b_1 + 80b_2) - 86400(4a_1 - 24a_2 + 560t)((8b_1 - 80b_2)(16a_2 - 1120t) + \\
& 32(16a_2 - 1120t)b_2 - 896(8b_1 - 80b_2t))(2x - 12b_1 + 60b_2) + 450(4a_1 - 24a_2 + 560t)^4(-17 + 28(8b_1 - 80b_2)^2 + 12(16a_2 - 1120t)^2) + 10800(4a_1 - 24a_2 + \\
& 560t)(-16a_2 + 2016t + 4(8b_1 - 80b_2)^2(16a_2 - 1120t) + 4(16a_2 - 1120t)^3) + \\
& 675(4a_1 - 24a_2 + 560t)^2(-3 + 16(8b_1 - 80b_2)^2 + 16(16a_2 - 1120t)^2) - 4096(8b_1 - 80b_2)b_2 - 114688(16a_2 - 1120t)t - 2764800(8b_1 - 80b_2)b_2)
\end{aligned}$$

and

$$\begin{aligned}
d(x, t) = & 2024 + 416179814400t^2 + 530841600b_2^2 + 356400(8b_1 - 80b_2)^2 + \\
& 518400(8b_1 - 80b_2)^4 + 874800(16a_2 - 1120t)^2 + 3720(4a_1 - 24a_2 + 560t)^8 + \\
& 120(4a_1 - 24a_2 + 560t)^{10} + 518400(16a_2 - 1120t)^4 + (1 + (2x - 12b_1 + 60b_2)^2 + \\
& (4a_1 - 24a_2 + 560t)^2)^6 + (-360(4a_1 - 24a_2 + 560t)^8(8b_1 - 80b_2) - 17280(4a_1 - \\
& 24a_2 + 560t)^5(8b_1 - 80b_2)(16a_2 - 1120t) - 1440(4a_1 - 24a_2 + 560t)^6(8b_1 - \\
& 240b_2) + 32400(4a_1 - 24a_2 + 560t)^4(8b_1 + 112b_2) + 64800(4a_1 - 24a_2 + \\
& 560t)^2(-40b_1 + 752b_2 + 4(8b_1 - 80b_2)^3 + 4(8b_1 - 80b_2)(16a_2 - 1120t)^2) - \\
& 777600(4a_1 - 24a_2 + 560t)((8b_1 - 80b_2)(16a_2 - 1120t) + 64(16a_2 - 1120t)b_2 - \\
& 1792(8b_1 - 80b_2)t) - 172800(4a_1 - 24a_2 + 560t)^3(3(8b_1 - 80b_2)(16a_2 - 1120t) + \\
& 32(16a_2 - 1120t)b_2 - 896(8b_1 - 80b_2)t) - 648000b_1 + 8553600b_2 + 259200(8b_1 - \\
& 80b_2)^3 + 1296000(8b_1 - 80b_2)(16a_2 - 1120t)^2 - 33177600(8b_1 - 80b_2)^2b_2 + \\
& 33177600(16a_2 - 1120t)^2b_2 - 1857945600(8b_1 - 80b_2)(16a_2 - 1120t)t(2x - \\
& 12b_1 + 60b_2) + 80(4a_1 - 24a_2 + 560t)^6(191 + 63(8b_1 - 80b_2)^2 + 27(16a_2 - \\
& 1120t)^2) + 21600(4a_1 - 24a_2 + 560t)^3(-368a_2 + 23072t + 4(8b_1 - 80b_2)^2(16a_2 - \\
& 1120t) + 4(16a_2 - 1120t)^3) + 120(8b_1 - 80b_2)(2x - 12b_1 + 60b_2)^9 + 46080b_2(2x - \\
& 12b_1 + 60b_2)^7 - 1161216000(16a_2 - 1120t)t + 240(4a_1 - 24a_2 + 560t)^4(599 + \\
& 135(8b_1 - 80b_2)^2 - 225(16a_2 - 1120t)^2 - 11520(8b_1 - 80b_2)b_2 - 322560(16a_2 - \\
& 1120t)t) - 16200(4a_1 - 24a_2 + 560t)(496a_2 - 52640t + 80(8b_1 - 80b_2)^2(16a_2 - \\
& 1120t) + 16(16a_2 - 1120t)^3 + 4096(8b_1 - 80b_2)(16a_2 - 1120t)b_2 - 57344(8b_1 - \\
& 80b_2)^2t + 57344(16a_2 - 1120t)^2t) + 24(4a_1 - 24a_2 + 560t)^2(3881 + 12150(8b_1 - \\
& 80b_2)^2 + 28350(16a_2 - 1120t)^2 + 691200(8b_1 - 80b_2)b_2 + 22118400b_2^2 + 17340825600t^2) + \\
& (-120(4a_1 - 24a_2 + 560t)^2 + 360(4a_1 - 24a_2 + 560t)(16a_2 - 1120t) + 120)(2x - \\
& 12b_1 + 60b_2)^8 + (480(4a_1 - 24a_2 + 560t)^2 - 240(4a_1 - 24a_2 + 560t)^4 + 960(4a_1 - \\
& 24a_2 + 560t)^3(16a_2 - 1120t) + 2320 + 2160(8b_1 - 80b_2)^2 + 5040(16a_2 - 1120t)^2 - \\
& 1440(4a_1 - 24a_2 + 560t)(64a_2 - 8960t))(2x - 12b_1 + 60b_2)^6 + (-720(4a_1 - \\
& 24a_2 + 560t)^4(8b_1 - 80b_2) - 17280(4a_1 - 24a_2 + 560t)(8b_1 - 80b_2)(16a_2 - \\
& 1120t) + 4320(4a_1 - 24a_2 + 560t)^2(8b_1 - 176b_2) - 51840b_1 + 103680b_2)(2x - \\
& 12b_1 + 60b_2)^5 + (-1440(4a_1 - 24a_2 + 560t)^4 + 720(4a_1 - 24a_2 + 560t)^5(16a_2 - \\
& 1120t) + 240(4a_1 - 24a_2 + 560t)^2(56 + 135(8b_1 - 80b_2)^2 - 45(16a_2 - 1120t)^2) + \\
& 32400(4a_1 - 24a_2 + 560t)(16a_2 - 2912t) + 7200(4a_1 - 24a_2 + 560t)^3(48a_2 - \\
& 4256t) + 3360 + 32400(8b_1 - 80b_2)^2 - 54000(16a_2 - 1120t)^2 + 2764800(8b_1 - \\
& 80b_2)b_2 + 77414400(16a_2 - 1120t)t(2x - 12b_1 + 60b_2)^4 + (-960(4a_1 - 24a_2 + \\
& 560t)^6(8b_1 - 80b_2) + 57600(4a_1 - 24a_2 + 560t)^3(8b_1 - 80b_2)(16a_2 - 1120t) - \\
& 43200(4a_1 - 24a_2 + 560t)^2(24b_1 - 272b_2) - 7200(4a_1 - 24a_2 + 560t)^4(48b_1 - \\
& 448b_2) + 345600b_1 - 5529600b_2 - 86400(8b_1 - 80b_2)^3 - 86400(8b_1 - 80b_2)(16a_2 - \\
& 1120t)^2 + 172800(4a_1 - 24a_2 + 560t)((8b_1 - 80b_2)(16a_2 - 1120t) - 32(16a_2 - \\
& 1120t)b_2 + 896(8b_1 - 80b_2)t))(2x - 12b_1 + 60b_2)^3 + (13440(4a_1 - 24a_2 + \\
& 560t)^6 + 240(4a_1 - 24a_2 + 560t)^8 - 240(4a_1 - 24a_2 + 560t)^4(-326 + 45(8b_1 -
\end{aligned}$$

$80 b_2)^2 - 135 (16 a_2 - 1120 t)^2) + 480 (4 a_1 - 24 a_2 + 560 t)^2(-76 + 135 (8 b_1 - 80 b_2)^2 + 1215 (16 a_2 - 1120 t)^2) - 129600 (4 a_1 - 24 a_2 + 560 t)^3(32 a_2 - 1344 t) - 12960 (4 a_1 - 24 a_2 + 560 t)^5(32 a_2 - 1344 t) - 64800 (4 a_1 - 24 a_2 + 560 t)(-96 a_2 + 11200 t + 4 (8 b_1 - 80 b_2)^2(16 a_2 - 1120 t) + 4 (16 a_2 - 1120 t)^3) + 12144 - 97200 (8 b_1 - 80 b_2)^2 + 32400 (16 a_2 - 1120 t)^2 + 530841600 b_2^2 - 464486400 (16 a_2 - 1120 t)t + 416179814400 t^2)(2 x - 12 b_1 + 60 b_2)^2 - 2160 (4 a_1 - 24 a_2 + 560 t)^5(240 a_2 - 47264 t) - 1440 (4 a_1 - 24 a_2 + 560 t)^7(80 a_2 - 6496 t) - 120 (4 a_1 - 24 a_2 + 560 t)^9(16 a_2 - 1120 t) + 1036800 (8 b_1 - 80 b_2)^2(16 a_2 - 1120 t)^2 - 24883200 (8 b_1 - 80 b_2)b_2$

is a solution to the (NLS5) equation (1).