



**HAL**  
open science

## Evolution of the contact network during tilting cycles of a granular pile under gravity

Stéphanie Deboeuf, Jean-Pierre Vilotte, Anne Mangeney, Lydie Staron,  
Olivier Dauchot

► **To cite this version:**

Stéphanie Deboeuf, Jean-Pierre Vilotte, Anne Mangeney, Lydie Staron, Olivier Dauchot. Evolution of the contact network during tilting cycles of a granular pile under gravity. International Conference on Powders & Grains 2005, Jul 2005, Stuttgart, Germany. 10.1201/NOE0415383486 . hal-04712548

**HAL Id: hal-04712548**

**<https://hal.science/hal-04712548v1>**

Submitted on 27 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# Evolution of the contact network during tilting cycles of a granular pile under gravity

S. Deboeuf, J.-P. Vilotte & A. Mangeney

*Institut de Physique du Globe de Paris, France*

L. Staron

*Department of Applied Mathematics and Theoretical Physics, Cambridge, UK*

O. Dauchot

*Commissariat à l'Energie Atomique, Saclay, France*

**ABSTRACT:** We perform discrete numerical simulations of a granular pile undergoing quasi-static tilting cycles in the gravity field. The volumetric deformation and the granular fabric exhibit hysteretic evolutions with the slope of the free surface. When exploring the range of metastable slopes -between the angle of repose and the angle of avalanche-, the contact network is strongly affected, as evidenced by the qualitative change of the hysteresis. A specific contribution to the observed hysteresis of the weak contacts is underlined: they carry the memory of the proximity of the slope destabilization.

## 1 INTRODUCTION

When inclined in the gravity field, a granular slope starts flowing when the surface slope reaches the avalanche angle  $\theta_a$ , and comes to rest afterwards at the angle of repose  $\theta_r < \theta_a$ . In between these two angles, the slope is metastable: avalanches can be triggered by finite amplitude perturbations (Daerr & Douady 1999). Recent numerical studies investigated the mechanisms of destabilization of granular piles during continuous load in the gravity field (Staron et al. 2002). Far before the avalanche, the authors observe the occurrence of local instabilities leading to the compaction of the pile. At the approach of the stability limit, they evidence dilatancy simultaneously with the percolation of clusters of critical contacts, that are essentially weak contacts. These results lead them to identify the major role of spatial correlations in the range of metastable slopes  $[\theta_r; \theta_a]$ . Similar correlations were also experimentally observed in Deboeuf et al. (2003) and Kabla et al. (2004).

The metastability of the slope can be seen as a signature of a major modification of the internal state of the granular packing. In order to investigate the nature of these modifications, and to study the influence of this metastability on the following evolution, we simulate by contact dynamics the quasi-static cyclic evolution of a granular pile tilted in the gravity field towards the slopes  $[\theta_r; \theta_a]$ . Details of the numerical procedure are

given in section 2. An analysis of the contact network in section 3 shows that the volumetric strain and the granular fabric evolutions are hysteretic. The shape of the hysteresis is strongly affected when the pile slope  $\theta$  explores the metastable range  $[\theta_r; \theta_a]$ . In section 4 we observe the contributions of the strong and weak contacts and we evidence the specific role of memory played by the latter.

## 2 NUMERICAL SET-UP

The discrete numerical simulations are performed using the contact dynamics method (Moreau 1994). The grains are perfectly rigid, and interact via frictional contacts governed by a Coulomb friction law ( $\mu = 0.5$ ). The granular pile is made of a two dimensional rectangular box filled with 4000 circular grains of mean diameter  $D$ , with a thickness of  $\approx 35D$ , and a width of  $\approx 120D$ . The solicitation consists in quasi-static rotations in the gravity field at a rate of  $0.001^\circ$  per time step. Initially, the slope increases from  $\theta = 0^\circ$  to a maximal inclination angle  $\theta_{rev}$ . At that stage, two successive cycles  $[-\theta_{rev}; \theta_{rev}]$  are performed. All the simulations are averaged over 50 granular piles differing in the initial grains arrangements, and showing characteristic angles  $\theta_r \approx 15^\circ$  and  $\theta_a \approx 21^\circ$ .

To study whether exploring the metastable regime influences the evolution of the granular pile, two values




Figure 1. Measure of the surface slope of the pile during the rotation cycles of amplitude  $\theta_{rev} = 10^\circ$  and  $\theta_{rev} = 18^\circ$ .

of the maximal inclination angle  $\theta_{rev}$  are investigated. Figure 1 shows the time evolution of the pile slope averaged over 50 simulations. The smallest amplitude is chosen in order to keep inclination below the metastable range:  $\theta_{rev} = 10^\circ < \theta_r$ . The largest amplitude is chosen such that the metastable regime is explored during each cycle, still below the minimal avalanche angle to prevent the pile from avalanching:  $\theta_{rev} = 18^\circ \in [\theta_r; \theta_a]$ .

### 3 EVOLUTION OF THE CONTACT NETWORK

#### 3.1 Volumetric strains of the granular pile

During each loading and unloading phase, local instabilities occur intermittently within the granular pile. The effect of these rearrangements of the grains is apparent when plotting the volumetric strain  $\epsilon_V = (V - V_0)/V_0$ , where  $V$  is the volume of the pile and  $V_0$  its initial volume, as a function of the slope of the pile (Fig. 2).

Irrespective of the value of  $\theta_{rev}$ , cycles produce an overall densification of the pile. Note that the consolidation is of the same order ( $10^{-3}$ ) as via avalanching from Barker & Mehta (2000). For small cycles ( $\theta_{rev} = 10^\circ$ ), the granular pile is always contracting:  $\epsilon_V$  decreases. On the contrary, for large cycles ( $\theta_{rev} = 18^\circ$ ), the granular pile exhibits compaction and dilatancy stages. Indeed for slopes in the metastable range  $[\theta_r; \theta_a]$ , the rearrangements cause the granular pile to dilate:  $\epsilon_V$  increases, as already observed by Staron *et al.* Interestingly, the dilatancy stage occurring when  $\theta \in [\theta_r; \theta_a]$ , changes dramatically the overall behavior of the pile over the complete solicitation: the compaction is twice more efficient for large cycles than for small ones.

#### 3.2 Coordination number of the grains

For both values of the amplitude  $\theta_{rev}$ , the coordination number averaged over all the grains  $Z$ , *i.e.* the mean




Figure 2. Estimation of the volumetric strain as a function of the pile slope (same symbols as figure 1).




Figure 3. Coordination number of the grains as a function of the pile slope (same symbols as figure 1).

number of contacts per grains, varies less than 1% (Fig. 3). Nevertheless, it exhibits a weak hysteresis according to the sense of rotation:  $Z$  tends to decrease during load, but to increase during unload.

Small and large cycles are identified by two different states: the value of the coordination number depends on  $\theta_{rev}$ . During the small cycles,  $Z$  remains approximately at its initial value, in contrast with the large cycles, for which a variation of the coordination number occurs when exploring the metastable range of slopes  $[\theta_r; \theta_a]$  enhances the susceptibility for grains to rearrange and deeply modifies the state of the granular packing according to its initial one.

#### 3.3 Granular fabric of the pile

To analyze further the micro-mechanical state of the granular pile, we investigate the statistics of the




Figure 4. Anisotropic intensity of the contact network as a function of the pile slope (same symbols as figure 1).




Figure 5. Anisotropic direction of the contact network as a function of the pile slope (same symbols as figure 1).

orientation of contacts. To do so, we compute the fabric tensor  $\mathbf{t}$  using the following definition:

$$t_{ij} = \frac{1}{n^c} \sum_{\alpha=1}^{n^c} n_i^\alpha n_j^\alpha, \quad (1)$$

where  $i$  and  $j$  denote the space dimension,  $n$  is the unity vector normal to the contact surface, and  $n^c$  is the total number of contacts  $\alpha$ . The anisotropy of the granular pile is analyzed with respect to the slope in terms of the anisotropic intensity  $\Lambda$  ( $= 2 \times$  the deviatoric component of  $\mathbf{t}$ ) and the direction of anisotropy  $\Phi$  with respect to the normal direction to the free surface (Fig. 4 and 5).

During the small cycles, a rather isotropic state is observed, and only small variations with  $\theta$  are observed ( $\Lambda \simeq 0.05$ ): the granular fabric is only weakly affected by the solicitation and remains close to the initial one. On the contrary, for large cycles, the granular fabric

evolves significantly up to  $\Lambda \simeq 0.12$ . Note that during large cycles, the asymmetry of the initial state is removed on the granular fabric, by contrast with small ones.

Irrespective of the value of  $\theta_{rev}$ , the direction of anisotropy tends to approach that of the free surface. For large inclinations,  $\Phi$  tends towards  $\pm 45^\circ$ . This behavior corresponds with the mechanisms of creation of contacts in the direction of compressive stress, and the loss of contacts in the direction of extension. Indeed in the particular case of a hydrostatic infinite slope inclined in the gravity field, continuous media mechanics predicts that the principal direction of stress is  $\pm 45^\circ$  with respect to the free surface.

Besides these general observations, both  $10^\circ$  and  $18^\circ$ -cycles exhibit hysteresis in the evolution of the granular fabric. Furthermore when the granular pile explores the metastable regime, the shape and the amplitude of the hysteresis are dramatically changed, pointing out the peculiar effect of this regime on the granular packing evolution. In particular, the fabric evolves less rapidly during unloading than during loading.

Altogether, this makes the state of the pile at an angle  $\theta$  depend strongly on the history: the previous proximity of the destabilization is apparent on the observed characteristics. Knowing that dense granular media exhibit two contact subnetworks, strong and weak, depending on the intensity of the force transmitted at the contacts (Radjai et al. 1998), the role of the respective contributions of the two subnetworks is now investigated.

#### 4 BI-MODAL RESPONSE OF THE FABRIC

A vertical gradient of the contact forces is related to the gravity field, such that the force intensity averaged over  $h$ -altitude contacts obeys:  $\langle f \rangle(h) \propto h$ . Therefore a  $h$ -altitude contact is defined as strong (resp. weak) if it transmits a normal force larger (resp. smaller) than the normal force averaged over all  $h$ -altitude contacts. The statistical analysis of the fabric is now restricted to strong and to weak contacts, from which the anisotropic direction of the strong ( $\Phi^s$ ) and the weak subnetwork ( $\Phi^w$ ) are calculated.

The evolution of  $\Phi^s$  as a function of the pile slope exhibits also hysteresis (Fig. 6). However, by contrast with all previous observations, the shape of the hysteresis is the same for both small and large cycles. Exploring the metastable regime does not affect the strong contact subnetwork.

On the contrary, the evolution of  $\Phi^w$  as a function of the pile slope (Fig. 7) is very different according to the cycle amplitude  $\theta_{rev}$ . For small cycles ( $\theta_{rev} = 10^\circ$ ),  $\Phi^w$  exhibits almost no hysteresis, and the hysteretic behavior of the pile (see Fig. 5) is therefore governed by the strong contacts only. During small cycles,  $\Phi^w$  remains




Figure 6. Fabric anisotropic direction of strong contacts as a function of the pile slope (same symbols as figure 1).




Figure 7. Fabric anisotropic direction of weak contacts as a function of the pile slope (same symbols as figure 1).

approximately always normal to  $\Phi^s$ , as observed by Staron *et al.* in the case of continuous load. On the contrary, for large cycles, a significant hysteretic behavior is observed in the evolution of  $\Phi^w$ . It consists in an early rotation of  $\Phi^w$ , which leads to a remarkable orientation of the fabric of the two contact subnetworks:  $\Phi^s$  and  $\Phi^w$  are now aligned during a large part of the cycle, when exploring the metastable regime.

Two conclusions can be drawn from these observations:

- For small cycles, the hysteretic behavior of the pile can be seen as a result of the strong contact subnetwork contribution;

- For large cycles, the weak contact subnetwork becomes strongly affected, and the hysteretic behavior of the whole pile is strongly modified despite the identical response of the strong subnetwork.

## 5 CONCLUSIONS

When performing quasi-static tilting cycles in the gravity field, the overall behavior of a granular pile is towards densification at constant coordination number. However differences of consolidation are observed between small and large cycles, due to the dilatancy stage occurring in the metastable range of slopes, that tends to enhance rearrangements of grains. Accordingly the evolution of volumetric strains on geomaterial samples could be measured for the detection of previous proximities of the destabilization. Further analyses confirm the peculiar effect of the metastable regime on the grain packing organization: the hysteretic evolution of the fabric is deeply modified when exploring the metastable regime. Finally, this hysteretic behavior is the result of a complex interplay between the strong and weak contact subnetworks. Such a specific interaction has to be considered when investigating the mechanical properties. For instance, it suggests to distinguish strong and weak contacts when appreciating stability limit analyses.

## REFERENCES

- Barker, G. & Mehta, A. 2000. Avalanches at rough surfaces. *Phys. Rev. E* 61(6).
- Daerr, A. & Douady, S. 1999. Two types of avalanche behaviour in granular media. *Nature* 399.
- Deboeuf, S., Bertin, E.M., Lajeunesse, E. & Dauchot, O. 2003. Jamming transition of a granular pile below the angle of repose. *Eur. Phys. J. B* 36.
- Kabla, A., Debregeas, G., di Meglio, J.-M., & Senden, T. J. 2004. X-ray observation of micro-failures in granular piles approaching an avalanche. in prep.
- Moreau, J.-J. 1994. Some numerical methods in multibody dynamics: Application to granular materials. *European Journal of Mechanics A/Solids*: 93–114.
- Radjai, F., Wolf, D.E., Jean, M. & Moreau, J.-J. 1998. Bimodal character of stress transmission in granular packings. *Phys. Rev. Lett.* 80(1).
- Staron, L., Vilotte, J.-P., & Radjai, F. 2002. Preavalanche instabilities in a granular pile. *Phys. Rev. Lett.* 89(20).