

Adults construct their representation of pain in babies cries with experience

Siloé Corvin, Alexis Koutseff, David Reby, Roland Peyron, Nicolas Mathevon,
Camille Fauchon

▶ To cite this version:

Siloé Corvin, Alexis Koutseff, David Reby, Roland Peyron, Nicolas Mathevon, et al.. Adults construct their representation of pain in babies cries with experience. 12th Congress of the European Pain Federation EFIC, Apr 2022, Dublin, Ireland. hal-04719265

HAL Id: hal-04719265 https://hal.science/hal-04719265v1

Submitted on 3 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Adults construct their representation of pain in babies cries with experience

Siloé Corvin^{1,2}, Alexis Koutseff², David Reby², Roland Peyron¹, Nicolas Mathevon², Camille Fauchon¹




¹NeuroPain / CRNL, University of Saint-Etienne, CNRS, Inserm, Saint-Etienne, France ²Equipe de Neuro-Ethologie Sensorielle ENES / CRNL, University of Saint-Etienne, CNRS, Inserm, Saint-Etienne, France.

Contact: Siloé Corvin, siloe.corvin@inserm.fr; Camille Fauchon, camille.fauchon@univ-st-etienne.fr

P360

Background: babies cries convey information through acoustic differences



Individual characteristics

Familiar babies are recognized by parents¹ and non-parents²

Experience and exposure improve this ability

Emotional and physiological state


Pain cries are longer, louder, harsher and rougher \wp Parents identify a pain state by listening to their baby's cries³

Does an individual's experience to take care of babies and familiarity with a given baby improve caregivers' ability to recognize pain in cries?

Experimental design

Online experiment

234 participants (133 women) with various experience in babies care



Training with discomfort cries familiar baby

Context recognition accuracy

Bayesian mixed model

75%

50%

75%

50%


75%

50%

75%

50%


PAIN



"Was it a cry of discomfort or pain?"

Test with discomfort and pain cries familiar and unknown babies





Five levels of experience

No experience (no previous contact with babies; N = 50)

Moderate experience (babysitting, younger siblings; N = 51)

Strong experience
parents
(with > 5 years-old baby; N = 50)

Strong experience

parents

(with < 2 years-old baby; N = 52)

Strong experience

non parents

(professional caregivers; N = 31)

Ability to identify cries context

Poor recognition of cries context

Effective recognition of pain cries for a familiar baby

Effective recognition of cries context for a familiar baby and of bath context for an unknown baby

Effective recognition of cries

context for both a familiar

and an unknown baby

Effective recognition of cries context for a familiar baby and of bath context for an

unknown baby

DIS discomfort cry

PAIN pain cry

unknown baby

familiar baby

50 % = chance level for context recognition (discomfort or pain)

Conclusion

Previous **experience** with babies **facilitates** the correct **identification** of acoustic features coding for pain in baby cries.


Only parents with younger babies were able to generalize and identify crying contexts for an unknown baby, even professional caregivers were not.



Scan the QR code to test a short version of the experiment.







References: 1 Gustafsson et al., 2013, Nat.Commun.; 2 Bouchet et al., 2020, Proc.Biol.Sci.; 3 Koutseff et al., 2017, Bioacoustics