



HAL
open science

Higher abundance of disturbance-favoured trees and shrubs in European temperate woodlands prior to the late-quaternary extinction of megafauna

Elena A Pearce, Florence Mazier, Ralph Fyfe, Charles W Davison, Signe Normand, Maria-antonia Serge, Paolo Scussolini, Jens-christian Svenning

► To cite this version:

Elena A Pearce, Florence Mazier, Ralph Fyfe, Charles W Davison, Signe Normand, et al.. Higher abundance of disturbance-favoured trees and shrubs in European temperate woodlands prior to the late-quaternary extinction of megafauna. *Journal of Ecology*, 2024, 10.1111/1365-2745.14422 . hal-04719507

HAL Id: hal-04719507

<https://hal.science/hal-04719507v1>

Submitted on 3 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

RESEARCH ARTICLE

Higher abundance of disturbance-favoured trees and shrubs in European temperate woodlands prior to the late-quaternary extinction of megafauna

Elena A. Pearce¹  | Florence Mazier²  | Ralph Fyfe³ | Charles W. Davison¹  |
Signe Normand⁴  | Maria-Antonia Serge²  | Paolo Scussolini⁵  |
Jens-Christian Svenning¹ 

¹Center for Ecological Dynamics in a Novel Biosphere (ECONOVO), Department of Biology, Aarhus University, Aarhus C, Denmark

²Department of Environmental Geography, CNRS UMR GEODE 5602, University Toulouse Jean Jaurès, Toulouse, France

³School of Geography, Earth and Environmental Sciences, University of Plymouth, Plymouth, UK

⁴Center for Sustainable Landscapes Under Global Change (SustainScapes) & Center for Landscape Research in Sustainable Agricultural Futures, Department of Biology, Aarhus University, Aarhus C, Denmark

⁵Institute for Environmental Studies, Vrije Universiteit Amsterdam, Amsterdam, Netherlands

Correspondence

Elena A. Pearce
Email: elena.pearce@bio.au.dk

Funding information

Danmarks Grundforskningsfond, Grant/Award Number: DNR173; Villum Fonden, Grant/Award Number: 16549; Danmarks Frie Forskningsfond, Grant/Award Number: 0135-00225B; Horizon 2020 Research and Innovation Programme, Grant/Award Number: 813904

Handling Editor: Guillaume de Lafontaine

Abstract

1. Large herbivorous mammals strongly influence vegetation structure by creating and maintaining open areas and causing disturbance within closed woody habitats. The herbivores alive today in Europe are only a small remnant of the large species that existed in high diversity and abundance before modern humans. The extinction of so many large herbivores during the last 50,000 years, and the loss of megaherbivores (body weight ≥ 1000 kg) from most of the continent before the Pleistocene–Holocene transition, is likely to have had cascading effects on vegetation structure and composition.
2. To evaluate these effects within the European temperate forest biome, we examine the abundance change of three important European woody taxa (deciduous oaks, *Quercus* spp.; hazel, *Corylus* [mainly *C. avellana*]; and yew, *Taxus baccata*) before and after the late-Quaternary downgrading of the region's large herbivore fauna. These taxa are disturbance-favoured, depend on canopy openings for regeneration, and tend to decline in closed dense forests. *Quercus* and *Corylus* may thrive in systems affected by megafauna herbivory or fire, whilst *Taxus* is fire-sensitive but can thrive in grazed systems. Using pollen-based reconstructions (REVEALS), we investigated the proportional cover of these three focal taxa in the Last Interglacial (129,000–116,000 before present [BP]) and early–mid-Holocene (8700–5700 BP).
3. We found that woodlands in the Last Interglacial exhibited higher cover of *Corylus* and *Taxus* relative to the Holocene, with the former reaching very high percentage cover; meanwhile, *Quercus* had a consistent, moderately high percentage cover in both periods. Furthermore, we found that the cover of *Corylus* and *Taxus* appeared to be influenced more by unmeasured, non-climatic factors than *Quercus*.
4. **Synthesis:** The abundance of *Taxus* suggests a limited role of fire; whilst the observed levels of *Taxus*, *Corylus* and *Quercus* align with the potential influence of

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2024 The Author(s). *Journal of Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

megafauna herbivory; however, a direct quantitative link remains to be established. Our results suggest that vegetation was structurally and compositionally affected by differences in disturbance regimes between the Last Interglacial and early-mid-Holocene, with the loss of diverse disturbance regimes likely contributing to the divergence of Holocene vegetation from long-term ecological baselines.

KEYWORDS

disturbance regimes, European forests, forest structure, megafauna, reference conditions

1 | INTRODUCTION

Large herbivorous mammals (hereafter, 'megafauna') have strong interactions with vegetation. Megafauna consume vegetation in large quantities, thereby altering plant biomass, community composition and structure (Pringle et al., 2023). Megafauna also cause physical disturbance to ecosystems by trampling, debarking and rooting (Pringle et al., 2023). Due to these processes, megafauna play an important role in creating and maintaining open areas. This is the case in the temperate forest biome, where closed woody vegetation would otherwise dominate under mild climatic conditions (Svenning, 2002). Today, megafauna assemblages are a fraction of their past size (Smith et al., 2018). Larger species existed in much greater diversity and abundance before the arrival of modern humans ~50,000 years ago (Bergman et al., 2023; Smith et al., 2018). The extinction of so many megafauna species likely had cascading effects on the abundance, structure and composition of vegetation communities (Bakker et al., 2016; Svenning et al., 2024). Extant smaller species such as *Capreolus capreolus* (roe deer), *Cervus elaphus* (red deer) and *Sus scrofa* (wild boar) are unlikely to have produced functionally equivalent vegetation interactions (Pringle et al., 2023). In the European temperate forest biome, pollen-based reconstructions have revealed high percentages of open and light-woodland vegetation before modern humans (Pearce et al., 2023). Climate and other environmental processes, such as soil moisture availability (le Roux et al., 2013) and degree of continentality (Giesecke et al., 2008), may have influenced vegetation patterns at this time. However, it is likely that megafauna assemblages, and their associated disturbance regimes, had a major influence on vegetation structure, composition and species interactions (Davoli et al., 2023; Svenning et al., 2024).

Many trees and shrubs depend on disturbance for reproduction, to reduce competition from other taxa and to provide open, sunlit conditions (Vera, 2000). However, different disturbance mechanisms can have diverging effects. For example, *Pinus* (pine), *Betula* (birch) and *Salix* (willow) are positively related to fire events (Molinari et al., 2020), whereas *Quercus* (oak) and *Corylus* (hazel) are fire-tolerant but equally benefit from canopy openings driven by fire or large grazers (Molinari et al., 2020; Vera, 2000). Meanwhile, *Taxus* (yew) is sensitive to fire but can thrive in grazed systems (Busing et al., 1995; Omarova & Asadulaev, 2016). Evidence suggests that intense fire regimes may not have been widespread in Europe

before the arrival of modern humans (Milner et al., 2016; Sandom, Faurby, et al., 2014). Conversely, the megafauna-rich conditions of past European landscapes may have promoted taxa such as *Quercus*, *Corylus* and *Taxus* (Pearce et al., 2023; Svenning & Magård, 1999).

Deciduous *Quercus* (deciduous oaks, with *Quercus robur* and *Q. petraea* being the most widespread and the only species in northern parts of Europe; hereafter 'Quercus') is a light-demanding taxon that thrives in disturbed, transitional habitats (Bobiec et al., 2018). *Quercus* seed dispersal and recruitment occur mostly in dynamic, mosaic landscapes; the taxon fails to regenerate under dense canopies in modern European forests (Bobiec et al., 2018). Abundant *Corylus* (hazel; largely *Corylus avellana*: common hazel; hereafter 'Corylus') is usually indicative of scrub woodland subject to ongoing disturbance (Vera, 2000). *Corylus* may thrive under the canopy of lightly shading trees such as *Quercus* and *Fraxinus excelsior* (common ash) (Coppins & Coppins, 2003); however, *Corylus* seedlings require high-light conditions and *Corylus* pollen production is enhanced in open and sunny sites (Bégeot, 1998; Vera, 2000). As a result, it is considered a ready coloniser of open landscapes (Kollmann & Schill, 1996; Vera, 2000) and fails to regenerate under a dense canopy (Vera, 2000). *Taxus baccata* (common yew; hereafter 'Taxus') is usually considered a shade-tolerant taxon, but also thrives in full sun (Linares, 2013). However, in modern forests, *Taxus* depends on canopy openings for reproduction, regeneration, growth and survival and declines under a dense canopy of shading trees (Linares, 2013; Svenning & Magård, 1999). *Taxus* requires substantial light for optimal growth, experiencing significant reductions in growth rates, biomass allocation and reproductive success under shaded conditions (Iszkuło et al., 2012; Perrin & Mitchell, 2013). Limited light availability in dense canopies, dominated by faster- and taller-growing species such as *Abies alba* (silver fir) and *Fagus sylvatica* (beech), hampers *Taxus* growth, survival and regeneration (Iszkuło et al., 2012; Perrin & Mitchell, 2013). Consequently, *Taxus* populations decline in dense forests where light is scarce (Iszkuło et al., 2012; Perrin & Mitchell, 2013). Additionally, *Taxus* is highly toxic to many large herbivores, except for deer and possibly certain other ruminants (Cortinovis & Caloni, 2015; Thomas & Polwart, 2003) and therefore may outcompete species more susceptible to browsing (Dhar et al., 2007; García & Ramón Obeso, 2003). On the other hand, *Taxus* is vulnerable to fire due to its thin bark and difficulty regenerating, particularly in dry environments (Mola et al., 2014; Thomas & Polwart, 2003) and its slow

growth and low seedling recruitment further delay population recovery post-fire (Thomas & Polwart, 2003). Therefore, *Taxus* would likely have been favoured in past ecosystems with high levels of herbivory provided by diverse megafauna (Davoli et al., 2023) but would have been disfavoured by frequent fires.

Quantifying vegetation communities before and after the late-Quaternary megafauna extinctions is challenging. The extinctions occurred with the arrival of *Homo sapiens* ~50,000BP, during the last glacial period (Bergman et al., 2023; Svenning et al., 2024), when forests and their tree species had strongly reduced distributions and abundances due to the cold, dry climate (Svenning et al., 2008; Willis & van Andel, 2004). Modern conservation practices are increasingly focused on restoring ecological function (Perino et al., 2019) and the role of long-term data (e.g. on functional baselines) has been recognised for almost 20 years (Willis et al., 2005, 2010). Pre-degradation baselines can provide unique ecological insights into systems before human-induced degradation, supplying modern restoration efforts with important context. However, we note that reconciling these insights with modern restoration efforts requires acknowledging contemporary ecological complexities, such as changing climate regimes and other anthropogenic influences, such as wildfire initiation, which may necessitate additional considerations beyond pre-degradation baselines alone. To establish a pre-degradation baseline for current and future systems, warm interglacial conditions are most relevant. The Last Interglacial in Europe (Eemian; ~129,000–116,000BP) is the most recent interglacial period prior to the Holocene. It occurred before the late-Quaternary megafauna extinctions and had a faunal structure broadly comparable to the preceding 10 million years or more (Blanco et al., 2021; Croitor & Brugal, 2010). This period is geologically recent enough to limit evolutionary changes, as most extant species were already present (Van Kolschoten, 2000). Finally, the Last Interglacial is climatically comparable to the warming Holocene, particularly in its central mesocratic phases, and has been described as a testbed for assessing environmental responses and climate feedbacks under conditions warmer than the pre-industrial benchmark (Kalis et al., 2003; Kühl et al., 2007; Salonen et al., 2018). Therefore, the Last Interglacial presents a long-term representative baseline for vegetation structure and dynamics under warm conditions prior to the late-Quaternary faunal downgrading (Smith et al., 2018).

Pollen-based vegetation reconstruction using the REVEALS model (Sugita, 2007) has shown that *Quercus* and *Corylus* were present in high abundance during the Last Interglacial in Europe (Pearce et al., 2023). Studies using raw pollen data from the Last Interglacial have shown a *Corylus* peak of up to 50% (Rychel et al., 2014; Suchora et al., 2022). Past abundances of *Quercus* and *Corylus* are better known for the Holocene, where REVEALS modelling shows 1%–15% *Quercus* cover and 10%–15% mean *Corylus* cover at 8200–5700BP (Githumbi et al., 2022; Serge et al., 2023). *Taxus* was present during the Last Interglacial to a varying degree (3%–15% in Central Europe; Malkiewicz, 2018; Schläfli et al., 2021). In the Holocene, *Taxus* pollen is infrequent and cover is considered low (2%–6%; Deforce & Bastiaens, 2007; Pérez Díaz et al., 2013). However, in both the Holocene and the Last Interglacial, *Taxus* cover has only

been examined using raw pollen percentages, which are less robust than REVEALS modelling for reconstructing past cover (Anderson et al., 2006; Sugita, 2007).

In this study, we compare the abundances of *Quercus*, *Corylus* and *Taxus* to assess how their populations changed before and after the late-Quaternary megafauna extinctions in the temperate forest biome. We quantify percentage cover for each taxon in the Last Interglacial and early-mid-Holocene using the latest REVEALS reconstructions (Pearce et al., 2023; Serge et al., 2023). Whilst direct testing of megafauna impact on vegetation was unfeasible in this study, we assess the extent to which observed abundance differences relate to climatic variations and estimate the residual variance. We were specifically interested in answering the following research questions: (1) How does the percentage cover of *Quercus*, *Corylus* and *Taxus* differ between the Last Interglacial and the early-mid-Holocene (before the onset of agriculture)? (2) How far can these differences be explained by climatic differences between the two periods? Overall, we expected that *Quercus*, *Corylus* and *Taxus* would be less abundant in the early-mid-Holocene than in the Last Interglacial as they depend on disturbed, transitional habitats for regeneration (Bobiec et al., 2018; Coppins & Coppins, 2003; Svenning & Magård, 1999), although human-enhanced fire regimes (Milner et al., 2016) may have compensated for the megafauna decline in the case of *Quercus* and *Corylus*. Furthermore, if changes in the abundance of these taxa were caused by human-linked changes in megafauna herbivory (Svenning et al., 2024), we expect that climate variables will have poor predictive ability. Together, our research questions will shed light on the pre-degradation abundances of important European tree species and their potential drivers, with implications for restoration initiatives and forest management (Lindbladh et al., 2007; Palli et al., 2023).

2 | METHODS

We used the latest pollen-based reconstructions of deciduous *Quercus*, *Corylus* and *Taxus* from the Last Interglacial (Pearce et al., 2023) and the early-mid-Holocene (Serge et al., 2023). The early-mid-Holocene dataset was missing *Taxus* reconstructions, which we produced using the same protocols as Serge et al. (2023). The two datasets and our early-mid-Holocene *Taxus* reconstructions were produced using the REVEALS model (Sugita, 2007) and following the same protocols developed by the LandClim and PAGES Land-cover6K projects (Gaillard et al., 2010; Githumbi et al., 2022). The REVEALS model transforms pollen count data to produce quantitative reconstructions of regional vegetation (Sugita, 2007). It overcomes the nonlinearity of the pollen-vegetation relationship by accounting for relative pollen productivity (RPP), dispersal and deposition differences between taxa. It is important to note that accurate RPP estimates for *Taxus* do not yet exist. In the absence of this, the estimate for *Juniperus* is used as a comparable substitution (Kuneš et al., 2011; Pearce et al., 2023). Whilst there therefore may be more uncertainty regarding the absolute percentages of *Taxus*,

this study is concerned with the relative abundance of *Taxus* in the Holocene and the Last Interglacial. For this comparison, it is most important that our method of measuring *Taxus* cover is consistent across time periods.

The REVEALS model was developed to transform pollen data from large lakes but can also produce regional vegetation cover estimates from multiple small-sized lakes and bogs (Trondman et al., 2016) and floodplains (Hoevers et al., 2022). The REVEALS model has been extensively tested and validated in Europe (Abraham et al., 2014; Hellman et al., 2008; Hjelle et al., 2018) and is robust to variations in site selection, sampling design and parameter values (Mazier, 2012). Pollen selection, harmonisation and REVEALS model parameterisation are comparable and available from Pearce et al. (2023) and Serge et al. (2023).

We focused on the distributions of *Quercus*, *Corylus* and *Taxus* in the temperate forest biome (Sensu Pearce et al., 2023). We also included pollen-based reconstructions from the adjoining sub-Arctic and Mediterranean biomes, to test for a Europe-wide effect, but only report these in the Supporting Information. *Quercus*, *Corylus* and *Taxus* are dependent on disturbance regimes that open the forest canopy for regeneration, often declining in dense, closed forests (Bobiec et al., 2018; Coppins & Coppins, 2003; Linares, 2013). Furthermore, the three taxa spread early at the beginning of interglacial periods and so are not strongly dispersal limited (Svenning et al., 2008). We focused on the mesocratic phases of vegetation development in both the Last Interglacial (Lang, 1994) and the Holocene (Birks, 1986). The mesocratic is the temperate, middle phase of an interglacial and thus is least affected by preceding and proceeding glacial conditions. In a typical interglacial cycle (i.e. one absent of pressure from modern humans), the mesocratic reflects the climatic optimum (Caspers et al., 2002) and maximum vegetation biomass development (Birks & Birks, 2004).

For the Last Interglacial, the mesocratic was identified by increases in *Quercus*, *Corylus*, *Taxus* and *Tilia* (first half of the mesocratic: Mesocratic I), followed by *Carpinus* and *Picea* (second half of the mesocratic: Mesocratic II) (Lang, 1994). Visual interpretation found that the vegetation community composition of Last Interglacial Mesocratic I was most comparable to the Holocene mesocratic, as there is no strong expansion of *Picea* and *Carpinus*. Furthermore, the Last Interglacial was a shorter interglacial period than the Holocene, with higher vegetation turnover (Felde et al., 2020). We therefore focus our analyses on Mesocratic I in the Last Interglacial. To test whether our choice to split the mesocratic affected our results, we analyse the full mesocratic period (including Mesocratic II: *Carpinus/Picea* phase) in the Supporting Information.

To assess the impact of megafauna extinction and other early, pre-agricultural human impacts on each taxon, we looked at *Quercus*, *Corylus* and *Taxus* abundance before the onset of substantial agriculture in the Holocene. We used REVEALS reconstructions of percentage *Cerealia* (members of the Poaceae family that are well-known indicators of agricultural activities) cover (Serge et al., 2023) as a proxy for the anthropogenic conversion of past landscapes to cropland. *Cerealia* cover began increasing exponentially at 5700

BP. As a result, we defined the mesocratic in the Holocene as encompassing 8700–5700BP, before the rise in *Cerealia* but after the protocratic phase and the associated disturbances of the preceding glacial period (Birks, 1986). For the main analysis, we used the reconstructions of our three focal taxa for the selected mesocratic phases in each time period (Last Interglacial: Mesocratic I; Holocene: 8700–5700BP; Table S1). We also included reconstructions that divided the mesocratic phases into individual time windows (Holocene: $n=6$; Last Interglacial, $n=2$) to provide more detail (Table S1).

We ensured our definition was restricted to periods where our three focal taxa were widespread in the temperate forest biome so that abundance variation would represent local ecological factors rather than large-scale biogeography. We captured early increases in *Quercus* and *Taxus*; however, we note that the early Holocene rise in *Corylus* precedes the mesocratic phase. *Corylus* expanded rapidly in the Holocene protocratic, particularly in Northwest Europe (Birks, 1986; Birks et al., 2016). As a result, our definition of the mesocratic in the Holocene misses the initial rise in *Corylus* cover. To test whether this exclusion impacted our results, we include a separate comparison of *Corylus* cover between the Holocene protocratic and Last Interglacial mesocratic in the Supporting Information.

Pollen-based reconstruction using REVEALS provides regional vegetation cover for 1-degree by 1-degree grid cells. A mean percentage cover estimate and mean standard error (SE) are given for each taxon per grid cell (Pearce et al., 2023; Serge et al., 2023). We averaged these values across grid cells to provide mean cover estimates for the temperate forest biome (Holocene: $n=342$; Last Interglacial: $n=49$) and Europe (Holocene: $n=539$; Last Interglacial: $n=66$; Supporting Information). The REVEALS model calculates uncertainty as the SEs derived from the sum of the within- and between-site variation in the REVEALS results in the grid cell (Githumbi et al., 2022). We used a coefficient of variation (SE/REVEALS estimate) to report SEs. When SE was greater than the REVEALS estimate, we considered the estimate unreliable.

2.1 | Climate data and zones

We used results from equilibrium simulations of climate, at 127ka (Last Interglacial) and 6ka (Holocene), executed according to the PMIP4 protocol (Otto-Bliesner et al., 2017). We used the results of the GISS-E2-1-G (Kelley et al., 2020) earth system model, as in Pearce et al. (2023). Temperature and precipitation fields were downscaled to a resolution of 5km and bias-corrected, as in Pearce et al. (2023). We derived bioclimatic variables, as in WorldClim (Fick & Hijmans, 2017), for each time period using the 'dismo' package in R (Hijmans et al., 2022). Using this data, the temperate forest biome was defined using Köppen–Geiger climate classifications, produced as in Beck et al. (2018) (Figure S1). Köppen–Geiger climate classifications were grouped into oceanic, continental, sub-Arctic, arid and Mediterranean categories following Pearce et al. (2023, Table S2). We defined the temperate forest biome as lowland grid cells that had an oceanic or continental climate in the Last Interglacial and/or

Holocene (Table S2). Lowland grid cells were defined as those where the standard deviation of elevation was <500 m.

2.2 | Statistical analyses

Where we had data for a given grid cell in both the Holocene mesocratic and the Last Interglacial mesocratic, we applied the paired Wilcoxon test for non-parametric data to compare percentage cover ($n=29$ paired grid cells). Using paired grid cells is more robust for measuring vegetation changes between the two time periods. However, the sample size is much smaller than using all grid cells and paired grid cells were mostly limited to Central Europe.

We used beta regression to assess the spatial, temporal and environmental drivers of each taxon across all grid cells in the temperate forest biome, as beta regression is appropriate for modelling continuous proportion data (Douma & Weedon, 2019). This is less robust than the paired analysis, as the sample sizes vary, but is more likely to capture Europe-wide effects. We selected explanatory variables based on known drivers of the distributions of each taxon from the literature (Berti & Svenning, 2020; Normand et al., 2011; Svenning, 2002). We included the above-mentioned bioclimatic variables and latitude (to assess disequilibrium dynamics following glaciation; Normand et al., 2011). We also included the standard deviation (SD) of elevation (Fischer et al., 2008) to assess the role of terrain roughness/herbivore accessibility on each taxon (c.f. Berti & Svenning, 2020). We tested for an effect of time period on each taxon by including time period (Last Interglacial; Holocene) as an explanatory factor. We used a pairwise comparison (z-test) of estimated marginal means from beta regression to test for a difference between the two time periods. We used backwards stepwise selection to exclude explanatory variables based on high variance inflation factors (>5) and correlation coefficients (>0.2).

In total, we ran four beta regression models. To look at the effect of time period whilst accounting for climate and other environmental factors, the first model (hereafter, 'the full model') contained six explanatory variables after variable selection: time period (Last Interglacial; Holocene), mean temperature of the wettest quarter ($^{\circ}\text{C}$), precipitation seasonality (coefficient of variation; $^{\circ}\text{C}$), precipitation of warmest quarter (mm), precipitation of coldest quarter (mm) and SD of elevation (m). To look at the relative importance of climate compared with time period, we removed time period from the full model to see if the model accuracy changed. A large drop in model accuracy after removing the time period variable from the full model would suggest that unmeasured, non-climatic variables account for a large part of explained variation (i.e., are important). For the third and fourth models, we ran the full model separately for the Last Interglacial and the Holocene to investigate any differences in climatic or environmental drivers between the two periods. We did not include interaction terms in any model, as we did not expect the physiological relationship between climate and vegetation to change between the Last Interglacial and the Holocene.

For the paired and beta regression analyses, we only used data from the temperate forest biome. Results for all grid cells (Europe-wide analyses) and for individual time windows (Table S3) are presented in the Supporting Information.

2.3 | Sensitivity analyses

To assess the robustness of our findings, we performed several sensitivity analyses. Due to *Abies* being susceptible to bias in vegetation cover estimates—especially in larger sedimentation basins—because of its high pollen fall speed and the limitations of GPM-based pollen productivity estimates (Abraham et al., 2014), we tested whether adjusting the RPP for *Abies* would influence our results for *Quercus*, *Corylus* and *Taxus*. Additionally, we tested the effects of excluding small sites and potentially local taxa (Cyperaceae) from the REVEALS reconstructions, as these factors may not align with the model's assumptions and could influence regional estimates. Lastly, we compared megafauna disturbance between the Last Interglacial and Holocene periods by comparing weighted mean grazing pressure values for herbaceous taxa in each period (Midolo et al., 2023). Detailed results for these analyses are presented in the Supporting Information.

3 | RESULTS

3.1 | *Corylus* (hazel)

We found considerably higher *Corylus* cover during the Last Interglacial mesocratic compared with the Holocene mesocratic in the European temperate forest biome. Paired analysis of grid cells using Wilcoxon testing showed strong evidence for a difference in *Corylus* cover between the two time periods ($p=0.002$; Figures 1 and 2; $n=29$ paired grid cells). In paired grid cells, cover was lower in the Holocene by an absolute mean of $14.0\% \pm 21.8$. This represents a relative decrease of 38.9% from the mean value of 36.0% in the Last Interglacial. *Corylus* cover was lower in all paired continental grid cells and 72% of paired oceanic grid cells (Figure S2).

Across all grid cells, the Last Interglacial mesocratic had $36.8\% \pm 17.6$ SD mean cover by *Corylus*, whereas the Holocene mesocratic had $16.9\% \pm 13.0$ mean cover (Figure 3). A pairwise comparison (z-test) of estimated marginal means from beta regression found very strong evidence for lower *Corylus* cover in the Holocene mesocratic than the Last Interglacial mesocratic (estimate = -0.19 , $z = -6.40$, $p < 0.0001$; all Holocene and Last Interglacial grid cells in the temperate forest biome). There was strong evidence that *Corylus* had lower cover in all time windows in the Holocene mesocratic than in Mesocratic I, but no evidence for a difference between the Holocene time windows and Mesocratic II (Table S3; Figure S3).

Corylus abundance was only partially explained by climatic variation between the two time periods. The full beta regression model




FIGURE 1 *Corylus*, *Quercus* and *Taxus* percentage cover in paired grid cells for the Last Interglacial and the early-mid-Holocene. Grid cells are depicted by points, which are joined by grey lines to show direction of change between the two time periods.




FIGURE 2 Change in mean *Corylus*, *Quercus* and *Taxus* cover in paired grid cells ($n=29$) in the Last Interglacial and Holocene (all time windows). Time windows in the Last Interglacial are given as phases relating to vegetation cover, whereas Holocene time windows have been dated (see Section 2). The darker grey box depicts the Last Glacial Period (LGP), for which we did not collect data. Lighter grey boxes depict the mesocratic phase of each time period, which we used for all analyses. Top panel: Mean and standard deviation (SD; shaded area) of *Corylus*, *Quercus* and *Taxus* percentage cover (%). Bottom panel: Cover of each taxon as a proportion of the maximum cover reached per taxon.

explained 22.5% of the variation in the data (pseudo R^2) and 13.1% when time period was removed (Table S4). The Last Interglacial model explained 12.7% of the variation in the data and the Holocene model explained 19.4% of the variation in the data (Table S4). In the Last Interglacial model, we found no evidence of any environmental

or climatic driver of *Corylus* cover (Figure S4). In the Holocene model, we found very strong evidence that *Corylus* cover was linked to precipitation seasonality (estimate = -0.054 , $p < 0.0001$) and moderate evidence that cover was linked to precipitation of the coldest quarter (estimate = 0.002 , $p = 0.020$; Figure S4).




FIGURE 3 REVEALS estimates of *Corylus* (top panel), *Quercus* (middle panel) and *Taxus* (bottom panel) percentage cover in the Last Interglacial mesocratic (left) and early-mid-Holocene mesocratic (right). Each square is a regional grid cell of $1^{\circ} \times 1^{\circ}$. Darker colours show greater cover by the given taxon. Paired grid cells are shown by a black outline. White/black circles inside each grid cell represent the coefficient of variation (standard error/REVEALS estimate). When $SE \geq$ REVEALS estimate, the circle fills the entire grid cell and the estimate is considered unreliable. Note the difference in scales of percentage cover between taxa.

3.2 | *Quercus* (oak)

Overall, there was no difference in *Quercus* abundance between the Last Interglacial and the Holocene. Paired analysis of grid cells showed no evidence of a difference in *Quercus* cover between the two time periods ($p=0.798$; Figures 1 and 2; $n=29$ paired grid cells). *Quercus* cover was lower in the Holocene in 80% of paired continental grid cells and 28% of paired oceanic grid cells (Figure S2).

Across all grid cells, the Last Interglacial mesocratic had $9.6\% \pm 6.4$ mean cover by *Quercus* ($n=47$), whereas the Holocene mesocratic had $5.6\% \pm 5.9$ mean cover ($n=292$; Figure 3). A pairwise comparison (z-test) of estimated marginal means from beta regression found no evidence for a difference in *Quercus* cover between the Holocene and Last Interglacial (Estimate=0.006, $z=-3.71$, $p=0.368$). When we compared the Holocene and Last Interglacial mesocratic time windows individually, we found evidence for higher percentage cover in the Holocene mesocratic time windows than in the Last Interglacial Mesocratic II, but no difference to Mesocratic I (Table S3; Figure S3).

The full beta regression model explained 24.3% of the variation in the data (pseudo R^2) and 24.1% when time period was removed, suggesting that climate was the most dominant driver that we measured (Table S4). The Last Interglacial model explained 27.1% of the variation in the data and the Holocene model explained 25.3% of the variation in the data (Table S4). In the Last Interglacial model, there was strong evidence that *Quercus* cover was linked to precipitation seasonality (coefficient of variation; estimate=0.038, $p=0.001$) and weak to moderate evidence that *Quercus* cover was linked to precipitation of the warmest quarter (estimate=-0.004, $p=0.053$; Figure S5). In the Holocene mesocratic model, we found very strong evidence that *Quercus* cover was linked to precipitation seasonality (estimate=-0.043, $p<0.0001$), mean temperature of the wettest quarter (estimate=0.077, $p<0.0001$), precipitation of the warmest quarter (estimate=-0.004, $p<0.0001$) and precipitation of the coldest quarter (estimate=0.003, $p<0.0001$; Figure S5).

3.3 | *Taxus* (yew)

We found much higher *Taxus* cover during the Last Interglacial mesocratic compared with the Holocene mesocratic. Paired analysis of grid cells using Wilcoxon testing showed strong evidence for a difference in *Taxus* cover between the two time periods ($p=0.002$; Figures 1 and 2; $n=29$ paired grid cells). In paired grid cells, cover was lower in the Holocene by an absolute mean of $1.1\% \pm 2.0$. This represents a relative decrease of 88.4% from the mean value of 1.3% in the Last Interglacial. *Taxus* cover was lower in all paired continental grid cells, as well as in 67% of paired oceanic grid cells (Figure S2).

Across all grid cells, the Last Interglacial mesocratic had $1.1\% \pm 1.8$ mean cover whereas the Holocene mesocratic had $0.02\% \pm 0.1$ mean cover (Figure 3). A pairwise comparison (z-test) of estimated marginal means from beta regression found very strong evidence for a lower *Taxus* cover in the Holocene mesocratic compared with the Last

Interglacial mesocratic (estimate=-0.004, $z=-7.22$, $p<0.0001$). At every time window in the Holocene mesocratic, there was very strong evidence that *Taxus* had lower cover than in both time windows in the Last Interglacial mesocratic (Table S3; Figure S3).

Taxus abundance was only partially explained by climatic variation between the two time periods. The full model explained 39.4% of the variation in the data (pseudo R^2) and 17.8% when time period was removed (Table S4). The Last Interglacial model explained 12.4% of the variation in the data and the Holocene model explained 8.6% of the variation in the data (Table S1). In the Last Interglacial model, beta regression found strong evidence that *Taxus* cover was linked to precipitation of the warmest quarter (estimate=-0.007, $p=0.001$) and precipitation of the coldest quarter (estimate=0.004, $p=0.004$) and weak to moderate evidence *Taxus* cover was linked to the mean temperature of the wettest quarter (estimate=0.044, $p=0.055$; Figure S6). In the Holocene model, we found strong evidence that *Taxus* cover was linked to precipitation seasonality (estimate=-0.008, $p=0.006$; Figure S6).


4 | DISCUSSION

Our analysis of the pre-degradation abundances of *Quercus*, *Corylus* and *Taxus* before and after the late-Quaternary megafauna extinctions revealed two principal findings. First, woodlands in the Last Interglacial of temperate Europe exhibited higher cover of *Corylus* and *Taxus* relative to the early-mid-Holocene, with the former reaching a very high percentage cover; meanwhile, *Quercus* had a consistent, moderately high percentage cover in both periods. Second, we found that the cover of *Corylus* and *Taxus* appeared to be influenced by unmeasured, non-climatic factors that differed between the two time periods, as the abundance of these two taxa was explained better by models that accounted for time period. Together, these findings suggest that, as a result of different drivers, forest structure differed between the two periods and from the classical, closed-canopy expectations (Figure 4).


Our findings are inconsistent with classic vegetation succession theory (Hilmers et al., 2018; Lienard et al., 2015). Within an interglacial period, vegetation succession is considered light-driven, as the dominant vegetation shifts from light-demanding to shade-tolerant taxa and forests form closed canopies with greater competition for light (Ellenberg, 1988). Therefore, *Quercus* and *Corylus*, as light-demanding taxa that rely on open and semi-open vegetation, should decline as they are out-competed by shade-tolerant species (Ellenberg, 1988). Indeed, earlier work concluded that declines in *Quercus* and *Corylus* pollen percentages indicate that much of the landscape was closed forest in the Last Interglacial (Svenning, 2002). Whilst phases of high *Quercus* and *Corylus* are a recognised feature of the early Last Interglacial (Suchora et al., 2022) and could be attributed to early vegetation succession, our REVEALS-based findings indicate that a high abundance of both taxa was a more persistent feature of Europe's temperate forest biome. We found that *Quercus* and particularly *Corylus* were abundant throughout

FIGURE 4 Examples of open and semi-open vegetation in Europe with regenerating (left) and established (right) *Corylus avellana* (top), *Quercus robur* (middle) and *Taxus baccata* (bottom). Photo credits: 1: Andrew Curtis, Wikimedia Commons, licensed under CC BY-SA 2.0; 2: Hazel Coppice at Torr Mhuilinn by M J Richardson, Wikimedia Commons, licensed under CC BY-SA 2.0; 3-5: Jens-Christian Svenning; Michael Geiser, iNaturalist (observation no. 115965559), licensed under CC BY-NC 4.0.

Regenerating



Established



both the Last Interglacial mesocratic phases as well as in the pre-agricultural Holocene. Whilst *Corylus* did decline in the second half of the Last Interglacial mesocratic, its cover remained much higher than would be considered possible under a closed canopy (Coppins & Coppins, 2003). Instead, it is possible that the *Corylus* persisted as the dominant taxa for thousands of years, forming a stable and long-term community (Coppins & Coppins, 2003).

Habitat disturbance resulting in open vegetation may enable *Corylus* to outcompete taller-growing, shade-producing trees (Hicks, 2022). *Corylus* pollen production is enhanced in more open settings (Bégeot, 1998) and female flowers are three times more productive in direct sunlight than in shady conditions (Azarenko et al., 1997). Furthermore, *Corylus* has a relatively deep rooting system and a strong capacity for regeneration, providing it with an advantage in post-fire vegetation recovery dynamics (Tallantire, 2002). Similarly, *Quercus* thrives in disturbed open and semi-open habitats, but not under closed canopies (Bobiec et al., 2018). Even with moderate shade, *Quercus* growth is slower and saplings are less likely to survive (Jones, 1959). In contrast, *Quercus* has a strong capacity for resprouting after damage from herbivory (Bobiec et al., 2018) or fire (Jones, 1959), particularly in open sites that enable higher rates of photosynthesis (Bobiec et al., 2018).

Whilst *Taxus* can tolerate some shade, it often declines in closed forest environments due to its poor competitive ability and the shading from taller- and faster-growing species, which

limits its growth and regeneration (Iszkuło et al., 2012; Perrin & Mitchell, 2013). Passive conservation approaches, such as passive rewilding or creating untouched forests are often not suitable for maintaining *Taxus* populations and active management strategies, including thinning competing species, are reported to be necessary to support its survival (Dobrowolska et al., 2017; Iszkuło et al., 2012). *Taxus* can withstand repeated pruning, continues growth under severe browsing pressure and is highly poisonous to many extant mammal species, even in small amounts (Cortinovis & Caloni, 2015). Deer are considered an exception and do browse *Taxus* in modern systems where their populations are not regulated by carnivores; at high population levels, deer can strongly depress *Taxus baccata* (Myserud & Østbye, 2004), as well as other *Taxus* species (Windels & Flaspohler, 2011). However, such mesoherbivores are subject to strong predation effects and are expected to have had reduced abundance, in contrast to bigger herbivores, in the complex pre-extinction megafauna assemblages (Svenning et al., 2024). Importantly, non-ruminants are particularly sensitive to *Taxus* toxicity and produce little browsing damage as a result (Thomas & Polwart, 2003). *Taxus* is therefore likely to have good defences against the diverse fauna of large and mega-herbivores that were widespread in Europe in the Last Interglacial (Davoli et al., 2023). This effect may be strengthened by the ability of *Taxus* seedlings to thrive in grazed systems via herbivore-mediated nurse shrubs (García & Ramón Obeso, 2003). Conversely, *Taxus* is sensitive to fire

due to its thin bark and difficulty regenerating in dry environments (Calvia et al., 2023; Mola et al., 2014; Thomas & Polwart, 2003), which is likely to have affected its abundance in the Holocene.

Corylus and *Taxus* were both less abundant in the Holocene mesocratic than in the Last Interglacial. Divergent natural disturbance regimes between the two periods may help to explain these findings. Whilst direct testing of the impact of megafauna or other non-climatic disturbance agents was not feasible and this study does not provide direct evidence of their role, we estimated residual variance by considering climatic and environmental effects. Even after accounting for these variables, the residual variation in the data suggests that unexplained factors played a substantial role in driving vegetation openness. Climatic differences between the Last Interglacial and the Holocene did not account for a substantial proportion of the variation in *Corylus* and *Taxus* abundance. The high importance of the 'time period' factor in our analysis indicates that other factors influencing abundance varied between these periods and were not captured by the explanatory variables. In contrast, *Quercus* abundance, which remained consistent across both periods, was mostly explained by climatic variables in our models. If climatic differences between the periods had strongly affected vegetation, we would expect *Quercus* cover to have changed more than *Corylus* and *Taxus* cover given that it was more explained by climate variables; however, *Quercus* cover changed the least. Overall, the inconsistent climate relations between taxa across different time periods suggest that other disturbance regimes likely played a role in driving open vegetation. We suggest a high level of herbivory by a diverse, abundant megafauna in the Last Interglacial and anthropogenic ecosystem management in the early-mid-Holocene are the most plausible candidates.

Megafauna abundance, species richness and functional diversity declined severely in the LGP, particularly amongst larger species (Davoli et al., 2023; Sandom, Ejrnæs, et al., 2014; Schowanek et al., 2021; Smith et al., 2018). Given the role of larger megafauna in creating and maintaining open and semi-open vegetation, defaunation is likely to have had cascading effects on forest community composition (Bakker et al., 2016). Our results for *Corylus* and *Taxus* are consistent with this hypothesis. Interestingly, however, *Quercus* is usually considered to benefit most from open, disturbed landscapes (Bobiec et al., 2018), yet, we found no evidence of a difference in *Quercus* cover; abundances were similar in the Last Interglacial and the Holocene mesocratic periods. This indicates that temperate forests in the Holocene mesocratic also had substantial openness (Serge et al., 2023), as in the Last Interglacial, but this was likely driven by factors other than megafauna due to their lower numbers in the Holocene (Bergman et al., 2023; Smith et al., 2018). We suggest that there may be an interplay between the effects of megafauna decline and additional factors that influence open and semi-open vegetation, which resulted in *Corylus* and *Taxus* being less abundant in the Holocene mesocratic.

Fire regimes were more frequent and intense in the Holocene than in the Last Interglacial (Lawson et al., 2010), which may have maintained open and semi-open vegetation. Increased fire

activity was likely due to direct fire management by *Homo sapiens* (Ellis et al., 2021), though it is also possible that the late-Quaternary megafauna extinctions contributed, as megafauna reduce flammable biomass (Karp et al., 2021). Increased fire activity, but less herbivory, in the Holocene compared with the Last Interglacial may have affected *Corylus* abundance by promoting more fire-tolerant taxa like *Pinus* (pine) and *Betula* (birch) (Molinari et al., 2020). *Pinus* and *Betula* may not have had a competitive advantage in the Last Interglacial as, though they are light-demanding, they are less tolerant of herbivory (Amsten et al., 2021).

In the early-mid-Holocene, high *Corylus* levels could have been promoted by *H. sapiens*, as hazelnuts were an important part of the Mesolithic diet (Bishop et al., 2023; Jacomet & Vandorpe, 2022; Jahns & Wolters, 2021). Deliberate coppicing of *Corylus* could promote flowering and therefore increase pollen production (Waller et al., 2012); however, the surviving browsing megafauna could have had the same effect (Vera, 2000). Moreover, whilst the human consumption of hazelnuts in the Mesolithic is well-documented, there is limited evidence of deliberate coppicing (Bishop et al., 2015). Finally, we found that *Corylus* presence was higher in the Last Interglacial than in the early Holocene. Although studies of archaeological sites have found charred hazelnut shells, indicating their consumption by *Homo neanderthalensis* (Neanderthals) (Salazar-García et al., 2013; Richter, 2016), the smaller population size of Neanderthals compared with later *H. sapiens* (Prüfer et al., 2014) is unlikely to have impacted *Corylus* abundance so substantially. This suggests that factors other than human activity played a role in promoting *Corylus* abundance, with megafauna likely having the strongest effects in the Last Interglacial and potentially also playing a role in the early-mid-Holocene (Vera, 2000).

Our findings show that disturbance-favoured *Corylus* and *Taxus* existed in greater abundance in the Last Interglacial, before the late-Quaternary megafauna extinctions. Interestingly, climate was not a primary driver of these abundance variations, emphasising the influence of unaccounted disturbance regimes in shaping past vegetation dynamics and opening an important avenue for future research. We conclude that the results presented here support active disturbance regimes in the Last Interglacial and are consistent with these being caused by megafauna herbivory rather than fire, given the higher abundance of *Taxus*. Megafauna existed in great abundance in Europe and world-wide for more than 30 million years (Smith et al., 2018). Our results indicate that the abundances of *Corylus* and *Taxus* in the early-mid-Holocene diverged from long-term baselines, which may have important implications for forest management and restoration initiatives (Bobiec et al., 2018; Mysterud & Østbye, 2004; Svenning & Magård, 1999). Modern management practices, characterised by either low densities of larger herbivores or high densities of meso-herbivores, do not reflect the greater diversity of herbivory modes and likely increased heterogeneity in herbivory that shaped past ecosystems (Atkinson et al., 2024). Consequently, trophic rewilding with a more diversified megafauna likely better aligns with long-term ecological conditions, promoting more resilient and functionally diverse woodland ecosystems.

AUTHOR CONTRIBUTIONS

Jens-Christian Svenning, Elena A. Pearce, Florence Mazier, Ralph Fyfe and Signe Normand designed the research, Florence Mazier, Maria-Antonia Serge and Paolo Scussolini contributed data; Elena A. Pearce and Charles W. Davison analysed the data; Elena A. Pearce wrote the paper, with input from all co-authors.

ACKNOWLEDGEMENTS

This work was supported by the project TERRANOVA, the European Landscape Learning Initiative, which received funding from the European Union's Horizon 2020 research and innovation program under Marie Skłodowska-Curie grant agreement no. 813904. The output reflects only the views of the authors and the European Union cannot be held responsible for the use which may be made of the information contained therein. We thank VILLUM FONDEN for economic support via J.-C.S' VILLUM Investigator project 'Biodiversity Dynamics in a Changing World' (grant 16549). We also consider this work a contribution to Centre for Ecological Dynamics in a Novel Biosphere (ECONOVO), funded by the Danish National Research Foundation (grant DNRF173 to J.-C.S.) and the MegaComplexity project, funded by Independent Research Fund Denmark | Natural Sciences (grant 0135-00225B to J.-C.S.).

CONFLICT OF INTEREST STATEMENT

The authors declare they have no conflicting interests.








PEER REVIEW

The peer review history for this article is available at <https://www.webofscience.com/api/gateway/wos/peer-review/10.1111/1365-2745.14422>.

DATA AVAILABILITY STATEMENT

The data used in this study are available from data.InDoRES: <https://data.indores.fr/dataset.xhtml?persistentId=doi:10.48579/PRO/J5GZUO> (Holocene; Serge, 2023) and OSF: <https://osf.io/nejbz/> (Last Interglacial; DOI: [10.17605/OSF.IO/NEJBZ](https://doi.org/10.17605/OSF.IO/NEJBZ); Pearce et al., 2024).

ORCID

Elena A. Pearce  <https://orcid.org/0000-0003-1635-0243>
 Florence Mazier  <https://orcid.org/0000-0003-2643-0925>
 Charles W. Davison  <https://orcid.org/0000-0001-9709-0951>
 Signe Normand  <https://orcid.org/0000-0002-8782-4154>
 Maria-Antonia Serge  <https://orcid.org/0000-0001-5506-9732>
 Paolo Scussolini  <https://orcid.org/0000-0001-6208-2169>
 Jens-Christian Svenning  <https://orcid.org/0000-0002-3415-0862>

REFERENCES

- Abraham, V., Oušková, V., & Kuneš, P. (2014). Present-day vegetation helps quantifying past land cover in selected regions of The Czech Republic. *PLoS One*, 9(6), e100117. <https://doi.org/10.1371/journal.pone.0100117>
- Amsten, K., Croomsigt, J. P. G. M., Kuijper, D. P. J., Loberg, J. M., Churski, M., & Niklasson, M. (2021). Fire- and herbivory-driven consumer control in a savanna-like temperate wood-pasture: An experimental approach. *Journal of Ecology*, 109(12), 4103–4114. <https://doi.org/10.1111/1365-2745.13783>
- Anderson, N. J., Bugmann, H., Dearing, J. A., & Gaillard, M.-J. (2006). Linking palaeoenvironmental data and models to understand the past and to predict the future. *Trends in Ecology & Evolution*, 21(12), 696–704. <https://doi.org/10.1016/j.tree.2006.09.005>
- Atkinson, J., Gallagher, R., Czyżewski, S., Kerr, M., Trepel, J., Buitenwerf, R., & Svenning, J.-C. (2024). Integrating functional traits into trophic rewilding science. *Journal of Ecology*, 112(5), 936–953. <https://doi.org/10.1111/1365-2745.14307>
- Azarenko, A. N., McCluskey, R. L., & Hampson, C. R. (1997). Proceedings of the 4th international symposium on hazelnut. *Acta Horticulturae*, 445, 179–183.
- Bakker, E. S., Gill, J. L., Johnson, C. N., Vera, F. W. M., Sandom, C. J., Asner, G. P., & Svenning, J.-C. (2016). Combining paleo-data and modern enclosure experiments to assess the impact of megafauna extinctions on woody vegetation. *Proceedings of the National Academy of Sciences of the United States of America*, 113(4), 847–855. <https://doi.org/10.1073/pnas.1502545112>
- Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., & Wood, E. F. (2018). Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Scientific Data*, 5(1), Article 1. <https://doi.org/10.1038/sdata.2018.214>
- Bégeot, C. (1998). Pollen behaviour of hazelnut (*Corylus avellana*), an indicator of human impacts? One example along a transect in Southern French Jura. *Acta Botanica Gallica*.
- Bergman, J., Pedersen, R. Ø., Lundgren, E. J., Lemoine, R. T., Monsarrat, S., Pearce, E. A., Schierup, M. H., & Svenning, J.-C. (2023). Worldwide late Pleistocene and early Holocene population declines in extant megafauna are associated with *Homo sapiens* expansion rather than climate change. *Nature Communications*, 14(1), Article 1. <https://doi.org/10.1038/s41467-023-43426-5>
- Berti, E., & Svenning, J.-C. (2020). Megafauna extinctions have reduced biotic connectivity worldwide. *Global Ecology and Biogeography*, 29(12), 2131–2142. <https://doi.org/10.1111/geb.13182>
- Birks, H., Felde, V. A., & Seddon, A. W. (2016). Biodiversity trends within the Holocene. *The Holocene*, 26(6), 994–1001. <https://doi.org/10.1177/0959683615622568>
- Birks, H. H., & Birks, H. J. B. (2004). The rise and fall of forests. *Science*, 305(5683), 484–485. <https://doi.org/10.1126/science.1101357>
- Birks, H. J. B. (1986). Late-quaternary biotic changes in terrestrial and lacustrine environments, with particular reference to north-west Europe. In B. E. Berglund (Ed.), *Handbook of Holocene palaeoecology and palaeohydrology* (pp. 3–52). John Wiley & Sons.
- Bishop, R. R., Church, M. J., & Rowley-Conwy, P. A. (2015). Firewood, food and human niche construction: The potential role of Mesolithic hunter-gatherers in actively structuring Scotland's woodlands. *Quaternary Science Reviews*, 108, 51–75. <https://doi.org/10.1016/j.quascirev.2014.11.004>
- Bishop, R. R., Kubiak-Martens, L., Warren, G. M., & Church, M. J. (2023). Getting to the root of the problem: New evidence for the use of plant root foods in Mesolithic hunter-gatherer subsistence in Europe. *Vegetation History and Archaeobotany*, 32(1), 65–83. <https://doi.org/10.1007/s00334-022-00882-1>
- Blanco, F., Calatayud, J., Martín-Perea, D. M., Domingo, M. S., Menéndez, I., Müller, J., Fernández, M. H., & Cantalapiedra, J. L. (2021). Punctuated ecological equilibrium in mammal communities over evolutionary time scales. *Science*, 372(6539), 300–303. <https://doi.org/10.1126/science.abd5110>
- Bobiec, A., Reif, A., & Öllerer, K. (2018). Seeing the oakscapes beyond the forest: A landscape approach to the oak regeneration in Europe. *Landscape Ecology*, 33(4), 513–528. <https://doi.org/10.1007/s10980-018-0619-y>
- Busing, R. T., Halpern, C. B., & Spies, T. A. (1995). Ecology of Pacific yew (*Taxus brevifolia*) in Western Oregon and Washington. *Conservation Biology*, 9(5), 1199–1207.

- Calvia, G., Casula, P., Farris, E., Fenu, G., Fantini, S., & Bacchetta, G. (2023). Shrub cover and soil moisture affect *Taxus baccata* L. regeneration at its southern range. *Plants*, 12(9), Article 9. <https://doi.org/10.3390/plants12091819>
- Caspers, G., Merkt, J., Müller, H., & Freund, H. (2002). The Eemian interglaciation in northwestern Germany. *Quaternary Research*, 58(1), 49–52. <https://doi.org/10.1006/qres.2002.2341>
- Coppins, A. M., & Coppins, B. J. (2003). Atlantic Hazelwoods—A neglected habitat? *Botanical Journal of Scotland*, 55(1), 149–160. <https://doi.org/10.1080/03746600308685056>
- Cortinovis, C., & Caloni, F. (2015). Alkaloid-containing plants poisonous to cattle and horses in Europe. *Toxins*, 7(12), Article 12. <https://doi.org/10.3390/toxins7124884>
- Croitor, R., & Brugal, J.-P. (2010). Ecological and evolutionary dynamics of the carnivore community in Europe during the last 3 million years. *Quaternary International*, 212(2), 98–108. <https://doi.org/10.1016/j.quaint.2009.06.001>
- Davoli, M., Monsarrat, S., Pedersen, R., Scussolini, P., Karger, D. N., Normand, S., & Svenning, J.-C. (2023). *Megafauna diversity and functional declines in Europe from the Last Interglacial (LIG) to the present* (p. 2023.06.12.544580). *bioRxiv* <https://doi.org/10.1101/2023.06.12.544580>
- Deforce, K., & Bastiaens, J. (2007). The Holocene history of *Taxus baccata* (yew) in Belgium and neighbouring regions. *Belgian Journal of Botany*, 140(2), 222–237.
- Dhar, A., Ruprecht, H., Klumpp, R., & Vacik, H. (2007). Comparison of ecological condition and conservation status of English yew population in two Austrian gene conservation forests. *Journal of Forestry Research*, 18(3), 181–186. <https://doi.org/10.1007/s11676-007-0037-5>
- Dobrowolska, D., Niemczyk, M., & Olszowska, G. (2017). The influence of stand structure on European yew *Taxus baccata* populations in its natural habitats in Central Poland. *Polish Journal of Ecology*, 65(3), 369–384. <https://doi.org/10.3161/15052249PJE2017.65.3.005>
- Douma, J. C., & Weedon, J. T. (2019). Analysing continuous proportions in ecology and evolution: A practical introduction to beta and Dirichlet regression. *Methods in Ecology and Evolution*, 10(9), 1412–1430. <https://doi.org/10.1111/2041-210X.13234>
- Ellenberg, H. (1988). *Vegetation ecology of Central Europe*. Cambridge University Press.
- Ellis, E. C., Gauthier, N., Klein Goldewijk, K., Bliege Bird, R., Boivin, N., Díaz, S., Fuller, D. Q., Gill, J. L., Kaplan, J. O., Kingston, N., Locke, H., McMichael, C. N. H., Ranco, D., Rick, T. C., Shaw, M. R., Stephens, L., Svenning, J.-C., & Watson, J. E. M. (2021). People have shaped most of terrestrial nature for at least 12,000 years. *Proceedings of the National Academy of Sciences of the United States of America*, 118(17), e2023483118. <https://doi.org/10.1073/pnas.2023483118>
- Felde, V. A., Flantua, S. G. A., Jenks, C. R., Benito, B. M., de Beaulieu, J.-L., Kuneš, P., Magri, D., Nalepka, D., Risebrobakken, B., ter Braak, C. J. F., Allen, J. R. M., Granaszewski, W., Helmens, K. F., Huntley, B., Kondratienė, O., Kalniņa, L., Kupryjanowicz, M., Malkiewicz, M., Milner, A. M., ... Birks, H. J. B. (2020). Compositional turnover and variation in Eemian pollen sequences in Europe. *Vegetation History and Archaeobotany*, 29(1), 101–109. <https://doi.org/10.1007/s00334-019-00726-5>
- Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, 37(12), 4302–4315. <https://doi.org/10.1002/joc.5086>
- Fischer, G., Nachtergaele, F., Prieler, S., van Velthuisen, H. T., Verelst, L., & Wiberg, D. (2008). *Global agro-ecological zones assessment for agriculture* (GAEZ 2008). IIASA, FAO.
- Gaillard, M.-J., Sugita, S., Mazier, F., Trondman, A.-K., Broström, A., Hickler, T., Kaplan, J. O., Kjellström, E., Kokfelt, U., Kuneš, P., Lemmen, C., Miller, P., Olofsson, J., Poska, A., Rundgren, M., Smith, B., Strandberg, G., Fyfe, R., Nielsen, A. B., ... Seppä, H. (2010). Holocene land-cover reconstructions for studies on land cover-climate feedbacks. *Climate of the Past*, 6(4), 483–499. <https://doi.org/10.5194/cp-6-483-2010>
- García, D., & Ramón Obeso, J. (2003). Facilitation by herbivore-mediated nurse plants in a threatened tree, *Taxus baccata*: Local effects and landscape level consistency. *Ecography*, 26(6), 739–750. <https://doi.org/10.1111/j.0906-7590.2003.03601.x>
- Giesecke, T., Bjune, A. E., Chiverrell, R. C., Seppä, H., Ojala, A. E. K., & Birks, H. J. B. (2008). Exploring Holocene continentality changes in Fennoscandia using present and past tree distributions. *Quaternary Science Reviews*, 27(13), 1296–1308. <https://doi.org/10.1016/j.quascirev.2008.03.008>
- Githumbi, E., Fyfe, R., Gaillard, M.-J., Trondman, A.-K., Mazier, F., Nielsen, A.-B., Poska, A., Sugita, S., Woodbridge, J., Azuara, J., Feurdean, A., Grindean, R., Lebreton, V., Marquer, L., Nebout-Combourieu, N., Stančikaitė, M., Tañău, I., Tonkov, S., Shumilovskikh, L., & LandClimII data contributors. (2022). European pollen-based REVEALS land-cover reconstructions for the Holocene: Methodology, mapping and potentials. *Earth System Science Data*, 14(4), 1581–1619. <https://doi.org/10.5194/essd-14-1581-2022>
- Hellman, S., Gaillard, M.-J., Broström, A., & Sugita, S. (2008). The REVEALS model, a new tool to estimate past regional plant abundance from pollen data in large lakes: Validation in southern Sweden. *Journal of Quaternary Science*, 23(1), 21–42. <https://doi.org/10.1002/jqs.1126>
- Hicks, D. (2022). Biological Flora of Britain and Ireland: *Corylus avellana*. *Journal of Ecology*, 110(12), 3053–3089. <https://doi.org/10.1111/1365-2745.14008>
- Hijmans, R. J., Phillips, S., & Elith, J. L. (2022). *dismo: Species distribution modeling* (1.3–9) [Computer software]. <https://CRAN.R-project.org/package=dismo>
- Hilmers, T., Friess, N., Bäessler, C., Heurich, M., Brandl, R., Pretzsch, H., Seidl, R., & Müller, J. (2018). Biodiversity along temperate forest succession. *Journal of Applied Ecology*, 55(6), 2756–2766. <https://doi.org/10.1111/1365-2664.13238>
- Hjelle, K. L., Halvorsen, L. S., Prøsch-Danielsen, L., Sugita, S., Paus, A., Kaland, P. E., Mehl, I. K., Overland, A., Danielsen, R., Høeg, H. I., & Midtbø, I. (2018). Long-term changes in regional vegetation cover along the west coast of southern Norway: The importance of human impact. *Journal of Vegetation Science*, 29(3), 404–415. <https://doi.org/10.1111/jvs.12626>
- Hoevers, R., Broothaerts, N., & Verstraeten, G. (2022). The potential of REVEALS-based vegetation reconstructions using pollen records from alluvial floodplains. *Vegetation History and Archaeobotany*, 31(5), 525–540. <https://doi.org/10.1007/s00334-022-00866-1>
- Iszkuło, G., Didukh, Y., Giertych, M. J., Jasińska, A. K., Sobierajska, K., & Szmyt, J. (2012). Weak competitive ability may explain decline of *Taxus baccata*. *Annals of Forest Science*, 69(6), 705–712. <https://doi.org/10.1007/s13595-012-0193-4>
- Jacomet, S., & Vandorpe, P. (2022). The search for a needle in a haystack—New studies on plant use during the Mesolithic in south-west Central Europe. *Journal of Archaeological Science: Reports*, 41, 103308. <https://doi.org/10.1016/j.jasrep.2021.103308>
- Jahns, S., & Wolters, S. (2021). *Mesolithic and early neolithic in Brandenburg from the botanical point of view* (pp. 205–230). Mesolithikum oder Neolithikum.
- Jones, E. W. (1959). *Quercus* L. *Journal of Ecology*, 47(1), 169–222. <https://doi.org/10.2307/2257253>
- Kalis, A. J., Merkt, J., & Wunderlich, J. (2003). Environmental changes during the Holocene climatic optimum in central Europe—Human impact and natural causes. *Quaternary Science Reviews*, 22(1), 33–79. [https://doi.org/10.1016/S0277-3791\(02\)00181-6](https://doi.org/10.1016/S0277-3791(02)00181-6)
- Karp, A. T., Faith, J. T., Marlon, J. R., & Staver, A. C. (2021). Global response of fire activity to late quaternary grazer extinctions. *Science*, 374(6571), 1145–1148. <https://doi.org/10.1126/science.abj1580>
- Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russqell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R.,

- Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., ... Yao, M. (2020). GISS-E2.1: Configurations and climatology. *Journal of Advances in Modeling Earth Systems*, 12(8), e2019MS002025. <https://doi.org/10.1029/2019MS002025>
- Kollmann, J., & Schill, H.-P. (1996). Spatial patterns of dispersal, seed predation and germination during colonization of abandoned grassland by *Quercus petraea* and *Corylus avellana*. *Vegetatio*, 125(2), 193–205. <https://doi.org/10.1007/BF00044651>
- Kühl, N., Litt, T., Schölzel, C., & Hense, A. (2007). Eemian and early Weichselian temperature and precipitation variability in northern Germany. *Quaternary Science Reviews*, 26(25), 3311–3317. <https://doi.org/10.1016/j.quascirev.2007.10.004>
- Kuneš, P., Odgaard, B. V., & Gaillard, M.-J. (2011). Soil phosphorus as a control of productivity and openness in temperate interglacial forest ecosystems. *Journal of Biogeography*, 38, 2150–2164.
- Lang, G. (1994). *Quartäre vegetationsgeschichte Europas: Methoden und ergebnisse*. Gustav Fischer Verlag Jena.
- Lawson, I., Venevsky, S., Sitch, S., Tzedakis, P., Roucoux, K., & Frogley, M. (2010). What is a natural wildfire regime in the Mediterranean? A comparison of Holocene and Eemian fire history at Ioannina. *Greece*, 12, 15350.
- le Roux, P. C., Aalto, J., & Luoto, M. (2013). Soil moisture's underestimated role in climate change impact modelling in low-energy systems. *Global Change Biology*, 19(10), 2965–2975. <https://doi.org/10.1111/gcb.12286>
- Lienard, J., Florescu, I., & Strigul, N. (2015). An appraisal of the classic forest succession paradigm with the shade tolerance index. *PLoS One*, 10(2), e0117138. <https://doi.org/10.1371/journal.pone.0117138>
- Linares, J. C. (2013). Shifting limiting factors for population dynamics and conservation status of the endangered English yew (*Taxus baccata* L., Taxaceae). *Forest Ecology and Management*, 291, 119–127. <https://doi.org/10.1016/j.foreco.2012.11.009>
- Lindbladh, M., Brunet, J., Hannon, G., Niklasson, M., Eliasson, P., Eriksson, G., & Ekstrand, A. (2007). Forest history as a basis for ecosystem restoration—A multidisciplinary case study in a south Swedish temperate landscape. *Restoration Ecology*, 15(2), 284–295. <https://doi.org/10.1111/j.1526-100X.2007.00211.x>
- Malkiewicz, M. (2018). Pollen-based vegetation and climate reconstruction of the Eemian sequence from Buntowo, N Poland. *Quaternary International*, 467, 54–61. <https://doi.org/10.1016/j.quaint.2016.07.006>
- Mazier, F. (2012). Testing the effect of site selection and parameter setting on REVEALS-model estimates of plant abundance using the Czech quaternary palynological database. *Review of Palaeobotany and Palynology*, 187, 38–49. <https://doi.org/10.1016/j.revpalbo.2012.07.017>
- Midolo, G., Herben, T., Axmanová, I., Marcenò, C., Pätzsch, R., Bruelheide, H., Karger, D. N., Ačić, S., Bergamini, A., Bergmeier, E., Biurrun, I., Bonari, G., Čarni, A., Chiarucci, A., De Sanctis, M., Demina, O., Dengler, J., Dziuba, T., Fanelli, G., ... Chytrý, M. (2023). Disturbance indicator values for European plants. *Global Ecology and Biogeography*, 32(1), 24–34. <https://doi.org/10.1111/geb.13603>
- Milner, A. M., Roucoux, K. H., Collier, R. E. L., Müller, U. C., Pross, J., & Tzedakis, P. C. (2016). Vegetation responses to abrupt climatic changes during the last interglacial complex (marine isotope stage 5) at Tenaghi Philippon, NE Greece. *Quaternary Science Reviews*, 154, 169–181. <https://doi.org/10.1016/j.quascirev.2016.10.016>
- Mola, J. M., Varner, J. M., Jules, E. S., & Spector, T. (2014). Altered community flammability in Florida's apalachicola ravines and implications for the persistence of the endangered conifer *Torreya taxifolia*. *PLoS One*, 9(8), e103933. <https://doi.org/10.1371/journal.pone.0103933>
- Molinari, C., Carcaillet, C., Bradshaw, R. H. W., Hannon, G. E., & Lehsten, V. (2020). Fire-vegetation interactions during the last 11,000 years in boreal and cold temperate forests of Fennoscandia. *Quaternary Science Reviews*, 241, 106408. <https://doi.org/10.1016/j.quascirev.2020.106408>
- Mysterud, A., & Østbye, E. (2004). Roe deer (*Capreolus capreolus*) browsing pressure affects yew (*Taxus baccata*) recruitment within nature reserves in Norway. *Biological Conservation*, 120(4), 545–548. <https://doi.org/10.1016/j.biocon.2004.03.027>
- Normand, S., Ricklefs, R. E., Skov, F., Bladt, J., Tackenberg, O., & Svenning, J.-C. (2011). Postglacial migration supplements climate in determining plant species ranges in Europe. *Proceedings of the Royal Society B: Biological Sciences*, 278(1725), 3644–3653. <https://doi.org/10.1098/rspb.2010.2769>
- Omarova, P. K., & Asadulaev, Z. M. (2016). Post-fire dynamics of vegetation in beech-yew woodland in piedmonts of Dagestan. *Lesovedenie*, 3, 209–215.
- Otto-Bliesner, B. L., Braconnot, P., Harrison, S. P., Lunt, D. J., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Capron, E., Carlson, A. E., Dutton, A., Fischer, H., Goelzer, H., Govin, A., Haywood, A., Joos, F., LeGrande, A. N., Lipscomb, W. H., Lohmann, G., Mahowald, N., ... Zhang, Q. (2017). The PMIP4 contribution to CMIP6—Part 2: Two interglacials, scientific objective and experimental design for Holocene and Last Interglacial simulations. *Geoscientific Model Development*, 10(11), 3979–4003. <https://doi.org/10.5194/gmd-10-3979-2017>
- Palli, J., Mensing, S. A., Schoolman, E. M., Solano, F., & Piovesan, G. (2023). Historical ecology identifies long-term rewilding strategy for conserving Mediterranean mountain forests in south Italy. *Ecological Applications*, 33(2), e2758. <https://doi.org/10.1002/eap.2758>
- Pearce, E. A., Mazier, F. F., Normand, S., Fyfe, R., Andrieu, V., Bakels, C., Balwierz, Z., Bińka, K., Boreham, S., Borisova, O., Brostrom, A., & Svenning, J.-C. (2023). Highly heterogeneous vegetation characterized the temperate forest biome before Homo sapiens. *Science Advances*, 9, eadi9135. <https://doi.org/10.1126/sciadv.adi9135>
- Pearce, E. A., Mazier, F. F., Normand, S., Fyfe, R., Andrieu, V., Bakels, C., Balwierz, Z., Bińka, K., Boreham, S., Borisova, O. K., Brostrom, A., de Beaulieu, J.-L., Gao, C., González-Sampérez, P., Granoszewski, W., Hrynowiecka, A., Kołaczek, P., Kuneš, P., Magri, D., ... Svenning, J.-C. (2024). Dataset: Substantial light woodland and open vegetation characterized the temperate forest biome before Homo sapiens. <https://doi.org/10.17605/OSF.IO/NEJBZ>
- Pérez Díaz, S., López Sáez, J. A., Ruiz Alonso, M., Zapata, L., & Abel Schaad, D. (2013). *Holocene history of Taxus baccata in the Basque Mountains (Northern Iberian Peninsula)*. Universidad Complutense de Madrid (UCM). https://doi.org/10.5209/rev_LAZA.2013.v34.n1.41933
- Perino, A., Pereira, H. M., Navarro, L. M., Fernández, N., Bullock, J. M., Ceaușu, S., Cortés-Avizanda, A., Van Klink, R., Kuemmerle, T., Lomba, A., Pe'er, G., Plieninger, T., Benayas, J. M. R., Sandom, C. J., Svenning, J.-C., & Wheeler, H. C. (2019). Rewilding complex ecosystems. *Science*, 364(6438), eaav5570. <https://doi.org/10.1126/science.aav5570>
- Perrin, P. M., & Mitchell, F. J. G. (2013). Effects of shade on growth, biomass allocation and leaf morphology in European yew (*Taxus baccata* L.). *European Journal of Forest Research*, 132(2), 211–218. <https://doi.org/10.1007/s10342-012-0668-8>
- Pringle, R. M., Abraham, J. O., Anderson, T. M., Coverdale, T. C., Davies, A. B., Dutton, C. L., Gaylard, A., Goheen, J. R., Holdo, R. M., Hutchinson, M. C., Kimuyu, D. M., Long, R. A., Subalusky, A. L., & Veldhuis, M. P. (2023). Impacts of large herbivores on terrestrial ecosystems. *Current Biology*, 33(11), R584–R610. <https://doi.org/10.1016/j.cub.2023.04.024>
- Prüfer, K., Racimo, F., Patterson, N., Jay, F., Sankararaman, S., Sawyer, S., Heinze, A., Renaud, G., Sudmant, P. H., de Filippo, C., Li, H., Mallick, S., Dannemann, M., Fu, Q., Kircher, M., Kuhlwilm, M., Lachmann, M., Meyer, M., Ongyerth, M., ... Pääbo, S. (2014). The complete genome sequence of a Neanderthal from the Altai Mountains. *Nature*, 505(7481), 43–49. <https://doi.org/10.1038/nature12886>
- Richter, J. (2016). Leave at the height of the party: A critical review of the middle Paleolithic in Western Central Europe from its beginnings to its rapid decline. *Quaternary International*, 411, 107–128. <https://doi.org/10.1016/j.quaint.2016.01.018>

- Rychel, J., Karasiewicz, M. T., Krześlak, I., Marks, L., Noryśkiewicz, B., & Woronko, B. (2014). Paleogeography of the environment in north-eastern Poland recorded in an Eemian sedimentary basin, based on the example of the Jafówka site. *Quaternary International*, 328–329, 60–73. <https://doi.org/10.1016/j.quaint.2013.09.018>
- Salazar-García, D. C., Power, R. C., Sanchis Serra, A., Villaverde, V., Walker, M. J., & Henry, A. G. (2013). Neanderthal diets in central and southeastern Mediterranean Iberia. *Quaternary International*, 318, 3–18. <https://doi.org/10.1016/j.quaint.2013.06.007>
- Salonen, J. S., Helmens, K. F., Brendryen, J., Kuosmanen, N., Väiliranta, M., Goring, S., Korpela, M., Kylander, M., Philip, A., Plikk, A., Renssen, H., & Luoto, M. (2018). Abrupt high-latitude climate events and decoupled seasonal trends during the Eemian. *Nature Communications*, 9, 2851. <https://doi.org/10.1038/s41467-018-05314-1>
- Sandom, C., Faurby, S., Sandel, B., & Svenning, J.-C. (2014). Global late quaternary megafauna extinctions linked to humans, not climate change. *Proceedings of the Royal Society B: Biological Sciences*, 281(1787), 20133254. <https://doi.org/10.1098/rspb.2013.3254>
- Sandom, C. J., Ejrnæs, R., Hansen, M. D. D., & Svenning, J.-C. (2014). High herbivore density associated with vegetation diversity in interglacial ecosystems. *Proceedings of the National Academy of Sciences of the United States of America*, 111(11), 4162–4167. <https://doi.org/10.1073/pnas.1311041111>
- Schläfli, P., Gobet, E., van Leeuwen, J. F. N., Vescovi, E., Schwenk, M. A., Bandou, D., Douillet, G. A., Schlunegger, F., & Tinner, W. (2021). Palynological investigations reveal Eemian interglacial vegetation dynamics at Spiezberg, Bernese Alps, Switzerland. *Quaternary Science Reviews*, 263, 106975. <https://doi.org/10.1016/j.quascirev.2021.106975>
- Schowaneck, S. D., Davis, M., Lundgren, E. J., Middleton, O., Rowan, J., Pedersen, R. Ø., Ramp, D., Sandom, C. J., & Svenning, J.-C. (2021). Reintroducing extirpated herbivores could partially reverse the late quaternary decline of large and grazing species. *Global Ecology and Biogeography*, 30(4), 896–908. <https://doi.org/10.1111/geb.13264>
- Serge, M. A., Mazier, F., Fyfe, R., Gaillard, M.-J., Klein, T., Lagnoux, A., Galop, D., Githumbi, E., Mindrescu, M., Nielsen, A. B., Trondman, A.-K., Poska, A., Sugita, S., & Woodbridge, J. (2023). Testing the effect of relative pollen productivity on the REVEALS model: A validated reconstruction of Europe-wide Holocene vegetation. *Land*, 12(5), Article 5. <https://doi.org/10.3390/land12050986>
- Serge, M. A. (2023). *Spatially extensive and temporally continuous three REVEALS pollen-based vegetation reconstructions in Europe over the Holocene*. (data.InDoRES, V2, UNF:6:6ucpBOTL4da0GLu2sG+C4Q== [fileUNF]) <https://doi.org/10.48579/PRO/J5GZUO>
- Smith, F. A., Smith, R. E. E., Lyons, S. K., & Payne, J. L. (2018). Body size downgrading of mammals over the late quaternary. *Science*, 360(6386), 310–313. <https://doi.org/10.1126/science.aao5987>
- Suchora, M., Kultys, K., Stachowicz-Rybka, R., Pidek, I. A., Hrynowiecka, A., Terpiłowski, S., Łabęcka, K., & Źarski, M. (2022). Palaeoecological record of long Eemian series from Kozłów (Central Poland) with reference to palaeoclimatic and palaeohydrological interpretation. *Quaternary International*, 632, 36–50. <https://doi.org/10.1016/j.quaint.2022.02.022>
- Sugita, S. (2007). Theory of quantitative reconstruction of vegetation I: Pollen from large sites REVEALS regional vegetation composition—Shinya Sugita, 2007. *The Holocene* <http://journals.sagepub.com/doi/10.1177/0959683607075837>, 17, 229–241.
- Svenning, J.-C. (2002). A review of natural vegetation openness in north-western Europe. *Biological Conservation*, 104(2), 133–148. [https://doi.org/10.1016/S0006-3207\(01\)00162-8](https://doi.org/10.1016/S0006-3207(01)00162-8)
- Svenning, J.-C., Lemoine, R. T., Bergman, J., Buitenwerf, R., Roux, E. L., Lundgren, E., Mungi, N., & Pedersen, R. Ø. (2024). The late-quaternary megafauna extinctions: Patterns, causes, ecological consequences and implications for ecosystem management in the Anthropocene. *Cambridge Prisms: Extinction*, 2, e5. <https://doi.org/10.1017/ext.2024.4>
- Svenning, J.-C., & Magård, E. (1999). Population ecology and conservation status of the last natural population of English yew *Taxus baccata* in Denmark. *Biological Conservation*, 88(2), 173–182. [https://doi.org/10.1016/S0006-3207\(98\)00106-2](https://doi.org/10.1016/S0006-3207(98)00106-2)
- Svenning, J.-C., Normand, S., & Kageyama, M. (2008). Glacial refugia of temperate trees in Europe: Insights from species distribution modelling. *Journal of Ecology*, 96(6), 1117–1127. <https://doi.org/10.1111/j.1365-2745.2008.01422.x>
- Tallantire, P. A. (2002). The early-Holocene spread of hazel (*Corylus avellana* L.) in Europe north and west of the Alps: An ecological hypothesis. *The Holocene*, 12(1), 81–96. <https://doi.org/10.1191/0959683602hl523rr>
- Thomas, P. A., & Polwart, A. (2003). *Taxus baccata* L. *Journal of Ecology*, 91(3), 489–524. <https://doi.org/10.1046/j.1365-2745.2003.00783.x>
- Trondman, A.-K., Gaillard, M.-J., Sugita, S., Björkman, L., Greisman, A., Hultberg, T., Lagerås, P., Lindbladh, M., & Mazier, F. (2016). Are pollen records from small sites appropriate for REVEALS model-based quantitative reconstructions of past regional vegetation? An empirical test in southern Sweden. *Vegetation History and Archaeobotany*, 25(2), 131–151. <https://doi.org/10.1007/s00334-015-0536-9>
- Van Kolfschoten, T. (2000). The Eemian mammal fauna of central Europe. *Netherlands Journal of Geosciences*, 79(2–3), 269–281. <https://doi.org/10.1017/S0016774600021752>
- Vera, F. W. M. (2000). *Grazing ecology and forest history*. Wallingford, UK: CABI Publishing. <https://doi.org/10.1079/9780851994420.0000>
- Waller, M., Grant, M. J., & Bunting, M. J. (2012). Modern pollen studies from coppiced woodlands and their implications for the detection of woodland management in Holocene pollen records. *Review of Palaeobotany and Palynology*, 187, 11–28. <https://doi.org/10.1016/j.revpalbo.2012.08.008>
- Willis, K. J., Bailey, R. M., Bhagwat, S. A., & Birks, H. J. B. (2010). Biodiversity baselines, thresholds and resilience: Testing predictions and assumptions using palaeoecological data. *Trends in Ecology & Evolution*, 25(10), 583–591. <https://doi.org/10.1016/j.tree.2010.07.006>
- Willis, K. J., Gillson, L., Brncic, T. M., & Figueroa-Rangel, B. L. (2005). Providing baselines for biodiversity measurement. *Trends in Ecology & Evolution*, 20(3), 107–108. <https://doi.org/10.1016/j.tree.2004.12.003>
- Willis, K. J., & van Andel, T. H. (2004). Trees or no trees? The environments of central and eastern Europe during the Last Glaciation. *Quaternary Science Reviews*, 23(23), 2369–2387. <https://doi.org/10.1016/j.quascirev.2004.06.002>
- Windels, S. K., & Flaspohler, D. J. (2011). The ecology of Canada yew (*Taxus canadensis* Marsh.): A review. *Botany*, 89(1), 1–17. <https://doi.org/10.1139/B10-084>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

Table S1. REVEALS time windows (TW) and years before present (BP; Holocene only) grouped to the interglacial phases used in this study (Last Interglacial mesocratic and Holocene mesocratic).

Table S2. Main climate zones covered by the Last Interglacial and Holocene pollen cores, grouped according to their description in the paper.

Table S3. Pairwise comparisons (z-test) of the estimated marginal means from beta regression modelling for each taxon across every time window.

Table S4. Model output (estimates and *p* values) for the four beta regression models per taxa.

Figure S1. Köppen-Geiger climate classification for the Last Interglacial (127k) and Holocene (6k). We reconstructed the classifications as in Beck et al. (2018), from monthly temperature and precipitation data from the GISS-E2-1-G earth system model (Kelley et al., 2020). For further methods and choice of model, see “climate data and zones” in main methods.

Figure S2. Maps showing the difference in percentage cover of *Quercus*, *Corylus* and *Taxus* for paired grid cells, from the Last Interglacial to the Holocene. More red colours show a decrease in cover, whereas more blue colours show an increase.

Figure S3. Boxplots of taxa percentage cover in the temperate forest biome (paired grid cells), at every time window, for the Last Interglacial (Protocratic to Telocratic) and the Holocene (11,700–(-65) BP).

Figure S4. Beta regression predictions of *Corylus* cover for variables where we found evidence of an effect in the Last Interglacial or Holocene: Mean Temperature of the Wettest Quarter (°C), Precipitation Seasonality (mm) and Precipitation of the Warmest Quarter (mm) and Precipitation of the Coldest Quarter (mm).

Figure S5. Beta regression predictions of *Quercus* cover for variables where we found evidence of an effect in the Last Interglacial or Holocene: Mean Temperature of the Wettest Quarter (°C), Precipitation Seasonality (mm) and Precipitation of the Warmest Quarter (mm) and Precipitation of the Coldest Quarter (mm).

Figure S6. Beta regression predictions of *Taxus* cover for variables where we found evidence of an effect in the Last Interglacial or Holocene: Mean Temperature of the Wettest Quarter (°C), Precipitation Seasonality (mm) and Precipitation of the Warmest Quarter (mm) and Precipitation of the Coldest Quarter (mm).

How to cite this article: Pearce, E. A., Mazier, F., Fyfe, R., Davison, C. W., Normand, S., Serge, M.-A., Scussolini, P., & Svenning, J.-C. (2024). Higher abundance of disturbance-favoured trees and shrubs in European temperate woodlands prior to the late-quaternary extinction of megafauna. *Journal of Ecology*, 00, 1–15. <https://doi.org/10.1111/1365-2745.14422>