



**HAL**  
open science

# CHIROPTICAL PROPERTIES OF ENANTIOPURE CRYPTOPHANES

T. Brotin

► **To cite this version:**

T. Brotin. CHIROPTICAL PROPERTIES OF ENANTIOPURE CRYPTOPHANES. 8th conference on vibrational SPECTROSCOPY, Aug 2024, BOCHUM, Germany. hal-04720285

**HAL Id: hal-04720285**

**<https://hal.science/hal-04720285v1>**

Submitted on 3 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# CHIROPTICAL PROPERTIES OF ENANTIOPURE CRYPTOPHANES

BROTIN Thierry

*Laboratoire de chimie de l'ENS LYON (CNRS, UMR 5182). École Normale Supérieure  
de Lyon, 46 Allée d'Italie, 69364 Lyon Cedex 07, FRANCE  
[tbrotin@ens-lyon.fr](mailto:tbrotin@ens-lyon.fr)*

*Cryptophane* is the common name for hollow macrocyclic compounds first studied by Collet and his collaborators.<sup>[1]</sup> Interestingly, these molecules can encapsulate a wide range of substrates (cations, anions, small molecules) in organic or aqueous solvents, depending on the substituents grafted onto the cryptophane backbone.<sup>[2]</sup> The vast majority of cryptophanes are also chiral compounds.<sup>[3]</sup> Thus, chiroptical techniques are valuable tools for characterizing these molecules.<sup>[4-5]</sup> In this presentation, we will give an overview of the chiroptical techniques (ECD, VCD, ROA, X-ray, Polarimetry) used to characterize cryptophane derivatives. Through several examples, we will show how chiroptical techniques can be used to determine the absolute configuration of cryptophanes, as well as providing valuable information on the conformation of the three linkers. Particular attention will be paid to Vibrational Circular Dichroism (VCD) and Raman Optical Activity (ROA) techniques.




Figure. Schematic representation of a cryptophane skeleton with the names of the various chiroptical techniques used to characterize these compounds.

## References

1. J. Gabard, A. Collet, *J. Chem. Soc. Chem. Comm.* (1981) 1137–1139
2. T. Brotin, J.-P. Dutasta, *Chem. Rev.* **109** (2009) 88–130
3. O. Baydoun, T. Buffeteau, T. Brotin, *Chirality* **33** (2021) 562–596
4. T. Brotin, D. Cavagnat, J.-P. Dutasta, T. Buffeteau, *J. Am. Chem. Soc.* **128** (2006) 5533–5540
5. T. Brotin, N. Daugey, J. Kapitan, N. Vanthuyne, M. Jean, E. Jeanneau, T. Buffeteau, *J. Org. Chem.* **88** (2023) 4829–4832