



HAL
open science

Shrinking body size may not provide meaningful thermoregulatory benefits in a warmer world

Andreas Nord, Elin Persson, Joshua Tabh, Elisa Thoral

► **To cite this version:**

Andreas Nord, Elin Persson, Joshua Tabh, Elisa Thoral. Shrinking body size may not provide meaningful thermoregulatory benefits in a warmer world. *Nature Ecology & Evolution*, 2024, 8 (3), pp.387-389. 10.1038/s41559-023-02307-2 . hal-04725251

HAL Id: hal-04725251

<https://hal.science/hal-04725251v1>

Submitted on 10 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Shrinking body size may not provide meaningful thermoregulatory benefits in a warmer world

Received: 17 July 2023

Andreas Nord  , Elin Persson , Joshua K. R. Tabh  & Elisa Thoräl 

Accepted: 13 December 2023

Published online: 15 January 2024

 Check for updatesARISING FROM C. Youngflesh et al. *Nature Ecology and Evolution* <https://doi.org/10.1038/s41559-022-01893-x> (2022)

In a recent and impressive analysis of avian morphological data (covering >250,000 male birds from 105 species), Youngflesh et al.¹ report that birds breeding in North America have become significantly lighter over the past three decades, coincident with rising breeding season temperatures. Because these observations recapitulate predictions under Bergmann's rule (that is, that the body size of congeners and conspecifics is usually larger near the cooler poles than near the warmer Equator), the authors argue a thermoregulatory benefit to their occurrence under the assumption that smaller-bodied animals have lower cooling costs than larger-bodied animals in a warming world. We agree that warmer environments during reproduction may well explain avian body size declines. However, we question whether (1) changes in heat balance attributed to these declines are sufficiently large to explain observed size reductions, and (2) increased thermoregulatory efficiency during the short windows where reproduction occurs is always relevant, particularly when tenancy in breeding-ground temperatures is short (that is, among migrants). Generalizability of these results may be limited further by sex-specific enquiry, which overlooks the possibility of divergent selection on body size in males and females under climate warming. In this commentary, we expand on these concerns hoping to instigate discussion on knowledge gaps that need closing if we wish to better understand the proximate and ultimate drivers of shapeshifting animals in a changing world.

Data presented by Youngflesh et al.¹ corroborate several recent reports detailing how animals are changing body size and shape in parallel with warming temperatures^{2–4}. These collective observations are notable, and we share the view of Youngflesh et al.¹ that such temporal trends must be described and understood from both evolutionary and applied points of view^{5,6}. We also agree that smaller animals are theoretically better equipped to withstand higher ambient temperatures in thermal environments characterized by net heat loss (that is, where the animal remains warmer than its surroundings), since their higher surface area to volume ratios increase the proportion of integument usable for heat transfer. However, the reported mean body size shifts in this and other studies are small (here, -0.56% ¹; Fig. 1a) and so, according to currently available allometric relationships, yield

near-negligible effects on mean body surface area (-0.38%) and surface area to volume ratios ($+0.19\%$) across species (Fig. 1b). Such minor changes in morphology hold little thermoregulatory bearing, with conservative heat balance models (that is, assuming both complete and static plumage cover) suggesting a mean increase in heat flux ($W\ cm^{-2}$) of around 0.14% (Fig. 1b). The estimated effects on evaporative water loss ($+0.10\%$), total heat loss rate (that is, thermal conductance; $+0.27\%$), basal heat production (that is, basal metabolic rate; -0.39%), upper critical temperature (-0.05%) and the metabolic response to warmth (-0.48%) are similarly weak (Fig. 1b,c). Even for higher estimates of body mass change (for example, -2.6% on average in ref. 4), thermoregulatory benefits are still likely to be limited (for example, $+0.60\%$ in heat flux, $+0.42\%$ in evaporative water loss, $+1.10\%$ in thermal conductance and -1.56% in basal metabolic rate). By comparison, a typical physiological thermoregulatory response may lead to a 5- to 15-fold increase in evaporative water loss upon heat exposure⁷. In all, it seems doubtful that the observed shifts in body mass or size reported by Youngflesh et al.¹ and elsewhere will hold any significant bearing over the degree to which species can respond to thermoregulatory pressures caused by warming temperatures. In fact, available data suggest that, interspecifically, maximum tolerable air temperature in passerines has positive allometry⁷, and reducing plumage depth by as little as 1% would over-double the heat dissipation effect ascribed to body shrinkage in ref. 1 (that is, 0.39% reduction in heat flux relative to 0.14%). We therefore caution against such adaptationist conclusions in the absence of empirical evidence of thermoregulatory benefits.

Second, the authors base inference on data obtained during Northern Hemisphere springs and summers (1 May to 31 July). While this in itself is perfectly reasonable, the overwhelming majority of the species studied were migrants (66%, or 69 of 105 species) or partial migrants (22%, 23 of 105)⁸, suggesting short tenancy in at least some study regions, or even measurement during migratory stop-over. In these contexts, regressing morphology against local ambient temperature appears problematic. Specifically, conceptualization and hypothesis testing become difficult when the precise temperatures to which the birds are responding to morphologically are unknown,




Fig. 1 | For most bird species, temporal changes in body masses are too small to contribute biologically meaningful shifts in heat production and dissipation, suggesting thermoregulatory benefit is an unlikely outcome of shapeshift under climate warming. **a**, Data on mean shifts in body masses from Youngflesh et al.¹. **b**, These data were used to predict changes in body proportions, heat flux and several thermo-physiological variables of relevance for how a bird exchanges heat with its environment. We predicted body surface area from body mass²⁰ and then assumed body densities of 1.1 g cm^{-3} (ref. 21) to estimate effects on surface area to volume ratios. Shifts in surface area were then used to estimate the change in sensible ('dry') heat loss using biophysical equations published previously²² and assuming still air conditions, no conductive heat loss (that is, no contact heat loss between the integument and the environment), static skin surface temperature of 41°C (that is,

a representative small-bird body temperature²³), a soft tissue thermal conductivity of $0.5 \text{ W m}^{-1}\text{C}^{-1}$ (ref. 22), a plumage conductivity of $0.0272 \text{ W m}^{-1}\text{C}^{-1}$ (ref. 22) and a plumage depth equalling $1/3 \times \log_{10}$ of body mass in cm. Latent ('wet') heat loss (that is, evaporative water loss) was estimated based on previously published data⁷, and total mass-specific heat dissipation rate (that is, thermal conductance) was predicted based on Aschoff's allometry for daytime conditions²⁴. **c**, Body size data were also used to estimate changes in basal heat production (that is, basal metabolic rate), upper critical temperature (that is, the temperature above which heat dissipation mechanisms must be engaged to prevent increasing body temperature) and the slope of the relationship between resting metabolic rate and air temperature above the upper critical temperature, using allometric relationships in refs. 7, 25. Solid plotting symbols show mean trait levels and bars denote the 89% quantile intervals.

even if migrants may track their thermal environment across the annual cycle⁹. The problem of spatial asynchrony arguably lessens if shrinking body sizes reflect plastic effects of developmental temperature on final morphometry (as suggested by Youngflesh et al. and others¹⁰), but recent theory still contends that broad generalizations of warming environments during development and decreasing size are likely naïve¹¹. In any case, spatial asynchrony between sites of measurement and sites of operation remains a concern if size declines reflect selection, for example, via temperature-mediated, size-dependent mortality¹. For this reason, we suggest life history parameters be carefully considered in future modelling efforts to shed light on the issues outlined above.

Last, inference in the study by Youngflesh et al.¹ is based only on male birds with the rationale that reproductive status (that is, gravidity) renders true female mass uncertain (for example, ref. 12). Removing

females from the dataset might well be appropriate statistically but raises concerns about the generality of conclusions if any selection for size is sex-specific. For example, larger females can lay larger eggs that produce larger chicks¹³, and so, may sire offspring with higher survival^{13,14}. It could therefore be expected that changes in body size in response to warming may be constrained in females relative to males. In line with this, a recent mammalian study found that while males and females responded qualitatively similarly to developmental temperature manipulations, the effect of body size was markedly larger when males were modelled alone compared to when measurements of both sexes were combined¹⁵. Other studies report that male and female birds show differential thermal sensitivity of reproductive investment¹⁶. While we understand that exclusion of females from the current data set may have been a necessity for modelling, we suggest that future studies acknowledge the possibility that natural selection for body size

differs between the sexes, instead of considering only the less variable sex or collapsing the data (for example, refs. 3,4).

By now, several ambitious studies have demonstrated that wild animals show concerted changes in body mass, size and shape in conjunction with warming temperatures. Evidence suggests that such shapeshifting probably stems from plasticity more than it reflects any adaptive evolutionary responses to changing thermal environments^{17–19}. Whether such plasticity is adaptive, non-adaptive or neutral remains unclear. However, predictions based on currently available allometries suggests that even if a reduction in body mass mediates changes to heat production and dissipation, the predicted shifts in mean trait levels are often too small to provide meaningful thermoregulatory value (Fig. 1), especially when compared against those accrued by plastic, physiological responses to cold and warmth. For this reason, the precise role of declining size toward thermal balance under climate warming in birds—or any endotherms—is speculative and should remain acknowledged as such until further evaluated. We suggest future research focus be placed on how size declinations occur in the warmth (for example, via selection, or plasticity) and why it matters by evaluating empirically the energetic and fitness benefits of reducing body size under climate warming by observed degrees.

References

- Youngflesh, C., Saracco, J. F., Siegel, R. B. & Tingley, M. W. Abiotic conditions shape spatial and temporal morphological variation in North American birds. *Nat. Ecol. Evol.* **6**, 1860–1870 (2022).
- Ryding, S., Klaassen, M., Tattersall, G. J., Gardner, J. L. & Symonds, M. R. E. Shape-shifting: changing animal morphologies as a response to climatic warming. *Trends Ecol. Evol.* **36**, 1036–1048 (2021).
- Zimova, M. et al. Body size predicts the rate of contemporary morphological change in birds. *Proc. Natl Acad. Sci. USA* **120**, e2206971120 (2023).
- Weeks, B. C. et al. Shared morphological consequences of global warming in North American migratory birds. *Ecol. Lett.* **23**, 316–325 (2020).
- Kronfeld-Schor, N. Conservation physiology. *Temperature* **1**, 94–95 (2014).
- Stillman, J. H. Heat waves, the new normal: summertime temperature extremes will impact animals, ecosystems, and human communities. *Physiology* **34**, 86–100 (2019).
- McKechnie, A. E., Gerson, A. R. & Wolf, B. O. Thermoregulation in desert birds: scaling and phylogenetic variation in heat tolerance and evaporative cooling. *J. Exp. Biol.* **224**, jeb229211 (2021).
- Billerman, S. M., Keeney, B. K., Rodewald, P. G. & Schulenberg, T. S. *Birds of the World* (Cornell Lab of Ornithology; 2022); <https://birdsoftheworld.org/bow/home>
- Somveille, M., Manica, A. & Rodrigues, A. S. L. Where the wild birds go: explaining the differences in migratory destinations across terrestrial bird species. *Ecography* **42**, 225–236 (2019).
- Weeks, B. C. et al. Temperature, size and developmental plasticity in birds. *Biol. Lett.* **18**, 20220357 (2022).
- Tabh, J. K. R. & Nord, A. Temperature-dependent developmental plasticity and its effects on Allen's and Bergmann's rules in endotherms. *Integr. Comp. Biol.* **63**, 758–771 (2023).
- Meijer, T., Möhring, F. J. & Trillmich, F. Annual and daily variation in body mass and fat of starlings *Sturnus vulgaris*. *J. Avian Biol.* **25**, 98–104 (1994).
- Ronget, V. et al. Causes and consequences of variation in offspring body mass: meta-analyses in birds and mammals. *Biol. Rev.* **93**, 1–27 (2018).
- Perrins, C. M. Eggs, egg formation and the timing of breeding. *Ibis* **138**, 2–15 (1996).
- Ballinger, M. A. & Nachman, M. W. The contribution of genetic and environmental effects to Bergmann's rule and Allen's rule in house mice. *Am. Nat.* **199**, 691–704 (2022).
- Schou, M. F. et al. Extreme temperatures compromise male and female fertility in a large desert bird. *Nat. Commun.* **12**, 666 (2021).
- Teplitsky, C., Mills, J. A., Alho, J. S., Yarrall, J. W. & Merilä, J. Bergmann's rule and climate change revisited: disentangling environmental and genetic responses in a wild bird population. *Proc. Natl Acad. Sci. USA* **105**, 13492–13496 (2008).
- Bosco, L. et al. Increasing winter temperatures explain body size decrease in wintering bird populations of Northern Europe – but response patterns vary along the spatioclimatic gradient. *Glob. Ecol. Biogeogr.* **32**, 2100–2110 (2023).
- Siepielski, A. M. et al. No evidence that warmer temperatures are associated with selection for smaller body sizes. *Proc. R. Soc. Lond. B* **286**, 20191332 (2019).
- Walsberg, G. E. The relationship of the external surface area of birds to skin surface area and body mass. *J. Exp. Biol.* **76**, 185–189 (1978).
- Heymsfield, S. B. et al. Chemical determination of human body density in vivo: relevance to hydrodensitometry. *Am. J. Clin. Nutr.* **50**, 1282–1289 (1989).
- Porter, W. P. & Kearney, M. Size, shape, and the thermal niche of endotherms. *Proc. Natl Acad. Sci. USA* **106**, 19666–19672 (2009).
- Prinzinger, R., Pressmar, A. & Schleucher, E. Body temperature in birds. *Comp. Biochem. Physiol. A* **99**, 499–506 (1991).
- Aschoff, J. Thermal conductance in mammals and birds: its dependence on body size and circadian phase. *Comp. Biochem. Physiol. A* **69**, 611–619 (1981).
- McKechnie, A. E. & Wolf, B. O. The allometry of avian basal metabolic rate: good predictions need good data. *Physiol. Biochem. Zool.* **77**, 502–521 (2004).

Acknowledgements

A.N. was supported by the Swedish Research Council (grant 2020-04686). E.P. was supported by the Crafoord foundation (grants 20211007 and 20221018) and a PhD studentship from Lund University. J.K.R.T. was supported by a postdoctoral grant from the Wenner-Gren Foundation (grant UPD2021-0038). E.T. was supported by a postdoctoral grant from the Carl Trygger Foundation (grant CTS21: 1173).

Author contributions

This commentary resulted from a lab meeting at the Department of Biology, Lund University, Sweden. A.N., E.P., J.K.R.T. and E.T. conceptualized the paper and drafted the outline. A.N. wrote the first complete draft, which was revised in equal measure by E.P., J.K.R.T., E.T. and A.N.; J.K.R.T. and A.N. constructed the heat balance model, J.K.R.T. performed the modelling; J.K.R.T. and A.N. developed the graphic material; A.N. procured funding. Author names are listed in alphabetical order to reflect equal contributions.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Andreas Nord.

Peer review information *Nature Ecology & Evolution* thanks the anonymous reviewers for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2024