



HAL
open science

Shaping non-reciprocal caustic spin-wave beams

V. Vlaminck, Loic Temdie, Vincent Castel, Benjamin Jungfleisch, Dinesh Wagle, Mojtaba Taghipour Kaffash, Daniel Stoeffler, Yves Henry, H Majjad, R Bernard, et al.

► **To cite this version:**

V. Vlaminck, Loic Temdie, Vincent Castel, Benjamin Jungfleisch, Dinesh Wagle, et al.. Shaping non-reciprocal caustic spin-wave beams. 2024 International Conference on Microwave Magnetics, Jun 2024, Oakland, MI, United States. hal-04726626

HAL Id: hal-04726626

<https://hal.science/hal-04726626v1>

Submitted on 8 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SHAPING NON-RECIPROCAL CAUSTIC SPIN-WAVE BEAMS

VINCENT VLAMINCK^{1,2*}, LOIC TEMDIE^{1,2}, VINCENT CASTEL^{1,2}, BENJAMIN JUNGFLAISCH³, DINESH WAGLE³, MOJTABA TAGHIPOUR KAFFASH³, DANIEL STOEFFLER⁴, YVES HENRY⁴, H. MAJJAD⁴, R. BERNARD⁴, MATTHIEU BAILLEUL⁴

IIMT Atlantique, Microwave Dpt., CS 83818, 29238 Brest, France

2Lab-STICC (UMR 6285), CNRS, Technopole Brest-Iroise, 29238 Brest Cedex 3, France

3University of Delaware, Dept. of Physics and Astronomy, Newark, Delaware, USA

4IPCMS – Institut de Physique et Chimie des matériaux de Strasbourg, France

Mastering spin waves interference at the sub-micron scale is central for the development of wave-based computing applications such as reservoir computing [1], holographic memory, or spectral analysis. The complexity of spin dynamics, inherently due to its dependence to numerous parameters, and the intricacy of magnon-magnon interactions, requires heavy computational methods, which can limit the scope of study. In this context, we developed an efficient tool to study the near-field diffraction (NFD) patterns of spin wave in homogeneous out-of-plane magnetized thin films for arbitrary distribution of excitation fields [2, 3]. In this communication, we present the adaptation of our NFD model to in-plane magnetized thin films, and reveal how caustic spin-wave beams can be directly emitted from a sharply constricted stripline. We show in particular the importance of not only meeting the suitable caustic conditions of field and frequency (see Fig. 1), but also having of a seemingly punctual source for shaping the caustic beam. Using spatially resolved micro-focused Brillouin light spectroscopy on various constriction sizes, we satisfyingly verify our predictions, and explore further the steerability as well as the caustic beams properties.

This work, which have important implications for the development magnonic interferometric devices, was supported by the French ANR project “MagFunc”, the U.S. Department of Energy under Award DE SC0020308, and the Transatlantic Research Partnership, a program of FACE Foundation and the French Embassy.




Figure 1. Slowness curve at 7.9 GHz for a 50 nm thick YIG film magnetized along y-axis under a bias field of 200mT.




Figure 2. Corresponding NFD simulations for both field orientations.

- 1 A. Papp *et al.* (2021) Characterization of nonlinear spin-wave interference by reservoir-computing metrics”, *Applied Physics Letters* **119**, 112403.
- 2 V. Vlaminc *et al.* (2023) Spin wave diffraction model for perpendicularly magnetized films. *Journal of Applied Physics* **133**, 053903.
- 3 L. Temdie *et al.* (2024) Probing Spin Wave Diffraction Patterns of Curved Antennas, *Phys. Rev. Applied* **21**, 014032.