



HAL
open science

A Lower Bound on the Critical Momentum of an Impurity in a Bose-Einstein Condensate

Benjamin Hinrichs, Jonas Lampart

► **To cite this version:**

Benjamin Hinrichs, Jonas Lampart. A Lower Bound on the Critical Momentum of an Impurity in a Bose-Einstein Condensate. 2024. hal-04727588

HAL Id: hal-04727588

<https://hal.science/hal-04727588v1>

Preprint submitted on 9 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A LOWER BOUND ON THE CRITICAL MOMENTUM OF AN IMPURITY IN A BOSE–EINSTEIN CONDENSATE

BENJAMIN HINRICHS AND JONAS LAMPART

ABSTRACT. In the Bogoliubov–Fröhlich model, we prove that an impurity immersed in a Bose–Einstein condensate forms a stable quasi-particle when the total momentum is less than its mass times the speed of sound. The system thus exhibits superfluid behavior, as this quasi-particle does not experience friction. We do not assume any infrared or ultraviolet regularization of the model, which contains massless excitations and point-like interactions.

1. INTRODUCTION

An impurity in a Bose–Einstein condensate will create excitations out of the ground state and may form a quasi-particle, called the Bose polaron, consisting of the particle and a surrounding cloud of excitations. The system is of interest in physics as the impurity can reveal properties of the condensate, such as superfluidity. Moreover, Bose–Einstein condensates are finely controllable experimental platforms from which one hopes to learn about polaron physics in solids by analogy.

The Bogoliubov–Fröhlich Hamiltonian is an effective model for such a system, in which the particle is linearly coupled to Bogoliubov’s excitation field. This model is relevant if the interaction between the particle and the bosons is sufficiently weak to not significantly impact the condensate [GD16], though there is some debate in the physics literature on what effects this model can or cannot capture [CLB15, GSSD17]. Recent mathematical results prove that it provides an accurate description of the system in certain mean-field [MS20, LP22] and dilute [LT] regimes.

In this letter we start from the translation-invariant Bogoliubov–Fröhlich Hamiltonian in \mathbb{R}^3 and prove that the Bose polaron is stable when the total momentum is less than the impurity mass times the speed of sound. Mathematically, this corresponds to proving that the Hamiltonian at fixed total momentum has an eigenvalue at the bottom of its spectrum. Since the excitations in this model are massless, this eigenvalue is always embedded in the essential spectrum. One expects that beyond some critical momentum this eigenvalue disappears and the system exhibits a Cherenkov transition. That is, the polaron would radiate sound waves, thereby slowing down to a stable state of smaller momentum. This has been validated numerically in [SSG⁺21a, SSG⁺21b], but there does not seem to be a mathematical proof of such a statement.

The dichotomy of stability at small velocities and a friction effect at high velocities has been studied in a model of a classical particle interacting with sound waves in the series of works [FGS11, FGS12, EG12, EFG⁺13, FG14a, FG14b], and later in [Lég20]. This model can be related to the Bose polaron system in a mean-field regime with a heavy impurity [DFPP14]. A simplified model is obtained by decoupling the directions of propagation of the particle and the waves, which limits the back-reaction of the field on the particle. Such a model was studied in [BDB02] in the quantum mechanical and [DBFS17] in the classical setting.

1.1. The Bogoliubov–Fröhlich Polaron. The Bogoliubov–Fröhlich Hamiltonian is characterized by the dispersion relation of the field of excitations, or phonons, and the form factor of the interaction. The dispersion relation is

$$(1.1) \quad \omega(k) := \sqrt{c^2|k|^2 + \xi^2|k|^4},$$

where $c > 0$ is the speed of sound and $\xi = 1/(2m_B)$, for the mass m_B of the bosons in the gas. The form factor of the particle-phonon interaction is

$$(1.2) \quad v_\Lambda(k) := \mathbf{g} \mathbf{1}_{|k| < \Lambda} \sqrt{|k|^2/\omega(k)},$$

where Λ is an ultraviolet cutoff (that may take the value infinity) and \mathbf{g} is a coupling constant, whose value will not be important in our analysis.

Our model is then realized as a selfadjoint lower-semibounded Hamiltonian on the bosonic Fock space \mathcal{F} over $L^2(\mathbb{R}^3)$. We use the standard notation a_k, a_k^* for the creation and annihilation operators on \mathcal{F} in the sense of operator-valued distributions. As usual, writing $d\Gamma(f) = \int f(k)a_k^*a_k dk$ and $\varphi(g) = \int (g(k)^*a_k + g(k)a_k^*)dk$ for second quantization and field operators, respectively, we define the Bogoliubov–Fröhlich Hamiltonian at momentum $P \in \mathbb{R}^3$ with cutoff $\Lambda < \infty$ by

$$(1.3) \quad H_\Lambda(P) := \frac{1}{2}(P - d\Gamma(\hat{p}))^2 + d\Gamma(\omega) + \varphi(v_\Lambda),$$

where $\hat{p} = (\hat{p}_1, \hat{p}_2, \hat{p}_3)$ denotes the vector of multiplication operators on $L^2(\mathbb{R}^3)$ given by $\hat{p}_i f(k) = k_i f(k)$. Note that we have set the impurity mass equal to one, keeping \mathbf{c}, ξ and \mathbf{g} as the model parameters. Moreover, we may choose $\mathbf{g} \geq 0$ without loss of generality, as the models with different signs (phases) in the coupling are unitarily equivalent via $e^{i\arg(\mathbf{g})d\Gamma(1)}$.

By the Kato–Rellich theorem, it follows from standard estimates that $H_\Lambda(P)$ is a selfadjoint lower-semibounded operator with $\mathcal{D}(H_\Lambda(P)) = \mathcal{D}(H_0(0)) = \mathcal{D}(d\Gamma(\hat{p})) \cap \mathcal{D}(d\Gamma(\omega))$ for all $\Lambda < \infty$, since $v_\Lambda, \omega^{-1/2}v_\Lambda \in L^2(\mathbb{R}^3)$.

For $\Lambda = \infty$, $H(P)$ is defined by the following renormalization result from [Lam20]. For the convenience of the reader, we sketch the proof in Section 2.

Proposition 1.1. *There exist $(\Sigma_\Lambda)_{\Lambda \geq 0} \subset \mathbb{R}$ and, for all $P \in \mathbb{R}^3$, a selfadjoint lower-semibounded operator $H(P)$ (given in Theorem 2.1) such that $H_\Lambda(P) - \Sigma_\Lambda$ converges to $H(P)$ as $\Lambda \rightarrow \infty$ in the norm resolvent sense.*

Proof. The statement is that of Theorem 2.1 with $\kappa = 0$. □

We are interested in studying the critical momentum of the operator $H(P)$.

1.2. Critical Momentum. As described earlier in this introduction, the polaron may become unstable for large momentum. We define the the critical momentum as

$$(1.4) \quad P_* := \sup\{P \in \mathbb{R}^3 : \inf \sigma(H(P)) \text{ is an eigenvalue of } H(P)\}.$$

The main result of this article can now easily be stated.

Theorem 1.2. *For any coupling constant $\mathbf{g} \geq 0$ and any speed of sound $\mathbf{c} > 0$, we have*

$$P_* \geq \mathbf{c}.$$

Proof. The statement is an immediate corollary of Theorem 3.1. □

Remark 1.3. It would be interesting to show that P_* is finite, and that there is a unique transition, i.e., $\inf \sigma(H(P))$ is an eigenvalue for all $P < P_*$. The numerical study [SSG⁺21a] supports this picture and indicates that the second derivative of E has a jump at P_* . Moreover, [SSG⁺21a, Fig.3(e)] suggests that P_* increases from \mathbf{c} to infinity as $\mathbf{g} \rightarrow \infty$.

Remark 1.4. An interpretation of the statement is to think of $P_* = m_*\mathbf{c}$, where m_* is the effective mass of the polaron (compare [SSG⁺21b]). Then we have shown $m_* \geq 1$, meaning that the quasi-particle is heavier than the impurity of mass one, in agreement with the picture that the particle is dressed by a cloud of phonons, increasing its effective mass. This definition of an effective mass is, of course, different from the common definition by the curvature $\partial_{|P|}^2 E(P)|_{P=0}$ at zero, but one may still expect similar qualitative behavior, see [Sei21, DS20] for a discussion of the latter quantity in the Fröhlich polaron model.

In order to prove Theorem 1.2, we have to deal with both an ultraviolet and an infrared problem. The first is due to the fact that $v_\infty \notin L^2(\mathbb{R}^3)$ (and also $\omega^{-1/2}v_\infty \notin L^2(\mathbb{R}^3)$). Using the method of interior boundary conditions, we can nevertheless describe the Bogoliubov–Fröhlich Hamiltonian, in particular its domain, without any ultraviolet regularization. This goes back to a recent article by the second author [Lam20], building on techniques developed for the related Fröhlich and Nelson models in [LS19] and improved on in the subsequent articles [Sch19, Lam19, Sch21, Lam23].

The infrared problem is due to the fact that $\omega(0) = 0$, which entails that $H(P)$ does not have a spectral gap and $\inf \sigma(H(P)) = \inf \sigma_{\text{ess}}(H(P))$. For massive polaron models, i.e., models satisfying $\text{ess inf } \omega > 0$, the existence of ground states is well known, see for example [Frö73, Spo88, DG99]. In the massless case, one distinguishes between the infrared singular case $v/\omega \notin L^2(\mathbb{R}^3)$ and the infrared regular one $v/\omega \in L^2(\mathbb{R}^3)$. In the infrared singular case, e.g., the famous Nelson model [Nel64], absence of ground states at arbitrary total momentum (and all non-zero couplings) has been shown, cf. [Frö73, HH08, Dam20, DH22]. In our case, however, the model is infrared-regular, as can be easily checked. There exists a variety of perturbative methods to prove existence of ground states in such a case for small values of the

total momentum and the coupling constant, e.g., operator theoretic renormalization [BFS98], iterated perturbation theory [Piz03, DP14] and functional integration methods [Spo98, BHL⁺02]. In particular, in presence of an ultraviolet cutoff $\Lambda < \infty$ and for small coupling, the fact that $P_* > 0$ follows from [DP14, Proposition 1.1]. Hence, we extend the existence of ground state to the case without ultraviolet cutoff, arbitrary coupling and a larger set of total momenta. The method we use in our proof is an adaption of a compactness argument first applied in [GLL01] and subsequently employed in the study of various models, e.g., the spin boson model [HHS21], the Nelson model [HM22, HHS23] and the Pauli–Fierz model [Mat16, HS23]. The general strategy is to introduce an artificial boson mass $\kappa > 0$, and then prove that the set of ground states with $\kappa \rightarrow 0$ is pre-compact and provides a minimizing sequence for $H(P)$.

In the remainder of this letter, we sketch the renormalization procedure leading to Proposition 1.1 in Section 2 and give the proof of Theorem 1.2 in Section 3.

2. RENORMALIZATION AND PROPERTIES OF THE BOGOLIUBOV–FRÖHLICH POLARON

In this section, we sketch the proof of Proposition 1.1, by reviewing the renormalization method employed in [Lam21]. The key idea is to identify a divergent and P -independent contribution Σ_Λ to $\inf \sigma(H_\Lambda(P))$. This contribution is of the form

$$(2.1) \quad \Sigma_\Lambda = e_1 \Lambda + e_2 \log \Lambda + \mathcal{O}(1).$$

The two divergent contributions of different orders arise in a two-step procedure of rewriting $H_\Lambda(P)$.

Throughout this section, we assume $P \in \mathbb{R}^3$ to be fixed. We emphasize that most of the defined objects, except for the contributions to Σ_Λ , do have a P -dependence. We now fix some parameter $\mu > 0$ and define

$$(2.2) \quad G_\Lambda = -(a(v_\Lambda)(H_0(P) + \mu)^{-1})^*.$$

Employing that $\omega^{-1}v \in L^2(\mathbb{R}^3)$, one can show that G_Λ is a bounded operator, including the case $\Lambda = \infty$, see the proof of Theorem 2.1 below for more details. Further, for $\Lambda < \infty$, we have the simple identity

$$(2.3) \quad H_\Lambda(P) = (1 - G_\Lambda^*)(H_0(P) + \mu)(1 - G_\Lambda) - G_\Lambda^*(H_0(P) + \mu)G_\Lambda - \mu,$$

which follows by expanding the product. The first singular contribution is contained in the term

$$(2.4) \quad -G_\Lambda^*(H_0(P) + \mu)G_\Lambda = -a(v_\Lambda)(H_0(P) + \mu)^{-1}a^*(v_\Lambda)$$

To make it explicit, we will put the creation and annihilation operators in this expression in normal order. With the pull-through formula $a_k(H_0(P) + \mu)^{-1} = (H_0(P - k) + \omega(k) + \mu)^{-1}a_k$, which holds by inspection on every n -particle sector of \mathcal{F} (see for example [BFS98, Lemma IV.8]), we find

$$(2.5) \quad \begin{aligned} -a(v_\Lambda)(H_0(P) + \mu)^{-1}a^*(v_\Lambda) &= -\int a_k \frac{v_\Lambda(k)v_\Lambda(\ell)}{H_0(P) + \mu} a_\ell^* dk d\ell \\ &= -\int \frac{v_\Lambda(k)^2}{H_0(P - k) + \omega(k) + \mu} dk - \int a_\ell^* \frac{v_\Lambda(k)v_\Lambda(\ell)}{H_0(P - k - \ell) + \omega(k) + \omega(\ell) + \mu} a_k dk d\ell. \end{aligned}$$

With this order of a_ℓ^*, a_k , the second term will be well defined also for $\Lambda = \infty$ as an unbounded operator, since the decay of $a_k \Psi$ in k for an element Ψ of its domain will make the integral convergent. For the first term this is not the case, and we will need to first subtract its divergent contribution to take $\Lambda \rightarrow \infty$. This can be chosen as

$$(2.6) \quad \Sigma_\Lambda^{(1)} = -\int \frac{v_\Lambda(k)^2}{\frac{1}{2}k^2 + \omega(k)} dk,$$

which has a divergence proportional to Λ since $v_\Lambda(k)$ is of order one for large k . We then define $T_{\Lambda,1} = \Theta_{\Lambda,1,0} + \Theta_{\Lambda,1,1}$, where $\Theta_{\Lambda,1,0} = \theta_{\Lambda,1,0}(d\Gamma(\hat{p}), d\Gamma(\omega))$ is a multiplication operator in the momentum representation and $\Theta_{\Lambda,1,1} = \int a_\ell^* \theta_{\Lambda,1,1}(d\Gamma(\hat{p}), d\Gamma(\omega), k, \ell) a_k dk d\ell$ is an integral operator, with

$$(2.7) \quad \begin{aligned} \theta_{\Lambda,1,0}(p, \eta) &= -\int \left(\frac{v_\Lambda(k)^2}{\frac{1}{2}(P - p - k)^2 + \eta + \omega(k) + \mu} - \frac{v_\Lambda(k)^2}{\frac{1}{2}k^2 + \omega(k)} \right) dk \\ \theta_{\Lambda,1,1}(p, \eta, k, \ell) &= -\frac{v_\Lambda(k)v_\Lambda(\ell)}{\frac{1}{2}(P - p - k - \ell)^2 + \eta + \omega(k) + \omega(\ell) + \mu}. \end{aligned}$$

Now, T_Λ makes sense also for $\Lambda = \infty$, since the integral defining $\theta_{\Lambda,1,0}$ has a limit for $\Lambda \rightarrow \infty$, and hence we could try to employ the identity

$$(2.8) \quad G_\Lambda^*(H_0(P) + \mu)G_\Lambda - \Sigma_\Lambda^{(1)} = T_\Lambda$$

to define the second term in (2.3) for a definition of $H(P)$. This is not enough, however, to remove the cutoff completely, since the (form) domain of the first term $(1 - G_\infty^*)(H_0(P) + \mu)(1 - G_\infty)$ is not contained in the (form) domain of T_∞ . To remedy this issue, we include T_Λ with the free operator.

For $\Lambda \in \mathbb{R}_+ \cup \{\infty\}$, let

$$(2.9) \quad \tilde{G}_\Lambda = -(a(v_\Lambda)(H_0(P) + T_\Lambda + \mu)^{-1})^*.$$

Then we can write a similar identity to (2.3) for $\Lambda < \infty$, explicitly

$$(2.10) \quad H_\Lambda = (1 - \tilde{G}_\Lambda^*)(H_0(P) + T_\Lambda + \mu)(1 - \tilde{G}_\Lambda) - a(v_\Lambda)(H_0(P) + T_\Lambda + \mu)^{-1}a^*(v_\Lambda) - T_\Lambda - \mu.$$

Expanding the resolvent gives

$$(2.11) \quad \begin{aligned} & -a(v_\Lambda)(H_0(P) + T_\Lambda + \mu)^{-1}a^*(v_\Lambda) - T_\Lambda \\ &= \Sigma_\Lambda^{(1)} + a(v_\Lambda)(H_0(P) + T_\Lambda + \mu)^{-1}T_\Lambda(H_0(P) + \mu)^{-1}a^*(v_\Lambda) \\ &= \Sigma_\Lambda^{(1)} + a(v_\Lambda)(H_0(P) + \mu)^{-1}T_\Lambda(H_0(P) + \mu)^{-1}a^*(v_\Lambda) \\ &\quad - a(v_\Lambda)(H_0(P) + \mu)^{-1}T_\Lambda(H_0(P) + T_\Lambda + \mu)^{-1}T_\Lambda(H_0(P) + \mu)^{-1}a^*(v_\Lambda). \end{aligned}$$

The term in the last line is regular in the case $\Lambda = \infty$ and will be treated as a remainder, while the first still contains the logarithmic divergence. To extract this, we proceed as before and put the creation and annihilation operators in normal order. However, there is now also the possibility of picking up a commutator between the operators in $\Theta_{\Lambda,1,1}$ and the outer creation/annihilation operators. With this in mind, the term with no remaining creation and annihilation operators reads

$$\int \frac{v_\Lambda(k)^2 \theta_{\Lambda,1,0}(d\Gamma(\hat{p}) + k, d\Gamma(\omega) + \omega(k))}{(H_0(P - k) + \omega(k) + \mu)^2} dk - \int \frac{v_\Lambda(k)v_\Lambda(\ell)\theta_{\Lambda,1,1}(d\Gamma(\hat{p}), d\Gamma(\omega), k, \ell)}{(H_0(P - k) + \omega(k) + \mu)(H_0(P - \ell) + \omega(\ell) + \mu)} dk d\ell.$$

These integrals have a logarithmic divergence as $\Lambda \rightarrow \infty$, captured by

$$(2.12) \quad \begin{aligned} \Sigma_\Lambda^{(2)} &= \int \frac{v_\Lambda(k)^2 \theta_{\Lambda,1,0}(k, \omega(k))}{(\frac{1}{2}k^2 + \omega(k))^2} dk \\ &\quad - \int \frac{v_\Lambda(k)^2 v_\Lambda(\ell)^2}{(\frac{1}{2}k^2 + \omega(k))(\frac{1}{2}(k + \ell)^2 + \omega(k) + \omega(\ell))(\frac{1}{2}\ell^2 + \omega(\ell))} dk d\ell. \end{aligned}$$

After subtracting this, we define $\tilde{T}_\Lambda = \Theta_{\Lambda,2,0} + \Theta_{\Lambda,2,1} + \Theta_{\Lambda,2,2}$, where $\Theta_{\Lambda,2,0}$ is a multiplication operator of the same type as $\Theta_{\Lambda,1,0}$, and $\Theta_{\Lambda,2,1}, \Theta_{\Lambda,2,2}$ are integrals with one, respectively two, remaining creation and annihilation operators. The expression for $\Theta_{\Lambda,2,0}$ is given by

$$(2.13) \quad \begin{aligned} \theta_{\Lambda,2,0}(p, \eta) &= \int \frac{v_\Lambda(k)^2 \theta_{\Lambda,1,0}(p + k, \eta + \omega(k))}{(\frac{1}{2}(P - p - k)^2 + \eta + \omega(k) + \mu)^2} dl \\ &\quad + \int \frac{v_\Lambda(k)v_\Lambda(\ell)\theta_{\Lambda,1,1}(p, \eta, k, \ell)}{(\frac{1}{2}(P - p - k)^2 + \eta + \omega(k) + \mu)(\frac{1}{2}(P - p - \ell)^2 + \eta + \omega(\ell) + \mu)} dk d\ell - \Sigma_\Lambda^{(2)}, \end{aligned}$$

where we observe that $\Sigma_\Lambda^{(2)}$ is simply the value of the integrals at $P = p = \eta = \mu = 0$. The integral operators have the kernels

$$(2.14) \quad \begin{aligned} \theta_{\Lambda,2,1}(p, \eta, k, \ell) &= \frac{v_\Lambda(k)v_\Lambda(\ell)\theta_{\Lambda,1,0}(p + k + \ell, \eta + \omega(k) + \omega(\ell))}{(\frac{1}{2}(P - p - k)^2 + \eta + \omega(k) + \mu)(\frac{1}{2}(P - p - \ell)^2 + \eta + \omega(\ell) + \mu)} \\ &\quad + \int \frac{v_\Lambda(\xi)^2 \theta_{\Lambda,1,1}(p + \xi, \eta + \omega(\xi), k, \ell)}{(\frac{1}{2}(P - p - \xi)^2 + \eta + \omega(\xi) + \mu)^2} dk d\ell \end{aligned}$$

$$(2.15) \quad \theta_{\Lambda,2,1}(p, \eta, k_1, k_2, \ell_1, \ell_2) = \frac{v_\Lambda(k_1)v_\Lambda(\ell_1)\theta_{\Lambda,1,1}(p + k_1 + \ell_1, \eta + \omega(k_1) + \omega(\ell_1), k_2, \ell_2)}{(\frac{1}{2}(P - p - k_1)^2 + \eta + \omega(k_1) + \mu)(\frac{1}{2}(P - p - \ell_1)^2 + \eta + \omega(\ell_1) + \mu)}.$$

Again, the definition of \tilde{T}_Λ may be extended to $\Lambda = \infty$ since these functions are defined also for this value. Finally, the definition of the remainder term reads

$$(2.16) \quad \begin{aligned} R_\Lambda &= -a(v_\Lambda)(H_0(P) + \mu)^{-1}T_\Lambda(H_0(P) + T_\Lambda + \mu)^{-1}T_\Lambda(H_0(P) + \mu)^{-1}a^*(v_\Lambda) \\ &= G_\Lambda^* T_\Lambda(H_0(P) + T_\Lambda + \mu)^{-1}T_\Lambda G_\Lambda, \end{aligned}$$

which defines a bounded operator also for $\Lambda = \infty$.

Since we require an infrared regularization in Section 3 additionally to the ultraviolet one provided by the cutoff Λ , we directly consider the family of operators $H_{\kappa,\Lambda}$ given by (1.3) with ω replaced by $\omega_\kappa = \omega + \kappa$, i.e.,

$$(2.17) \quad H_{\kappa,\Lambda}(P) := H_\Lambda(P) + \kappa N \quad \text{for } \kappa \geq 0, \Lambda \in \mathbb{R}_+,$$

where $N = d\Gamma(1)$ is the particle number operator as usual. Note that $\mathcal{D}(H_{\kappa,0}(P)) = \mathcal{D}(H_{0,0}(0)) \cap \mathcal{D}(\kappa N)$, so we simply denote this domain by $\mathcal{D}(H_{\kappa,0})$. For $\Lambda < \infty$, (2.17) immediately defines a selfadjoint lower-semibounded operator on $\mathcal{D}(H_{\kappa,0})$, since $v_\Lambda \in L^2(\mathbb{R}^3)$.

The preceding discussion applies in the same way with ω_κ , yielding objects $T_\Lambda = T_{\kappa,\Lambda}$, $\tilde{G}_\Lambda = \tilde{G}_{\kappa,\Lambda}$, $\tilde{T}_\Lambda = \tilde{T}_{\kappa,\Lambda}$, whose dependence on κ we will not make explicit. Proposition 1.1 is now a consequence of the following theorem for $\kappa = 0$.

Theorem 2.1 ([Lam21]). *Let $\kappa \geq 0$ and let $\tilde{G}_\infty, T_\infty, \tilde{T}_\infty, R_\infty$ be defined by (2.9), (2.7), (2.13)–(2.15), and (2.16) with $\omega = \omega_\kappa$ respectively. The operator*

$$H_{\kappa,\infty}(P) = (1 - \tilde{G}_\infty^*)(H_{\kappa,0}(P) + T_\infty + \mu)(1 - \tilde{G}_\infty) + \tilde{T}_\infty + R_\infty - \mu$$

$$\mathcal{D}(H_{\kappa,\infty}(P)) = \{\psi \in \mathcal{F} : (1 - \tilde{G}_\infty)\psi \in \mathcal{D}(H_{\kappa,0})\}$$

is selfadjoint and bounded from below. We have the convergence

$$H_{\kappa,\Lambda}(P) - \Sigma_\Lambda^{(1)} - \Sigma_\Lambda^{(2)} \rightarrow H_{\kappa,\infty}(P)$$

in norm resolvent sense, where $\Sigma_\Lambda^{(i)}$, $i = 1, 2$ are defined as in (2.6) and (2.12).

Sketch of the proof. We give a short outline of the proof with references to key technical lemmas for the convenience of the reader. Throughout this proof, $\kappa \geq 0$ is fixed.

The first step is to prove that

$$(2.18) \quad \|T_\Lambda \psi\| \leq C \|(H_{\kappa,0}(P) + 1)^{1/2} \psi\| \quad \text{for } \Lambda \in \mathbb{R}_+ \cup \{\infty\},$$

$$\|(T_\Lambda - T_\infty) \psi\| \leq C_\Lambda \|(H_{\kappa,0}(P) + 1)^{1/2+\varepsilon} \psi\|,$$

with $\varepsilon > 0$ and $\lim_{\Lambda \rightarrow \infty} C_\Lambda = 0$ (the part $\Theta_{\Lambda,1,0}$ can be bounded by an elementary calculation; concerning $\Theta_{\Lambda,1,1}$, see [Lam19, Lem.17] and [Lam23, Lemma B.2] for proofs in the case $\kappa > 0$ that are easily adapted to $\kappa = 0$).

Using this, one shows that for $\Lambda \in \mathbb{R}_+ \cup \{\infty\}$, \tilde{G}_Λ are bounded operators on \mathcal{F} , satisfying

$$(2.19) \quad \|(H_{\kappa,0}(P) + \mu)^s \tilde{G}_\Lambda\| \leq C, \quad (H_{\kappa,0}(P) + \mu)^s (\tilde{G}_\Lambda - \tilde{G}_\infty) \xrightarrow{\Lambda \rightarrow \infty} 0 \quad \text{for } 0 \leq s < 1/4.$$

This follows easily from the bound $\|a(f)d\Gamma(\eta)^{-1/2}\| \leq \|f/\eta\|$, $v/\omega \in L^2(\mathbb{R}^d)$ and the fact that T_Λ is an infinitesimal perturbation of $H_{\kappa,0}(P)$.

In particular, for μ large enough, $\|\tilde{G}_\Lambda\| < 1$, so $1 - \tilde{G}_\Lambda$ is invertible. This shows that $\mathcal{D}(H_{\kappa,\infty}(P))$ is dense and combined with (2.18), the operator

$$(2.20) \quad K := (1 - \tilde{G}_\infty^*)(H_{\kappa,0}(P) + T_\infty + \mu)(1 - \tilde{G}_\infty)$$

is selfadjoint and bounded from below on this domain. The terms $\tilde{T}_\infty, R_\infty$ will be treated as perturbations of K . For R_∞ , boundedness follows directly from the properties of T_Λ and \tilde{G}_Λ .

For \tilde{T}_Λ one can again show

$$(2.21) \quad \|\tilde{T}_\Lambda \psi\| \leq C \|(H_{\kappa,0}(P) + 1)^\varepsilon \psi\|,$$

$$(2.22) \quad \|(\tilde{T}_\Lambda - \tilde{T}_\infty) \psi\| \leq C_\Lambda \|(H_{\kappa,0}(P) + 1)^\varepsilon \psi\|$$

for $\varepsilon > 0$ and $\lim_{\Lambda \rightarrow \infty} C_\Lambda = 0$ (cf. [Lam23, Lemma B.2], [Lam19, Lem.19]). This implies that

$$(2.23) \quad \|\tilde{T}_\infty \psi\| \leq \|\tilde{T}_\infty(1 - \tilde{G}_\infty)\psi\| + \|\tilde{T}_\infty \tilde{G}_\infty \psi\|$$

$$\leq C(\|(H_{\kappa,0}(P) + 1)^\varepsilon(1 - \tilde{G}_\infty)\psi\| + \|(H_{\kappa,0}(P) + 1)^\varepsilon \tilde{G}_\infty \psi\|)$$

$$\leq \delta \|K_\kappa \psi\| + C_\delta \|\psi\|$$

for any $\delta > 0$. Thus $H_{\kappa,\infty}(P)$ is selfadjoint by the Kato–Rellich theorem.

Convergence of resolvents follows from the identity (2.10), the resolvent formula and the convergence properties of $T_\Lambda, \tilde{G}_\Lambda$ already mentioned. \square

From the proof, we also obtain the following Lemma 2.2, which relates the domains of $H(P)$ and N . It will be important to our proof of Theorem 1.2 in the next section.

Lemma 2.2. *For any $P \in \mathbb{R}^3$, the subspace $\mathcal{D}(N) \cap \mathcal{D}(H_{0,\infty}(P))$ is a core for $H_{0,\infty}(P)$. Further, $\mathcal{D}(H_{\kappa,\infty}(P)) = \mathcal{D}(N) \cap \mathcal{D}(H_{0,\infty}(P))$ for all $\kappa > 0$ and $H_{\kappa,\infty}(P) = H_{0,\infty}(P) + \kappa N$.*

Proof. From Theorem 2.1, we know that $\mathcal{D}(H_{\kappa,\infty}(P)) = (1 - \tilde{G}_\infty)^{-1} \mathcal{D}(H_{\kappa,0})$. Moreover, for any core \mathcal{C} of $H_{\kappa,0}(P)$, $(1 - \tilde{G}_\infty)^{-1} \mathcal{C}$ is a core for $H_{\kappa,\infty}(P)$, since $(1 - \tilde{G}_\infty)^{-1} : \mathcal{D}(H_{\kappa,0}) \rightarrow \mathcal{D}(K)$ is continuous for the graph norms. Hence, to prove the domain statements, it suffices to show $(1 - \tilde{G}_\infty) \mathcal{D}(N) = \mathcal{D}(N)$. This follows from the observation $N\tilde{G}_\infty = \tilde{G}_\infty(N + 1)$, implying

$$(2.24) \quad \|N(1 - \tilde{G}_\infty)\psi\| \leq \|N\psi\| + \|\tilde{G}_\infty\| \|(N + 1)\psi\|,$$

$$(2.25) \quad \|N\psi\| \leq \|N(1 - \tilde{G}_\infty)\psi\| + \|N\tilde{G}_\infty\psi\| \leq \|N(1 - \tilde{G}_\infty)\psi\| + \|\tilde{G}_\infty\| \|(N + 1)\psi\|,$$

from where we conclude using $\|\tilde{G}_\infty\| < 1$. Moreover,

$$(2.26) \quad H_{\kappa,\infty}(P) = H_{0,\infty}(P) + \kappa N$$

holds since both sides are the weak graph limit of $H_{\kappa,\Lambda}(P)$, by Theorem 2.1, [RS72, Thm.VIII.26], and the uniform bound $\|N\psi\| \leq C\|(H_{\kappa,\Lambda}(P) + \mu)\psi\|$. \square

Remark 2.3. One can observe that $\mathcal{D}(H(P)) \neq \mathcal{D}(H(P'))$ for $P \neq P'$. This is the case because

$$(2.27) \quad (H_0(P) + \mu)^{-1} d\Gamma(\hat{p})G_\infty,$$

which is proportional to the difference of G_∞ for two different values of P , does not map $\mathcal{D}(H_0)$ to itself. It does, however, map the form domain of $H_{\kappa,0}(P)$ to itself, so the operators with different total momenta still have comparable quadratic forms. Notwithstanding, this fact will not be used in our arguments.

3. EXISTENCE OF GROUND STATES

In this section, we prove the following theorem.

Theorem 3.1. *If $|P| < c$, then $\inf \sigma(H(P))$ is an eigenvalue of $H(P)$.*

To prove the statement, we approximate $H(P) = H_{0,\infty}(P)$ by the infrared regularized Hamiltonians $H_{\kappa,\infty}(P)$ with $\kappa > 0$ introduced in (2.17). Further, we write

$$(3.1) \quad E_{\kappa,\Lambda}(P) := \inf \sigma(H_{\kappa,\Lambda}(P)) \quad \text{for all } \kappa \geq 0, \Lambda \in \mathbb{R}_+ \cup \{\infty\}, P \in \mathbb{R}^3.$$

Let us first observe that the ground state energies converge, when removing any regularization.

Lemma 3.2. *For any fixed $\kappa \geq 0$, $\Lambda \in \mathbb{R}_+ \cup \{\infty\}$ and $P \in \mathbb{R}^3$, we have*

$$E_{\kappa,\infty} = \lim_{\sigma \rightarrow \infty} E_{\kappa,\sigma} \quad \text{and} \quad E_{0,\Lambda} = \lim_{\eta \downarrow 0} E_{\eta,\Lambda}.$$

Proof. The first statement is a consequence of the norm resolvent convergence established in Theorem 2.1 (cf. [dO09]). For the second statement, we observe that $\mathcal{D}(N) \cap \mathcal{D}(H_{0,\Lambda}(P))$ is a core for $H_{0,\Lambda}(P)$, by the Kato–Rellich theorem for $\Lambda < \infty$ and by Lemma 2.2 for $\Lambda = \infty$. Hence, picking any $\varepsilon > 0$, there exists $\varphi_\varepsilon \in \mathcal{D}(N) \cap \mathcal{D}(H_{0,\Lambda}(P))$ with $\|\varphi_\varepsilon\| = 1$ such that $\langle \varphi_\varepsilon, H_{0,\Lambda}(P)\varphi_\varepsilon \rangle < E_{0,\Lambda}(P) + \varepsilon$. Further employing that $H_{\eta,\Lambda}(P) - H_{0,\Lambda}(P) \geq 0$ (as a form inequality), again by (2.17) and Lemma 2.2, we find

$$E_{0,\Lambda}(P) \leq E_{\eta,\Lambda}(P) \leq \langle \varphi_\varepsilon, H_{\eta,\Lambda}(P)\varphi_\varepsilon \rangle = \langle \varphi_\varepsilon, H_{0,\Lambda}(P)\varphi_\varepsilon \rangle + \eta \langle \varphi_\varepsilon, N\varphi_\varepsilon \rangle \leq E_{0,\Lambda}(P) + \varepsilon + \eta \langle \varphi_\varepsilon, N\varphi_\varepsilon \rangle.$$

First taking $\eta \downarrow 0$ and then $\varepsilon \downarrow 0$ finishes the proof. \square

The mass term κN ensures the existence of a spectral gap for small enough P , as a consequence of the following well-known HVZ-type theorem [Fr73, Møl05].

Proposition 3.3 ([Møl05, Theorem 1.2]). *For all $\kappa > 0$, $\Lambda \in \mathbb{R}_+$, we have*

$$(3.2) \quad \inf \sigma_{\text{ess}}(H_{\kappa,\Lambda}(P)) = \inf_{\substack{k_1, \dots, k_n \in \mathbb{R}^3 \\ n \in \mathbb{N}}} [E_{\kappa,\Lambda}(P - k_1 - \dots - k_n) + \omega(k_1) + \dots + \omega(k_n) + n\kappa].$$

In view of the above proposition, we need to estimate the difference $E(P - k) - E(P)$. This can be done using simple convexity arguments, cf. [LMS07, HHS23].

Lemma 3.4. *Let $\kappa \geq 0$, $\Lambda \in \mathbb{R}_+ \cup \{\infty\}$ and $P, K \in \mathbb{R}^3$. Then*

$$E_{\kappa,\Lambda}(P - K) - E_{\kappa,\Lambda}(P) \geq -|K||P|.$$

Proof. First, we assume that $\kappa > 0$ and $\Lambda < \infty$ and prove the inequalities

$$(3.3) \quad 0 \leq E_{\kappa,\Lambda}(P) - E_{\kappa,\Lambda}(0) \leq \frac{1}{2}|P|^2 \quad \text{for all } P \in \mathbb{R}^3.$$

The first inequality goes back to Gross [Gro72], see [Hin22, Lemma 3.4] for a recent adaption which covers our case. Now, note that Proposition 3.3 combined with the first inequality yields

$$(3.4) \quad \inf \sigma_{\text{ess}}(H_{\kappa,\Lambda}(0)) \geq E_{\kappa,\Lambda}(0) + \kappa,$$

so $E_{\kappa,\Lambda}(0)$ is a discrete eigenvalue with corresponding normalized eigenvector $\psi_0 \in \mathcal{D}(H_{\kappa,\Lambda}(0)) = \mathcal{D}(H_{\kappa,\Lambda}(P))$. Then

$$E_{\kappa,\Lambda}(P) \leq \langle \psi_0, H_{\kappa,\Lambda}(P)\psi_0 \rangle = E_{\kappa,\Lambda}(0) + \frac{1}{2}|P|^2 + \langle \psi_0, P \cdot \text{d}\Gamma(\hat{p})\psi_0 \rangle.$$

This implies $\frac{1}{2}|P|^2 + \langle \psi_0, P \cdot \text{d}\Gamma(\hat{p})\psi_0 \rangle \geq 0$ for all $P \in \mathbb{R}^3$. Letting $P \rightarrow 0$, this yields $e \cdot \langle \psi_0, \text{d}\Gamma(\hat{p})\psi_0 \rangle \geq 0$ for all normalized $e \in \mathbb{R}^3$, whence $\langle \psi_0, \text{d}\Gamma(\hat{p})\psi_0 \rangle = 0$. This proves the upper bound in (3.3).

Clearly, the map

$$(3.5) \quad P \mapsto \frac{1}{2}P^2 - E_{\kappa,\Lambda}(P) = - \inf_{\psi \in \mathcal{D}(H_{\kappa,\Lambda}(P))} \langle \psi, (-P \cdot \text{d}\Gamma(\hat{p}) + \frac{1}{2}\text{d}\Gamma(\hat{p})^2 + H_{\kappa,\Lambda}(0))\psi \rangle$$

is convex, as a supremum over linear functions of P . By a general result on convex functions taking nonnegative values below the standard parabola (essentially the fact that such a function must lie below any segment that intersects its graph and is tangent to the parabola, cf. [LMS07, Appendix A] or [HHS23, Cor. A.6]), this gives for $\kappa > 0$, $\Lambda < \infty$

$$(3.6) \quad E_{\kappa,\Lambda}(P - K) - E_{\kappa,\Lambda}(P) \geq \begin{cases} -|K||P| + \frac{1}{2}|K|^2 & \text{if } |K| \leq |P|, \\ -\frac{1}{2}|P|^2 & \text{if } |K| > |P|. \end{cases}$$

In both cases the right hand side is larger than $-|K||P|$ as claimed. This proves the claim for $\kappa > 0$ and $\Lambda < \infty$. The general statement follows from the convergence results of Lemma 3.2. \square

Corollary 3.5. *If $|P| \leq c$, then $E_{\kappa,\Lambda}(P)$ is a discrete eigenvalue of $H_{\kappa,\Lambda}(P)$ for all $\kappa > 0$ and $\Lambda \in \mathbb{R}_+ \cup \{\infty\}$.*

Proof. First assume $\Lambda < \infty$. Combining the HVZ Theorem, Proposition 3.3, with Lemma 3.4 gives

$$(3.7) \quad \inf \sigma_{\text{ess}}(H_{\kappa,\Lambda}(P)) - E_{\kappa,\Lambda}(P) \geq \inf_{\substack{k_1, \dots, k_n \in \mathbb{R}^3 \\ n \in \mathbb{N}}} \left(\sum_{i=1}^n (\omega(k_i) + \kappa) - |P| \left| \sum_{i=1}^n k_i \right| \right).$$

Since the absolute value is subadditive and $\omega(k) \geq c|k|$, this is larger than κ for $|P| \leq c$, which proves the statement in the case $\Lambda < \infty$. The case $\Lambda = \infty$ directly follows from Theorem 2.1, since norm resolvent convergence implies convergence of $\inf \sigma_{\text{ess}}(H_{\kappa,\Lambda}(P))$ and $E_{\kappa,\Lambda}(P)$. \square

We now identify a compact set in Fock space containing the (normalized) ground states of $H_{\kappa,\Lambda}(P)$. To this end, we define

$$(3.8) \quad \mathcal{G}_r := \left\{ \psi \in \mathcal{F} : \|a_k \psi\| \leq \frac{r}{\sqrt{|k|} \vee |k|^2}, \|(a_{k+p} - a_k)\psi\| \leq \frac{r|p|}{|k|^2} \quad \text{for a.e. } k, p \in \mathbb{R}^3, |p| \leq \frac{1}{2}|k| \right\}.$$

Lemma 3.6. *For all $r > 0$, the set \mathcal{G}_r is pre-compact in \mathcal{F} .*

Proof. The elements of \mathcal{G}_r are localized by the first bound, and regular by the second. Conditions of this type are well-known to yield compactness, see [HHS23, Theorem 3.4] for a detailed proof. \square

Now, we prove that $\overline{\mathcal{G}_r}$ contains the ground states of $H_{\kappa,\Lambda}(P)$.

Proposition 3.7. *If $|P| < c$, there exist $r > 0$ (depending on $|P|$, \mathfrak{g} and c) such that for all $\kappa > 0$ and $\Lambda \in \mathbb{R}_+ \cup \{\infty\}$ and any normalized $\psi \in \mathcal{D}(H_{\kappa,\Lambda}(P))$ with $H_{\kappa,\Lambda}(P)\psi = E_{\kappa,\Lambda}(P)\psi$, we have $\psi \in \overline{\mathcal{G}_r}$.*

Proof. Throughout this proof, $r > 0$ denotes a (not necessarily fixed) constant solely depending on $|P|$, \mathfrak{g} and c . Especially, r is independent of κ or Λ . Further, fix $\kappa > 0$, $\Lambda < \infty$ and ψ as in the statement.

The starting point of our proof is the pull-through formula

$$(3.9) \quad a_k \psi = -v_\Lambda(k) R_{\kappa,\Lambda}(P, k) \psi \quad \text{with} \quad R_{\kappa,\Lambda}(P, k) := (H_{\kappa,\Lambda}(P - k) - E_{\kappa,\Lambda}(P) + \omega(k) + \kappa)^{-1},$$

which holds true for almost every $k \in \mathbb{R}^3$. To check this, compute using the commutation relations

$$(3.10) \quad 0 = a_k (H_{\kappa,\Lambda}(P) - E_{\kappa,\Lambda}(P)) \psi = v_\Lambda(k) \psi + (H_{\kappa,\Lambda}(P - k) + \omega(k) + \kappa - E_{\kappa,\Lambda}(P)) a_k \psi.$$

The formula then follows by applying $R_{\kappa,\Lambda}(P, k)$, which is well defined since $E_{\kappa,\Lambda}(P - k) \geq E_{\kappa,\Lambda}(P) - |k||P| \geq E_{\kappa,\Lambda}(P) - \omega(k)$, by Lemma 3.4 and the assumption $|P| \leq c$, see e.g. [Gér00, Dam20] for more details.

To estimate the resolvent, we use Lemma 3.4 to obtain the bounds

$$(3.11) \quad E_{\kappa,\Lambda}(P - k) - E_{\kappa,\Lambda}(P) + \omega(k) \geq \omega(k) - |P||k| \geq \begin{cases} (c - |P|)|k| & \text{for all } k \in \mathbb{R}^3, \\ \frac{\xi}{2}|k|^2 & \text{if } |k| > 2\xi^{-1}|P|. \end{cases}$$

Then using that $H_{\kappa,\Lambda}(P - k) \geq E_{\kappa,\Lambda}(P - k)$, it follows directly from the spectral theorem that

$$(3.12) \quad \|R_{\kappa,\Lambda}(P, k)\| \leq \begin{cases} ((c - |P|)|k|)^{-1} & \text{for all } k \in \mathbb{R}^3, \\ 2\xi^{-1}|k|^{-2} & \text{if } |k| > 2\xi^{-1}|P|. \end{cases}$$

Further employing $|v(k)| \leq \mathfrak{g}(\sqrt{|k|/c} \wedge 1)$ we find, for an appropriate choice of r ,

$$(3.13) \quad \|v_\Lambda(k)R_{\kappa,\Lambda}(P, k)\| \leq \frac{r}{\sqrt{|k|} \vee |k|^2},$$

which combined with (3.9) proves the desired bound on $a_k\psi$ in the definition (3.8).

Now let $|p| \leq \frac{1}{2}|k|$. The resolvent identity gives

$$(3.14) \quad \begin{aligned} a_{k+p}\psi - a_k\psi &= (v_\Lambda(k+p) - v_\Lambda(k))R_{\kappa,\Lambda}(P, k+p)\psi \\ &\quad + v_\Lambda(k)R_{\kappa,\Lambda}(P, k+p)[p \cdot (P - k - d\Gamma(\hat{p}))]R_{\kappa,\Lambda}(P, k)\psi. \end{aligned}$$

Since $|\nabla v_\Lambda(\ell)| \leq C\omega^{-1/2}(k)$ for $\frac{1}{2}|k| \leq \ell \leq \frac{3}{2}|k|$, cf. (1.2), using (3.12) yields

$$(3.15) \quad \|(v_\Lambda(k+p) - v_\Lambda(k))R_{\kappa,\Lambda}(P, k+p)\psi\| \leq \frac{8C|p|}{(c - |P|)|k|\sqrt{\omega(k)}} \leq \frac{r|p|}{|k|^2}.$$

As $\| |P - k - d\Gamma(\hat{p}) | R_{\kappa,\Lambda}(P, k) \| \leq 1$, (3.13) also implies

$$(3.16) \quad \|v_\Lambda(k)R_{\kappa,\Lambda}(P, k+p)[p \cdot (P - k - d\Gamma(\hat{p}))]R_{\kappa,\Lambda}(P, k)\psi\| \leq \frac{r|p|}{|k|^2}.$$

Combining (3.14), (3.15) and (3.16) and the definition (3.8), this proves that $\psi \in \mathcal{G}_r$ in the case $\Lambda < \infty$.

As a consequence of norm-resolvent convergence and the uniform gap estimate in Corollary 3.5, the spectral projections of $H_{\kappa,\Lambda}(P)$ converge to those of $H_{\kappa,\infty}(P)$ (cf. [RS72, Theorem VIII.23]), whence the ground states of $H_{\kappa,\infty}$ are contained in the closure $\overline{\mathcal{G}_r}$. \square

We conclude with the proof of our main result.

Proof of Theorem 3.1. Let ψ_κ denote any normalized ground state of $H_{\kappa,\infty}(P)$ for $\kappa > 0$. Since $\mathcal{D}(H_{\kappa,\infty}(P)) \subset \mathcal{D}(H(P))$, cf. Lemma 2.2, we find

$$(3.17) \quad 0 \leq \langle \psi_\kappa, (H(P) - E(P))\psi_\kappa \rangle \leq \langle \psi_\kappa, (H_{\kappa,\infty}(P) - E(P))\psi_\kappa \rangle = E_{\kappa,\infty}(P) - E_{0,\infty}(P) \xrightarrow{\kappa \downarrow 0} 0,$$

by Lemma 3.2. Further, since $(\psi_\kappa)_{\kappa > 0} \subset \overline{\mathcal{G}_r}$ by Proposition 3.7, which is compact by Lemma 3.6, there exists a zero sequence $(\kappa_n)_{n \in \mathbb{N}}$ such that the limit $\psi_\infty = \lim_{n \rightarrow \infty} \psi_{\kappa_n}$ exists. The estimate (3.17) then implies that $\psi_\infty \in \mathcal{D}(H(P)^{1/2})$, whence ψ_∞ is a minimizer of the closed quadratic form of $H(P)$, and thus an eigenvector, which finishes the proof. \square

ACKNOWLEDGEMENTS

BH acknowledges funding by the Ministry of Culture and Science of the State of North Rhine-Westphalia within the project ‘PhoQC’. JL thanks Christian Hainzl for discussions on the subject and acknowledges financial support through the EUR-EIPHI Graduate School (ANR-17-EURE-0002).

REFERENCES

- [BDB02] L. Bruneau and S. De Bièvre. A Hamiltonian Model for Linear Friction in a Homogeneous Medium. *Commun. Math. Phys.*, 229(3):511–542, 2002. doi:10.1007/s00220-002-0689-0.
- [BFS98] V. Bach, J. Fröhlich, and I. M. Sigal. Quantum Electrodynamics of Confined Nonrelativistic Particles. *Adv. Math.*, 137(2):299–395, 1998. doi:10.1006/aima.1998.1734.
- [BHL⁺02] V. Betz, F. Hiroshima, J. Lőrinczi, R. A. Minlos, and H. Spohn. Ground state properties of the Nelson Hamiltonian - A Gibbs measure-based approach. *Rev. Math. Phys.*, 14(02):173–198, 2002, arXiv:math-ph/0106015. doi:10.1142/S0129055X02001119.

- [CLB15] R. S. Christensen, J. Levinsen, and G. M. Bruun. Quasiparticle Properties of a Mobile Impurity in a Bose-Einstein Condensate. *Phys. Rev. Lett.*, 115(16):160401, 2015, arXiv:1503.06979. doi:10.1103/PhysRevLett.115.160401.
- [Dam20] T. N. Dam. Absence of Ground States in the Translation Invariant Massless Nelson Model. *Ann. Henri Poincaré*, 21(8):2655–2679, 2020, arXiv:1808.00088. doi:10.1007/s00023-020-00928-y.
- [DBFS17] S. De Bièvre, J. Faupin, and B. Schubnel. Spectral analysis of a model for quantum friction. *Rev. Math. Phys.*, 29(06):1750019, 2017, arXiv:1512.05704. doi:10.1142/S0129055X17500192.
- [DFPP14] D.-A. Deckert, J. Fröhlich, P. Pickl, and A. Pizzo. Effective Dynamics of a Tracer Particle Interacting with an Ideal Bose Gas. *Commun. Math. Phys.*, 328(2):597–624, 2014, arXiv:1210.0253. doi:10.1007/s00220-014-1987-z.
- [DG99] J. Dereziński and C. Gérard. Asymptotic Completeness in Quantum Field Theory. Massive Pauli-Fierz Hamiltonians. *Rev. Math. Phys.*, 11(4):383–450, 1999. doi:10.1142/S0129055X99000155.
- [DH22] T. N. Dam and B. Hinrichs. Absence of ground states in the renormalized massless translation-invariant Nelson model. *Rev. Math. Phys.*, 34(10):2250033, 2022, arXiv:1909.07661. doi:10.1142/S0129055X22500337.
- [dO09] C. R. de Oliveira. *Intermediate Spectral Theory and Quantum Dynamics*, volume 54 of *Progress in Mathematical Physics*. Birkhäuser, Basel, 2009. doi:10.1007/978-3-7643-8795-2.
- [DP14] W. Dybalski and A. Pizzo. Coulomb Scattering in the Massless Nelson model I. Foundations of Two-Electron Scattering. *J. Stat. Phys.*, 154(1):543–587, 2014, arXiv:1302.5001. doi:10.1007/s10955-013-0857-y.
- [DS20] W. Dybalski and H. Spohn. Effective Mass of the Polaron – Revisited. *Ann. Henri Poincaré*, 21(5):1573–1594, 2020, arXiv:1908.03432. doi:10.1007/s00023-020-00892-7.
- [EFG⁺13] D. Egli, J. Fröhlich, Z. Gang, A. Shao, and I. M. Sigal. Hamiltonian Dynamics of a Particle Interacting with a Wave Field. *Commun. Part. Diff. Eq.*, 38(12):2155–2198, 2013, arXiv:1211.6154. doi:10.1080/03605302.2013.816857.
- [EG12] D. Egli and Z. Gang. Some Hamiltonian models of friction II. *J. Math. Phys.*, 53(10):103707, 2012, arXiv:1108.5545. doi:10.1063/1.4757278.
- [FG14a] J. Fröhlich and Z. Gang. Ballistic motion of a tracer particle coupled to a Bose gas. *Adv. Math.*, 259:252–268, 2014, arXiv:1302.0429. doi:10.1016/j.aim.2014.03.014.
- [FG14b] J. Fröhlich and Z. Gang. Emission of Cherenkov radiation as a mechanism for Hamiltonian friction. *Adv. Math.*, 264:183–235, 2014, arXiv:1311.6377. doi:10.1016/j.aim.2014.07.013.
- [FGS11] J. Fröhlich, Z. Gang, and A. Soffer. Some Hamiltonian models of friction. *J. Math. Phys.*, 52(8):083508, 2011, arXiv:1011.1283. doi:10.1063/1.3619799.
- [FGS12] J. Fröhlich, Z. Gang, and A. Soffer. Friction in a Model of Hamiltonian Dynamics. *Commun. Math. Phys.*, 315(2):401–444, 2012, arXiv:1110.6550. doi:10.1007/s00220-012-1564-2.
- [Frö73] J. Fröhlich. On the infrared problem in a model of scalar electrons and massless scalar bosons. *Ann. de l'Inst. Henri Poincaré*, 19(1):1–103, 1973.
- [GD16] F. Grusdt and E. Demler. New theoretical approaches to Bose polarons. In *Quantum Matter at Ultralow Temperatures*, volume 191 of *Proceedings of the international school of physics "Enrico Fermi"*, pages 325–411, 2016, arXiv:1510.04934. doi:10.3254/978-1-61499-694-1-325.
- [Gér00] C. Gérard. On the Existence of Ground States for Massless Pauli-Fierz Hamiltonians. *Ann. Henri Poincaré*, 1(3):443–459, 2000. doi:10.1007/s000230050002.
- [GLL01] M. Griesemer, E. Lieb, and M. Loss. Ground states in non-relativistic quantum electrodynamics. *Invent. math.*, 145(3):557–595, 2001, arXiv:math-ph/0007014. doi:10.1007/s002220100159.
- [Gro72] L. Gross. Existence and uniqueness of physical ground states. *J. Funct. Anal.*, 10(1):52–109, 1972. doi:10.1016/0022-1236(72)90057-2.
- [GSSD17] F. Grusdt, R. Schmidt, Y. E. Shchadilova, and E. Demler. Strong-coupling Bose polarons in a Bose-Einstein condensate. *Phys. Rev. A*, 96(1):013607, 2017, arXiv:1704.02605. doi:10.1103/PhysRevA.96.013607.
- [HH08] D. Hasler and I. Herbst. Absence of Ground States for a Class of Translation Invariant Models of Non-relativistic QED. *Commun. Math. Phys.*, 279(3):769–787, 2008, arXiv:math-ph/0702096. doi:10.1007/s00220-008-0444-2.
- [HHS21] D. Hasler, B. Hinrichs, and O. Siebert. On Existence of Ground States in the Spin Boson Model. *Commun. Math. Phys.*, 388(1):419–433, 2021, arXiv:2102.13373. doi:10.1007/s00220-021-04185-w.
- [HHS23] D. Hasler, B. Hinrichs, and O. Siebert. Non-Fock Ground States in the Translation-Invariant Nelson Model Revisited Non-Perturbatively. Preprint, 2023, arXiv:2302.06998.
- [Hin22] B. Hinrichs. *Existence of Ground States for Infrared-Critical Models of Quantum Field Theory*. PhD thesis, Friedrich Schiller University Jena, 2022. doi:10.22032/dbt.51516.
- [HM22] F. Hiroshima and O. Matte. Ground states and their associated path measures in the renormalized Nelson model. *Rev. Math. Phys.*, 34(2):2250002, 2022, arXiv:1903.12024. doi:10.1142/S0129055X22500027.
- [HS23] D. Hasler and O. Siebert. Ground States for translationally invariant Pauli-Fierz Models at zero Momentum. *J. Funct. Anal.*, 284(1):109725, 2023, arXiv:2007.01250. doi:10.1016/j.jfa.2022.109725.
- [Lam19] J. Lampart. A Nonrelativistic Quantum Field Theory with Point Interactions in Three Dimensions. *Ann. Henri Poincaré*, 20(11):3509–3541, 2019, arXiv:1804.08295. doi:10.1007/s00023-019-00843-x.
- [Lam20] J. Lampart. The renormalized Bogoliubov-Fröhlich Hamiltonian. *J. Math. Phys.*, 61(10):101902, 2020, arXiv:1909.02430. doi:10.1063/5.0014217.
- [Lam21] J. Lampart. The Resolvent of the Nelson Hamiltonian Improves Positivity. *Math. Phys. Anal. Geom.*, 24(1):2, 2021, arXiv:2010.03235. doi:10.1007/s11040-021-09374-6.
- [Lam23] J. Lampart. Hamiltonians for Polaron Models with Subcritical Ultraviolet Singularities. *Ann. Henri Poincaré*, 24(8):2687–2728, 2023, arXiv:2203.07253. doi:10.1007/s00023-023-01285-2.
- [Lég20] T. Léger. Scattering for a particle interacting with a Bose gas. *Commun. Part. Diff. Eq.*, 45(10):1381–1413, 2020, arXiv:1902.03012. doi:10.1080/03605302.2020.1774896.
- [LMS07] M. Loss, T. Miyao, and H. Spohn. Lowest energy states in nonrelativistic QED: Atoms and ions in motion. *J. Funct. Anal.*, 243(2):353–393, 2007, arXiv:math-ph/0605005. doi:10.1016/j.jfa.2006.10.012.

- [LP22] J. Lampart and P. Pickl. Dynamics of a Tracer Particle Interacting with Excitations of a Bose–Einstein Condensate. *Ann. Henri Poincaré*, 23(8):2855–2876, 2022, arXiv:2011.14428. doi:10.1007/s00023-022-01153-5.
- [LS19] J. Lampart and J. Schmidt. On Nelson–Type Hamiltonians and Abstract Boundary Conditions. *Commun. Math. Phys.*, 367(2):629–663, 2019, arXiv:1803.00872. doi:10.1007/s00220-019-03294-x.
- [LT] J. Lampart and A. Triay. The excitation spectrum of a dilute Bose gas with an impurity. In preparation.
- [Mat16] O. Matte. Continuity properties of the semi-group and its integral kernel in non-relativistic QED. *Rev. Math. Phys.*, 28(05):1650011, 2016, arXiv:1512.04494. doi:10.1142/S0129055X16500112.
- [Møl05] J. S. Møller. The Translation Invariant Massive Nelson Model: I. The Bottom of the Spectrum. *Ann. Henri Poincaré*, 6(6):1091–1135, 2005. doi:10.1007/s00023-005-0234-8.
- [MS20] K. Myśliwy and R. Seiringer. Microscopic Derivation of the Fröhlich Hamiltonian for the Bose Polaron in the Mean-Field Limit. *Ann. Henri Poincaré*, 21(12):4003–4025, 2020, arXiv:2003.12371. doi:10.1007/s00023-020-00969-3.
- [Nel64] E. Nelson. Interaction of Nonrelativistic Particles with a Quantized Scalar Field. *J. Math. Phys.*, 5(9):1190–1197, 1964. doi:10.1063/1.1704225.
- [Piz03] A. Pizzo. One-particle (improper) States in Nelson’s Massless Model. *Ann. Henri Poincaré*, 4(3):439–486, 2003. doi:10.1007/s00023-003-0136-6.
- [RS72] M. Reed and B. Simon. *Functional Analysis*, volume 1 of *Methods of Modern Mathematical Physics*. Academic Press, San Diego, 1972.
- [Sch19] J. Schmidt. On a direct description of pseudorelativistic Nelson Hamiltonians. *J. Math. Phys.*, 60(10):102303, 2019, arXiv:1810.03313. doi:10.1063/1.5109640.
- [Sch21] J. Schmidt. The Massless Nelson Hamiltonian and its Domain. In A. Michelangeli, editor, *Mathematical Challenges of Zero-Range Physics*, volume 42 of *Springer INdAM Series*. Springer, 2021, arXiv:1901.05751. doi:10.1007/978-3-030-60453-0.
- [Sei21] R. Seiringer. The polaron at strong coupling. *Rev. Math. Phys.*, 33(1):2060012, 2021, arXiv:1912.12509. doi:10.1142/S0129055X20600120.
- [Spo88] H. Spohn. The polaron at large total momentum. *J. Phys. A*, 21(5):1199–1211, 1988. doi:10.1088/0305-4470/21/5/016.
- [Spo98] H. Spohn. Ground State of a Quantum Particle Coupled to a Scalar Bose Field. *Lett. Math. Phys.*, 44(1):9–16, 1998. doi:10.1023/A:1007473300274.
- [SSG⁺21a] K. Seetharam, Y. Shchadilova, F. Grusdt, M. Zvonarev, and E. Demler. Quantum Cherenkov transition of finite momentum Bose polarons. Preprint, 2021, arXiv:2109.12260.
- [SSG⁺21b] K. Seetharam, Y. Shchadilova, F. Grusdt, M. B. Zvonarev, and E. Demler. Dynamical Quantum Cherenkov Transition of Fast Impurities in Quantum Liquids. *Phys. Rev. Lett.*, 127:185302, 2021, arXiv:2101.00030. doi:10.1103/PhysRevLett.127.185302.

BENJAMIN HINRICHS, UNIVERSITÄT PADERBORN, INSTITUT FÜR MATHEMATIK, INSTITUT FÜR PHOTONISCHE QUANTENSYSTEME, WARBURGER STR. 100, 33098 PADERBORN, GERMANY

Email address: benjamin.hinrichs@math.upb.de

JONAS LAMPART, CNRS & LABORATOIRE INTERDISCIPLINAIRE CARNOT DE BOURGOGNE (UMR 6303), UNIVERSITÉ DE BOURGOGNE FRANCHE-COMTÉ, 9 AV. A. SAVARY, 21078 DIJON CEDEX, FRANCE.

Email address: jonas.lampart@u-bourgogne.fr