



**HAL**  
open science

## Assessing mutualistic metacommunity capacity by integrating spatial and interaction networks

Marc Ohlmann, François Munoz, François Massol, Wilfried Thuiller

### ► To cite this version:

Marc Ohlmann, François Munoz, François Massol, Wilfried Thuiller. Assessing mutualistic metacommunity capacity by integrating spatial and interaction networks. *Theoretical Population Biology*, 2024, 156, pp.22-39. 10.1016/j.tpb.2024.01.001 . hal-04729045

**HAL Id: hal-04729045**

**<https://hal.science/hal-04729045v1>**

Submitted on 9 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



<sup>21</sup> [//gitlab.com/marcohlmann/metacomunity\\_theory](https://gitlab.com/marcohlmann/metacomunity_theory)).

## 22 **Abstract**

23 We develop a spatially realistic model of mutualistic metacommunities that exploits the joint struc-  
24 ture of spatial and interaction networks. Assuming that all species have the same colonisation  
25 and extinction parameters, this model exhibits a sharp transition between stable non-null equilib-  
26 rium states and a global extinction state. This behaviour allows defining a threshold on coloni-  
27 sation/extinction parameters for the long-term metacommunity persistence. This threshold, the  
28 'metacommunity capacity', extends the metapopulation capacity concept and can be calculated  
29 from the spatial and interaction networks without needing to simulate the whole dynamics. In  
30 several applications we illustrate how the joint structure of the spatial and the interaction net-  
31 works affects metacommunity capacity. It results that a weakly modular spatial network and a  
32 power-law degree distribution of the interaction network provide the most favourable configuration  
33 for the long-term persistence of a mutualistic metacommunity. Our model that encodes several  
34 explicit ecological assumptions should pave the way for a larger exploration of spatially realistic  
35 metacommunity models involving multiple interaction types.

## 36 1 Introduction

37 A fundamental goal of predictive ecology is to forecast the dynamics of interacting species in a given  
38 region (Thuiller *et al.* 2013, Mouquet *et al.* 2015). Reaching such a goal has direct implications  
39 for biodiversity management and conservation and to anticipate or mitigate the effects of habitat  
40 destruction and global change on biodiversity.

41 Metapopulation models have long been used to characterise the dynamics of populations that  
42 can colonise, persist or go extinct in a given landscape configuration (Hanski & Ovaskainen 2003).  
43 This configuration is often summarised by a spatial network of suitable patches (Dale & Fortin  
44 2010; Hagen *et al.* 2012) that best represents habitat patchiness in both natural and human-altered  
45 ecosystems (Haddad *et al.* 2015). Levins (1969) devised a seminal model of species occupancy *i.e.*,  
46 the probability of presence of species populations across a landscape. In this model, a mean-field,  
47 deterministic differential equation model represented the population dynamics in fully connected  
48 patches, so that equilibrium occupancy depended on both a colonisation and an extinction pa-  
49 rameter. More than 30 years later, Etienne & Nagelkerke (2002) proposed a stochastic analogue  
50 of Levins' model and studied the links between the properties of the two models. Two sources  
51 of spatial heterogeneity can be embedded in metapopulation models: the heterogeneity on coloni-  
52 sation/extinction parameters among species (functional connectivity) and on the spatial network  
53 structure (structural connectivity) (Tischendorf & Fahrig 2000). The impact of structural con-  
54 nectivity on stationary occupancy (*e.g.*, Gilarranz & Bascompte 2012) underlines the influence of  
55 fragmentation on metapopulation persistence (Fahrig 2003, Fletcher Jr *et al.* 2018). Subsequent  
56 deterministic, spatially realistic models acknowledged variation of connectivity among nodes, and  
57 allowed quantifying analytically the viability of a metapopulation that depends on the mere struc-  
58 tural properties of the spatial network (Ovaskainen & Hanski 2001, Hanski & Ovaskainen 2003).  
59 The viability is defined through the metapopulation capacity, *i.e.*, a threshold on colonisation and  
60 extinction parameters above which the metapopulation can survive. This threshold is thus of prime  
61 importance in biological conservation (Groffman *et al.* 2006).

62 However, populations of a species are likely to interact with many other species within habitat

63 patches. These interactions should also affect the spatial coexistence of multiple metapopulations  
64 and their respective capacities (Thuiller *et al.* 2013). Metacommunity models are designed to assess  
65 the joint dynamics of multiple species in a habitat network (Leibold *et al.* 2004). While the structure  
66 of interaction networks is known to strongly influence biodiversity dynamics (Sole & Bascompte  
67 2007), most existing deterministic metacommunity models generally focused on global competition  
68 and competition-colonisation trade-off in fully connected patches (Tilman *et al.* 1997, Calcagno *et al.*  
69 2006), or sometimes in evenly connected patches (*e.g.*, lattice Amarasekare *et al.* 2004, Mouquet  
70 *et al.* 2011). Models focusing on other interaction types (*e.g.* facilitation, mutualistic and trophic)  
71 were developed for species-poor communities, homogeneous or lattice space (*i.e.* for few species  
72 Nee *et al.* 1997, Gravel & Massol 2020, homogeneous space Astegiano *et al.* 2015, lattice space Kéfi  
73 *et al.* 2007), preventing the study of complex networks and further generalisations.

74 Yet, stochastic models of interactions where species are either present or absent can encode  
75 mechanisms through specific rules, like having at least one prey to survive in the Trophic Theory  
76 of Island Biogeography (Gravel *et al.* 2011, Massol *et al.* 2017), or through increasing probability of  
77 presence depending on prey availability (Cazelles *et al.* 2016, Auclair *et al.* 2017). The latter model  
78 belongs to graphical models, a class of statistical models that represents conditional dependencies  
79 between species distributions using graphs. Using network-based metrics, these models can encode  
80 several mechanisms in terms of conditional probabilities of presence (Staniczenko *et al.* 2017).  
81 Nevertheless, these approaches still ignore the spatial structure of the environment.

82 So far, theoretical studies on the dynamics of metacommunities within a spatially explicit en-  
83 vironment and with biotic interactions have rarely considered how the dynamics jointly depend on  
84 graph properties of both interaction and spatial networks (*e.g.*, Amarasekare *et al.* 2004, but see  
85 Zhang *et al.* 2021), trophic interactions (Pillai *et al.* 2010, Brechtel *et al.* 2018, Gross *et al.* 2020  
86 but see Wang *et al.* 2021) or mutualistic interactions on a lattice (Filotas *et al.* 2010, Sardanyés  
87 *et al.* 2019). These models often elude the question of existence of a non-null equilibrium, and  
88 the metacommunity persistence is often assessed through tedious dynamic simulations or using  
89 strong approximations (Wang *et al.* 2021). If this approach provides points in the parameter space  
90 where the metacommunity persists, it neither maps regions of this space leading to persistence,

91 nor it demonstrates the existence of critical thresholds acting on metacommunity persistence as in  
92 metapopulation theory.

93 Interestingly, thresholds between local community persistence and extinction have already been  
94 identified in the case of positive interactions (Callaway 1997, Kéfi *et al.* 2016). For instance,  
95 mutualistic interactions play a major role in natural systems by conditioning coexistence (Valdovinos  
96 2019). Thébault & Fontaine (2010) showed that mutualistic networks generally have a nested  
97 architecture favouring persistence, and empirical surveys evidenced a truncated power-law of degree  
98 distribution (Bascompte & Jordano 2006, Vázquez *et al.* 2009, Bascompte 2009). Kéfi *et al.* 2007  
99 studied a metacommunity model with facilitation on a lattice space. However, no network-based  
100 model of spatially realistic, mutualistic metacommunities has been proposed so far. Such model  
101 should allow to test the joint impact of the structure of the spatial and interaction networks on the  
102 viability of a metacommunity and, potentially, allow to exhibit thresholds acting at the mutualistic  
103 metacommunity level. It should also reconcile the ongoing debate on the impact of the structure of  
104 the spatial network on metapopulations (Fletcher Jr *et al.* 2018).

105 In this paper, we explicitly model mutualistic interactions in an heterogeneous space using  
106 dynamic Bayesian networks (Auclair *et al.* 2017). We derive then a deterministic approximation  
107 and exhibit a threshold in metacommunity persistence assuming that all species have the same  
108 colonisation and extinction parameters. It defines an abrupt transition between stable coexistence  
109 and global metacommunity extinction. Our approach extends the computation of metapopula-  
110 tion capacity *sensu* Ovaskainen & Hanski to the case of mutualistic metacommunities with specific  
111 assumptions on colonisation and extinction functions. Using numerical methods, we show how  
112 metacommunity capacity relies on the structure of both mutualistic and spatial networks. Import-  
113 tantly, specific submodels can be derived to encode key ecological assumptions on extinction and  
114 colonisation. For these different ecological assumptions, we represent how spatial proximity of sites  
115 and mutualistic interactions modulate colonisation and/or extinction probability, and we compute  
116 metacommunity capacities. We finally explore the relationship between the degrees of the nodes of  
117 both spatial and interaction networks and species' occupancy at equilibrium. This allows extracting  
118 ecological relevant quantities on species among the sites (*e.g.*, mean occupancy) or in sites among

119 species (*e.g.*, species diversity, interaction network diversity). We thus quantify how metacommunity capacity is shaped by the joint structure of spatial and interaction networks.

121

## 122 2 Stochastic models of metacommunity dynamics using dynamic Bayesian networks

123

124 We first present a formalism that unifies stochastic spatially realistic metapopulation models and mainland-island models of biotic interactions in discrete time using Dynamic Bayesian Networks (DBNs). DBNs describe dependencies between random variables at different time steps through a bipartite directed graph, and represent stochastic models in which parameters are networks (Lähdesmäki & Shmulevich 2008, Koller & Friedman 2009). The network represents the influences between species distributions between two time steps. Once the structure of causal influences is fixed, several distributions can be associated to a given network structure through different parameterisations. These parameterisations represent interaction mechanisms that describe the effect of neighbour species or sites on the probability of presence of a given species at time  $t + 1$ . See Appendix for a more precise introduction on dynamic Bayesian networks and proof of the convergence of the different models.

135 The heterogeneous space is represented by a spatial network  $G_s = (V_s, E_s)$ , where  $V_s$  is the set of spatial vertices and  $E_s$  the set of spatial edges (linking unordered pairs of vertices). We assume that this network is undirected and connected, *i.e.*, considering two nodes  $u$  and  $v$  of  $G_s$ , there is always a path from  $u$  to  $v$ . Biotic interactions in the metacommunity are represented by an interaction network  $G_b = (V_b, E_b)$ , with  $V_b$  its set of vertices and  $E_b$  its set of edges, which we also assume undirected and connected. We note  $n = |V_s|$  and  $m = |V_b|$  (see Table 1 for notations).

141



Object	Name
$G_s$	Spatial network ( $n$ nodes)
$G_b$	Interaction network ( $m$ nodes)
$G_s^0$	Spatial network where edges have been deleted ( $n$ nodes)
$G_b^0$	Interaction network where edges have been deleted ( $m$ nodes)
$G_{s,b} = G_s \square G_b$	Cartesian product of the spatial and biotic interaction networks ( $n * m$ nodes)
$\mathbf{A}_s$	Adjacency matrix of the spatial network
$\mathbf{A}_b$	Adjacency matrix of the biotic interaction network
$\mathbf{A}_{s,b} = \mathbf{A}_s \otimes \mathbf{I}_m + \mathbf{I}_n \otimes \mathbf{A}_b$	Adjacency matrix of the Cartesian product network)
$G_c$	Colonisation network ( $n * m$ nodes)
$G_e$	Extinction network ( $n * m$ nodes)
$\mathbf{A}_c$	Adjacency matrix of the colonisation network
$\mathbf{A}_e$	Adjacency matrix of the extinction network
$\lambda_M$	Metacommunity persistence capacity
$\lambda_I$	Metacommunity invasion capacity
$\Lambda_s$	Dominant eigenvalue of the adjacency matrix of the spatial network
$\Lambda_b$	Dominant eigenvalue of the adjacency matrix of the biotic interaction network
$\Lambda_{s,b} = \Lambda_s + \Lambda_b$	Dominant eigenvalue of the adjacency matrix of the Cartesian product network

Table 1: Notations

## 142 2.1 Spatially realistic metapopulation model

143 We start by defining, using DBNs, a spatially realistic metapopulation model where populations  
144 of a single species colonise the spatial network  $G_s$ . Let  $X_i^t$  be a random variable associated to the  
145 presence of a population in a site  $i$  (*i.e.* the node  $v_i$  of  $G_s$ ) at time  $t$  ( $1 \leq i \leq n$ ,  $t \in \mathbb{N}^*$ , where  
146  $\mathbb{N}^*$  is the set of positive integers). We depict the dependency structure between the  $X_i^t$  using a  
147 DBN built from  $G_s$  (Fig. 1a). Defining the neighbours of  $v_i$  in  $G_s$  as  $N_s(i)$ , the parents of  $X_i^{t+1}$  in  
148 the DBN are  $\{X_i^t, \mathbf{X}_{N_s(i)}^t\}$ . This means that the presence of a population at time  $t + 1$  is causally  
149 influenced by the presence of a population at time  $t$  in site  $i$  and in sites adjacent to  $i$ . In this first  
150 model, no other variables or species influence the presence of a population in site  $i$  at time  $t + 1$ .  
151 Through conditional probabilities, the parameterisation encodes the way the presence or absence  
152 of a population in adjacent sites modulates the probability of presence of a population in the focal  
153 site. Here, we chose the same parameterisation as in [Gilarranz & Bascompte 2012](#).

$$\mathbb{P}(X_i^{t+1} = 1 | X_i^t, \mathbf{X}_{N_s(i)}^t) = (1 - (1 - c)^{\sum_{k \in N_s(i)} X_k^t})(1 - X_i^t) + (1 - e)X_i^t \quad (1)$$

154 where  $c$  and  $e$  are the respective colonisation ( $0 < c < 1$ ) and extinction ( $0 < e < 1$ ) parameters. In  
155 Eq. 1, the probability of presence grows with the number of occupied adjacent sites. Specifically,  
156 the probability that node  $i$  includes a population at time  $t + 1$  is  $1 - e$  if it had one at time  $t$ ,  
157 while the probability that node  $i$  is colonised between time  $t$  and time  $t + 1$  is equal to 1 minus  
158 the probability that all occupied neighbouring sites do not colonise node  $i$ , which happens with  
159 probability  $1 - c$  independently for each of these nodes.

160

## 161 2.2 A mainland-island model with biotic interactions

162 In this section, we present, using DBNs, a mainland-island model of species community where dif-  
163 ferent species colonise an island without any spatial structure but with a biotic interaction network  
164  $G_b$ .

165 Let  $X_j^t$  be the random variable associated to the presence of population of species  $j$  on the island. A  
166 DBN representing the dependency structure is built from  $G_b$  (Fig. 1a). Here, the DBN represents  
167 the network of species interactions as interactions affect colonisation and extinction probabilities  
168 on the island. Defining as  $N_{G_b}(j)$  the neighbours of  $v_j$  in  $G_b$ , the parents of  $X_j^{(t+1)}$  in the DBN are  
169  $\{X_j^t, \mathbf{X}_{N_{G_b}(j)}^t\}$ , meaning that the presence of species  $v_j$  and species that interact with  $v_j$  at time  $t$   
170 on the island, causally influences the presence of species  $v_j$  at time  $t + 1$ . Importantly, there is no  
171 other variables influencing the presence of a species  $v_j$  at time  $t + 1$ . We chose a parameterisation  
172 similar to [Auclair \*et al.\* 2017](#):

$$\mathbb{P}(X_j^{t+1} = 1 | X_j^t, \mathbf{X}_{N_{G_b}(j)}^t) = c(1 - X_j^t) + (1 - e(1 - \frac{\sum_{k \in N_{G_b}(j)} X_k^t}{1 + deg_{G_b}(j)}))X_j^t \quad (2)$$

173 where  $deg_{G_b}(j)$  is the degree of  $j$  in  $G_b$ . The probability of extinction (defined by Eq. 2) belongs  
174 to  $]0, 1[$  (Appendix). Although the dependency between species occurrences can encode any kind  
175 of interactions, we here focus on the mutualistic case by imposing an extinction function. In this  
176 case, the probability of extinction of a given species decreases with the number of species present  
177 that interact with the focal species.

## 179 2.3 Spatially realistic models of mutualistic metacommunities

180 Integrating the models from Section 2.1 and 2.2, we built a spatially explicit metacommunity model.  
 181 In this model, several species, interacting through  $G_b$ , are colonising the spatial network  $G_s$ . The  
 182 colonisation and extinction probabilities of population of a given species in a site are affected by  
 183 the presence of interacting species in the same site and presence of population of focal species in  
 184 neighbour sites. To do so, we used the Cartesian product of graphs that builds a network from  $G_b$   
 185 and  $G_s$  (Imrich & Klavzar 2000).

186 **Definition 1.** *The Cartesian product of  $G_s$  and  $G_b$ ,  $G_{s,b} = G_s \square G_b$  is the graph in which the set of*  
 187 *nodes is  $V_s \times V_b$ . A node of this graph is identified by a pair of nodes of  $G_s$  and  $G_b$ . Moreover, there*  
 188 *is an edge between  $(u_s, u_b)$  and  $(v_s, v_b)$  if  $(u_s = v_s$  and  $(u_b, v_b) \in E_b)$  or  $(u_b = v_b$  and  $(u_s, v_s) \in E_s)$ .*  
 189 *The first condition corresponds to the case where the two species are present at the same location*  
 190 *and interact with one another; the second condition, to the case where only one species is considered*  
 191 *and the two locations are linked by a spatial edge.*

192 The adjacency matrix,  $A_{s,b}$ , of  $G_{s,b}$  is

$$\mathbf{A}_{s,b} = \mathbf{A}_s \otimes \mathbf{I}_m + \mathbf{I}_n \otimes \mathbf{A}_b \quad (3)$$

193 where  $\mathbf{I}_m$  and  $\mathbf{I}_n$  denotes the identity matrices of dimension  $m$  and  $n$  and  $\otimes$  denotes the Kronecker  
 194 product of two matrices.

195 Let  $X_{ij}^t$  be the random variable associated to the presence of a population of species  $j$  in site  $i$   
 196 at time  $t$ . The dependency structure between the  $X_{ij}^t$  is depicted using a DBN that is built from  
 197  $G_{s,b}$  (Fig. 1a). Defining as  $N(i, j)$  the neighbours of  $(v_i, v_j)$  in  $G_{s,b}$ , the parents of  $X_{ij}^{t+1}$  in the  
 198 DBN are  $\{X_{ij}^t, \mathbf{X}_{N(i,j)}^t\}$ . This means that the presence of a population of species  $j$  in site  $i$  at time  
 199  $t + 1$  is causally influenced by the presence of population of the same species in adjacent sites at  
 200 time  $t$  and by the presence of populations of species that interact with  $j$  in the same site.

201 At this stage, it is crucial to define several submodels that formalise key ecological assumptions in




Figure 1: (a) Metapopulation model, mainland-island interaction model and metacommunity model. The second column represents the network associated to each model (spatial, interaction and product network). The third column represents the dynamic Bayesian network associated to each model that represents the causal influences of variables (presence of populations of a given species, species on the island, species in sites) at  $t$  on variables at  $t + 1$

(b) Simulating a dynamic in the combined effect model between two time steps. The nodes of the product network are either empty or occupied (grey: occupied, white: empty). For the sake of simplicity, the model here is turned deterministic ( $c = 1, e = 1$ ). To colonise a new node of the product network, species  $A$  and  $B$  must be both present in the same site and can colonise adjacent site only. The population of species  $B$  originally present in site  $c$  goes extinct since it does not co-occur with  $A$  at  $t$  whereas species  $A$  and  $B$  that co-occur in site  $a$  colonise the site  $b$ .

202 the product graph, using either the spatial network or the biotic interaction network to modulate  
 203 colonisation and extinction probability.

204 Let  $G_s^0$  be the network that has the same set of nodes as  $G_s$  but an empty set of edges, and let  
 205  $G_b^0$  be the network that has the same set of nodes as  $G_b$  but an empty set of edges. We introduce  
 206 then the colonisation network  $G_c$  ( $\mathbf{A}_c$  is its adjacency matrix) and the extinction network  $G_e$  ( $\mathbf{A}_e$   
 207 is its adjacency matrix). These networks modulate the colonisation and extinction probability in  
 208 the different submodels. We build two submodels from a given product graph (Fig. 2) :

- 209 • a Levins type submodel, where both the spatial and biotic interaction networks modulate the  
 210 colonisation probability ( $G_c = G_s \square G_b$ ), while the extinction probability is constant ( $G_e =$   
 211  $G_s^0 \square G_b^0$ )
- 212 • a combined effect submodel, where both the spatial and the biotic interaction networks modu-  
 213 late the colonisation probability ( $G_c = G_s \square G_b$ ), and the biotic interaction network modulates  
 214 the extinction probability ( $G_e = G_s^0 \square G_b$ )

215 For the two submodels, the conditional probabilities of colonisation and non-extinction are expressed  
 216 as:

$$\mathbb{P}(X_{ij}^{t+1} = 1 | X_{ij}^t = 0, \sum_{(k,l) \in N_{G_c}(i,j)} X_{kl}^t) = \epsilon + (1 - \epsilon) \left[ 1 - (1 - c)^{\sum_{(k,l) \in N_{G_c}(i,j)} X_{kl}^t} \right] \quad (4)$$

$$\mathbb{P}(X_{ij}^{t+1} = 1 | X_{ij}^t = 1, \sum_{(k,l) \in N_{G_e}(i,j)} X_{kl}^t) = 1 - e \left( 1 - \frac{\sum_{(k,l) \in N_{G_e}(i,j)} X_{kl}^t}{1 + \text{deg}_{G_e}((i,j))} \right) \quad (5)$$

218 where  $\epsilon \in ]0; 1[$  is a constant that guarantees the convergence of the model. This constant allows  
 219 colonisation from an external source, analogous to nodal self-infection in the epidemiology literature  
 220 (Van Mieghem & Cator 2012). The proposed metacommunity model is analogous to the open Levins  
 221 model, that better fits with data than the classic Levins model (Laroche *et al.* 2018).  $\text{deg}_{G_e}((i,j))$   
 222 is the degree of  $(v_i, v_j)$  in  $G_e$ , and  $N_{G_c}(i,j)$  (resp.  $N_{G_e}(i,j)$ ) denotes the neighbours of  $(v_i, v_j)$  in  
 223  $G_c$  (resp.  $G_e$ ). Fig. 1b shows a simplistic dynamics in the combined effect model. Computing the  
 224 stationary distribution is also intractable in the general case (since transition matrix is of dimension  
 225  $2^{nm}$ ), but, it is however possible to simulate the dynamics of the metacommunity as Gilarranz &

226 Bascompte (2012) did for metapopulation model. The code to sample in the stochastic model is  
 227 available on the gitlab repository ([https://gitlab.com/marcohlmann/metacommunity\\_theory](https://gitlab.com/marcohlmann/metacommunity_theory)).

### 228 3 The $nm$ -intertwined model

229 Since studying the stochastic model of Section 2.3 is intractable in the general case, we propose to  
 230 study deterministic models that approximate the stochastic models, referred to as the intertwined  
 231 model in the epidemiology literature (Van Mieghem 2011). We extended the spatially realistic  
 232 Levins model to a metacommunity model based on the product of spatial and interaction network  
 233 (Ovaskainen & Hanski 2001). The approximation is derived from Van Mieghem (2011) and Bianconi  
 234 (2018). The aim is to study the dynamics of mean occupancy of each species  $j$  in each site  $i$ , *i.e.*  
 235  $p_{ij}(t) = \mathbb{E}(X_{ij}^t) = \mathbb{P}(X_{ij}^t = 1)$  where  $\mathbb{E}(\cdot)$  denotes the expected value. For all  $i$  and  $j$ :

$$p_{ij}(t+1) = \mathbb{E}((1 - X_{ij}^t)(\epsilon + (1 - \epsilon)(1 - (1 - c)^{\sum_{(k,l) \in N_{G_c}(i,j)} X_{kl}^t}))) + \mathbb{E}((1 - e(1 - e)^{\sum_{(k,l) \in N_{G_e}(i,j)} X_{kl}^t})X_{ij}^t)) \quad (6)$$

236 Eq. 6 leads to a hierarchy of equations that cannot be solved (*i.e.* we need to consider  $\mathbb{E}(X_{1,1}^t, \dots, X_{m,n}^t)$ )  
 237 to find a solution to the system). In order to get moment closure, we assume that site occupancies  
 238 are independent. More precisely, for any sequence of indices  $n(1), n'(1); \dots, n(r), n'(r')$  :

$$\mathbb{E}(X_{n(1),n'(1)}^t, \dots, X_{n(r),n'(r')}^t) \simeq \mathbb{E}(X_{n(1),n'(1)}^t) \dots \mathbb{E}(X_{n(r),n'(r')}^t) \quad (7)$$

239 After some algebra, introducing a new single index  $v$  for the nodes of the product network and  
 240 assuming that  $c \ll 1, e \ll 1$  and  $\epsilon \ll c$  (see Appendix), it follows :

$$p_v(t+1) - p_v(t) = C_v(\mathbf{p}(t))(1 - p_v(t)) - E_v(\mathbf{p}(t))(p_v(t)) \quad (8)$$

241 where  $C_v(\mathbf{p}(t)) = c \sum_u [A_c]_{v,u} p_u(t)$  and  $E_v(\mathbf{p}(t)) = e(1 - \sum_u [A_e]_{v,u} p_u(t)/M_u)$  with  $M_u = 1 +$   
 242  $\text{deg}_{G_e}(u)$

243 This rewrites:

$$\mathbf{p}(\mathbf{t} + \mathbf{1}) - \mathbf{p}(\mathbf{t}) = c(\mathbf{A}_c \mathbf{p}(\mathbf{t})) \odot (\mathbf{1} - \mathbf{p}(\mathbf{t})) - e(\mathbf{1} - (\mathbf{D}_e + \mathbf{I}_{nm})^{-1} \mathbf{A}_e \mathbf{p}(\mathbf{t})) \odot \mathbf{p}(\mathbf{t}) \quad (9)$$

244 where  $\odot$  denotes the element-wise product,  $\mathbf{D}_e$  denotes the in-degree matrix of  $G_e$  and  $\mathbf{I}_{nm}$  denotes  
245 the identity matrix of dimension  $nm$ .

246 Eq. 8 is analogous to master equation of [Ovaskainen & Hanski \(2001\)](#). Now, to assess the viability  
247 of a given mutualistic metacommunity, we need to determine the equilibrium states and evaluate  
248 their local stability in function of  $c$  and  $e$  parameters.

### 249 3.1 Metapopulation capacity

250 In these spatially realistic metapopulation models, equilibrium state is either stable coexistence (all  
251 sites have non-null occupancy) or global extinction (all patches have null occupancy). Metapopula-  
252 tion capacities have thus been derived to assess both the persistence and the stability of metapopula-  
253 tions at equilibrium ([Hanski & Ovaskainen 2000](#), [Ovaskainen & Hanski 2001](#)). The metapopulation  
254 persistence capacity  $\lambda_M$  is a threshold between coexistence and global extinction (depending on  
255 the colonisation and extinction parameters), computable from the spatial network. Importantly,  
256 in spatially realistic metapopulation models, as soon as a population is present in a site, it can  
257 colonise gradually the entire spatial network.

258 More formally, in the metapopulation case,  $G_b$  is made of a single node, ( $m = 1$ ) and we assume  
259 that  $G_s$  is undirected and connected. We have:

$$\forall t \in \mathbb{N}^*, \mathbf{p}(\mathbf{t}) \in \bar{\Omega} = \{x \in \mathbb{R}^n, \forall i, 0 \leq x_i \leq 1\} \quad (10)$$

260 with the following assumptions on the colonisation functions (per site  $i$ ),  $C_i(\cdot)$ , and extinction  
261 functions,  $E_i(\cdot)$ :

- 262 • there is no external source of migrants

$$C_i(\mathbf{0}) = 0 \quad (11)$$

- 263 • the occupied sites make a positive contribution to the colonisation function of an empty site

$$\forall \mathbf{p} \in \Omega = \{x \in \mathbb{R}^n, \forall i, 0 < x_i < 1\}, C_i(\mathbf{p}) > 0 \quad (12)$$

264

$$\begin{cases} \frac{\partial C_i}{\partial p_j}(\mathbf{p}) \geq 0 & \text{for } i \neq j \\ \frac{\partial C_i}{\partial p_i}(\mathbf{p}) = 0 \end{cases} \quad (13)$$

- 265 • there is no mainland population, extinction rates are positive and, eventually, reduced by the  
266 presence of local populations

$$\forall p \in \bar{\Omega}, E_i(\mathbf{p}) > 0 \quad (14)$$

$$\begin{cases} \frac{\partial E_i}{\partial p_j} \leq 0 & \text{for } i \neq j \\ \frac{\partial E_i}{\partial p_i} = 0 \end{cases} \quad (15)$$

- 267 • Colonisation and extinction functions are smooth functions

$$C_i \in \mathcal{C}^1(\bar{\Omega}) \quad (16)$$

268

$$E_i \in \mathcal{C}^1(\bar{\Omega}) \quad (17)$$

269 Let:

$$g_i(\mathbf{p}) = \frac{eC_i(\mathbf{p})}{cE_i(\mathbf{p})} \quad (18)$$



270 The model is also assumed to be irreducible. Let  $\mathbf{J}$  be the matrix of dimension  $n \times n$  so that:

271

$$J_{ij} = \begin{cases} 1 & \text{if } \frac{\partial g_i}{\partial p_j}(\mathbf{p}) > 0, \mathbf{p} \in \bar{\Omega} \\ 0 & \text{otherwise} \end{cases} \quad (19)$$

272 We say that the model is irreducible if  $\mathbf{J}$  is irreducible, i.e., the graph that has  $\mathbf{J}$  as adjacency  
273 matrix is strongly connected.

274 In the case of the spatially realistic Levins model :

275 •  $C_i(\mathbf{p}) = c(\mathbf{A}_s \mathbf{p})_i$

276 •  $E(\mathbf{p}) = e$

277 where  $\mathbf{A}_s$  is the adjacency matrix of the spatial network. Then,

$$g_i(\mathbf{p}) = (\mathbf{A}_s \mathbf{p})_i \quad (20)$$

278 and the model is irreducible since  $\mathbf{A}_s$  is irreducible.

279 The metapopulation invasion capacity,  $\lambda_I$ , is defined as the dominant eigenvalue of the Jacobian  
280 matrix of  $g$  evaluated in  $\mathbf{p} = 0$ . It measures the stability of the equilibrium  $\mathbf{p} = 0$  that is the ability  
281 of a single population to invade the spatial network.

282

**Definition 2.** *The metapopulation persistence capacity,  $\lambda_M$ , is defined as:*

$$\lambda_M = \sup_{\mathbf{p} \in \Omega} h(\mathbf{p})$$

where

$$h(\mathbf{p}) = \min_i h_i(\mathbf{p})$$

and

$$h_i(\mathbf{p}) = g_i(\mathbf{p}) \frac{1 - p_i}{p_i}$$

283 We now present a weak version of the main theorem of [Ovaskainen & Hanski \(2001\)](#).

284 **Theorem 1.** (*Ovaskainen & Hanski*) *The deterministic metapopulation model has a non-trivial*  
285 *non-negative equilibrium state if and only if the threshold condition  $\lambda_M > \frac{c}{c}$  (if all the components*  
286 *of  $\mathbf{g}$  are concave) or  $\lambda_M \geq \frac{c}{c}$  (otherwise) is satisfied. Moreover, if the threshold condition is*  
287 *satisfied, the non-trivial non-negative equilibrium state is unique if all components of  $g$  are concave*  
288 *and equilibria are ordered otherwise.*

289  $\lambda_M$  is a threshold on the colonisation/extinction parameters that allows the metapopulation  
290 to persist. Importantly, if the metapopulation persists, the equilibrium points are interior (they  
291 belong to  $\Omega$ ), meaning that all occupancies are strictly positive.

292 Moreover, if all the components of  $\mathbf{g}$  are concave (it is the case for the spatially realistic Levins  
293 model), we have:

$$\lambda_M = \lambda_I = \Lambda_s \tag{21}$$

294 where  $\Lambda_s$  is the dominant eigenvalue of  $\mathbf{A}_s$ . If one component (or more) of  $\mathbf{g}$  is not concave, then  
295  $\lambda_I < \lambda_M$

### 296 **3.2 Extension to mutualistic metacommunity capacity**

297 In order to extend the metapopulation concept to metacommunity concept, we first consider a  
298 case with no interaction. Assuming that all species have the same colonisation and extinction  
299 parameters, since they also share the same spatial network, they have then the same metapopulation  
300 capacity even if their have independent dynamics. However, in this case, we cannot define a  
301 metacommunity capacity since the different metapopulations have independent dynamics: if a  
302 species is initially absent from the metacommunity, it will never colonise it. In this section, we show  
303 that adding mutualistic interactions to the metacommunity tangle the dynamics of the different  
304 metapopulations and allows defining a single threshold controlling the extinction of the entire  
305 metacommunity. Importantly, we show that this metacommunity capacity is higher than individual  
306 metapopulation capacities.

307 **3.2.1 The mutualistic metacommunity concept**

308 We extend metapopulation capacities from Section 3.1 to mutualistic metacommunity capacities in  
 309 the dynamical system defined by Eq. 9, using the product of the spatial network and the biotic  
 310 interaction network and specific assumptions on colonisation and extinction functions. Importantly,  
 311 in our spatially realistic mutualistic metacommunity model, as soon as a population of a given  
 312 species is present in a site, it can colonise gradually the entire spatial network and populations of  
 313 partner species will also colonise the spatial network thanks to this focal species. As a consequence,  
 314 the proposed mutualistic metacommunity model presents a sharp transition between coexistence (all  
 315 species have non-null occupancy in all sites) and global extinction (all species have null occupancy  
 316 in all sites).

317 For this model, the state space is:  $\Omega = \{x \in \mathbb{R}^{n*m}, \forall v \in \{1, \dots, n * m\} 0 < x_v < 1\}$

318 We have:

$$C_v(\mathbf{p}(t)) = c \sum_u [A_c]_{v,u} p_u(t) \quad (22)$$

319 and

$$E_v(\mathbf{p}(t)) = e(1 - \sum_u [A_e]_{v,u} p_u(t) / M_u) \quad (23)$$

320 In order to apply theorem 1 to the product network, we first verify assumptions on colonisation and  
 321 extinction functions (notice that index  $v$  represents a combination of a site and a species index).

- 322 • there is no external source of migrants

$$C_v(\mathbf{0}) = 0 \quad (24)$$

323 Notice that this assumption is only verified at order 1

- 324 • species occupying sites make a positive contribution to the colonisation function of an empty  
 325 site

$$\forall \mathbf{p} \in \Omega, C_v(\mathbf{p}) > 0 \quad (25)$$

326

$$\begin{cases} \frac{\partial C_u}{\partial p_u}(\mathbf{p}) \geq 0 & \text{for } u \neq v \\ \frac{\partial C_u}{\partial p_v}(\mathbf{p}) = 0 \end{cases} \quad (26)$$

327

Importantly, in both Levins type and combined effect model, since  $G_c = G_s \square G_b$ , an empty

328

site can be colonised by a given species if this species is present in neighbor sites or if species

329

that interact with the focal species are present in this site. By doing so, even if a species is

330

initially absent from the metacommunity, it can colonise it thanks to partner species.

331

- there is no mainland population, extinction rates are positive and reduced by the presence of

332

others species

$$\forall p \in \bar{\Omega}, E_v(\mathbf{p}) > 0 \quad (27)$$

333

$$\begin{cases} \frac{\partial E_i}{\partial p_j} \leq 0 & \text{for } i \neq j \\ \frac{\partial E_i}{\partial p_i} = 0 \end{cases} \quad (28)$$

334

Notice that, due to this assumption, we stick to the modelling of mutualistic metacommunity.

335

- Colonisation and extinction are smooth functions

$$C_v \in \mathcal{C}^1(\bar{\Omega}) \quad (29)$$

336

$$E_v \in \mathcal{C}^1(\bar{\Omega}) \quad (30)$$

337

Additionally:

338

**Proposition 1.** *The Levins type model and the combined effect model are irreducible on  $\bar{\Omega}$*

339

See proof in Appendix.

340

We then define metacommunity invasion capacity as the dominant eigenvalue of the jacobian matrix

341

of  $g$  evaluated in  $\mathbf{p} = 0$ .

342

**Definition 3.** *The mutualistic metacommunity persistence capacity,  $\lambda_M$ , is defined as:*

$$\lambda_M = \sup_{\mathbf{p} \in \Omega} h(\mathbf{p})$$

where

$$h(\mathbf{p}) = \min_v h_v(\mathbf{p})$$

and

$$h_v(\mathbf{p}) = g_v(p) \frac{1 - p_v}{p_v}$$

343 By applying theorem 1, a non-trivial equilibrium that the dynamical system has a non trivial  
 344 equilibrium if and only if  $\lambda_M > \frac{e}{c}$ .  $\lambda_M$  is then a threshold on the colonisation/extinction param-  
 345 eters that allows the mutualistic metacommunity to persist. Importantly, a non-trivial equilibrium  
 346 point is interior (it belongs to  $\Omega$ ), so each species in each site has a positive abundance at equilibrium.  
 347

348 **Proposition 2.** *For the Levins type model,  $\lambda_M = \lambda_I = \Lambda_s + \Lambda_b$*

349 The Levins type model is actually the spatially realistic metapopulation model with  $G_c =$   
 350  $G_s \square G_b$  as spatial network (the extinction is constant equals to  $e$ ). The dominant eigenvalue,  $\Lambda_c$ ,  
 351 of  $A_c$  is  $\Lambda_s + \Lambda_b$ . Consequently, for the Levins type submodel:

$$\lambda_M = \lambda_I = \Lambda_s + \Lambda_b \tag{31}$$

352 For the combined effect model,  $\lambda_I = \Lambda_s + \Lambda_b$  (see Appendix for proof). Notice that in the Levins  
 353 type model, both the biotic interaction and the spatial networks play interchangeable roles.

### 354 3.2.2 Computation of metacommunity capacity for the combined effect model

355 For the combined effect model, we computed the metacommunity capacity  $\lambda_M$  using Appendix  
 356 D of [Ovaskainen & Hanski 2001](#) and simulating annealing. We propose an implementation in R  
 357 and Python available at: [https://gitlab.com/marcohlmann/metacommunity\\_theory](https://gitlab.com/marcohlmann/metacommunity_theory). Only the




Figure 2: Top: Map of the different models, submodels and their parameters. Bottom: The four submodels associated to mutualistic metacommunity models, their assumptions, colonisation/extinction networks and metacommunity capacities

358 metapopulation or the metacommunity persistence capacity is really the focus for assessing viability.  
 359 For the sake of simplicity, we will thus use metacommunity capacity as metacommunity persistence  
 360 capacity in the rest of the text (unless specified otherwise)

## 361 4 Applications

### 362 4.1 Illustration

363 To illustrate the metacommunity capacity concept, we built a toy model (Fig. 3). We used a circu-  
 364 lar spatial network with 4 nodes (Fig. 3a) and a star shaped interaction network made of 4 nodes  
 365 (Fig. 3b), which could represent a plant species and its mutualistic mycorrhizal fungi species. The  
 366 Cartesian product is built from the spatial and the interaction networks (Fig. 3c). For the illustra-  
 367 tion, we derived the Levins type submodel dynamics. In this case, both metacommunity invasion  
 368 capacity  $\lambda_I$  and persistence capacity  $\lambda_M$  are equal to the dominant eigenvalue of the product of  
 369 the networks (3.73).  $\lambda_M$  defines the feasibility domain that is the portion of space where all species

370 have a non-null occupancy (see [Song et al. \(2018\)](#)) (Fig. 3d). We showed two possible outcomes of  
371 species occupancy dynamics (Fig. 3). One had a combination of colonisation and extinction values  
372 allowing metacommunity persistence, while the other had values outside the feasibility domain and  
373 yielded metacommunity extinction. Occupancies of persisting species converge toward two different  
374 values due to symmetries in the product network. Despite its simplicity, this toy model shows  
375 that we can predict the outcome of mutualistic metacommunity dynamics for any location of the  
376 parameter space, depending on the metacommunity capacity.

## 377 4.2 Structures of spatial and mutualistic interaction network jointly shape 378 the metacommunity capacity

379 We applied our model to investigate how the structure of the spatial and interaction networks shape  
380 the metacommunity capacity of a bipartite mutualistic system. To simulate landscape fragmenta-  
381 tion, we sampled two types of spatial networks while keeping constant the expected number of edges.  
382 We generated random spatial networks with 10 nodes in either Erdős-Renyi graphs (all edges are  
383 independent and identically distributed, with connectance  $C = 0.25$ ) or modular graphs using a  
384 block model ( $C = 0.25$ , more details in Appendix). We only kept connected spatial networks and  
385 used 15 replicates for each type of spatial network. Concerning the mutualistic network, we sampled  
386 two types of bipartite networks while keeping constant the number of edges. We generated random  
387 interaction networks with 14 nodes and 16 edges in either Erdős-Renyi graphs or networks with  
388 degree distribution shaped as a power-law of scaling parameter equals to 2. We used the function  
389 `sample_fitness_pl` implemented in the R package `igraph` ([Csardi & Nepusz 2006](#)). We only kept  
390 connected interaction networks and used 15 replicates per type of interaction network. We then  
391 computed the colonisation and extinction networks for each combination of spatial and interaction  
392 networks, so generating  $4 * 4 * 15 = 900$  different networks in total. This number of replicates was  
393 large enough to generate robust results (see Appendix). We first computed the metacommunity  
394 capacities for each combination of spatial and interaction networks to assess the viability range of  
395 the metapopulations. Then, we choose  $c$  and  $e$  parameters so that the mutualistic metacommunity  
396 persists and compared the stochastic model with its  $nm$ -intertwined deterministic approximation.




Figure 3: Toy model built from (a) a circular spatial network, (b) a star-shaped interaction network giving (c) the product network. The product network defines (d) the persistence and extinction domain. (e) Two trajectories sampled in and outside the persistence domain leading to persistence or extinction of the metacommunity



397 We also studied how species occupancy at equilibrium and aggregated quantities build from these  
398 occupancies (mean occupancy, species diversity) depend on node characteristics of both networks  
399 in the deterministic model.

#### 400 4.2.1 Computing metacommunity capacities

401 We computed the metacommunity capacity  $\lambda_M$  for the Levins type and combined effect submodels  
402 with the four combinations of networks structure (Fig. 4, Fig. S3). Despite known concerns on  
403 the ability to fit power-laws on small networks (Clauset *et al.* 2009, Stumpf & Porter 2012), we  
404 were able to statistically distinguish estimation of metacommunity capacity for almost all sampled  
405 combinations of structures (cf. Appendix). For both the Levins type and combined effect submodel,  
406 the metacommunity capacity decreased when the spatial network was modular and when the degree  
407 distribution was not a power-law. In this case, the modularity of the spatial network had a stronger  
408 impact on the metacommunity capacity than the structure of the mutualistic interaction network.  
409 Metacommunity capacity values were similar for the combined effect and Levins type model.

#### 410 4.2.2 Comparison between the stochastic model and the $nm$ -intertwined model

411 We compared the output of the stochastic metacommunity models with the  $nm$ -intertwined model  
412 for a given network combination with colonisation and extinction parameters chosen so that the  
413 metacommunity persist. We set  $c = e = 0.05$ ,  $\epsilon = 0.0005$  and used a spatial network with modularity  
414 of 0.36 and a mutualistic network with a degree distribution sampled in a power-law with parameter  
415 2. We sampled 1000 trajectories on 1500 time steps in the Levins type and the combined effect  
416 model and compared the mean stationary local occupancies and total occupancy with the prediction  
417 of the  $nm$ -intertwined model. For the combined effect model, the  $nm$ -intertwined model provides  
418 a accurate approximation of total occupancy (sum of occupancies of all species in all sites) at  
419 equilibrium (Fig. 5a). The deterministic model also provides a reasonable approximation of local  
420 occupancies (occupancy of each species in each site) compared to the mean values computed from  
421 the stationary distribution built from the stochastic simulations (Fig. 5b). We show this comparison  
422 for the Levins type model in Appendix (Fig. S4). Both the total occupancy and local occupancies




Figure 4: Assessing metacommunity persistence capacity in function of the structure of the spatial network (Erdős-Rényi/Modular) and the structure of the interaction network (Erdős-Rényi/Power law) for the combined effect model

423 are higher at equilibrium with the combined effect model compared to the Levins type model.

#### 424 4.2.3 A focus on species occupancies at equilibrium for a given network combination

425 Using the same parameters than the previous section, we simulated metacommunity dynamics and  
 426 studied how the occupancy at equilibrium of each node of the product network depends on its  
 427 degree for the combined effect model (see Appendix for the Levins type model). Additionally, we  
 428 studied the mean occupancy of species across sites, plus species and link diversity in each site.

429 We represented the occupancy of the nodes of the product network (that is the colonisation




Figure 5: Comparison between the stochastic metacommunity model and the *nm*-intertwined model for the combined effect model. (a) Comparison of the mean total occupancy dynamics averaged over 1000 replicates (solid line, the standard deviation is represented in grey) with the prediction of the equilibrium by the *nm*-intertwined model (dashed line) (b) Comparison of the mean local occupancies in the stationary distribution of the stochastic metacommunity model with the predicted values by the *nm*-intertwined model

430 network in this combined effect model) in function of their degree (Fig. 6). The occupancy of  
 431 the nodes of the product network (indexed by a species and a site) increased with the degree  
 432 of the nodes. Moreover, in this submodel, at a fixed node degree of the product network, the  
 433 occupancy decreased with the ratio of the degree of the site over the degree of the node of the  
 434 product network. This means that nodes of the product network that combined a generalist species  
 435 with a low-connected site have a higher occupancy at equilibrium compared to nodes that combined  
 436 a specialist species with a highly connected site. We observed the same patterns for the Levins  
 437 type model with lower occupancies (Appendix). From the occupancies at equilibrium, we then

438 computed, species  $\alpha$ -diversities in each site using the framework developed in (Ohlmann *et al.*,  
439 2019) with  $\eta = 2$  (Fig. 6). We observed a positive relationship between species  $\alpha$ -diversity and the  
440 degree of the nodes of the spatial network (Fig. 6). Mirroring the analysis on the spatial network,  
441 we represented the mean occupancy among the sites (Fig. 6) and observed a positive relationship  
442 between mean occupancy of a species and its degree in the biotic interaction network. For the  
443 Levins type model, we observed similar patterns except lower occupancies and mean occupancies  
444 per species (Fig. S5).

## 445 5 Discussion

446 In this paper, we proposed a stochastic spatially explicit model of mutualistic metacommunities  
447 that depends on the structure of spatial and biotic interaction networks, using Dynamic Bayesian  
448 Networks and graph products. Our stochastic model is built by integrating a metapopulation and  
449 a mainland-island interaction model (where species colonise an island influenced by a known meta-  
450 network). Spatial and interaction networks can modulate colonisation and extinction probabilities  
451 depending on the mechanisms that are encoded in the model. We proposed two sub-models but  
452 we encourage the implementation of other parameterisations or even interaction type since the  
453 stochastic model is highly flexible. The proposed mainland-island model is analogous to the trophic  
454 theory of island biogeography (TTIB, Gravel *et al.* 2011, Massol *et al.* 2017). However, in the TTIB,  
455 the interaction network must be a directed acyclic graph contrary to our mainland-island model  
456 where any network, even empty, can be used. The TTIB represents trophic interaction as energy  
457 flow from basal to non-basal species at a given time step whereas our stochastic interaction model  
458 represents population dynamics between two time steps, allowing feedback loops. The downside of  
459 this flexibility is the complexity and high-dimensionality of our stochastic model. However, network  
460 symmetries can be used to perform exact dimension reduction as in epidemics model (Simon *et al.*  
461 2011).

462 In order to further investigate properties of our mutualistic metacommunity model, we did a  
463 deterministic approximation to obtain the  $nm$ -intertwined model, named in reference to epidemics




Figure 6: Simulating the dynamics for a given spatial and biotic interaction network with the combined effect model. (a) Colonisation network whose size of the nodes is proportional to their degree and colour indicates the occupancy at equilibrium (grey: low occupancy, black: high occupancy) (b) Spatial network whose size of the nodes is proportional to their degree and colour indicates the species  $\alpha$ -diversity at equilibrium (grey: low  $\alpha$ -diversity, black: high  $\alpha$ -diversity) (c) Mutualistic interaction network whose size of the nodes is proportional to their degree and colour indicates the mean occupancy across the sites at equilibrium (grey: low mean occupancy, black: high mean occupancy) (d) Relationship between the occupancy at equilibrium and the degree of the node of the product network. Each point of the relationship (corresponding to a node of the product graph) is coloured according to the ratio of the degree of the site in the spatial network over the degree of the focal node in the colonisation network (e) Relationship between the species  $\alpha$ -diversity at equilibrium and the degree of the sites in the spatial network (f) Relationship between the mean occupancy at equilibrium and the degree of the species in the biotic interaction network

464 model (Van Mieghem 2011). It allows to keep track of interaction and spatial network structure  
465 in the deterministic model, contrary to mean field approximation in lattice based metacommunity  
466 models (Kéfi *et al.* 2007). We assume that species occupancies vary independently and we showed,  
467 using simulations, that it provides a reasonable approximation of our metacommunity stochastic  
468 model (Fig. 5). However, this approximation holds as soon as the metacommunity is far from  
469 the extinction threshold. Otherwise, pairwise correlation between species occupancies must be  
470 considered (pair approximation, *e.g.*, Kéfi *et al.* 2007) or even higher order correlation structure  
471 (Hiebeler & Millett 2011, Wuyts & Sieber 2022).

472 Assuming that all species have the same colonisation and extinction parameters, our determin-  
473 istic metacommunity model showed a sharp transition between states where the metacommunity  
474 persisted (*i.e.*, all species have non-null occupancy in all sites), and a state where the entire meta-  
475 community went extinct (*i.e.*, all species have null occupancy in all sites). The transition depended  
476 on the structure of the interaction and spatial networks and on colonisation and extinction pa-  
477 rameters. We defined the metacommunity capacity, a scalar quantity depending on the structure  
478 of both networks, as a threshold on colonisation/extinction parameters governing persistence of  
479 interacting species, thus extending the single-species concept of metapopulation capacity (Hanski  
480 & Ovaskainen 2000, Ovaskainen & Hanski 2001) to a metacommunity context. Importantly, strong  
481 assumptions on colonisation and extinction functions lead to the threshold behaviour of our model.  
482 We assume that both spatial and interaction networks contribute to species colonisation. By doing  
483 so, even if a species is absent from the metacommunity, populations of this species can colonise  
484 sites where partner species are present. This guarantees not to have prior invariant (except  $\mathbf{0}$ ) in  
485 the model, as for deterministic metapopulation model and leads to the threshold behaviour of the  
486 metacommunity model. We consider that mutualistic interactions help implantation of new species.  
487 This assumption is supported by the existence of foundation species that helps the metacommuni-  
488 ties to settle down (*e.g.*, cushion plants, Reid & Lortie 2012). In our mutualistic model, presence  
489 of a foundation species in the metacommunity will lead to colonisation of the metacommunity by  
490 partner species.

491 We extended the framework of metapopulation capacity to the case of a mutualistic meta-

492 community with a critical extinction threshold that is the same for all species belonging to the  
493 metacommunity. Importantly, in this model, even in the absence of biotic interactions, all species  
494 have the same metapopulation capacity (since they have the same spatial network and colonisa-  
495 tion and extinction parameters) leading to a the same extinction threshold for all species (even if  
496 they have independant dynamics). Adding mutualistic interactions tangle the different metapop-  
497 ulation dynamics and increase metapopulation capacity (that becomes metacommunity capacity)  
498 thus strengthening the metacommunity in regard to extinction. This conclusion is specific to the  
499 deterministic model, while local extinctions are still possible in our stochastic model. We showed  
500 that spatial and interaction networks jointly determine the metacommunity capacity (Fig. 4, Fig.  
501 S3). In other words, any viability statement on a metacommunity (like classic metapopulation  
502 viability statements, *e.g.*, [Bulman et al. 2007](#)) should be done using both networks, although we  
503 should keep in mind that the perceived spatial grain (*i.e.* nodes of the spatial network) and coloni-  
504 sation/extinction parameters might differ among species. Metacommunity capacity has important  
505 implications for biodiversity management (*e.g.*, for metapopulations [Groffman et al. 2006](#)), since it  
506 helps conservationists to forecast and thus prevent crossing critical thresholds to metacommunity  
507 extinction when facing habitat destruction, pollution or other alteration. Despite appealing proper-  
508 ties, our deterministic mutualistic metacommunity model is ecologically unrealistic since all species  
509 have stricly positive occupancies at equilibrium (in case of metacommunity persistence) ignoring so  
510 the possibility of local extinction due to environmental constrains or demographic stochasticity.

511 Our model of mutualistic metacommunity showed a sharp state-transition. Such abrupt transi-  
512 tions are known for community with positive interactions along environmental gradients ([Callaway](#)  
513 [1997](#), [Kéfi et al. 2016](#)). We somehow extended these known results for mutualistic metacommunities.  
514 Mutualistic interactions tangle individual metapopulation dynamics and strengthen metacommuni-  
515 nity in regards of extinction and, thanks to the proposed framework, we are able to quantify the  
516 gain in viability. If we assume that species have different colonisation and extinction parameters  
517 (depending on the environment for example) or that new species cannot colonise the metacommuni-  
518 nity thanks to mutualistic partners, we can no more apply [Ovaskainen & Hanski \(2001\)](#). We might  
519 expect intermediate equilibrium states (*i.e.*, states where only a subset of species goes extinct). Can

520 we extend the framework for other types of interactions? The assumptions on extinction functions  
521 in our model cannot represent non-mutualistic interactions and thus prevent its extension to com-  
522 petitive or multitrophic metacommunities. Regarding competition, competitive exclusion models  
523 in communities (Chesson 2000) and metacommunities (Calcagno *et al.* 2006) can lead to several  
524 intermediate states between coexistence and extinction of the entire metacommunity. However,  
525 competitive interactions along environmental gradients can induce dependencies between species,  
526 entailing alternative stable states (Liautaud *et al.* 2019). In the classic Lotka-Volterra deterministic  
527 model, conditions on trophic interaction network can lead to states where some of the species goes  
528 extinct but not the entire community (Takeuchi 1996; Bunin 2017). Wang *et al.* (2021) proposed  
529 a two species extension of metapopulation capacity with trophic interaction. They consider the  
530 metapopulation capacity for the prey and the predator separately. By approximating equilibrium  
531 prey occupancy, they compute predator metapopulation capacity. They extend the results to food  
532 chain in a hierarchical way. Contrary to the proposed framework, they do not propose a meta-  
533 community capacity but rather a set of metapopulation capacity that depends on each other in  
534 hierarchical way. It could be extended towards a trophic metacommunity model in a more general  
535 framework in several ways (Gross *et al.* 2020). However, predicting the outcome of these models  
536 from parameters only still poses tough challenges (Gross *et al.* 2020). In particular, this makes it  
537 difficult to establish critical thresholds for conservation science for competitive and trophic meta-  
538 communities. Nevertheless, we doubt that a single threshold value governs the fate of many species  
539 engaged in several types of interaction with each others as we believe that threshold phenomena  
540 occur in multi-interactions metacommunity. Our model should pave the way for a better under-  
541 standing of properties of spatially realistic trophic and competitive metacommunity models.

## 542 Acknowledgements

543 We dedicate this article to the memory of Marc Ohlmann, who tragically died in a mountain  
544 accident in June 2023. This was the last article of his thesis and one on which he spent the most  
545 energy. We thank Fabien Laroche for insightful comments and references on metapopulation and



546 metacommunity models. This research was funded by the French Agence Nationale de la Recherche  
 547 (ANR) through the GlobNet (ANR-16-CE02-0009) and EcoNet (ANR-18-CE02-0010) projects and  
 548 from ‘Investissement d’Avenir’ grants managed by the ANR (Montane: ANR-10-LAB-56).

## 549 **A Appendix: details on the model and simulations and proofs**

### 550 **A.1 Stochastic models of metacommunity dynamics using dynamic Bayesian** 551 **networks**

#### 552 **A.1.1 Bayesian networks and dynamic Bayesian networks**

553 Given a set of  $n$  random variables  $(X_1, \dots, X_n)$  (we note  $I = \{1, \dots, n\}$ ),

**Definition 4.** *Two random variables  $X_i$  and  $X_j$  are independent conditionally given  $\mathbf{X}_{I \setminus \{i,j\}}$  iff:*

$$\mathbb{P}(X_i, X_j | \mathbf{X}_{I \setminus \{i,j\}}) = \mathbb{P}(X_i | \mathbf{X}_{I \setminus \{i,j\}}) \mathbb{P}(X_j | \mathbf{X}_{I \setminus \{i,j\}})$$

554 Bayesian networks aim to map conditional independence statements using a Directed Acyclic Graph  
 555  $G$  (DAG). For a given node  $u$ , we note  $Pa_u(G)$  the set of nodes that are parents of  $u$ .

$$Pa_u(G) = \{v \in V, (v, u) \in E\} \tag{32}$$

556 The joint probability  $\mathbb{P}(\mathbf{X})$  factorises over  $G$  as :

$$\mathbb{P}(X_1, \dots, X_n) = \prod_i \mathbb{P}(X_i | \mathbf{X}_{Pa_i(G)}) \tag{33}$$

557 The factorisation gives the independence conditional statement according to the structure of the  
 558 DAG.

559 A particular case of Bayesian network consists in Dynamic Bayesian Networks (DBNs). Indexing  
 560 our previous  $n$  random variables by time  $t$ , a DBN describes the homogeneous dependencies be-  
 561 tween  $\{X_1^t, \dots, X_n^t\}$  and  $\{X_1^{t+1}, \dots, X_n^{t+1}\}$  using a directed bipartite network  $G_{bip}$  (we note  $\mathbf{A}_{bip}$  its

562 adjacency matrix). Importantly, as the structure of  $G_{bip}$  does not depend on  $t$ , it can be built  
 563 using an aggregated network  $G$  (we note  $\mathbf{A}$  its adjacency matrix) and a graph  $P_2$  (we note  $\mathbf{A}_2$  its  
 564 adjacency matrix) whose set of nodes is  $\{t, t + 1\}$  and set of edges is  $\{(t, t + 1)\}$ . We have

$$\mathbf{A}_{bip} = \mathbf{A}_2 \otimes (\mathbf{A} + \mathbf{I}_n) \quad (34)$$

565 where  $\mathbf{I}_n$  denotes the identity matrices of dimension  $n$ . We set  $\tilde{\mathbf{A}} = \mathbf{A} + \mathbf{I}_n$  and denotes  $\tilde{G}$  the  
 566 associated graph. The joint probability factorizes over  $G_{bip}$ :

$$\mathbb{P}(X_1^{t+1}, \dots, X_n^{t+1} | X_1^t, \dots, X_n^t) = \prod_i \mathbb{P}(X_i^{t+1} | \mathbf{X}_{Pa_i(\tilde{G})}) \quad (35)$$

### 567 A.1.2 Convergence properties of the stochastic models

568 The spatially realistic metapopulation model is a homogeneous Markov chain on  $\chi = \{0, 1\}^n$ . A  
 569 state of the metapopulation is a binary vector of length  $n$  indicating whether each site is occupied  
 570 or not. The dimension of the transition matrix is  $2^n * 2^n$  and the probability of transition between  
 571 a state  $s_k = (x_1, \dots, x_n)$  and  $s_l = (\tilde{x}_1, \dots, \tilde{x}_n)$  is

$$P_{k,l} = \mathbb{P}(X_1^{t+1} = \tilde{x}_1, \dots, X_n^{t+1} = \tilde{x}_n | X_1^t = x_1, \dots, X_n^t = x_n) \quad (36)$$

572 By applying conditional independence statements, we get:

$$P_{k,l} = \prod_i \mathbb{P}(X_i^{t+1} = \tilde{x}_i | X_i^t = x_i, \mathbf{X}_{N_s(i)}^t = (x_{N_s(i)})) \quad (37)$$

573  $\mathbf{0}$  is an absorbing state of the model. However, the model will reach a quasi-stationary distribution  
 574 (see [Darroch & Seneta 1965](#)) before extinction which gives a distribution of all possible states of the  
 575 metapopulation among sites. Getting extinction time and quasi-stationary distribution require to  
 576 compute eigenvectors and eigenvalues of  $\mathbf{P}$  that are intractable in the general case since  $\mathbf{P}$  is high-  
 577 dimensional. Using the ‘‘sampling from the past’’ algorithm ([Aldous \*et al.\*, 1988](#)) is an alternative  
 578 option to estimate the quasi-stationary distributions and associated eigenvalues, see e.g. [Schreiber](#)

579 *et al.* (2023) for an application of this technique in an ecological context.

580 The mainland-island model of species interaction is a homogeneous Markov chain on  $\chi = \{0, 1\}^m$   
 581 with no absorbing state. A state of the mainland-island model of species interaction is a binary  
 582 vector of length  $m$ , representing the composition of the community. The dimension of the transition  
 583 matrix is  $2^m * 2^m$  and the probability of transition between a state  $s_k = (x_1, \dots, x_m)$  and  $s_l =$   
 584  $(\tilde{x}_1, \dots, \tilde{x}_m)$  is

$$P_{k,l} = \prod_j \mathbb{P}(X_j^{t+1} = \tilde{x}_j | X_j^t = x_j, X_{N_b(j)}^t = (x_{N_b(j)})) \quad (38)$$

585 The chain converges towards a unique stationary distribution, a distribution of probability over all  
 586 possible species communities. However, as in the metapopulation case, computing the stationary  
 587 distribution is intractable in the general case since  $\mathbf{P}$  is high-dimensional. To summarise, in the  
 588 metapopulation model, the spatial network acts on the probability of colonisation, whereas in the  
 589 interaction model, the biotic network acts on the probability of extinction.

590

591 **Proposition 3.** *The stochastic spatially realistic metacommunity model converges towards a unique*  
 592 *stationary distribution*

593 In the stochastic spatially realistic models of mutualistic metacommunities, the transition matrix  
 594 of the chain is of dimension  $2^{mn} * 2^{mn}$ , encoding the probability of transition between a state  
 595  $\mathbf{s}_k = (x_{11}, \dots, x_{mn})$  of the metacommunity and a state  $\mathbf{s}_l = (\tilde{x}_{11}, \dots, \tilde{x}_{mn})$ , where  $x_{ij} \in \{0, 1\}$   
 596 describes the presence of a population of species  $i$  in site  $j$ . We note  $\mathbf{P}$  the transition matrix, the  
 597 probability of transition between  $\mathbf{s}_k$  and  $\mathbf{s}_l$  is :

$$P_{k,l} = \prod_{i,j} \mathbb{P}(X_{ij}^{t+1} = \tilde{x}_{ij} | X_{11}^t = x_{11}, \dots, X_{mn}^t = x_{mn}) \quad (39)$$

598 Moreover, we have:

$$\mathbb{P}(X_{i,j}^{t+1} = 1 | X_{i,j}^t = 0, \sum_{(k,l) \in N_{G_c}(i,j)} X_{k,l}^t) = \epsilon + (1 - \epsilon) * (1 - (1 - c)^{\sum_{(k,l) \in N_{G_c}(i,j)} X_{k,l}^t}) \quad (40)$$

599 with  $\epsilon \in ]0; 1[$  and  $c \in ]0; 1[$ . We have:

$$\mathbb{P}(\epsilon < \epsilon + (1 - \epsilon) * (1 - (1 - c)^{\sum_{(k,l) \in N_{G_c}(i,j)} X_{k,l}^t} < 1) = 1 \quad (41)$$

600 Moreover:

$$\mathbb{P}(X_{i,j}^{t+1} = 1 | X_{i,j}^t = 1, \sum_{(k,l) \in N_{G_e}(i,j)} X_{k,l}^t) = 1 - e(1 - \frac{\sum_{(k,l) \in N_{G_e}(i,j)} X_{k,l}^t}{1 + \text{deg}_{G_e}((i,j))}) \quad (42)$$

601 where  $e \in ]0; 1[$ . We have then:

$$\frac{e}{1 + \text{deg}_{G_e}((i,j))} < e(1 - \frac{\sum_{(k,l) \in N_{G_e}(i,j)} X_{k,l}^t}{1 + \text{deg}_{G_e}((i,j))}) < e \quad (43)$$

602 The probability of extinction is in  $]0; 1[$ .

603 Consequently :

$$\forall i \in \{1, \dots, n\}, \forall j \in \{1, \dots, m\}, \mathbb{P}(X_{i,j}^{t+1} = x_{i,j} | X_{1,1}^t = x_{1,1}, \dots, X_{m,n}^t = x_{m,n}) > 0 \quad (44)$$

604 It follows that  $\mathbf{P}$  is irreducible and aperiodic and  $(\mathbf{X}^t)^t$  converges towards a unique stationary  
 605 distribution. Importantly, in the stationary distribution, each species in each sites has a non-nul  
 606 probability of presence.

## 607 A.2 The $nm$ -intertwined model

608 The approximation is derived from [Bianconi \(2018\)](#) and [Van Mieghem \(2011\)](#). The aim is to study  
 609 the dynamics of occupancy of each species  $j$  in each site  $i$ :  $p_{ij}(t) = \mathbb{E}(X_{ij}^t)$ . For all  $i$  and  $j$ , we  
 610 have

$$p_{i,j}(t+1) = \mathbb{E}((1 - X_{i,j}^t)(\epsilon + (1 - \epsilon)(1 - (1 - c)^{\sum_{(k,l) \in N_{G_c}(i,j)} X_{k,l}^t}))) + \mathbb{E}(X_{i,j}^t(1 - e(1 - \frac{\sum_{(k,l) \in N_{G_e}(i,j)} X_{k,l}^t}{1 + \text{deg}_{G_e}((i,j))}))) \quad (45)$$

611 This approach leads to a hierarchy of equations that cannot be solved (i.e. we need to consider

612  $\mathbb{E}(X_{1,1}^t, \dots, X_{m,n}^t)$  to find a solution to the system). A drastic approximation consists in the mean  
 613 field approximation, for any sequence of indices  $n(1), n'(1); \dots, n(r), n'(r')$ , we assume :

$$\mathbb{E}(X_{n(1),n'(1)}^t, \dots, X_{n(r),n'(r')}^t) \simeq \mathbb{E}(X_{n(1),n'(1)}^t) \dots \mathbb{E}(X_{n(r),n'(r')}^t) \quad (46)$$

614

$$p_{i,j}(t+1) = (1-p_{i,j}(t))(\epsilon + (1-\epsilon)(1-(1-c)^{\sum_{(k,l) \in N_{G_c}(i,j)} p_{k,l}(t)})) + (1-e(1 - \frac{\sum_{(k,l) \in N_{G_e}(i,j)} p_{k,l}(t)}{1 + \text{deg}_{G_e}((i,j))}))p_{i,j}(t) \quad (47)$$

615 We assume that  $c = o(1)$ ,  $e = o(1)$  and  $\epsilon = o(c)$ , a Taylor expansion at order 1 with set  $M_{i,j} :=$   
 616  $1 + \text{deg}_{G_e}((i,j))$  leads to:

$$p_{i,j}(t+1) = (1-p_{i,j}(t))(\epsilon + (1-\epsilon)(c \sum_{(k,l) \in N_{G_c}(i,j)} p_{k,l}(t) + (1-e+e \sum_{(k,l) \in N_{G_e}(i,j)} p_{k,l}(t))/M_{i,j}))p_{i,j}(t) \quad (48)$$

617

$$p_{i,j}(t+1) = (1-p_{i,j}(t))(c \sum_{(k,l) \in N_{G_c}(i,j)} p_{k,l}(t) + (1-e+e \sum_{(k,l) \in N_{G_e}(i,j)} p_{k,l}(t))/M_{i,j})p_{i,j}(t) \quad (49)$$

618 We introduce a single index  $v$  for the nodes of the product networks and get:

$$p_v(t+1) = (1-p_v(t))(c \sum_{u \in N_{G_c}(v)} p_u(t) + ((1-e+e \sum_{u \in N_{G_e}(v)} p_u(t))/M_v)p_v(t) \quad (50)$$

619

$$p_v(t+1) - p_v(t) = (1-p_v(t))(c \sum_{u \in N_{G_c}(v)} p_u(t) - e((1 - \sum_{u \in N_{G_e}(v)} p_u(t))/M_v)p_v(t) \quad (51)$$

$$p_v(t+1) - p_v(t) = C_v(\mathbf{p}(t))(1-p_v(t)) - E_v(\mathbf{p}(t))p_v(t) \quad (52)$$

620 where  $C_v(\mathbf{p}(t)) = c \sum_u [A_c]_{v,u} p_u(t)$  and  $E_v(\mathbf{p}(t)) = e(1 - \sum_u [A_e]_{v,u} p_u(t)/M_v)$ .

621

$$\mathbf{p}(t+1) - \mathbf{p}(t) = c(\mathbf{A}_c \mathbf{p}(t)) \odot (\mathbf{1} - \mathbf{p}(t)) - e(\mathbf{1} - (\mathbf{D}_e + \mathbf{I}_{nm})^{-1} \mathbf{A}_e \mathbf{p}(t)) \odot \mathbf{p}(t) \quad (53)$$

622 where  $\odot$  denotes the element-wise product,  $D_e$  denotes the indegree matrix of  $G_e$  and  $\mathbf{I}_{nm}$  the  
 623 identity matrix of dimension  $nm$ .

### 624 A.3 Proof of proposition 1

625 We need to show that the two submodels are irreducible. Let  $\mathbf{J}$  be the matrix of dimension  $n \times n$   
 626 so that:

$$627 \quad J_{ij} = \begin{cases} 1 & \text{if } \frac{\partial g_i}{\partial p_j}(\mathbf{p}) > 0, \mathbf{p} \in \bar{\Omega} \\ 0 & \text{otherwise} \end{cases} \quad (54)$$

628 We say that the model is irreducible if  $\mathbf{J}$  is irreducible, i.e., the graph that has  $\mathbf{J}$  as adjacency  
 629 matrix is strongly connected.

630 Importantly, as pointed out in [Smith \(2008\)](#), we need to show that the models are irreducible on  
 631  $\bar{\Omega}$ , that is the interior of the domain but also its boundary.

632 We have:

$$g_v(\mathbf{p}) = \frac{\sum_u [A_c]_{v,u} p_u}{1 - \sum_u [A_e]_{v,u} p_u / M_v} \quad (55)$$

633 We first note that  $g_v$  is defined on  $\bar{\Omega}$  since:

$$|\sum_u [A_e]_{v,u} p_u / M_v| \leq |\sum_u [A_e]_{v,u} / M_v| < 1 \quad (56)$$

634 We have:

$$\frac{\partial g_v}{\partial p_u}(\mathbf{p}) = \frac{[A_c]_{v,u} (1 - \sum_k [A_e]_{v,k} p_k / M_v) + ([A_e]_{v,u} / M_v) \sum_k [A_c]_{v,k} p_k}{(1 - \sum_k [A_e]_{v,k} p_k / M_v)^2} \quad (57)$$

635

636 For the Levins type model:

$$\frac{\partial g_v}{\partial p_u}(\mathbf{p}) = [A_c]_{v,u} \quad (58)$$

637 And since,  $G_s$  and  $G_b$  are both strongly connected and  $G_c = G_s \square G_b$ ,  $G_c$  is also strongly connected  
 638 and  $\mathbf{J}$  is irreducible on  $\bar{\Omega}$ .

639

640 For the combined effect model, we note that  $E(G_e) \subset E(G_c)$ . It follows that  $[A^c]_{v,u} = 0 \implies$

641  $[A^e]_{v,u} = 0$  and  $[A^e]_{v,u} = 1 \implies [A^c]_{v,u} = 1$ . We have then:

$$\frac{\partial g_v}{\partial p_u}(\mathbf{p}) = \frac{[A_c]_{v,u} + (\sum_k [A_c]_{v,k} [A_e]_{v,u} p_k / M_v - [A_c]_{v,u} [A_e]_{v,k} p_k / M_v)}{(1 - \sum_k [A_e]_{v,k} p_k / M_v)^2} \quad (59)$$

642 • If  $[A_c]_{v,u} = 0$ , then  $[A_e]_{v,u} = 0$  and, for all  $k$ ,  $[A_c]_{v,u} [A_e]_{v,k} - [A_c]_{v,k} [A_e]_{v,u} = 0$ . It follows  
643 that  $\frac{\partial g_v}{\partial p_u}(\mathbf{p}) = 0$

644 • If  $[A_c]_{v,u} = 1$  and  $[A_e]_{v,u} = 0$ , then:

$$\frac{\partial g_v}{\partial p_u}(\mathbf{p}) = 1 - \sum_k [A_e]_{v,k} p_k / M_v \quad (60)$$

645 and  $\frac{\partial g_v}{\partial p_u}(\mathbf{p}) > 0$

646 • If  $[A_c]_{v,u} = 1$  and  $[A_e]_{v,u} = 1$ , then:

$$\frac{\partial g_v}{\partial p_u}(\mathbf{p}) = \frac{1 + (\sum_k [A_c]_{v,k} - [A_e]_{v,k}) / p_k M_v}{(1 - \sum_k [A_e]_{v,k} p_k / M_v)^2} \quad (61)$$

647 Since  $E(G_e) \subset E(G_c)$ , we have  $(\sum_k [A_c]_{v,k} - [A_e]_{v,k}) > 0$  and  $\frac{\partial g_v}{\partial p_u}(\mathbf{p}) > 0$ .

648 Consequently, for the combined effect model, then non-zero elements of  $\mathbf{J}$  are the non-zero elements  
649 of  $\mathbf{A}_c$ . Since  $G_c = G_s \square G_b$  it follows that  $\mathbf{J}$  is irreducible.

#### 650 **A.4 Computation of $\lambda_I$**

651 As provided in the main text, for the Levins type submodel,  $\lambda_I = \lambda_M = \Lambda_s + \Lambda_b$ . We now compute  
652 the  $\lambda_I$  for the three other submodels.

653 We first compute the Jacobian matrix of  $\mathbf{p} \mapsto g(\mathbf{p})$  evaluated in  $\mathbf{p} = \mathbf{0}$ . We have

$$\frac{\partial g_v}{\partial p_u}(\mathbf{0}) = [A_c]_{v,u} \quad (62)$$

654  $\lambda_I$  is the dominant eigenvalue of  $\left( \frac{\partial g_v}{\partial p_u}(\mathbf{0}) \right)_{u,v}$

655 • Combined effect submodel

656 For this submodel,  $A_c = A_s \otimes I_m + I_n \otimes A_b$ , it follows  $\lambda_I = \Lambda_s$

## 657 A.5 Computation of $\lambda_M$

658 In order to compute  $\lambda_M$  for the combined effect submodel, the separated effect model and the  
659 rescue effect submodel where the components of  $\mathbf{g}$  are not concave, we used a simulated annealing  
660 algorithm. We used the result of the iterative procedure described in Appendix D of [Ovaskainen &  
661 Hanski 2001](#) as starting point.

662 The code to compute the metacommunity capacity in the different models is available at: [https:  
663 //gitlab.com/marcohlmann/metacommunity\\_theory](https://gitlab.com/marcohlmann/metacommunity_theory).

664 We assessed the performance of the method on the Levins type model on the simulated data, since  
665 we know analytically the metacommunity capacity in this case. We used 20000 time steps on the  
666 900 different networks for the two submodels. The maximum is not reached (Fig. S1a) but there  
667 is a strong correlation (0.955) between the estimated metacommunity capacities and the theoretical  
668 metacommunity capacities (Fig. S1b), allowing so comparison of the metacommunity capacities  
669 among the different network structures.

## 670 B Appendix: detail on the simulation

### 671 B.1 Spatial networks

In order to mimic fragmentation of the landscape, we sampled spatial networks (10 nodes) using  
Erdős-Renyi model and a block model. For the Erdős-Renyi model, the probability of connection  
was  $C = 0.25$  and we kept connected networks only. For the block model, we partitioned in two






Figure S1: (a) Distribution of the relative error in the estimation of the metacommunity capacity (b) Relation between the metacommunity persistence capacity computing using a simulating annealing algorithm and the theoretical metacommunity capacity for the Levins type submodel

groups of equal sizes,  $p$  and  $q$ , with a matrix of probability of connection,  $\mathbf{\Pi}$ , given by:

$$\begin{pmatrix} p & q \\ \left(\frac{7C}{4} & \frac{C}{4}\right) p \\ \left(\frac{C}{4} & \frac{7C}{4}\right) q \end{pmatrix}$$

672 where  $C = 0.25$ .

673 The overall probability of connection in the network is :

$$\mathbb{P}(i \leftrightarrow j) = \sum_{k \in \{p, q\}, l \in \{p, q\}} Pr(i \leftrightarrow j | i \in k, j \in l) Pr(i \in k) Pr(j \in l) \quad (63)$$

$$\mathbb{P}(i \leftrightarrow j) = \frac{1}{4} \left( \frac{7C}{4} + \frac{C}{4} + \frac{C}{4} + \frac{7C}{4} \right) \quad (64)$$

$$\mathbb{P}(i \leftrightarrow j) = C \quad (65)$$

674 So the expected value of connectance for all spatial networks is the same despite different  
 675 modularity values (Fig. S2).




Figure S2: Distribution of the modularity of the spatial networks over the 15 replicates for the Erdős-Renyi structure and the modular structure

## 676 B.2 Biotic interaction networks

677 We first generated random undirected network with various shapes of the degree distribution using  
 678 the function `sample_fitness_pl` implemented in the R package `igraph` (Csardi & Nepusz, 2006). We  
 679 generated Erdős-Renyi networks and networks with a degree distribution given by a power-law.  
 680 We only kept connected networks. On the random network  $G$  sampled ( $\mathbf{A}$  is its adjacency matrix),  
 681 we build a bipartite network  $G_{bip}$  with adjacency matrix  $\mathbf{A}_{bip}$  as:

$$\mathbf{A}_{bip} = \mathbf{A}_2 \otimes (\mathbf{A} + \mathbf{I}_n) \quad (66)$$

682 where  $\mathbf{A}_2$  is the adjacency matrix of an undirected graph made of two nodes and a single edge  
683 between these two nodes. By doing so, all the sampled undirected bipartite networks are strongly  
684 connected.

### 685 B.3 Results

686 We simulated the dynamic (as presented in the main text for the combined effect submodel) for the  
687 Levins type model (Fig. S5, Fig. S6).




Figure S3: Assessing metacommunity persistence capacity in function of the structure of the spatial network (Erdős-Renyi/Modular) and the structure of the interaction network (Erdős-Renyi/Power law) for the Levins type model




Figure S4: Comparison between the stochastic metacommunity model and the  $nm$ -intertwined model for the Levins type model. (a) Comparison of the mean total occupancy dynamics averaged over 1000 replicates (solid line, the standard deviation is represented in grey) with the prediction of the equilibrium by the  $nm$ -intertwined model (dashed line) (b) Comparison of the mean local occupancies in the stationary distribution of the stochastic metacommunity model with the predicted values by the  $nm$ -intertwined model

#### 688 B.4 Robustness of metacommunity capacity estimation

689 We analysed the robustness of the estimation of  $\lambda_M$  for the four different structures for each  
 690 submodel. We described the distribution of  $\lambda_M$  (225 samples per combination of structure for  
 691 each model) using a boxplot (Fig. S7). Moreover, we used a Tukey test to estimate the confidence  
 692 intervals of the difference in mean metacommunity capacity per pairs of structures (Fig. S8). For  
 693 the Levins type and combined effect model, all differences in mean  $\lambda_M$  were statistically different  
 694 of 0. For the separated effect and rescue effect model, difference in mean  $\lambda_M$  of PL/E-E/E (PL:  
 695 Power-Law, E: Erdős-Renyi, M: Modular) and PL/M-E/M were statistically not different from 0.




Figure S5: Simulating the dynamics for a given spatial and biotic interaction network with the Levins type model. (a) Colonisation network whose size of the nodes is proportional to their degree and colour indicates the occupancy at equilibrium (grey: low occupancy, black: high occupancy) (b) Spatial network whose size of the nodes is proportional to their degree and colour indicates the species  $\alpha$ -diversity at equilibrium (grey: low  $\alpha$ -diversity, black: high  $\alpha$ -diversity) (c) Mutualistic interaction network whose size of the nodes is proportional to their degree and colour indicates the mean occupancy across the sites at equilibrium (grey: low mean occupancy, black: high mean occupancy) (d) Relationship between the occupancy at equilibrium and the degree of the node of the product network. Each point of the relationship (corresponding to a node of the product graph) is coloured according to the ratio of the degree of the site in the spatial network over the degree of the focal node in the colonisation network (e) Relationship between the species  $\alpha$ -diversity at equilibrium and the degree of the sites in the spatial network (f) Relationship between the mean occupancy at equilibrium and the degree of the species in the biotic interaction network




Figure S6: Aggregated statistics from occupancy at equilibrium for the combined effect submodel in the spatial network and the biotic interaction network. (a) Spatial network whose size of the nodes is proportional to their degree and colour indicates the  $\alpha$ -diversity at equilibrium (grey: low  $\alpha$ -diversity, black: high  $\alpha$ -diversity). (b) Relationship between the  $\alpha$ -diversity at equilibrium and the degree of the sites in the spatial network. (c) Biotic interaction network whose size of the nodes is proportional to their degree and colour indicates the mean occupancy across the sites at equilibrium (grey: low  $\alpha$ -diversity, black: high  $\alpha$ -diversity). (d) Relationship between the mean occupancy at equilibrium and the degree of the species in the biotic interaction network.

696 It means that, for these two models, whatever the structure of the spatial network (Modular or  
 697 Erdős-Renyi), mean  $\lambda_M$  was comparable for a power-law or Erdős-Renyi biotic interaction network.




Figure S7: Boxplot representing distributions of  $\lambda_M$  for each combination of structure in the Levins type and the combined effect model. E: Erdős-Renyi, PL: Power-Law, M: Modular




Figure S8: Tukey plot representing the confidence intervals of the difference in mean metacommunity capacity per pairs of structures. E: Erdős-Renyi, PL: Power-Law, M: Modular

## 698 References

- 699 Aldous, D., Flannery, B. & Palacios, J.L. (1988). Two applications of urn processes the fringe  
700 analysis of search trees and the simulation of quasi-stationary distributions of markov chains.  
701 *Probability in the engineering and informational sciences* 2, 293–307.
- 702 Amarasekare, P., Hoopes, M.F., Mouquet, N. & Holyoak, M. (2004). Mechanisms of coexistence in  
703 competitive metacommunities. *The American Naturalist* 164, 310–326.
- 704 Astegiano, J., Guimarães Jr, P.R., Cheptou, P.O., Vidal, M.M., Mandai, C.Y., Ashworth, L. &  
705 Massol, F. (2015). Persistence of plants and pollinators in the face of habitat loss: insights from  
706 trait-based metacommunity models. In: *Advances in ecological research*. Elsevier vol. 53 pp.  
707 201–257.
- 708 Auclair, E., Peyrard, N. & Sabbadin, R. (2017). Labeled dbn learning with community struc-  
709 ture knowledge. In: *Joint european conference on machine learning and knowledge discovery in*  
710 *databases*. Springer pp. 158–174.
- 711 Bascompte, J. (2009). Mutualistic networks. *Frontiers in Ecology and the Environment* 7, 429–436.
- 712 Bascompte, J. & Jordano, P. (2006). The structure of plant-animal mutualistic networks. *Ecological*  
713 *networks: linking structure to dynamics in food webs*. Oxford University Press, Oxford, UK pp.  
714 143–159.
- 715 Bianconi, G. (2018). *Multilayer Networks: Structure and Function*. Oxford University Press; p.60-  
716 62.
- 717 Brechtel, A., Gramlich, P., Ritterskamp, D., Drossel, B. & Gross, T. (2018). Master stability  
718 functions reveal diffusion-driven pattern formation in networks. *Physical Review E* 97, 032307.
- 719 Bulman, C.R., Wilson, R.J., Holt, A.R., Bravo, L.G., Early, R.I., Warren, M.S. & Thomas, C.D.  
720 (2007). Minimum viable metapopulation size, extinction debt, and the conservation of a declining  
721 species. *Ecological Applications* 17, 1460–1473.



- 722 Bunin, G. (2017). Ecological communities with Lotka-Volterra dynamics. *Physical Review E* 95,  
723 042414.
- 724 Calcagno, V., Mouquet, N., Jarne, P. & David, P. (2006). Coexistence in a metacommunity: the  
725 competition–colonization trade-off is not dead. *Ecology letters* 9, 897–907.
- 726 Callaway, R.M. (1997). Positive interactions in plant communities and the individualistic-continuum  
727 concept. *Oecologia* 112, 143–149.
- 728 Cazelles, K., Mouquet, N., Mouillot, D. & Gravel, D. (2016). On the integration of biotic interaction  
729 and environmental constraints at the biogeographical scale. *Ecography* 39, 921–931.
- 730 Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review of Ecology and*  
731 *Systematics* 31, 343–366.
- 732 Clauset, A., Shalizi, C.R. & Newman, M.E. (2009). Power-law distributions in empirical data.  
733 *SIAM review* 51, 661–703.
- 734 Csardi, G. & Nepusz, T. (2006). The igraph software package for complex network research.  
735 *InterJournal Complex Systems*, 1695.
- 736 Dale, M. & Fortin, M.J. (2010). From graphs to spatial graphs. *Annual Review of Ecology, Evolu-*  
737 *tion, and Systematics* 41.
- 738 Darroch, J.N. & Seneta, E. (1965). On quasi-stationary distributions in absorbing discrete-time  
739 finite markov chains. *Journal of Applied Probability* 2, 88–100.
- 740 Etienne, R.S. & Nagelkerke, C. (2002). Non-equilibria in small metapopulations: comparing the  
741 deterministic levins model with its stochastic counterpart. *Journal of Theoretical Biology* 219,  
742 463–478.
- 743 Fahrig, L. (2003). Effects of habitat fragmentation on biodiversity. *Annual review of ecology,*  
744 *evolution, and systematics* 34, 487–515.

- 745 Filotas, E., Grant, M., Parrott, L. & Rikvold, P.A. (2010). The effect of positive interactions on  
746 community structure in a multi-species metacommunity model along an environmental gradient.  
747 *Ecological Modelling* 221, 885–894.
- 748 Fletcher Jr, R.J., Didham, R.K., Banks-Leite, C., Barlow, J., Ewers, R.M., Rosindell, J., Holt,  
749 R.D., Gonzalez, A., Pardini, R., Damschen, E.I. *et al.* (2018). Is habitat fragmentation good for  
750 biodiversity? *Biological conservation* 226, 9–15.
- 751 Gilarranz, L.J. & Bascompte, J. (2012). Spatial network structure and metapopulation persistence.  
752 *Journal of Theoretical Biology* 297, 11–16.
- 753 Gravel, D. & Massol, F. (2020). Toward a general theory of metacommunity ecology. In: *Theoretical*  
754 *Ecology*. Oxford University Press.
- 755 Gravel, D., Massol, F., Canard, E., Mouillot, D. & Mouquet, N. (2011). Trophic theory of island  
756 biogeography. *Ecology Letters* 14, 1010–1016.
- 757 Groffman, P.M., Baron, J.S., Blett, T., Gold, A.J., Goodman, I., Gunderson, L.H., Levinson, B.M.,  
758 Palmer, M.A., Paerl, H.W., Peterson, G.D. *et al.* (2006). Ecological thresholds: the key to  
759 successful environmental management or an important concept with no practical application?  
760 *Ecosystems* 9, 1–13.
- 761 Gross, T., Allhoff, K.T., Blasius, B., Brose, U., Drossel, B., Fahimipour, A.K., Guill, C., Yeakel,  
762 J.D. & Zeng, F. (2020). Modern models of trophic meta-communities. *Philosophical Transactions*  
763 *of the Royal Society B* 375, 20190455.
- 764 Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A., Holt, R.D., Lovejoy, T.E.,  
765 Sexton, J.O., Austin, M.P., Collins, C.D. *et al.* (2015). Habitat fragmentation and its lasting  
766 impact on earth’s ecosystems. *Science advances* 1, e1500052.
- 767 Hagen, M., Kissling, W.D., Rasmussen, C., De Aguiar, M.A., Brown, L.E., Carstensen, D.W.,  
768 Alves-Dos-Santos, I., Dupont, Y.L., Edwards, F.K., Genini, J. *et al.* (2012). Biodiversity, species  
769 interactions and ecological networks in a fragmented world. 46, 89–210.

- 770 Hanski, I. & Ovaskainen, O. (2000). The metapopulation capacity of a fragmented landscape.  
771 *Nature* 404, 755.
- 772 Hanski, I. & Ovaskainen, O. (2003). Metapopulation theory for fragmented landscapes. *Theoretical*  
773 *population biology* 64, 119–127.
- 774 Hiebeler, D.E. & Millett, N.E. (2011). Pair and triplet approximation of a spatial lattice popula-  
775 tion model with multiscale dispersal using markov chains for estimating spatial autocorrelation.  
776 *Journal of theoretical biology* 279, 74–82.
- 777 Imrich, W. & Klavzar, S. (2000). *Product graphs: structure and recognition*. Wiley p.27.
- 778 Kéfi, S., Holmgren, M. & Scheffer, M. (2016). When can positive interactions cause alternative  
779 stable states in ecosystems? *Functional Ecology* 30, 88–97.
- 780 Kéfi, S., Rietkerk, M., van Baalen, M. & Loreau, M. (2007). Local facilitation, bistability and  
781 transitions in arid ecosystems. *Theoretical population biology* 71, 367–379.
- 782 Koller, D. & Friedman, N. (2009). *Probabilistic graphical models: principles and techniques*. MIT  
783 press.
- 784 Lähdesmäki, H. & Shmulevich, I. (2008). Learning the structure of dynamic bayesian networks  
785 from time series and steady state measurements. *Machine Learning* 71, 185–217.
- 786 Laroche, F., Paltto, H. & Ranius, T. (2018). Abundance-based detectability in a spatially-explicit  
787 metapopulation: a case study on a vulnerable beetle species in hollow trees. *Oecologia* 188,  
788 671–682.
- 789 Leibold, M.A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J.M., Hoopes, M.F., Holt, R.D.,  
790 Shurin, J.B., Law, R., Tilman, D. *et al.* (2004). The metacommunity concept: a framework for  
791 multi-scale community ecology. *Ecology letters* 7, 601–613.
- 792 Levins, R. (1969). Some demographic and genetic consequences of environmental heterogeneity for  
793 biological control. *American Entomologist* 15, 237–240.

- 794 Liataud, K., van Nes, E.H., Barbier, M., Scheffer, M. & Loreau, M. (2019). Superorganisms  
795 or loose collections of species? a unifying theory of community patterns along environmental  
796 gradients. *Ecology letters* 22, 1243–1252.
- 797 Massol, F., Dubart, M., Calcagno, V., Cazelles, K., Jacquet, C., Kéfi, S. & Gravel, D. (2017). Island  
798 biogeography of food webs. *Advances in Ecological Research* 56.
- 799 Mouquet, N., Lagadeuc, Y., Devictor, V., Doyen, L., Duputié, A., Eveillard, D., Faure, D., Garnier,  
800 E., Gimenez, O., Huneman, P. *et al.* (2015). Predictive ecology in a changing world. *Journal of*  
801 *Applied Ecology* 52, 1293–1310.
- 802 Mouquet, N., Matthiessen, B., Miller, T. & Gonzalez, A. (2011). Extinction debt in source-sink  
803 metacommunities. *PLoS One* 6, e17567.
- 804 Nee, S., Hassell, M.P. & May, R.M. (1997). Two-species metapopulation models. In: *Metapopulation*  
805 *biology*. Elsevier pp. 123–147.
- 806 Ohlmann, M., Miele, V., Dray, S., Chalmandrier, L., O’connor, L. & Thuiller, W. (2019). Diversity  
807 indices for ecological networks: a unifying framework using hill numbers. *Ecology letters* 22,  
808 737–747.
- 809 Ovaskainen, O. & Hanski, I. (2001). Spatially structured metapopulation models: global and local  
810 assessment of metapopulation capacity. *Theoretical population biology* 60, 281–302.
- 811 Pillai, P., Loreau, M. & Gonzalez, A. (2010). A patch-dynamic framework for food web metacom-  
812 munities. *Theoretical Ecology* 3, 223–237.
- 813 Reid, A.M. & Lortie, C.J. (2012). Cushion plants are foundation species with positive effects  
814 extending to higher trophic levels. *Ecosphere* 3, 1–18.
- 815 Sardanyés, J., Piñero, J. & Solé, R. (2019). Habitat loss-induced tipping points in metapopulations  
816 with facilitation. *Population Ecology* 61, 436–449.
- 817 Schreiber, S.J., Levine, J.M., Godoy, O., Kraft, N.J.B. & Hart, S.P. (2023). Does deterministic  
818 coexistence theory matter in a finite world? *Ecology* 104, e3838.

- 819 Simon, P.L., Taylor, M. & Kiss, I.Z. (2011). Exact epidemic models on graphs using graph-  
820 automorphism driven lumping. *Journal of mathematical biology* 62, 479–508.
- 821 Smith, H.L. (2008). *Monotone dynamical systems: an introduction to the theory of competitive and*  
822 *cooperative systems: an introduction to the theory of competitive and cooperative systems*. 41.  
823 American Mathematical Soc.
- 824 Sole, R.V. & Bascompte, J. (2007). Self organization in complex ecosystems.
- 825 Song, C., Rohr, R.P. & Saavedra, S. (2018). A guideline to study the feasibility domain of multi-  
826 trophic and changing ecological communities. *Journal of theoretical biology* 450, 30–36.
- 827 Staniczenko, P.P., Sivasubramaniam, P., Suttle, K.B. & Pearson, R.G. (2017). Linking macroecology  
828 and community ecology: refining predictions of species distributions using biotic interaction  
829 networks. *Ecology letters* 20, 693–707.
- 830 Stumpf, M.P. & Porter, M.A. (2012). Critical truths about power laws. *Science* 335, 665–666.
- 831 Takeuchi, Y. (1996). *Global dynamical properties of Lotka-Volterra systems*. World Scientific.
- 832 Thébault, E. & Fontaine, C. (2010). Stability of ecological communities and the architecture of  
833 mutualistic and trophic networks. *Science* 329, 853–856.
- 834 Thuiller, W., Münkemüller, T., Lavergne, S., Mouillot, D., Mouquet, N., Schiffrers, K. & Gravel, D.  
835 (2013). A road map for integrating eco-evolutionary processes into biodiversity models. *Ecology*  
836 *letters* 16, 94–105.
- 837 Tilman, D., Lehman, C.L. & Yin, C. (1997). Habitat destruction, dispersal, and deterministic  
838 extinction in competitive communities. *The American Naturalist* 149, 407–435.
- 839 Tischendorf, L. & Fahrig, L. (2000). On the usage and measurement of landscape connectivity.  
840 *Oikos* 90, 7–19.
- 841 Valdovinos, F.S. (2019). Mutualistic networks: moving closer to a predictive theory. *Ecology letters*  
842 22, 1517–1534.

- 843 Van Mieghem, P. (2011). The n-intertwined sis epidemic network model. *Computing* 93, 147–169.
- 844 Van Mieghem, P. & Cator, E. (2012). Epidemics in networks with nodal self-infection and the  
845 epidemic threshold. *Physical Review E* 86, 016116.
- 846 Vázquez, D.P., Chacoff, N.P. & Cagnolo, L. (2009). Evaluating multiple determinants of the struc-  
847 ture of plant–animal mutualistic networks. *Ecology* 90, 2039–2046.
- 848 Wang, S., Brose, U., van Nouhuys, S., Holt, R.D. & Loreau, M. (2021). Metapopulation capacity  
849 determines food chain length in fragmented landscapes. *Proceedings of the National Academy of*  
850 *Sciences* 118.
- 851 Wuyts, B. & Sieber, J. (2022). Mean-field models of dynamics on networks via moment closure:  
852 An automated procedure. *Physical Review E* 106, 054312.
- 853 Zhang, H., Bearup, D., Nijs, I., Wang, S., Barabás, G., Tao, Y. & Liao, J. (2021). Dispersal network  
854 heterogeneity promotes species coexistence in hierarchical competitive communities. *Ecology*  
855 *Letters* 24, 50–59.