



Past, current and future states of Green Lizard (*Lacerta viridis*) populations in Eastern Europe

Oksana Nekrasova, Oleksii Marushchak, Anastasiia Marushchak, Jean-Yves Georges

► To cite this version:

Oksana Nekrasova, Oleksii Marushchak, Anastasiia Marushchak, Jean-Yves Georges. Past, current and future states of Green Lizard (*Lacerta viridis*) populations in Eastern Europe. Biology of Lizard 2024, Rodeo, NM, USA, 24-27 July 2024, Jul 2024, Rodeo, United States. hal-04730029

HAL Id: hal-04730029

<https://hal.science/hal-04730029v1>

Submitted on 11 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



PAST, CURRENT AND FUTURE STATES OF GREEN LIZARD (LACERTA VIRIDIS) POPULATIONS IN EASTERN EUROPE

Oksana Nekrasova^{1,2,3}, Oleksii Marushchak^{1,3}, Anastasija Marushchak⁴, Mihails Puipins², Andris Čeirāns², Arturs Skute², Kathrin Theisinger^{5,6}, Jean-Yves Georges¹

¹ Université de Strasbourg, CNRS IPhC, UMR 7178, Strasbourg, France; ² Institute of Life Sciences and Technologies, Daugavpils University, Daugavpils, Latvia; ³ I. I. Schmalhausen Institute of Zoology NASU, Kyiv, Ukraine; ⁴ Kyiv Zoo, Kyiv, Ukraine; ⁵ Justus Liebig University Gießen, Institute for Insect Biotechnology, Gießen, Germany; ⁶ Fraunhofer Institute for Molecular Biology and Applied Ecology, Gießen, Germany

INTRODUCTION

The green lizard (*Lacerta viridis* Laurent 1768) is one of the species of big-sized reptiles (Reptilia) of a total length of up to 400 mm with natural range occupying territories from Italy to Turkey. This species requires special protection being included in Appendix II of the Bern Convention (1979), Annex IV of the Habitats Directive (1992) and IUCN protection lists (category Least Concern [LC]). It is also included in the 4th edition of the Red Data Book of Ukraine (category «Vulnerable»).

Over the last decades, climate change and various anthropogenic pressures have been reported to have greatest impacts on reptile populations worldwide. Moreover, thermophilic species such as *L. viridis* are protected due to their dramatic population trends especially in Eastern Europe.

MATERIALS AND METHODS

To study the prospects for the 2050 of green lizard in Europe, we used the GBIF database (n=1031), literature sources and personal observations. We used species distribution bioclimatic modeling Maxent software (25 replicates, test 25%) using the 35 CLIMond and 19 PaleoClim bioclimatic variables. To account for sampling bias, we used the nearest neighbor distance ('nbox' package in R; Osoño-Olvera et al., 2020) method for thinning the data. Occurrence points that were >0.1 units away from each other were removed to avoid errors due to spatial autocorrelation. Modeling and calculations were carried out using Maxent v3.3.3 software with 25 replicates (Phillips, 2005). Additional processed material from Ukrainian populations was included into the study in order to illustrate modern problems and trends face by the lizards, includes museum specimens from the 1960s (n = 50) and modern records (2012–2017, n=97).

PAST

By studying the morphology of both museum specimens from the 1960s (n = 50) and comparing with modern populations (2012–2017, n=97), we noticed an increase in number of anomalies of head foliosis in the most recent specimens (21.1%). Recent field censuses and observations in our field sites throughout Ukraine also revealed an increase in the occurrence of rare or so-called abnormal morphs – melanized forms, "meridionalis morphotype" (Odeia and Mykolajiv regions) and "leopard morphotype" (Cherkasy region), with visible signs of viral (Reoviridae) papillomas in the populations of southern Ukraine.



Figure 2. «Meridionalis morphotypes» and «leopard morphotypes» (photo by V. Golub) of *L. viridis* from southern and central Ukraine respectively.




Figure 3. Viral papillomas (Reoviridae) From *L. viridis* of southern populations in Ukraine. In a population of Sukhyi Lyman, Odessa, up to 15% of the detected lizards are have visual evidence of these viral papillomas.




Figure 4. Modelled current distribution of *L. viridis* in Europe based on the used bioclimatic variables for current situation (2000) and taking into account future climate changes (2050).
The research was partly funded by the project Enviro-R https://enviroscars.fr through the BioDivRestore ERA-Net COFUND programme and with the funding organizations Agence Nationale de la Recherche (ANR, France), Bundesministerium für Bildung und Forschung (BMBF, Germany), State Education Development Agency (VIAA, Latvia) National Science Center (NSC, Poland). Additionally the study was supported by the projects 16W015, PAUSE (ANR-23-PAUP-0074) and Nr.ZP-2021/1-0247 (Latvia)

CURRENT



Figure 5. Morphology of both museum specimens from the 1960s (n = 50) and comparing with modern populations (2012–2017, n=97), we noticed an increase in number of anomalies of head foliosis in the most recent specimens (21.1%). Recent field censuses and observations in our field sites throughout Ukraine also revealed an increase in the occurrence of rare or so-called abnormal morphs – melanized forms, "meridionalis morphotype" (Odeia and Mykolajiv regions) and "leopard morphotype" (Cherkasy region), with visible signs of viral (Reoviridae) papillomas in the populations of southern Ukraine.



Figure 6. Viral papillomas (Reoviridae) From *L. viridis* of southern populations in Ukraine. In a population of Sukhyi Lyman, Odessa, up to 15% of the detected lizards are have visual evidence of these viral papillomas.

FUTURE

Finally, by implementing bioclimatic models based in GIS, Maxent and Wordclim, we could predict a northward shift of the range of the species in Europe up to the Baltic countries by 2050. Consistently there is already almost a 2-fold decrease in the number of populations in the south of its range (Black Sea region). All these records collected in the recent decades indicate that green lizard populations require creation of local and international conservation programs for the protection of the species.




Figure 7. Modelled range shift of *L. viridis* in Europe based on the used bioclimatic variables for current situation (2000) and taking into account future climate changes (2050).
The research was partly funded by the project Enviro-R https://enviroscars.fr through the BioDivRestore ERA-Net COFUND programme and with the funding organizations Agence Nationale de la Recherche (ANR, France), Bundesministerium für Bildung und Forschung (BMBF, Germany), State Education Development Agency (VIAA, Latvia) National Science Center (NSC, Poland). Additionally the study was supported by the projects 16W015, PAUSE (ANR-23-PAUP-0074) and Nr.ZP-2021/1-0247 (Latvia)