



HAL
open science

Cutoff phenomenon for asymmetric zero range process with monotone rates

Ons Rameh

► **To cite this version:**

Ons Rameh. Cutoff phenomenon for asymmetric zero range process with monotone rates. 2024.
hal-04731272

HAL Id: hal-04731272

<https://hal.science/hal-04731272v1>

Preprint submitted on 10 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License

CUTOFF PHENOMENON FOR ASYMMETRIC ZERO RANGE PROCESS WITH MONOTONE RATES

ONS RAMEH

ABSTRACT. We investigate the mixing time of the asymmetric Zero Range process on the segment with a non-decreasing rate. We show that the cutoff holds in the totally asymmetric case with a convex flux, and also with a concave flux if the asymmetry is strong enough. We show that the mixing occurs when the macroscopic system reaches equilibrium. A key ingredient of the proof, of independent interest, is the hydrodynamic limit for irregular initial data.

CONTENTS

1. Introduction	1
2. Main Results	3
3. Preliminaries	6
4. Hydrodynamic limit on the segment	12
5. Proof of Lower and Upper bounds on the mixing time	17
6. Evolution of the left-most particle and the stack at N	21
Appendix A. Proof of Lemma 12	26
Appendix B. A technical lemma	27
List of symbols	27
References	28

1. INTRODUCTION

1.1. Model. The Zero range process on $I_N = \llbracket 1, N \rrbracket$ is a model describing the evolution of $k \geq 1$ indistinguishable particles hopping on the lattice I_N with a rate that depends on the number of the particles at the departure site and the direction of the jump. This model was introduced by Spitzer in 1970 [32] as a stochastic lattice gas with on-site interaction. Since then, it has been widely studied (see [15, 27, 18] and the reference therein).

Let us now explicitly describe the process. Define

$$\Omega_{N,k}^0 = \left\{ \eta \in \mathbb{N}^{\llbracket 1, N \rrbracket} : \sum_{i=1}^N \eta(i) = k \right\}. \quad (1)$$

the space of configurations of k particles on I_N .

Given $p \in (\frac{1}{2}, 1]$, $q = 1 - p$ and a rate function by $g : \mathbb{N} \mapsto [0, +\infty)$ a non-decreasing Lipschitz function such that $g(0) = 0 < g(1)$, an Asymmetric Zero Range process (ZRP (g, p)) is a continuous time Markov process on $\Omega_{N,k}^0$ with the following generator:

$$\mathcal{A}_N J(\eta) = \sum_{i=1}^{N-1} pg(\eta(i)) (J(\eta^{i,i+1}) - J(\eta)) + \sum_{i=2}^N qg(\eta(i)) (J(\eta^{i,i-1}) - J(\eta)) \quad (2)$$

where $\eta^{x,y}$ is the configuration obtained when one particle from x jumps to the site y and $J : \Omega_{N,k}^0 \mapsto \mathbb{R}$ is a test function.




FIGURE 1. A configuration of particles with $k = 11$ and $N = 8$. The crossed red arrows correspond to forbidden jumps.

Remark 1. When g is linear, the dynamic describes the evolution of k independent particles. Each particle jumps to the right with rate $pg(1)$ and to the left with rate $qg(1)$.

The process is irreducible and reversible with respect to its equilibrium measure

$$\pi_{N,k}(\eta) = \frac{1}{Z_{N,k}} \prod_{y=1}^N \frac{\left(\frac{p}{q}\right)^{\eta(y)y}}{g(\eta(y))!} \quad (3)$$

where $g(k)! = g(k)g(k-1) \cdots g(1)$. In particular, when $p = 1$, the equilibrium measure is a Dirac mass on the configuration $\vee_{N,k} = k\delta_N$.

1.2. Mixing time and Cutoff Phenomenon. Classical Markov process theory guarantees that the process converges to its equilibrium measure $\pi_{N,k}$ starting from any configuration. This article aims to estimate the speed at which the equilibrium is reached. The standard choice is the total-variation distance, as it allows for uniform control, following Diaconis' earlier work [14, 1]. We denote the worst-case total-variation distance by:

$$d_{N,k}(t) = \max_{\eta \in \Omega_{N,k}^0} \|P_t^\eta(\cdot) - \pi_{N,k}\|_{\text{TV}} \quad (4)$$

where P_t^η is the law of a ZRP at time t started at η .

For any level of precision $\theta > 0$, we define the mixing time of ZRP :

$$T_{\text{mix}}^{N,k}(\theta) := \inf\{t \geq 0; \quad d_{N,k}(t) \leq \theta\}. \quad (5)$$

In particular, we aim to describe the behaviour of the mixing time when the size of the system and the number of particles diverge to infinity. In this article, we work in the natural setting for a scaling limit, where $\frac{k}{N}$ converge to $\alpha \in \mathbb{R}_+$, a macroscopic finite density.

Of particular interest, is the dependence of the mixing time on the precision ε . We say that the system exhibits a *Cutoff phenomenon* if

$$\forall \theta \in (0, 1), \quad \frac{T_{\text{mix}}^{N,k}(\theta)}{T_{\text{mix}}^{N,k}(1/4)} \xrightarrow{N \rightarrow +\infty} 1.$$

In other words, the distance to the equilibrium stays around 1 and drops abruptly to 0. This remarkable phase transition was discovered in the context of card shuffling [14, 1]. We refer the reader to [26, Chapter 18] for more background on the Cutoff Phenomenon.

1.3. Previous Results. The Cutoff was investigated in various contexts, among which the simple exclusion process that describes the evolution of particles hopping on a lattice and constrained to not coexist on the same site. SEP's mixing time has been studied in both symmetric [34, 23, 22, 24] and (weakly) asymmetric [25, 8, 19, 20] settings. The articles [22, 19, 20, 21] underlined a notable link between the macroscopic relaxation to equilibrium (the Hydrodynamic limit) and the mixing time. In particular, it was proven that in the asymmetric setup, the macroscopic equilibrium is reached in a finite time and it corresponds exactly to the mixing time. This link was also emphasised for the symmetric SEP with reservoir [16] where the study of hydrodynamic limit allows for a precise description of the Cutoff profile.

ZRP can be seen as a generalization of SEP as it can be mapped into a simple exclusion process (SEP) when the rate function g is constant. Cutoff for ZRP was also shown when the rate function is constant on the complete graph [29]. For more general rate functions, it was only studied to the best of our knowledge on the complete graph [17, 33].

In the context of asymmetric ZRP, the hydrodynamic limit was thoroughly studied on \mathbb{Z}^d [30] for regular initial data. It was also established in the non-conservative case [3] and in a random environment [4]. In all these results the rate function is assumed nondecreasing to guarantee the attractivity of the process and to avoid the condensation phenomena (see for example [2]).

1.4. Our contribution. The main achievement of this paper is to prove that cutoff holds in the totally asymmetric convex flux, and in the concave flux if the asymmetry is strong enough (Theorems 2 and 3). Moreover, we show that it occurs exactly at the macroscopic equilibrium time, which highlights once again the link between hydrodynamic limit and mixing times. Our proof exploits the hydrodynamic limit of ZRP on the segment, which we generalise for irregular initial data (Theorem 1). As far as we know, the hydrodynamic limit for irregular initial data hasn't been studied before. We also give a precise description of the evolution of the position of the left-most particles and the stack of particles at position N .

1.5. Acknowledgment. The author warmly thanks Max Fathi for his advice throughout this project and for thoroughly reading the drafts. The author also kindly thanks Christophe Bahadoran, Cyril Imbert, Cyril Labbé, Hubert Lacoïn, Fraydoun Rezakhanlou, Ellen Saada, Justin Salez, Marielle Simon and Cristina Toninelli for helpful discussions. This work received funding from the Agence Nationale de la Recherche (ANR) Grant ANR-23-CE40-0003 (Project CONVIVIALITY), as well as funding from the Institut Universitaire de France.

2. MAIN RESULTS

Before we state the main theorems, we introduce some notation. For any $\varphi < \varphi^*$, we define:

$$Z(\varphi) = \sum_{k \geq 0} \frac{\varphi^k}{g(k)!} \quad \text{and} \quad \forall k \in \mathbb{N}, \bar{\nu}_\varphi(k) = \frac{1}{Z(\varphi)} \frac{\varphi^k}{g(k)!}$$

where φ^* is the radius of convergence of Z . Throughout the paper, we will always assume the following:

Assumption.

- g is non-decreasing,
- g is Lipschitz :

$$g^* := \sup_{k \geq 0} |g(k+1) - g(k)| < \infty,$$

- $\lim_{\varphi \uparrow \varphi^*} Z(\varphi) = \infty$.

The first assumption ensures the attractivity of the process and avoids potential condensation phenomena that may arise. The family of measure $(\bar{\nu}_\varphi)_\varphi$ can be thus parametrized by its expectation:

$$\forall \alpha \in \mathbb{R}_+, \nu(\alpha) = \bar{\nu}_{\Phi(\alpha)}$$

where Φ is the inverse function of $\varphi \rightarrow \mathbb{E}_{\bar{\nu}_\varphi}(X)$ and corresponds to the macroscopic flux for a density α . When Φ is convex, we also denote by Ψ its convex conjugate:

$$\Psi(x) = \sup_{\alpha} (\alpha x - \Phi(\alpha)).$$

If Φ is concave, Ψ denotes the convex conjugate of $(-\Phi)$ instead. We recall that the convexity (resp. concavity) of g implies that Φ is strictly convex (resp. strictly concave) [5, Proposition 3.1].

Let $\mathcal{M}_+([0, 1])$ be the space of non-negative measure on $[0, 1]$ endowed with the weak topology. For any sequence of configurations $(\eta_{N,0})_N \in \Omega_{N,k}^0$, we define the empirical measure associated with the process $(\eta_{N,t})$:

$$\rho_{N,t}(dy) = \frac{1}{N} \sum_{i \in I_N} \eta_{N,t}(i) \delta_{i/N}(dy) \in \mathcal{M}_+([0, 1])$$

where $(\eta_{N,t})$ is ZRP started at $(\eta_{N,0})_N$. Notice that $(\rho_{N,t})_t$ belongs to the $\mathbb{D}(\mathbb{R}_+, \mathcal{M}_+([0, 1]))$ the set of cadlag functions with values in $\mathcal{M}_+([0, 1])$ endowed with the Skorokhod topology. This allows us to simultaneously embed the configurations with different sizes in a common space.

In the context of ZRP, it is intuitively clear that the worst initial condition should correspond to $\wedge_{N,k} = k\delta_1$ where all particles are on the first site. From a macroscopic scale, it corresponds to imposing $\alpha\delta_0$ as initial data where α would be the macroscopically observed mass. The following theorem gives the hydrodynamic limit of ZRP on the segment for general irregular initial data.

Theorem 1. *Assume that Φ is either strictly convex or strictly concave. Given $u_0 \in \mathcal{M}_+([0, 1])$ with total mass α , let U be the unique Barron–Jensen viscosity solution to*

$$\begin{cases} \partial_t U + \Phi(\partial_x U) = 0, & t \geq 0, x \in \mathbb{R}, \\ U(0, \cdot) = \int_{[0, \cdot]} u_0(dy). \end{cases} \quad (\text{HJ})$$

For any $p > \frac{1}{2}$, if $(\rho_N(0, dy))_N$ weakly converge to $u_0(dx)$, the sequence $(t \rightarrow \rho_{N, \frac{Nt}{p-q}})_N$ converge in law in $\mathbb{D}(\mathbb{R}_+, \mathcal{M}_+(\mathbb{R}))$ to a measure-valued function $t \rightarrow v(t, dx)$. Moreover,

$$\forall t > 0, \forall x \in (0, 1], v(t, [0, x]) = \max(U, \alpha \mathbf{1}_{\mathbb{R}_+ \times [1, +\infty)})^*.$$

Let us underline that we chose to speed up the process by $\frac{N}{p-q}$ to ensure the limit does not depend on p . The precise definition of Barron–Jensen viscosity solution of the Hamilton–Jacobi equation (HJ) is delayed to subsection 3.3. The limit is usually determined (see [31, 15]) as the unique entropic solution to the scalar conservation law:

$$\begin{cases} \partial_t u + \partial_x \Phi(u) = 0, & t \geq 0, x \in \mathbb{R}, \\ u(0, \cdot) = u_0. \end{cases} \quad (\text{CL})$$

Both notions of solutions are equivalent when $u_0 \in L^1 \cap L^\infty$ for convex or concave flux Φ [28]. Beyond the difficulties raised by the presence of boundaries, the irregularity of initial data with eventually non-bounded rate jumps required the use of Hamilton–Jacobi equations

to uniquely identify the limit v . Note that although there exists a notion of measure-valued solution of scalar conservation law, ensuring well-posedness in a more general framework (see [13, 9, 10, 11]), adapting the uniqueness criterion to our situation seems more challenging. Regarding the boundary conditions, we extensively use the attractivity of the process and the fact that the characteristics point towards the right.

For any $\alpha \geq 0$, let U^α be the Barron–Jensen viscosity solution associated to $u_0^\alpha = \alpha\delta_0$. One can in particular remark that the macroscopic system reaches equilibrium in finite time. We define for any $\alpha \geq 0$ the macroscopic equilibrium time

$$T_{eq,p,\alpha} = \frac{1}{p-q} \inf\{t > 0, U^\alpha(t, dx) \leq \alpha \mathbb{1}_{x>1}\}. \quad (6)$$

If the macroscopic system has not reached the equilibrium, the microscopic system is also still far from it. Consequently, the macroscopic equilibrium time provides the following lower-bound on the mixing time.

Theorem 2. *Assume that Φ is either strictly convex or strictly concave. For any $p > \frac{1}{2}$,*

$$\forall \alpha \in \mathbb{R}_+, \forall \theta \in (0, 1), \quad \liminf_{\substack{N \rightarrow \infty \\ k/N \rightarrow \alpha}} \frac{T_{mix}^{N,k}(\theta)}{N} \geq T_{eq,p,\alpha}.$$

Our main result is that the lower-bound is sharp in the totally asymmetric convex case and the concave setting if the asymmetry is strong enough.

Theorem 3.

1. *For $p = 1$ and assuming that Φ is strictly convex and $g^* = g(1)$,*

$$\forall \alpha \in \mathbb{R}_+, \forall \theta \in (0, 1), \quad \lim_{\substack{N \rightarrow \infty \\ k/N \rightarrow \alpha}} \frac{T_{mix}^{N,k}(\theta)}{N} = T_{eq,1,\alpha} = \frac{1}{g(1)}.$$

2. *Assume that Φ is strictly concave and bounded by \bar{g} . Assume $p > \frac{1}{2}, \alpha > 0$ are such that*

$$\frac{pg(1) - q\bar{g}}{p-q} > \max \left\{ \frac{\Phi \left(\Psi' \left(\Psi^{-1} \left(\frac{\alpha}{T_{eq,1,\alpha}} \right) \right) \right)}{\Psi' \left(\Psi^{-1} \left(\frac{\alpha}{T_{eq,1,\alpha}} \right) \right)}, \Phi \left((-\Phi')^{-1} \left(\frac{-1}{T_{eq,1,\alpha}} \right) \right) \right\}. \quad (7)$$

Then,

$$\forall \alpha \in \mathbb{R}_+, \forall \theta \in (0, 1), \quad \lim_{\substack{N \rightarrow \infty \\ k/N \rightarrow \alpha}} \frac{T_{mix}^{N,k}(\theta)}{N} = T_{eq,p,\alpha}.$$

Clearly, it implies that the ZRP in this framework exhibits a Cutoff since the limit does not depend on θ . As was already the case in the ASEP [19], the hydrodynamic limit does not provide precise enough information to prove the matching upper-bound. An easy way to see it is by considering $1\delta_1 + (k-1)\delta_N$ as an initial condition where macroscopic equilibrium is achieved and microscopically we need to wait for $(p-q)g(1)N$.

The main idea is to control precisely:

- $L_{N,k}$: the position of the left-most particle
- $S_{N,k}$: the number of particles at the position N

and prove that they do follow their macroscopic analogue $(\mathcal{L}_p^\alpha, \mathcal{S}_p^\alpha)$, predicted by the hydrodynamic limit. Once these two statistics reach their equilibrium state, the system rapidly mixes. At this level of generality, $\pi_{N,k}$, \mathcal{L}_p^α and \mathcal{S}_p^α are not explicit, making the problem delicate. We also emphasise that ZRP do not have a particle-hole duality, unlike ASEP. Thus, the study of

$L_{N,k}$ and $S_{N,k}$ need to be conducted separately. The technical condition (7) is due to a comparison to ASEP case which lacks precision. Indeed, condition (7) requires that the maximum macroscopic speed and flux achieved by the ZRP be below the critical level $pg(1) - q\bar{g}$ of the ASEP, to which we are comparing (see remarks 9 and 10). We believe that the result should hold in a more general context.

Remark 2. *The condition (7) is verified on a wider class of ZRP than just for the constant rate. Indeed, we observe that when α converge to 0, $\Phi\left((-\Phi')^{-1}\left(\frac{-1}{T_{eq,1,\alpha}}\right)\right) \rightarrow 0$. Hence, one can fix α_{\max} such that*

$$\Phi\left((-\Phi')^{-1}\left(\frac{-1}{T_{eq,1,\alpha}}\right)\right) < g(1).$$

Thus, when $p = 1$, the condition (7) is trivially fulfilled for any $\alpha \leq \alpha_{\max}$. Consequently,

$\exists \alpha_{\max}, \forall 0 < \alpha \leq \alpha_{\max}, \exists p_{\min} \in (1/2, 1), \forall p \geq p_{\min}$ such that the condition (7) is fulfilled.

2.1. Organisation of the paper. The rest of the article is organized as follows. In Subsections 3.1 and 3.2, we recall the attractivity property of ZRP and the mapping to the exclusion process when the rate is constant. Then, we recall the definition of the Barron–Jensen viscosity solution in Subsection 3.3 and generalize the hydrodynamic limit on \mathbb{Z} with irregular initial data in Subsection 3.4. In Section 4, we prove the hydrodynamic limit of ZRP on the segment. The proof of Theorems 2 and 3 occupies the whole Section 5. Section 6 is devoted to the control of the position of the left-most particle and the stack at position N when the flux is concave. A list of symbols used is inducted at the end of the Appendix.

3. PRELIMINARIES

3.1. Coupling, attractivity, and orders. Throughout the article, we shall frequently need to couple ZRP on different graphs and rate functions. For convenience, we introduce a generic form that serves as a standard framework, to which all specific cases can be reduced. Given $g_+, g_- : \mathbb{N} \times \mathbb{Z} \mapsto \mathbb{R}_+$, we define the following generator:

$$\mathcal{A}_{g_+,g_-} J(\eta) = \sum_{i \in \mathbb{Z}} g_+(\eta(i), i) (J(\eta^{i,i+1}) - J(\eta)) + g_-(\eta(i), i) (J(\eta^{i,i-1}) - J(\eta)), \quad (8)$$

where g_+ (resp. g_-) is the rate of jumps to the right (resp. left) from position i . Let us mention that any ZRP generator on any sub-graph of \mathbb{Z} could be written in this generic form.

It can be more convenient to compare their evolution by representing a configuration η through its height function. For any $\eta \in \mathbb{N}^{\mathbb{Z}}$, we define the height function by

$$h_\eta(j) = \sum_{i=1}^j \eta(i), \quad \forall j \in \mathbb{Z}. \quad (9)$$

We denote by $\Omega_{\mathbb{Z}}$ the set of height functions corresponding to configurations on \mathbb{Z} . It is endowed with the order defined by

$$\{h \geq \tilde{h}\} \Leftrightarrow \{\forall i \in \mathbb{Z}, h(i) \geq \tilde{h}(i)\}.$$

As the asymmetry induces a bias to the right, we can understand this partial order as a comparison of the evolution of the processes, where h is farther than \tilde{h} from the equilibrium.

Remark 3. *In the case of ZRP on the segment, we will denote the set of height functions by $\Omega_{N,k} = h(\Omega_{N,k}^0)$. In particular, the configuration $\vee_{N,k} = k\delta_N$ is the minimal element and $\wedge_{N,k} = k\delta_1$ to the maximal element. Due to the bias to the right, the configuration $\wedge_{N,k}$ corresponds to the worst initial condition.*

The following lemma provides a straightforward criterion to compare different processes, which is a slight generalization of the usual attractive coupling.

Lemma 1 (Attractivity). *Given $g_+, g_-, \tilde{g}_+, \tilde{g}_- : \mathbb{N} \times \mathbb{Z} \mapsto \mathbb{R}_+$ such that:*

$$\forall i \in \mathbb{Z}, \forall l \geq m, \quad \tilde{g}_+(l, i) \geq g_+(m, i) \text{ and } \tilde{g}_-(m, i) \leq g_-(l, i). \quad (10)$$

For any $\tilde{h}_0 \leq h_0 \in \Omega_{\mathbb{Z}}$, there exists a coupling such that almost surely:

$$\forall t \geq 0, \tilde{h}_t \leq h_t$$

where h (resp. \tilde{h}) follows the dynamic \mathcal{A}_{g_+, g_-} (resp. $\mathcal{A}_{\tilde{g}_+, \tilde{g}_-}$).

Proof. We introduce a coupling via the following Markov generator:

$$\tilde{\mathcal{A}}J(\eta, \tilde{\eta}) = \sum_{i \in \mathbb{Z}} \mathcal{G}_{i,+} + \mathcal{G}_{i,-} \quad (\mathcal{C})$$

where

$$\begin{aligned} \mathcal{G}_{k,\pm} J(\eta, \tilde{\eta}) &= (g_{\pm}(\eta(i), i) \wedge \tilde{g}_{\pm}(\tilde{\eta}(i), i)) (J(\eta^{i,i\pm 1}, \tilde{\eta}^{i,i\pm 1}) - J(\eta, \tilde{\eta})) \\ &\quad + [g_{\pm}(\eta(i), i) - \tilde{g}_{\pm}(\tilde{\eta}(i), i)]_{\pm} (J(\eta^{i,i\pm 1}, \tilde{\eta}) - J(\eta, \tilde{\eta})) \\ &\quad + [\tilde{g}_{\pm}(\tilde{\eta}(i), i) - g_{\pm}(\eta(i), i)]_{\pm} (J(\eta, \tilde{\eta}^{i,i\pm 1}) - J(\eta, \tilde{\eta})). \end{aligned}$$

As usual, we want particles for the two processes to jump simultaneously as much as possible. We denote by

$$\mathcal{E} = \{(h, \tilde{h}) \in \Omega_{\mathbb{Z}}^2, h \geq \tilde{h}\}.$$

It remains to show that if $h \geq \tilde{h}$ then $\tilde{\mathcal{A}}\mathbb{1}_{\mathcal{E}}(h, \tilde{h}) \geq 0$. We remark that the jump from $i \in \mathbb{Z}$ that may contribute negatively are:

- A jump to the right of an η -particle without an $\tilde{\eta}$ -particle jump when $h(i) = \tilde{h}(i)$. In such a case, $\eta(i) \leq \tilde{\eta}(i)$ which implies that $g_+(\eta(i), i) \leq \tilde{g}_+(\tilde{\eta}(i), i)$.
- A jump to the left of an $\tilde{\eta}$ -particle without an η -particle jump when $h(i-1) = \tilde{h}(i-1)$. In such a case, $\eta(i) \geq \tilde{\eta}(i)$ which implies that $g_-(\eta(i), i) \geq \tilde{g}_-(\tilde{\eta}(i), i)$.

In both cases, the corresponding term in the generator is null which concludes the proof. \square

Remark 4. *As g is non-decreasing, we recover the attractivity of ZRP :*

For any $\tilde{h}_0 \leq h_0$, there exists a coupling such that almost surely for any $t \geq 0$, $\tilde{h}_t \leq h_t$. One can see the attractivity as the analogue of the L^1 -contraction of entropy solution to (CL). It can also be seen as a microscopic counterpart of the Hamilton-Jacobi's maximum principle. The attractivity is the key argument in the proof of the hydrodynamic limit of the ZRP in [30],[18, Chapter 6] and also for Theorem 1.

We will also seek to transfer information about the macroscopic behaviour of ZRP on \mathbb{Z} to the ZRP on subintervals. In the spirit of [19, Lemma 18] and [3, Lemma 3.3], the same coupling (\mathcal{C}) guarantees a comparison of both processes for short times and far from the border. The proof is identical that in the references but we include it for completeness.

Lemma 2 (Comparison far from borders). *Given $a < b \in \mathbb{Z} \cup \{\pm\infty\}$, $\beta \in \mathbb{R}_+$, let η be a ZRP (g, p) on \mathbb{Z} and $\tilde{\eta}$ a ZRP (g, p) on $[[aN, bN]]$ coupled according to (\mathcal{C}).*

There exists a constant $C > 0$ such that for any $a', b' \in \mathbb{R}$ such that $a < a' < b' < b$

$$\lim_{N \rightarrow \infty} \sup_{\substack{\eta_0 = \tilde{\eta}_0 \in \mathbb{N}^{\mathbb{Z}} \\ \sum_i \tilde{\eta}_0(i) \leq \beta N}} \sup_{s \in [0, C\varepsilon]} \mathbb{E} \left[\frac{1}{N} \sum_{i=a'N}^{b'N} \left| \eta_{\frac{sN}{p-q}}(i) - \tilde{\eta}_{\frac{sN}{p-q}}(i) \right| \right] = 0,$$

where $\varepsilon = \min(b - b', a' - a)$.

Proof of Lemma 2. Given such $a', b' \in \mathbb{R}$. We introduce

- $H : \mathbb{R} \mapsto [0, 1]$ smooth, non-increasing, equal to 1 on \mathbb{R}_- and to 0 on $[1, \infty)$.
- $q : \mathbb{R} \mapsto [0, 1]$ equal to 0 on $[a', b']$, to 1 on $(-\infty, a' - \frac{2\varepsilon}{3}] \cup [b' + \frac{2\varepsilon}{3}, \infty)$ and such that $\|q'\|_\infty \leq 3\varepsilon^{-1}$.

We denote by $J(t, x) = H\left(\frac{6g^*}{(p-q)\varepsilon}t + q(x)\right)$. Then, there exists $c, C > 0$ such that $J(t, x) > c$ for any $(t, x) \in [0, C\varepsilon] \times [a', b']$ which implies:

$$\mathbb{E} \left(\frac{1}{N} \sum_{i=a'N}^{b'N} \left| \eta_{\frac{tN}{p-q}}(i) - \tilde{\eta}_{\frac{tN}{p-q}}(i) \right| \right) \leq \mathbb{E} \left(\frac{1}{cN} \sum_{i \in \mathbb{Z}} J \left(t, \frac{i}{N} \right) \left| \eta_{\frac{tN}{p-q}}(i) - \tilde{\eta}_{\frac{tN}{p-q}}(i) \right| \right).$$

Since $\eta_0 = \tilde{\eta}_0$, we have:

$$\sum_{i \in \mathbb{Z}} J \left(t, \frac{i}{N} \right) \left| \eta_{\frac{tN}{p-q}}(i) - \tilde{\eta}_{\frac{tN}{p-q}}(i) \right| = M_t + I_t \quad (11)$$

where M is a martingale with $M_0 = 0$ and

$$I_t = \int_0^t ds \left[\sum_{i \in \mathbb{Z}} \partial_t J \left(s, \frac{i}{N} \right) \left| \eta_{\frac{sN}{p-q}}(i) - \tilde{\eta}_{\frac{sN}{p-q}}(i) \right| + \frac{N}{p-q} J \left(s, \frac{i}{N} \right) \tilde{\mathcal{A}} \left| \eta_{\frac{sN}{p-q}}(i) - \tilde{\eta}_{\frac{sN}{p-q}}(i) \right| \right].$$

As J is equal to 0 as soon as $x \notin [a' - \frac{2\varepsilon}{3}, b' + \frac{2\varepsilon}{3}]$, we obtain:

$$\begin{aligned} & \sum_{i \in \mathbb{Z}} J \left(\frac{s}{N}, \frac{i}{N} \right) \tilde{\mathcal{A}} \left| \eta_s(i) - \tilde{\eta}_s(i) \right| \\ & \leq \sum_{i \in \mathbb{Z}} p \left[J \left(s, \frac{i+1}{N} \right) - J \left(s, \frac{i}{N} \right) \right] \left| \eta_{\frac{sN}{p-q}}(i) - \tilde{\eta}_{\frac{sN}{p-q}}(i) \right| \text{ord}_{i,i+1} \\ & \quad + \sum_{i \in \mathbb{Z}} q \left[J \left(s, \frac{i-1}{N} \right) - J \left(s, \frac{i}{N} \right) \right] \left| \eta_{\frac{sN}{p-q}}(i) - \tilde{\eta}_{\frac{sN}{p-q}}(i) \right| \text{ord}_{i,i-1}, \end{aligned}$$

where $\text{ord}_{i,j} = \mathbb{1}_{(\eta(i) - \tilde{\eta}(i))(\eta(j) - \tilde{\eta}(j)) \geq 0}$. Since the number of particles is bounded by βN and J is smooth, there exists $C_{J,\beta} > 0$ such that for any $t \leq C\varepsilon$

$$I_t \leq C_{J,\beta} + \int_0^t ds \sum_{i \in \mathbb{Z}} \left(\partial_t J + \frac{2g^*}{p-q} |\partial_x J| \right) \left(\frac{s}{N}, \frac{i}{N} \right) \left| \eta_{\frac{sN}{p-q}}(i) - \tilde{\eta}_{\frac{sN}{p-q}}(i) \right|.$$

Finally, we observe that $\partial_t J + \frac{2g^*}{p-q} |\partial_x J| \leq 0$ which implies for any $t \in [0, C\varepsilon]$:

$$\mathbb{E} \left(\frac{1}{cN} \sum_{i \in \mathbb{Z}} J \left(t, \frac{i}{N} \right) \left| \eta_{\frac{tN}{p-q}}(i) - \tilde{\eta}_{\frac{tN}{p-q}}(i) \right| \right) \leq \frac{C_{J,\beta}}{N},$$

and concludes the proof. \square

3.2. Link with the exclusion process. Let us recall the generator of the asymmetric exclusion process (ASEP) on a segment:

$$\mathcal{A}_N^{\text{ex}} J(\eta) = \sum_{i=1}^{N-1} p\eta(i)[1-\eta(i+1)] (J(\eta^{i,i+1}) - J(\eta)) + \sum_{i=2}^N q\eta(i)[1-\eta(i-1)] (J(\eta^{i,i-1}) - J(\eta)).$$

For any configuration $\xi \in \Omega_{N,k}^{0,\text{ex}} = \left\{ \xi \in \{0, 1\}^{I_{N+k-1}}, \sum_{i=1}^{N+k-1} \xi_i = k \right\}$, we define $E(\xi) \in \Omega_{N,k}^0$ according the following rules:

- Label $(z_i)_{i=1}^N$ the position of empty-sites on the configuration ξ . Fix $z_0 = 0$ and $z_N = N + k + 1$

- On each site i , put $z_i - z_{i-1} - 1$ particles.




FIGURE 2. The exclusion configuration corresponding to the figure (Fig.1.1)

It is easy to see that E is a bijection from $\Omega_{N,k}^{0,\text{ex}}$ to $\Omega_{N,k}^0$. Moreover, it is well-known that ZRP is in bijection with ASEP on a segment I_{N+k-1} with k particles if the rate function is constant [15, 12].

Lemma 3 (Link between ZRP and SEP). *If $(\eta_t)_{t \geq 0}$ is a ZRP with a rate function $g = \mathbb{1}_{x \geq 1}$ started at η_0 , then process $(E^{-1}(\eta_t))_t$ evolves according to $\mathcal{A}_N^{\text{ex}}$ started at $E^{-1}(\eta_0)$.*

3.3. Hamilton Jacobi Equation. In this subsection, we recall uniqueness and stability properties of the Hamilton-Jacobi equation on $\bar{\Omega} = [0, +\infty) \times \mathbb{R}$

$$\begin{cases} \partial_t U + \Phi(\partial_x U) = 0, & t, x \in \Omega \\ U(0, \cdot) = U_0, \end{cases} \quad (\text{HJ})$$

where $\Omega = (0, +\infty) \times \mathbb{R}$. Let us assume that Φ is strictly convex.

Barron and Jensen introduced in [7] a notion of viscosity solution adapted for discontinuous initial data.

Definition 1. *A bounded lower semicontinuous function is a Barron–Jensen solution (BJ) iff for all $\varphi \in C^1(\bar{\Omega})$ and at each maximum point $(t_0, x_0) \in \Omega$ of $U - \varphi$, we have*

$$\partial_t \varphi(t_0, x_0) + \Phi(\partial_x \varphi(t_0, x_0)) = 0. \quad (\text{BJ})$$

Remark 5. *The Barron–Jensen notion of viscosity solution is also called the “bilateral solution” and is equivalent to the Crandall–Lions viscosity solution if Φ is convex and $U \in W_{loc}^{1,\infty}(\Omega)$ [7, Theorem 2.3].*

This notion ensures the uniqueness of the solution to (HJ).

Theorem 4 ([6, Theorem 5.14]). *Assume that U_0 is a l.s.c bounded function. There exists at most one l.s.c function U which satisfies both (BJ) and*

$$u(x, 0) = u_0 \quad \text{on } \mathbb{R}^N.$$

It is also well-known that if the initial data U_0 is Lipschitz, the viscosity solution is given by the Hopf–Lax formula:

$$U(t, x) = \inf_{y \in \mathbb{R}} \left(U_0(y) + t \Psi \left(\frac{x - y}{t} \right) \right) \quad (12)$$

where we recall that Ψ is the convex conjugate function of Φ . Demengel and Serre proved in [13] that the Hopf–Lax formula is stable.

Proposition 1 ([13, Proposition 4.1]). *Given a sequence $(u_{0,n})$ bounded in $L^1(\mathbb{R})$ and converging vaguely to a measure $u_{0,\infty}$, let*

$$\forall n \in \mathbb{N} \cup \{\infty\}, \quad U_n(t, x) = \inf_{y \in \mathbb{R}} \left(\int_{(-\infty, y)} u_{0,n}(z) dz + t \Psi \left(\frac{x - y}{t} \right) \right).$$

Then, for any $t \geq 0, x \in \text{Cont}(U(t))$, $(U_n(t, x))$ converge to $U_\infty(t, x)$ defined by the Hopf–Lax formula.

In the reference, the convergence is stated to be point-wise which is not the case even for $t = 0$. Nonetheless, the proof yields the convergence at continuity points. On the other hand, Barron and Jensen proved the stability of their solutions in [7, Corollary 3.9]. The proof is provided for a bounded Hamiltonian but can easily be extended to our framework. Thus, we have the following corollary:

Corollary 1. *Given $u_0 \in \mathcal{M}_+(\mathbb{R})$ with finite mass, the unique Barron–Jensen solution U is the function given by the Hopf–Lax formula (12) where*

$$U(0, x) = \int_{(-\infty, x)} u_0(dy) \quad \text{on } \mathbb{R}.$$

Since in our case Φ' is non-negative, the characteristics always point towards the right direction. The uniqueness result can be generalized by specifying the influence domain.

Lemma 4. *Assume that U_0, V_0 are l.s.c bounded functions and that there exists $C > 0$ such that:*

$$U_0(x) = V_0(x), \quad \forall x \in (-\infty, C]$$

then, $U(t, x) = V(t, x)$ for all $x < C + t\Phi'(0)$.

Proof. For any $t > 0, x \in \mathbb{R}$, we observe that if y is such that $\frac{x-y}{t} \geq \Phi'(0)$ then $\Psi\left(\frac{x-y}{t}\right) = 0$. So, the infimum can be taken on the smaller set $y \in (-\infty, x - \Phi'(0)t]$. This implies the lemma. \square

Remark 6. *We emphasize that all these results can be adapted in the case of a strictly concave flux by replacing l.s.c with u.s.c and maximum by minimum. It is sufficient to consider $V(t, x) = U_0(\infty) - U(t, -x)$. The Hopf–Lax formula becomes:*

$$U(t, x) = \sup_{y \in \mathbb{R}} \left(\int_{(-\infty, y]} u_0(dz) - t\Psi\left(\frac{x-y}{t}\right) \right), \quad (13)$$

where Ψ is the convex conjugate of $-\Phi$.

3.4. Hydrodynamic limit on \mathbb{Z} . In this subsection, we consider the ZRP on \mathbb{Z} . Let $\mathcal{M}_+(\mathbb{R})$ be the space of non-negative measures on \mathbb{R} endowed with the weak topology.

We consider $(\eta_{N,0})_N \in \mathbb{N}^{\mathbb{Z}}$ a sequence of configurations and we define the empirical measure associated with the process $\eta_{N,t}$:

$$\rho_{N,t}(dy) = \frac{1}{N} \sum_{i \in \mathbb{Z}} \eta_{N,t}(i) \delta_{i/N}(dy).$$

Note that we extended the empirical measure to the hole line \mathbb{Z} .

Theorem 5 (Hydrodynamic limit on \mathbb{Z}). *Given $u_0 \in \mathcal{M}_+(\mathbb{R})$ with a finite mass, we consider the Barron–Jensen viscosity solution to*

$$\begin{cases} \partial_t U + \Phi(\partial_x U) = 0, & t, x \in \Omega \\ U(0, \cdot) = U_0, \end{cases} \quad (14)$$

where U_0 is the cumulative distribution function of u_0 . Let $(\rho_N(0, dy))_N$ be a sequence weakly convergent in law to $u_0(dx)$.

We assume that there exists $A > 0$ such that:

$$\lim_{N \rightarrow \infty} \mathbb{P}(\rho_N(0, \mathbb{R} \setminus [-A, A]) = 0) = 1. \quad (15)$$

Then, the sequence $(t \rightarrow \rho_{N, \frac{Nt}{p-q}})_N$ converge in law in $\mathbb{D}(\mathbb{R}_+, \mathcal{M}_+(\mathbb{R}))$ to a measure-valued function $t \rightarrow u(t, dx)$. More over, for each $t \geq 0$, the c.d.f of $(u_t(dx))$ is $(U_t)^*$, the u.s.c envelope of U_t .

Rezakhanlou [30] has proved the hydrodynamic limit when the initial configuration is sampled according to a local equilibrium $\otimes_{i \in \mathbb{Z}^{\nu}} (u_{i,N})$ where

$$\lim_{N \rightarrow \infty} \int_{|x| \leq \ell} |u_{[xN],N} - u_0(x)| dx = 0$$

and u_0 has a bounded density with respect to the Lebesgue measure. Actually, in [30], g is assumed to be bounded. We underline that one can adapt the proof [18, Theorem 0.3, Chapter 8] to generalise the result to non-bounded g when $(\rho_{N,0})$ are compactly supported. Our theorem generalizes the hydrodynamic limit to irregular initial data, as we want to consider the macroscopic behaviour for $u_0 = \delta_0$.

Let us denote \mathcal{Q}_N the law of $t \mapsto \rho_{N, \frac{Nt}{p-q}}$. The proof is based on a tightness and uniqueness argument.

Lemma 5 (Tightness). *The sequence $(\mathcal{Q}_N)_N$ is tight. Moreover, if \mathcal{Q} is any limit point of $(\mathcal{Q}_N)_N$, then for \mathcal{Q} almost every u , u is weakly continuous in t .*

The proof of this lemma is a standard argument similar to the tightness proof on the segment (Lemma 6) and we omit it.

Proof of Theorem 5. Let \mathcal{Q}^* be a limit point of \mathcal{Q}_N . The idea is to trap the process η between two processes with more regular initial data. For any $\varepsilon > 0$, we introduce the following functions

$$\begin{aligned} \underline{U}_0^\varepsilon(x) &= \inf_y \left\{ U_0(y) + \frac{|x - \varepsilon - y|^2}{\varepsilon^2} - \varepsilon \right\}_+, \\ \bar{U}_0^\varepsilon(x) &= \iota_\varepsilon(x) + \sup_y \left\{ U_0(y) - \frac{|x + \varepsilon - y|^2}{\varepsilon^2} \right\}. \end{aligned}$$

where ι_ε is a smooth function equal to 0 on $(-\infty, -A - 2]$ and ε on $[-A - 1, \infty)$. We denote their associated unique viscosity solution by $\underline{U}^\varepsilon$ and \bar{U}^ε . We also denote by

$$\inf_\varepsilon \bar{U}^\varepsilon = \bar{U} \quad \text{et} \quad \sup_\varepsilon \underline{U}^\varepsilon = \underline{U}$$

For any $\varepsilon > 0$, $\underline{U}_0^\varepsilon$ and \bar{U}_0^ε are Lipschitz, non-decreasing, bounded, and converge to zero when x goes to $-\infty$. We can thus consider two ZRP with the following initial configurations:

$$\bar{\xi}_0^\varepsilon \sim \otimes_{i \in \mathbb{Z}^{\nu}} \left(N \left[\bar{U}_{0, \frac{i}{N} + \frac{1}{2N}}^\varepsilon - \bar{U}_{0, \frac{i}{N} - \frac{1}{2N}}^\varepsilon \right] \right) \quad \text{and} \quad \underline{\xi}_0^\varepsilon \sim \otimes_{i \in \mathbb{Z}^{\nu}} \left(N \left[\underline{U}_{0, \frac{i}{N} + \frac{1}{2N}}^\varepsilon - \underline{U}_{0, \frac{i}{N} - \frac{1}{2N}}^\varepsilon \right] \right).$$

By [30, Theorem 1.3], the hydrodynamic limit of $\underline{\xi}^\varepsilon$ (resp. $\bar{\xi}^\varepsilon$) corresponds to the solution of HJ with initial data $\underline{U}_0^\varepsilon$ (resp. \bar{U}_0^ε).

Let us denote by \mathcal{E}_N , the event where for any $x \in \mathbb{R}$,

$$\begin{aligned} U_0 \left(x - \frac{\varepsilon}{2} \right) - \frac{\varepsilon}{2} &\leq h(Nx) \leq U_0 \left(x + \frac{\varepsilon}{2} \right) + \frac{\varepsilon}{2}; \\ \bar{h}_0^\varepsilon(Nx) &\geq \bar{U}_0^\varepsilon \left(x - \frac{\varepsilon}{2} \right) - \frac{\varepsilon}{2}; \\ \underline{h}_0^\varepsilon(Nx) &\leq \underline{U}_0^\varepsilon \left(x + \frac{\varepsilon}{2} \right) + \frac{\varepsilon}{2}. \end{aligned}$$

By using the Levy distance, we obtain that $\mathcal{Q}_N(\mathcal{E}_N)$ converges to 1. Thanks to the choice of $\bar{U}_0^\varepsilon, \underline{U}_0^\varepsilon$ and the assumption (15), we obtain on the event \mathcal{E}_N :

$$\underline{h}_0^\varepsilon \leq h_0 \leq \bar{h}_0^\varepsilon.$$

A similar coupling as (\mathcal{E}) ensures for any $t \geq 0, \varepsilon > 0$, and almost surely on \mathcal{E}_N ,

$$\underline{h}_t^\varepsilon \leq h_t \leq \bar{h}_t^\varepsilon.$$

By using the hydrodynamic limit of $\underline{\xi}^\varepsilon$ and $\bar{\xi}^\varepsilon$, we obtain for any $t \geq 0$,

$$\mathcal{Q}^* \text{-a.s.}, \quad \forall x \in \text{Cont}(U_t), \quad \underline{U}_t^\varepsilon \leq U_t \leq \bar{U}_t^\varepsilon \quad (16)$$

where $\text{Cont}(U_t)$ are the continuity point of U_t . By making ε go to 0, we have for any $t \geq 0$:

$$\mathcal{Q}^* \text{-a.s.}, \quad \forall x \in \text{Cont}(U_t), \quad \underline{U}_t \leq U_t \leq \bar{U}_t. \quad (17)$$

By Proposition 1, $\underline{U}(t, x) = \bar{U}_*(t, x)$ on their continuity point. Thus, we obtain $\forall t \geq 0$,

$$\mathcal{Q}^* \text{-a.s.}, \quad \forall x \in \text{Cont}(U_t), \quad U_t = \underline{U}_t$$

which uniquely determines \mathcal{Q}^* . \square

4. HYDRODYNAMIC LIMIT ON THE SEGMENT

Let $(\eta_{N,0}) \in \Omega_{N,k}$ such that the associated empirical measure $(\rho_{N,0})$ weakly converge to $u_0 \in \mathcal{M}_+([0, 1])$. The proof is also a tightness–uniqueness argument. We denote by \mathcal{Q}_N the law of $t \rightarrow \rho_{N, \frac{N}{p-q}t}$.

Lemma 6 (Tightness). *The sequence $(\mathcal{Q}_N)_N$ is tight. Moreover, if \mathcal{Q} is any limit point of $(\mathcal{Q}_N)_N$, then for \mathcal{Q} -almost every u , u is weakly continuous in t .*

The main argument is the point-wise convergence of the cumulative distribution functions.

Proposition 2 (Concentration on BJ solution). *Given $(\eta_{N,0}) \in \Omega_{N,k}$ such that the associated empirical measure $\rho_{N,0}$ weakly converge to $u_0 \in \mathcal{M}_+([0, 1])$, we consider U the unique Barron–Jensen viscosity solution associated to (HJ) where U_0 is the c.d.f of u_0 .*

For any $t \geq 0, \delta > 0$ and $y \in (0, 1) \cap \text{Cont}(U_t)$, we obtain

$$\lim_{N \rightarrow \infty} \mathbb{P} \left(\left| \frac{1}{N} h_{N, \frac{Nt}{p-q}}(yN) - U(t, y) \right| \geq \delta \right) = 1.$$

The proof of Lemma 6 and Proposition 2 are deferred to the next subsections.

Proof of Theorem 1. Given \mathcal{Q}^* a limit point of \mathcal{Q}^N , the previous lemma implies that for any $t \geq 0$,

$$\mathcal{Q}^* \text{-a.s.}, \quad y \in (0, 1) \cap \text{Cont}(U_t), \quad \int_{-\infty}^y \pi_t^*(dy) = U(t, y)$$

The total number of particles is conserved. Thus, for any $t \geq 0$,

$$\mathcal{Q}^* \text{-a.s.}, \quad \pi_t^*([0, 1]) = \alpha$$

which uniquely determine \mathcal{Q}^* . \square

4.1. Proof of Lemma 6: By [18, Proposition 1.7], it is sufficient to show that for any $J \in \mathcal{C}^\infty([0, 1])$, the sequence $(t \rightarrow \langle \rho_{N, \frac{N}{p-q}t}, J \rangle)$ is tight. Since the total mass of the empirical measure is bounded, it remains to prove that for any $T > 0$:

$$\overline{\lim}_{\delta \downarrow 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{E} \left[\sup_{s, t \leq T, |t-s| \leq \delta} \left| \langle \rho_{N, \frac{N}{p-q}t} - \rho_{N, \frac{N}{p-q}s}, J \rangle \right| \right] = 0 \quad (18)$$

Let $f(\eta) = \frac{1}{N} \sum_{x=1}^N J\left(\frac{x}{N}\right) \eta(x)$. We consider the associated martingale $(M_{N,t})$ such that for any $t \geq 0$

$$f\left(\eta_{Nt, \frac{Nt}{p-q}}\right) - f\left(\eta_{N, \frac{Ns}{p-q}}\right) = \int_{\frac{Ns}{p-q}}^{\frac{Nt}{p-q}} \mathcal{A}_N f(\eta_{N,r}) dr + M_{N, \frac{Nt}{p-q}} - M_{N, \frac{Ns}{p-q}}. \quad (19)$$

A direct computation shows that

$$\begin{aligned}
& \left| \int_{\frac{Ns}{p-q}}^{\frac{Nt}{p-q}} \mathcal{A}_N f(\eta_{N,r}) dr \right| \\
&= \left| \int_{\frac{N}{p-q}t}^{\frac{Nt}{p-q}} \frac{1}{N} \sum_{x=1}^{N-1} (pg(\eta_r(x)) - qpg(\eta_r(x+1))) \left(J\left(\frac{x+1}{N}\right) - J\left(\frac{x}{N}\right) \right) dr \right| \\
&\leq \|\nabla J\|_\infty g^* \delta \frac{k}{N}.
\end{aligned} \tag{20}$$

On the other hand, we obtain by Doob inequality:

$$\begin{aligned}
\mathbb{E} \left(\sup_{s \leq T} M_{N, \frac{Ns}{p-q}}^2 \right) &\leq \mathbb{E} \left(M_{N, \frac{NT}{p-q}}^2 \right) \\
&= \mathbb{E} \left(\int_0^{\frac{NT}{p-q}} \mathcal{A}_N f(\eta_{N,r})^2 - 2f(\eta_{N,r}) \mathcal{A}_N f(\eta_{N,r}) \right) \\
&= \frac{1}{N^2} \mathbb{E} \left(\int_0^{\frac{NT}{p-q}} \sum_{x=1}^{N-1} (pg(\eta_r(x)) + qg(\eta_r(x+1))) \left(J\left(\frac{x+1}{N}\right) - J\left(\frac{x}{N}\right) \right)^2 \right) \\
&\leq \frac{kC_{J,T,g}}{(p-q)N^3}.
\end{aligned} \tag{21}$$

Clearly, (20) and (21) imply (18). \square

4.2. Proof of Proposition 2 (Lower bound). The idea is to compare the process on the segment to the process on \mathbb{Z} through two different couplings.

Given $t^*, \delta, \varepsilon > 0$, we fix $R = \frac{6g^*t^*}{p-q}$. We denote by C the constant given by Lemma 2 for $a = 0$ and $b = +\infty$. As the macroscopic comparisons are valid only over a period of $\frac{NC\varepsilon}{(p-q)}$, we introduce a time subdivision $(t_i)_{i \in \mathbb{N}} = (\frac{C\varepsilon}{2}i)_{i \in \mathbb{N}}$. For convenience, let us denote $(T_i)_{i \in \mathbb{N}} = (\frac{CN\varepsilon}{2(p-q)}i)_{i \in \mathbb{N}}$.

We introduce a first process $\tilde{\eta}$ on the subgraph \mathbb{N} started at $\tilde{\eta}_0 = \eta(\cdot - \lfloor 2\varepsilon N \rfloor)$ which we couple with the ZRP process η on the segment following (\mathcal{C}) . For any $t \geq 0$,

$$\tilde{h}_{N,t} \leq h_{N,t}, \quad \text{a.s.} \tag{22}$$

Then, we define $\hat{\eta}$ a ZRP on \mathbb{Z} coupled with $\tilde{\eta}$ according to (\mathcal{C}) on $[T_i, T_{i+1})$ and updated at each T_i by imposing $\hat{\eta} = \tilde{\eta}$. We claim that for any $t \leq 2t^*$, the sequence $(\hat{\pi}(t, dy))_N$ converge in law for the weak topology to the measure $\partial_x[U_t^*(\cdot - 2\varepsilon)]$.

Thanks to Theorem 5, it is sufficient to prove the claim for each update $t_i \leq 2t^*$. We argue by induction and we assume that the claim holds at t_{i-1} .

Let us denote $\mathcal{E} = \{\forall s \leq t^*, \tilde{h}(s, RN) = k\}$. Thanks to Lemma 1, we can couple $\tilde{\eta}$ with k independent particles jumping only to the right with rate pg^* . Thus,

$$\mathbb{P}(\mathcal{E}) \geq 1 - \mathbb{P} \left(\max_{i=1, \dots, k} X_i \geq RN \right),$$

where X_i are independent Poisson variables with parameters $\frac{2g^*Nt^*}{p-q}$. Lemma 13 implies that r.h.s term converges to 1.

Given a bounded continuous test function J , we obtain on the event \mathcal{E} by the triangular inequality:

$$\left| \frac{1}{N} \sum_{x \in \mathbb{Z}} J\left(\frac{x}{N}\right) \hat{\eta}([T_i]^+, x) - \int J(x + 2\varepsilon) \partial_x U_{t_i}^*(dx) \right| \leq A_{i,N} + B_{i,N} + C_{i,N} + D_{i,N}, \quad (23)$$

where

$$\begin{aligned} A_{i,N} &= \frac{2\|J\|_\infty}{N} \sum_{x=\varepsilon N}^{RN} |\tilde{\eta}(T_i, x) - \hat{\eta}([T_i]^-, x)| \\ B_{i,N} &= \frac{k}{N} \sup_{x \leq \varepsilon} |J(x) - J(0)| \\ C_{i,N} &= \left| \frac{1}{N} \sum_{x=\varepsilon N}^{RN} J(x/N) \hat{\eta}([T_i]^-, x) - \int_{-\varepsilon}^{R-2\varepsilon} J(x + 2\varepsilon) \partial_x U_{t_i}^*(dx) \right| \\ D_{i,N} &= J(0) \left| \frac{1}{N} \sum_{x=\varepsilon N}^{RN} \hat{\eta}([T_i]^-, x) - \frac{k}{N} \right|. \end{aligned}$$

First, $(A_{i,N})$ converges to 0 in expectation by Lemma 2. The second term vanishes by continuity of J in 0. By the induction assumption, $(C_{i,N})$ and $(D_{i,N})$ converge in probability to zero which concludes the proof of the claim.

Consequently, for any $t \leq 2t^*$ and $x \in \mathbb{R}$ such that $x - 2\varepsilon \in \text{Cont}(U_t)$:

$$\begin{aligned} \mathbb{P} \left(\left| \tilde{h}_{\frac{Nt}{p-q}}(Nx) - U(t, x - 2\varepsilon) \right| \geq \frac{\delta}{2} \right) &\leq \mathbb{P}(\mathcal{E}^c) + \frac{16}{\varepsilon} \mathbb{E} \left[\frac{1}{N} \sum_{x \in \llbracket \varepsilon N, RN \rrbracket} |\tilde{\eta}_{\frac{Nt}{p-q}}(x) - \hat{\eta}_{\frac{Nt}{p-q}}(x)| \right] \\ &+ \mathbb{P} \left(\left| \sum_{j=\varepsilon N}^{RN} \hat{\eta}_{\frac{Nt}{p-q}}(j) - \alpha \right| \geq \frac{\delta}{8} \right) + \mathbb{P} \left(\left| \hat{h}_{\frac{Nt}{p-q}}(Nx) - U(t, x - 2\varepsilon) \right| \geq \frac{\delta}{8} \right). \end{aligned}$$

where \mathcal{E}^c the complementary event. The third and fourth right-hand side terms converge to zero thanks to the hydrodynamic limit of $\hat{\eta}$. Lemma 2 guarantees the convergence of the second term. Thus,

$$\mathbb{P} \left(\left| \tilde{h}_{\frac{Nt}{p-q}}(Nx) - U(t, x - 2\varepsilon) \right| \geq \frac{\delta}{2} \right) \xrightarrow{N \rightarrow \infty} 0.$$

By (22), we obtain

$$\lim_{N \rightarrow \infty} \mathbb{P} \left(\frac{1}{N} h_{N,t}(xN) \geq U(t, x - 2\varepsilon) - \frac{\delta}{2} \right) = 1.$$

Finally, $\text{Cont}(U_t)$ is dense and for any $x \in \text{Cont}(U_t)$, we can choose $\varepsilon > 0$ such that

$$x - 2\varepsilon \in \text{Cont}(U_t) \quad \text{and} \quad U(t, x - 2\varepsilon) \geq U(t, x) - \frac{\delta}{2}.$$

Thus,

$$\lim_{N \rightarrow \infty} \mathbb{P} \left(\frac{1}{N} h_{N,t}(xN) \geq U(t, x) - \delta \right) = 1, \quad \forall \delta > 0.$$

□

4.3. Proof of Proposition 2 (Upper bound). The proof follows a similar approach to the lower bound proof, but we need to be cautious at the boundary at N . Thanks to Lemma 1, it is sufficient to prove the same result on the subgraph $\llbracket -\infty, N \rrbracket$.

Lemma 7. *Let $\tilde{\eta}_N$ a ZRP on $\llbracket -\infty, N \rrbracket$ started at $\eta_{N,0}$. For any $t, \delta > 0$ and $y \in (0, 1) \cap \text{Cont}(U_t)$:*

$$\lim_{\substack{N \rightarrow \infty \\ k/N \rightarrow \alpha}} \mathbb{P} \left(\frac{1}{N} \tilde{h}_{N, \frac{Nt}{p-q}}(yN) \leq U(t, y) + \delta \right) = 1.$$

Proof of Lemma 7. Given $\delta, t^* > 0$ and $\varepsilon > 0$. We fix $R = \frac{6g^*t^*}{p-q}$ and C the constant provided by the Lemma 2 for $a = -\infty$ and $b = 1$. As above, we introduce a time subdivision $(t_i)_{i \in \mathbb{N}} = (\frac{C\varepsilon}{2}i)_{i \in \mathbb{N}}$ and $(T_i)_{i \in \mathbb{N}} = (\frac{CN\varepsilon}{2(p-q)}i)_{i \in \mathbb{N}}$.

Whenever we use Lemma 2, we lose information about the density profile between $(1 - \varepsilon)N$ and N . Unlike the lower bound, this part may contain a non-negligible number of particles. To get around this, we introduce a delayed version at each t_i where we gather the particles at a well-chosen point before $N(1 - \varepsilon)$. This is still sufficient as we are looking for an upper bound. More precisely, let us define the function V^d by:

- For any $t \in [t_0, t_1)$, $V_t^d = U_t$ where U_t is the Barron–Jensen viscosity solution to (HJ) with U_0 as initial condition.
- For any $i \in \mathbb{N}$, there exists $y_i \in \text{Cont } V^d(t_i^-) \cap (1 - 2\varepsilon, 1 - \varepsilon)$

$$\begin{cases} V^d(x, t_i) = V^d(x, t_i^-) & \text{if } x \leq y_i, \\ V^d(x, t_i) = \alpha & \text{if } x > y_i. \end{cases}$$

For any $t \in (t_i, t_{i+1}]$, V^d is equal to the Barron–Jensen viscosity solution started at $V^d(t_i)$.

As the updates at each t_i affect only the function on a support $[1 - 2\varepsilon, +\infty)$, Lemma 4 implies that $V^d = U$ for any $t \geq 0$ and $x \leq 1 - 4\varepsilon$.

We also define the microscopic analog $\tilde{\eta}^d$ to V^d on $\llbracket -\infty, N \rrbracket$

- For any $t \in [0, T_1)$, $\tilde{\eta}^d$ is set to be equal to $\tilde{\eta}$.
- For any $i \in \mathbb{N}$, conditionally on $\sigma(\tilde{\eta}(s), s \leq T_i)$

$$\begin{cases} \tilde{\eta}^d(x, T_i) = \tilde{\eta}^d(x, T_i^-) & \text{if } x < \lfloor Ny_i \rfloor, \\ \tilde{\eta}^d(x, T_i) = \sum_{y=\lfloor Ny_i \rfloor}^N \tilde{\eta}^d(y, T_i^-) & \text{if } x = \lfloor Ny_i \rfloor, \\ \tilde{\eta}^d(x, T_i) = 0 & \text{if } x > \lfloor Ny_i \rfloor. \end{cases}$$

This modification delays the evolution. More precisely, $\tilde{h}^d(T_i^+, \cdot) \geq \tilde{h}(T_i^+, \cdot)$. For any $t \in (T_i, T_{i+1}]$, both processes evolve according to (\mathcal{C}) to respect the order of height functions.

We finally introduce a ZRP $\hat{\eta}^d$ evolving on \mathbb{Z} coupled with $\tilde{\eta}^d$ according to \mathcal{C} and updated on each T_i as for $\hat{\eta}^d$. We claim that

Lemma 8. *For any $t \leq 2t^*$, the sequence $\left(\hat{\pi}^d \left(\frac{Nt}{p-q}, dy \right) \right)_N$ converges in law (in the weak topology) to the measure $\partial_x V_t^*$.*

We defer the proof of this lemma to the end of this sub-section. For any $t \leq 2t^*$, $x \in \text{Cont}(U_t) \cap (-\infty, 1 - 4\varepsilon)$:

$$\begin{aligned} \mathbb{P} \left(\left| \tilde{h}_{\frac{Nt}{p-q}}^d(Nx) - U(t, x) \right| \geq \delta \right) &\leq \mathbb{P} \left(\tilde{h}_{\frac{Nt}{p-q}}^d(-RN) \geq \frac{\delta N}{3} \right) \\ &+ \frac{3}{\delta} \mathbb{E} \left[\frac{1}{N} \sum_{j \in \llbracket -RN, N(1-\varepsilon) \rrbracket} \left| \tilde{\eta}_{\frac{Nt}{p-q}}^d(j) - \hat{\eta}_{\frac{Nt}{p-q}}^d(j) \right| \right] + \mathbb{P} \left(\left| \hat{\eta}_{\frac{Nt}{p-q}}^d(\llbracket -RN, Nx \rrbracket) - U(t, x) \right| \geq \frac{\delta}{3} \right). \end{aligned}$$

The second right-hand side term vanishes as N goes to infinity thanks to Lemma 2 and Lemma 8 implies the convergence to zero of the third term.

By Lemma 1, we can couple the process $\tilde{\eta}^d$ with a process of k independent particles that jumps only to the left with a rate qg^* . Thanks to this coupling, we obtain for any $s \leq \frac{2t^*}{p-q}$,

$$\mathbb{P} \left(\tilde{h}_{sN}^d(-RN) \geq 1 \right) \leq \mathbb{P} \left(\max_{i=1, \dots, k} X_i \geq RN \right), \quad (24)$$

where $(X_i)_{i=1}^k$ are independent Poisson variables with parameter $\frac{2qg^*Nt^*}{p-q}$. Lemma 13 implies for any $t \geq 0$ and $x \in \text{Cont}(U_t) \cap (-\infty, 1 - 4\varepsilon)$:

$$\lim_{N \rightarrow \infty} \mathbb{P} \left(\left| \tilde{h}_{\frac{Nt}{p-q}}^d(Nx) - U(t, x) \right| \geq \delta \right) = 0.$$

Since $\tilde{h}^d \geq \tilde{h}$, we conclude the proof, up to Lemma 8. \square

We turn to prove the last lemma.

Proof of Lemma 8. We first recall that for any $i \leq i^*$, $\{\tilde{h}_{T_i^+}^d(-RN) \geq 1\}$ occurs with probability converging to one. Thanks to Theorem 5, it remains to prove the convergence for each update $t_i \leq 2t^*$.

As above, we argue by induction and we assume that the lemma holds at t_{i-1} . Given a bounded continuous test function J , we obtain almost surely on $\{\tilde{h}_{T_i^+}^d(-RN) \geq -RN\}$:

$$\left| \frac{1}{N} \sum_{x \in \mathbb{Z}} J\left(\frac{x}{N}\right) \hat{\eta}^d(T_i, x) - \int J(x) \partial_x V_{t_i}^*(dx) \right| \leq A_{i,N} + B_{i,N} + C_{i,N} + D_{i,N}, \quad (25)$$

where

$$\begin{aligned} A_{i,N} &= \frac{2\|J\|_\infty}{N} \sum_{x \in \llbracket -RN, N(1-\varepsilon) \rrbracket} \left| \tilde{\eta}^d([T_i]^- , x) - \hat{\eta}^d([T_i]^- , x) \right| \\ B_{i,N} &= \left| J\left(\frac{\lfloor Ny_i \rfloor}{N}\right) \frac{k}{N} - J(y_i) \alpha \right| \\ C_{i,N} &= \left| \frac{1}{N} \sum_{x=-NR}^{y_i N} J(x/N) \hat{\eta}^d([T_i]^- , x) - \int_{-R}^{y_i} J(x) \partial_x V_{t_i}^*(dx) \right| \\ D_{i,N} &= \left| \frac{J\left(\frac{\lfloor Ny_i \rfloor}{N}\right)}{N} \sum_{x=-NR}^{y_i N} \hat{\eta}^d([T_i]^- , x) - J(y_i) \int_{-R}^{y_i} \partial_x V_{t_i}^*(dx) \right|. \end{aligned}$$

First, $(A_{i,N})$ converges to 0 in expectation by Lemma 2. The second term converges to 0 by continuity of J . By the induction assumption and Theorem 5, we obtain the convergence in

law of $(\hat{\pi}^d (T_i^-, dy))_N$ to $\partial_x V_{t_i^-}$. Thus, $(C_{i,N})$ and $(D_{i,N})$ converge to 0 in probability as $V_{t_i^-}$ is continuous in y_i and $-R$. \square

5. PROOF OF LOWER AND UPPER BOUNDS ON THE MIXING TIME

In this section, we prove the main theorems on the mixing time.

Let us recall that $\wedge_{N,k} = k\delta_1$ and $\vee_{N,k} = k\delta_N$. For the sake of clarity, we will omit the indices when it is clear from the context. As mentioned, \wedge corresponds to the configuration farthest from the equilibrium. By Theorem 1, its macroscopic density evolution is described by U^α the unique Barron–Jensen solution of (HJ) and its macroscopic equilibrium time $T_{eq,p,\alpha}$ corresponds to the mixing time.

5.1. Lower bound of the mixing time. The proof of the lower bound relies on the hydrodynamic limit of the ZRP on the segment and uses the position of the left-most particle as a distinctive statistic.

Given a configuration $\eta \in \Omega_{N,k}^0$, we denote by $\ell(\eta)$ the position of the left-most particle. The following lemma controls $\ell(\eta)$ when η is sampled according to the equilibrium measure $\pi_{N,k}$. The proof is the analogue of [19, Lemma 11] and [25, Proposition 11].

Lemma 9 (Concentration of $\ell(\eta)$). *There exists C_g depending on g, p such that for any $N, k, \delta \in \mathbb{N}$:*

$$\pi_{N,k}(\ell(\eta) \leq N - \delta) \leq C_g g(k) \left(\frac{p}{q}\right)^{-\delta}. \quad (26)$$

Proof. We observe that

$$\frac{\pi_{N,k}(\eta^{i,N})}{\pi_{N,k}(\eta)} = \left(\frac{p}{q}\right)^{N-i} \frac{g(\eta(i))}{g(\eta(N) + 1)} \geq \frac{g(1)}{g(k)} \left(\frac{p}{q}\right)^{N-i},$$

where $\eta^{i,N}$ is the configuration obtained by moving a particle from a position i is moved to the position N . Thus,

$$\pi_{N,k}(\ell(\eta) = i) = \sum_{\eta, \ell(\eta)=i} \pi_{N,k}(\eta) \leq \frac{g(k)}{g(1)} \left(\frac{p}{q}\right)^{-(N-i)} \sum_{\eta, \ell(\eta)=i} \pi_{N,k}(\eta^{i,N}) \leq \frac{g(k)}{g(1)} \left(\frac{p}{q}\right)^{-(N-i)}.$$

We obtain the lemma by summing over $i \leq N - \delta$. \square

We can now prove Theorem 2.

Proof of Theorem 2. As mentioned above, it is natural to consider as initial condition \wedge . We obtain for any $\delta > 0$,

$$\begin{aligned} d_{N,k}((T_{eq,p,\alpha} - \delta)N) &\geq \|P_{(T_{eq,p,\alpha} - \delta)N}^\wedge - \pi_{N,k}\|_{\text{TV}} \\ &\geq \mathbb{P}_{(T_{eq,p,\alpha} - \delta)N}^\wedge(\ell(\eta) \leq N - \varepsilon N) - \pi_{N,k}(\ell(\eta) \geq N - \varepsilon N). \end{aligned}$$

where P_t^\wedge is the law of the process started at \wedge . Since

$$T_{eq,p,\alpha} = \frac{1}{p-q} \inf\{t > 0, U^\alpha(t, dx) \leq \alpha \mathbf{1}_{x>1}\}, \quad (27)$$

the hydrodynamic limit on the segment (Theorem 1) implies that for any $\delta > 0$, there exists $\varepsilon \in (0, 1)$,

$$\lim_{\substack{N \rightarrow \infty \\ k/N \rightarrow \alpha}} \mathbb{P}_{(T_{eq,p,\alpha} - \delta)N}^\wedge(\ell(\eta) \leq N - \varepsilon N) = 1.$$

Lemma 9 concludes the proof. \square

Remark 7. *Let us emphasise that the same method would apply for g of polynomial growth if the hydrodynamic limit holds.*

5.2. Upper bound: totally asymmetric jumps and convex flux. Let us first observe that as $p = 1$, the system is mixed when it reaches the configuration \vee . Thanks to the attractivity, we have in particular

$$\forall t \geq 0, d_{N,k}(t) = P_t^\wedge (h(N-1) > 0).$$

On the other hand, the Hopf–Lax formula written (12) as:

$$U^\alpha(x, t) = \inf \left\{ \inf_{z \leq \frac{x}{t}} [\alpha + t\Psi(z)]; \inf_{z \geq \frac{x}{t}} [t\Psi(z)] \right\}$$

implies that the macroscopic equilibrium is reached at

$$T_{eq,1,\alpha} = \frac{1}{\Phi'(0)} = \frac{1}{g(1)}.$$

As $T_{eq,1,\alpha}$ depends only on the value of g in 1, the idea is to couple through (\mathcal{C}) with the linear rate function, the simplest and slowest convex rate function.

We consider a ZRP (η^{ind}) started at \wedge and a linear rate function $g_{\text{ind}} : x \rightarrow g(1)x$. Since $g \geq g_{\text{ind}}$, we couple both processes according to (\mathcal{C}) to obtain almost surely for any $t > 0$:

$$h_t^{\text{ind}} \geq h_t.$$

We recall that ZRP with a linear rate corresponds to independent random walks. Consequently, we obtain for any $t \geq 0$,

$$d_{N,k}(t) = \mathbb{P}(h_t(N-1) > 0) \leq \mathbb{P}(h_t^{\text{ind}}(N-1) > 0) = 1 - \mathbb{P}(\mathcal{P}(tg(1)) \geq N)^k.$$

If t is fixed to $N \left(\frac{1}{g(1)} + \varepsilon \right)$, the limit is zero, which yields the first assertion of Theorem 3.

5.3. Upper-bound: Concave flux. In this subsection, we will work within the framework of Theorem 3.2. The main idea is to show that the leftmost particle and the number of particles at position N approach their equilibrium state around $NT_{eq,p,\alpha}$. Once there, the system will mix rapidly. Let us denote by:

$$L_{N,k} = \ell(\wedge_t), \quad \text{and} \quad S_{N,k} = \wedge_t(N), \quad (28)$$

where $(\wedge_t)_t$ is a ZRP (g, p) started from the configuration \wedge . We also introduce their macroscopic analogue:

$$\begin{aligned} \forall t \geq 0, \quad \mathcal{L}_p^\alpha(t) &= \inf \{x \in \mathbb{R}, U^\alpha((2p-1)t, x) > 0\}, \\ \mathcal{S}_p^\alpha(t) &= \alpha - U^\alpha((2p-1)t, 1^-). \end{aligned} \quad (29)$$

The macroscopic equilibrium time could be written as:

$$T_{eq,p,\alpha} = (\mathcal{L}_p^\alpha)^{-1}(1) = (\mathcal{S}_p^\alpha)^{-1}(\alpha).$$

The following lemma ensures that the microscopic behaviour corresponds to the expected macroscopic behaviour.

Proposition 3 (Macroscopic behaviour of $L_{N,k}$ and $S_{N,k}$). *For any $t \leq T_{eq,p,\alpha}$, we have the following convergence in probability:*

$$\frac{1}{N} S_{N,k}(Nt) \xrightarrow[k/N \rightarrow \alpha]{N \rightarrow \infty} \mathcal{S}_p^\alpha(t) \quad \text{and} \quad \frac{1}{N} L_{N,k}(Nt) \xrightarrow[k/N \rightarrow \alpha]{N \rightarrow \infty} \mathcal{L}_p^\alpha(t).$$

The proof of this lemma is deferred to Section 6. As in the case of the exclusion process [19, Proposition 9], we emphasise that the hydrodynamic limit of (\wedge_t) provides only the upper-bound on $L_{N,k}$ and $S_{N,k}$: For any $\delta > 0$

$$\forall t \leq T_{eq,p,\alpha}, \quad \lim_{N \rightarrow \infty} \mathbb{P} \left(\frac{S_{N,k}(Nt)}{N} \leq \mathcal{L}_p^\alpha(t) + \delta \text{ and } \lim_{N \rightarrow \infty} \frac{L_{N,k}(Nt)}{N} \leq \mathcal{L}_p^\alpha(t) + \delta \right) = 1.$$

Let us focus on the proof of the second statement of 3.

Proof of Theorem 3.2. By triangular inequality, we have for any $t \geq 0$:

$$d_{N,k}(t) \leq 2 \max_{h,h'} \left\| P_t^h - P_t^{h'} \right\|_{\text{TV}}.$$

The attractivity of the ZRP permits a first simplification. Indeed, starting for any h, h' , we can trap their evolution between (\wedge_t) and (\vee_t) thanks to a similar coupling to (\mathcal{C}) :

$$d_{N,k}(t) \leq 2 \max_{h,h'} \left\| P_t^h - P_t^{h'} \right\|_{\text{TV}} \leq 2\mathbb{P}(\wedge_t \neq \vee_t).$$

Then, once the process (\wedge_t) reaches the minimal state \vee , the processes $(\wedge_s)_s$ and $(\vee_s)_s$ become identical:

$$d_{N,k}(t) \leq 2\mathbb{P}(\forall s \leq t, \wedge_s \neq \vee).$$

For any $\varepsilon > 0$, we define

$$\mathcal{E}_{N,k}^\varepsilon = \{ \eta \in \Omega_{N,k}^0, \ell(\eta) \geq N(1 - \varepsilon) \text{ and } \eta(N) \geq (1 - \varepsilon)k \}.$$

We also denote by \wedge^ε the maximal element of the set $\mathcal{E}_{N,k}^\varepsilon$. For any $\delta > 0$, we obtain

$$\begin{aligned} d_{N,k}(N [T_{eq,p,\alpha} + \delta]) &\leq 2\mathbb{P}(\forall t \leq N [T_{eq,p,\alpha} + \delta], \wedge_t \neq \vee) \\ &\leq 2\mathbb{P}(\forall t \leq N [T_{eq,p,\alpha} + \delta], \wedge_t \neq \vee \ \& \ \wedge_{NT_{eq,p,\alpha}} \in \mathcal{E}_{N,k}^\varepsilon) + 2\mathbb{P}(\wedge_{NT_{eq,p,\alpha}} \notin \mathcal{E}_{N,k}^\varepsilon) \\ &\leq 2\mathbb{P}(\forall t \leq \delta N, \wedge_t^\varepsilon \neq \vee) + 2\mathbb{P}(\wedge_{NT_{eq,p,\alpha}} \notin \mathcal{E}_{N,k}^\varepsilon), \end{aligned}$$

where (\wedge_t^ε) is a ZRP started at \wedge^ε .

Since $T_{eq,p,\alpha} = (\mathcal{L}_p^\alpha)^{-1}(1) = (\mathcal{L}_p^\alpha)^{-1}(\alpha)$, Proposition 3 implies that

$$\mathbb{P}(\wedge_{NT_{eq,p,\alpha}} \notin \mathcal{E}_{N,k}^\varepsilon) \xrightarrow[k/N \rightarrow \alpha]{N \rightarrow \infty} 0.$$

We compare the system to an exclusion process to control the last term. We denote by $\bar{g} = \max_k g(k)$. Let us introduce two ZRP $(\tilde{\wedge}_t^\varepsilon)$ (resp. $(\tilde{\vee}_t)$) with a modified rate function $\tilde{g} = (pg(1) + q\bar{g})\mathbf{1}_{\geq 1}$ and a bias $\tilde{p} = \frac{pg(1)}{pg(1) + q\bar{g}}$ started at \wedge^ε (resp. \vee). We couple these three processes with (\mathcal{C}) :

$$\forall t \geq 0, \quad h_{\tilde{\wedge}_t} \geq \max(h_{\wedge_t}, h_{\tilde{\vee}_t}) \quad \text{a.s.}$$

We obtain that:

$$\begin{aligned} \mathbb{P}(\forall t \leq \delta N, \wedge_t^\varepsilon \neq \vee) &\leq \mathbb{P}(\forall t \leq \delta N, \tilde{\wedge}_t^\varepsilon \neq \vee) \\ &\leq \mathbb{P}(\tilde{\wedge}_{\frac{\delta}{2}N}^\varepsilon \neq \tilde{\vee}_{\frac{\delta}{2}N}) + \mathbb{P}\left(\forall t \in \left[\frac{\delta}{2}N, \delta N\right], \tilde{\vee}_t \neq \vee\right). \end{aligned}$$

We recall that Lemma 3 implies that the processes $(E^{-1}(\tilde{\vee}_t))$ and $(E^{-1}(\tilde{\wedge}_t^\varepsilon))$ evolve according to an accelerated exclusion process. So, we can use the same argument as in [19, Theorem 2] and [20, Lemma 12] since we are reduced to an exclusion process.

In particular, there exists an increasing function $f_{N,k}$ on $\Omega_{N,k}$ and such that:

$$\mathbb{E}[f_{N,k}(h'_t) - f_{N,k}(h_t)] \leq e^{-\varepsilon t} (f_{N,k}(h') - f_{N,k}(h)),$$

where h, h' are ZRP (\tilde{g}, \tilde{p}) coupled according to (\mathcal{C}) and ϱ is a constant (see [20, Equation 45]). We obtain by Markov's inequality and [20, Lemma 12]:

$$\mathbb{P}\left(\tilde{\Lambda}_{\frac{\delta}{2}N}^\varepsilon \neq \tilde{V}_{\frac{\delta}{2}N}\right) \leq \frac{\mathbb{E}\left(f_{N,k}\left(h_{\tilde{\Lambda}_{\frac{\delta}{2}N}^\varepsilon}\right) - f_{N,k}\left(h_{\tilde{V}_{\frac{\delta}{2}N}}\right)\right)}{\min_{h \leq h'} [(f_{N,k}(h') - f_{N,k}(h))]} \leq CN \left[\frac{pg(1)}{q\bar{g}}\right]^{\varepsilon N} e^{-\frac{\delta}{2}\rho N}.$$

Thus, for a fixed $\delta > 0$, we can choose ε small enough to obtain:

$$\mathbb{P}\left(\tilde{\Lambda}_{\frac{\delta}{2}N}^\varepsilon \neq \tilde{V}_{\frac{\delta}{2}N}\right) \xrightarrow[k/N \rightarrow \alpha]{N \rightarrow \infty} 0.$$

Finally, the Markov inequality implies:

$$\mathbb{P}\left(\forall t \in \left[\frac{\delta}{2}N, \delta N\right], \tilde{V}_t \neq \vee\right) \leq \frac{2\mathbb{E}_\vee(\tilde{\tau}_\vee)}{\delta N} = \frac{2}{\delta N \tilde{\pi}_{N,k}(\vee)},$$

where $\tilde{\tau}_\vee$ an excursion's duration from the configuration \vee and $\tilde{\pi}_{N,k}$ the equilibrium measure of ZRP (\tilde{g}, \tilde{p}) . It is then sufficient to lower-bound $\tilde{\pi}_{N,k}(\vee)$ by a constant. For any $\eta \in \Omega_{N,k}^0$, we have

$$\tilde{\pi}_{N,k}(\vee) \geq \tilde{\pi}_{N,k}(\eta)$$

Thus, for any $\Delta > 0$,

$$\tilde{\pi}_{N,k}(\vee) \# \left\{ \eta \in \Omega_{N,k}^0, \ell(\eta) + \eta(N) \geq N + k - \Delta \right\} \geq \tilde{\pi}_{N,k}(\ell(\eta) + \eta(N) \geq N + k - \Delta)$$

Due to [19, Lemma 11], we can choose Δ independent from N, k such that

$$\tilde{\pi}_{N,k}(\ell(\eta) + \eta(N) \geq N + k - \Delta) \geq \frac{1}{2}.$$

We conclude by observing that

$$\# \left\{ \eta \in \Omega_{N,k}^0, \ell(\eta) + \eta(N) \geq N + k - \Delta \right\} \leq \sum_{i=1}^{\Delta} \binom{\Delta}{i}.$$

□

Remark 8 (Pre-Cutoff). We emphasise that by using the same approach, we obtain the pre-cutoff for asymmetric ZRP as long as $pg(1) > q\bar{g}$ and Φ strictly concave (see [26, Chapter 18] for more background on the Pre-Cutoff). Indeed, we can couple the ZRP to ZRP (\tilde{g}, \tilde{p}) where $\tilde{g} = (pg(1) + q\bar{g})\mathbb{1}_{\geq 1}$ and $\tilde{p} = \frac{pg(1)}{pg(1) + q\bar{g}}$. By using Lemma 3 and [19, Proposition 9], we have for any $\varepsilon > 0$:

$$\mathbb{P}\left(\wedge_{N\tilde{T}_{eq,p,\alpha}} \notin \mathcal{E}_{N,k}^\varepsilon\right) \xrightarrow[k/N \rightarrow \alpha]{N \rightarrow \infty} 0,$$

where $\tilde{T}_{eq,p,\alpha}$ is associated to ZRP (\tilde{g}, \tilde{p}) . Then, the same method yields for any $\theta > 0$,

$$T_{eq,p,\alpha} \leq \liminf_{\substack{N \rightarrow \infty \\ k/N \rightarrow \alpha}} \frac{T_{mix}^{N,k}(\theta)}{N} \leq \limsup_{\substack{N \rightarrow \infty \\ k/N \rightarrow \alpha}} \frac{T_{mix}^{N,k}(\theta)}{N} \leq \tilde{T}_{eq,p,\alpha}$$

6. EVOLUTION OF THE LEFT–MOST PARTICLE AND THE STACK AT \mathbb{N}

In the sequel, we prove Proposition 3 and assume that the flux Φ is strictly concave and bounded by \bar{g} . We recall that

$$L_{N,k} = \ell(\wedge_t) \quad \text{and} \quad S_{N,k} = \wedge_t(N) \quad (30)$$

where $(\wedge_t)_t$ is a ZRP(g, p) started from the configuration \wedge .

Let us first describe the behaviour of \mathcal{L}_p^α and \mathcal{S}_p^α . Thanks to the Hopf–Lax formula (13), we can explicitly compute U^α .

$$U^\alpha(t, x) = \begin{cases} \max(0, \alpha - t\Psi(-\frac{x}{t})) & \text{if } x \leq g(1)t, \\ \alpha & \text{if } x > g(1)t. \end{cases} \quad (31)$$

This implies that $\mathcal{L}_p^\alpha(t) = \alpha \mathcal{L}\left(\frac{(2p-1)t}{\alpha}\right)$ and $\mathcal{S}_p^\alpha(t) = \mathcal{S}((2p-1)t)$ where

$$\mathcal{L}(t) = \begin{cases} 0 & \text{if } t \leq \frac{1}{\Psi(0)}, \\ -t\Psi^{-1}\left(\frac{1}{t}\right) & \text{otherwise} \end{cases}, \quad \mathcal{S}(t) = \begin{cases} 0 & \text{if } t \leq \frac{1}{g(1)}, \\ t\Psi\left(-\frac{1}{t}\right) & \text{otherwise.} \end{cases} \quad (32)$$

These functions are continuous, increasing and strictly convex. Indeed, we can easily compute their derivatives:

$$\mathcal{L}'(t) = \frac{\Phi(\Psi'(\Psi^{-1}(1/t)))}{\Psi'(\Psi^{-1}(1/t))}, \quad \forall t \geq \frac{1}{\Psi(0)}, \quad \mathcal{S}'(t) = \Phi\left(\Psi'\left(\frac{-1}{t}\right)\right), \quad \forall t \geq \frac{1}{g(1)}.$$

The proof is now divided into two subsections: 6.1 for the lower–bound on $(L_{N,k})$ and 6.2 for the lower–bound on $(S_{N,k})$. Unlike for the exclusion process, we do not have a particle-hole symmetry, so we need to prove both estimates.

6.1. Convergence to \mathcal{L}_p^α . Let us start by noting that the fastest density is the null density. Thus, if we suppose that a few particles are left behind $\lfloor N\mathcal{L}_p^\alpha \rfloor$, they will travel at a higher speed and eventually catch up. The proof will rely on this observation. We first prove a partial version of the Proposition.

Lemma 10. *Given $\varepsilon > 0$, let $t^*(\varepsilon) = (\mathcal{L}_p^\alpha)^{-1}(6g^*q\varepsilon)$. For any $\kappa \in \mathbb{N}$ such that $t^*(\varepsilon) + (\kappa - 1)\varepsilon \leq T_{eq,p,\alpha}$ and any $\gamma > 0$,*

$$\lim_{\substack{N \rightarrow \infty \\ \kappa/N \rightarrow \alpha}} \mathbb{P}\left(L_{N,k}[Nt^*(\varepsilon) + \kappa\varepsilon N] \geq N\left[\mathcal{L}_p^\alpha\left((\kappa - 1)\varepsilon + \frac{\alpha}{\lambda_2(2p-1)}\right) - \gamma\right]\right) = 1 \quad (H_{\kappa,1})$$

where $\lambda_2 = \Psi(0)$.

Proof. Given $\varepsilon > 0$, we introduce the associated time subdivision $T_\kappa = N[t^*(\varepsilon) + \kappa\varepsilon]$. For convenience, we denote $W_{\kappa,\varepsilon} = \mathcal{L}_p^\alpha\left(\kappa\varepsilon + \frac{\alpha}{\lambda_2(p-q)}\right)$ and $R_{\kappa,\varepsilon} = \mathcal{L}_p^\alpha(t^*(\varepsilon) + \kappa\varepsilon)$.

We will prove $(H_{\kappa,1})$ by induction on κ . Let us sketch the idea of the proof. At T_κ , the hydrodynamic limit implies that there are only $o(N)$ particles between $NW_{\kappa,\varepsilon}$ and $NR_{\kappa,\varepsilon}$. In the totally asymmetric case, we can follow the same approach as in the convex case. Indeed, let us remark that when density converges to zero, the asymptotic speed is $g(1)$. So, we can compare to a ZRP with a constant jump rate fixed at $g(1)$ and use the analogous result proven by Labbé & Lacoïn [19, Proposition 9] for ASEP using the bijection between the two processes. If $p \neq 1$, strict concavity of Φ allows for an improvement of the result.




FIGURE 3. The configuration $\tilde{\eta}$ at $t = T_\kappa$ zoomed out by a factor N . The red dashed line corresponds to the added wall.

For $\kappa = 1$, the result is trivial. Assume $(H_{\kappa,1})$ holds and that $t^*(\varepsilon) + \kappa\varepsilon \leq T_{e_{q,p,\alpha}}$. Given $\gamma > 0$, we introduce three parameters $\gamma_1, \gamma_2, \beta > 0$ that we will choose later.

We define a process $\tilde{\eta}$ (figure 6.1) initialized at time T_κ by

$$\tilde{\eta}_{T_\kappa}(x) = \begin{cases} 0 & \text{if } x \leq \lceil NR_{\kappa,\varepsilon} - N\gamma_2 \rceil \ \& \ x \neq \lceil NW_{\kappa-1,\varepsilon} - N\gamma_1 \rceil, \\ \beta N & \text{if } x = \lceil NW_{\kappa-1,\varepsilon} - N\gamma_1 \rceil, \\ \wedge_{T_\kappa}(x) & \text{if } x > \lceil NR_{\kappa,\varepsilon} - N\gamma_2 \rceil. \end{cases}$$

Regarding the dynamics, we impose that no particles cross through the site $\lfloor NW_{\kappa,\varepsilon} \rfloor$. The evolution on the sites after $\lfloor NW_{\kappa,\varepsilon} \rfloor$ will follow a ZRP(g, p) dynamics. However, the particles on the sites before $\lfloor NW_{\kappa,\varepsilon} \rfloor$ evolve with a modified rate function $\tilde{g} = (pg(1) + q\bar{g})\mathbb{1}_{\geq 1}$ and a bias $\tilde{p} = \frac{pg(1)}{pg(1) + q\bar{g}}$. Let us remark that the generator could be written in the generic form (8). For convenience, (h_t) (resp. (\tilde{h}_t)) denotes the height function associated to (\wedge_t) (resp. $(\tilde{\eta}_t)$). We denote by

$$\mathcal{E} = \{L_{N,k}(T_\kappa) \geq N \lceil W_{\kappa-1,\varepsilon} - \gamma_1 \rceil\} \cap \{h_{T_\kappa}(\lceil NR_{\kappa,\varepsilon} - N\gamma_2 \rceil) \leq \beta N\}.$$

Thanks to Theorem 1 applied to (\wedge_t) , we have:

$$\lim_{N \rightarrow +\infty} \mathbb{P}(h_{T_\kappa}(\lceil NR_{\kappa,\varepsilon} - N\gamma_2 \rceil) \leq \beta N) = 1.$$

Therefore, the induction assumption $(H_{\kappa,1})$ implies that \mathcal{E} occurs with probability converging to one.

On the event \mathcal{E} , we observe that the height functions are ordered:

$$\tilde{h}_{T_\kappa} \geq h_{T_\kappa}.$$

Since $\tilde{p}\tilde{g} \leq pg$ and $\tilde{q}\tilde{g} = q\bar{g} \leq qg$, the processes $(\tilde{\eta}_t)$ and (\wedge_t) coupled through (\mathcal{C}) verify almost surely on \mathcal{E}

$$\forall \tau \geq t \geq T_\kappa, \quad \tilde{h}_t \geq h_t,$$

where

$$\tau = \inf \{s \geq T_\kappa, \quad \tilde{\eta}_s(\lfloor NW_{\kappa,\varepsilon} \rfloor + 1) \geq 1\},$$

the first time that the $\tilde{\eta}$ -particles beyond the wall attempt to cross it.

We obtain the following inequality

$$\begin{aligned} \mathbb{P}(L_{N,k}(T_{\kappa+1}) \leq N \lceil W_{\kappa,\varepsilon} - \gamma \rceil) \\ \leq \mathbb{P}(\{\ell(\tilde{\eta})(T_{\kappa+1}) \leq N \lceil W_{\kappa,\varepsilon} - \gamma \rceil\} \cap \mathcal{E}) + \mathbb{P}(\mathcal{E}^c) + \mathbb{P}(\tau \leq N\varepsilon). \end{aligned} \quad (33)$$

Regarding the third term of (33), we couple through (\mathcal{C}) to a process where the particles on sites $i > \lfloor NR_{\kappa,\varepsilon} \rfloor$ jump to the left independently from each other with rate $qg(1)$. If $\tau \leq N\varepsilon$,

at least one of the particles had jumped more than $\lfloor NR_{\kappa,\varepsilon} - N\gamma_2 \rfloor - \lceil NW_{\kappa,\varepsilon} \rceil - 1$. We obtain that

$$\mathbb{P}(\tau \leq \varepsilon) \leq \mathbb{P}\left(\max_k X_i \geq \lfloor NR_{\kappa,\varepsilon} - N\gamma_2 \rfloor - \lceil NW_{\kappa,\varepsilon} \rceil - 1\right),$$

where $(X_i)_{i=1}^k$ are independent Poisson variable with parameter $qg^*\varepsilon N$. As \mathcal{L}_p^α is convex,

$$R_{\kappa,\varepsilon} - W_{\kappa,\varepsilon} \geq R_{0,\varepsilon} - W_{0,\varepsilon} \geq 6\varepsilon g^* q.$$

If we fix $\gamma_2 \leq \varepsilon g^* q$, Lemma 13 implies that $\mathbb{P}(\tau \leq N\varepsilon)$ converge to zero.

It remains to control the first term of (33). The process restricted to $I_{\lceil NW_{\kappa,\varepsilon} \rceil}$ behaves like a ZRP (\tilde{g}, \tilde{p}) on a segment with the following features:

$$\tilde{N} = \lceil NW_{\kappa,\varepsilon} \rceil, \quad \tilde{k} = \beta N,$$

and initialised by $\xi = \beta N \delta_{\lceil NW_{\kappa-1,\varepsilon} - N\gamma_1 \rceil}$. Thanks to Lemma 3 and [19, Proposition 13], we obtain

$$\forall t \geq \frac{(\sqrt{\beta} + \sqrt{W_{\kappa,\varepsilon} - W_{\kappa-1,\varepsilon} + \gamma_1})^2}{pg(1) - q\bar{g}}, \quad \lim_{N \rightarrow \infty} \mathbb{P}(\ell(\tilde{\eta})(Nt + T_\kappa) \geq N[W_{\kappa,\varepsilon} - \gamma]) = 1.$$

To conclude, we should choose β, γ_1 verifying:

$$\varepsilon \geq \frac{(\sqrt{\beta} + \sqrt{W_{\kappa,\varepsilon} - W_{\kappa-1,\varepsilon} + \gamma_1})^2}{pg(1) - q\bar{g}}. \quad (34)$$

As \mathcal{L}_p^α is convex, we have

$$W_{\kappa,\varepsilon} - W_{\kappa-1,\varepsilon} \leq \varepsilon (\mathcal{L}_p^\alpha)'(T_{eq,p,\alpha}) = \varepsilon(p-q)\mathcal{L}'(\mathcal{L}^{-1}(1/\alpha))$$

and since we assumed

$$\frac{pg(1) - q\bar{g}}{p-q} > \mathcal{L}'(\mathcal{L}^{-1}(1/\alpha)),$$

we can choose β, γ_1 small enough to verify (34) which concludes the proof. \square

Remark 9. The condition (7) is due to the comparison to the exclusion process, which lacks precision. Indeed, we require that the maximal speed attained $(\mathcal{L}_p^\alpha)'(T_{eq,p,\alpha})$ is less than the highest speed reachable by ZRP (\tilde{g}, \tilde{p}) . The same method would imply a more general solution if one could control $L_{N,k}$ for small densities.

We can now finish the proof of the lower-bound on $L_{N,k}$.

Proof of Proposition 3 (Part 1). Given $t \leq T_{eq,p,\alpha}$ and $\varepsilon, \gamma > 0$, we fix

$$\kappa = \left\lfloor \frac{t - t^*(\varepsilon)}{\varepsilon} + 1 \right\rfloor - 1.$$

As \mathcal{L}_p^α is non-decreasing,

$$\mathcal{L}_p^\alpha \left((\kappa - 1)\varepsilon + \frac{\alpha}{\lambda_2(p-q)} \right) \geq \mathcal{L}_p^\alpha \left(t - t^*(\varepsilon) - 2\varepsilon + \frac{\alpha}{\lambda_2(p-q)} \right).$$

Since $t^*(\varepsilon)$ converge to $\frac{\alpha}{\lambda_2(p-q)}$ and \mathcal{L}_p^α is continuous, there exist $\varepsilon_1(\gamma)$ such that for all $\varepsilon < \varepsilon_1(\gamma)$,

$$\mathcal{L}_p^\alpha \left((\kappa - 1)\varepsilon + \frac{\alpha}{\lambda_2(p-q)} \right) \geq \mathcal{L}_p^\alpha(t) - \gamma/3.$$

It follows from the previous lemma that:

$$\lim_{\substack{N \rightarrow \infty \\ k/N \rightarrow \alpha}} \mathbb{P} \left(L_{N,k}(t^*(\varepsilon)N + \kappa\varepsilon N) \geq N \left[\mathcal{L}_p^\alpha(t) - 2\gamma/3 \right] \right) = 1.$$

Finally, we observe that

$$t - \varepsilon \leq t^*(\varepsilon) + \kappa^*\varepsilon \leq t.$$

During $N(t - t^*(\varepsilon) + \kappa^*\varepsilon)$, $L_{N,k}$ cannot move any further than $N\gamma/3$. Indeed, between $t^*(\varepsilon)N + \kappa\varepsilon N$ and t , we couple (\wedge_t) by (\mathcal{C}) to the process where particles jump independently to the left with rate $qg(1)$. Thus,

$$\begin{aligned} & \mathbb{P} \left(L_{N,k}(tN) \geq N \left[\mathcal{L}_p^\alpha(t) - \gamma \right] \right) \\ & \geq \mathbb{P} \left(L_{N,k}(t^*(\varepsilon)N + \kappa\varepsilon N) \geq N \left[\mathcal{L}_p^\alpha(t) - 2\gamma/3 \right] \right) - \mathbb{P} \left(\max_k X_i \geq \frac{N\gamma}{3} \right), \end{aligned}$$

where $(X_i)_{i=1}^k$ are independent Poisson variable with parameter $N(t^*(\varepsilon) + \kappa^*\varepsilon - t)$. Lemma 13 implies the result once ε is small enough. \square

6.2. Convergence to \mathcal{S}_p^α . Let us start with a lemma that guarantees a partial result.

Lemma 11. *Given $\varepsilon > 0$, let $t^*(\varepsilon) = (\mathcal{S}_p^\alpha)^{-1}(6\bar{g}q\varepsilon)$. For any $\kappa \in \mathbb{N}$ such that $t^*(\varepsilon) + (\kappa - 1)\varepsilon \leq T_{eq,p,\alpha}$ and any $\gamma > 0$,*

$$\lim_{\substack{N \rightarrow \infty \\ k/N \rightarrow \alpha}} \mathbb{P} \left(S_{N,k}[Nt^*(\varepsilon) + \kappa\varepsilon N] \geq N \left[\mathcal{S}_p^\alpha \left((\kappa - 1)\varepsilon + \frac{1}{g(1)(2p-1)} \right) - \gamma \right] \right) = 1 \quad (H_{\kappa,2})$$

Using the same argument as in Lemma 10 to derive the convergence of $L_{N,k}$, one can show that Lemma 11 implies the convergence of $S_{N,k}$ in probability.

Proof. Given $\varepsilon > 0$, we introduce the associated time subdivision $T_\kappa = N[t^*(\varepsilon) + \kappa\varepsilon]$. For convenience, we denote $W_{\kappa,\varepsilon} = \mathcal{S}_p^\alpha \left(\kappa\varepsilon + \frac{\alpha}{\lambda_2(p-q)} \right)$ and $R_{\kappa,\varepsilon} = \mathcal{S}_p^\alpha(t^*(\varepsilon) + \kappa\varepsilon)$.

We also prove $(H_{\kappa,2})$ by induction on κ . Let us sketch the proof. The hydrodynamic limit on the segment implies that on a $o(N)$ window near N , there are $NR_{\kappa,\varepsilon}$. The strategy consists of slowing down the $NW_{\kappa,\varepsilon}$ first particles by comparing them to a ZRP with a constant rate. Consequently, we can use the analogous result proven by Labbé & Lacoïn [19, Proposition 9] to control the flux through N .

For $\kappa = 1$, the result is trivial. Assume $(H_{\kappa,2})$ holds and that $t^*(\varepsilon) + \kappa\varepsilon \leq T_{eq,p,\alpha}$. Given $\gamma > 0$, we introduce three parameters $\gamma_1, \gamma_2, \beta > 0$ that we will choose later.

We define a coloured process $\tilde{\eta} = \tilde{\eta}^b + \tilde{\eta}^w$ (figure 6.2) where we denoted the black particles




FIGURE 4. The configuration $\tilde{\eta}$ at T_κ zoomed out by a factor N .

by $\tilde{\eta}^b$ and $\tilde{\eta}^w$ the white particles. We initialize them at time T_κ by:

$$\tilde{\eta}^b(x, T_\kappa) = \begin{cases} \wedge_{T_\kappa}(x) & \text{if } x \leq N - \lceil N\beta \rceil, \\ k - h_{T_\kappa}(N - \lceil N\beta \rceil - 1) - NW_{\kappa, \varepsilon} & \text{if } x = N - \lceil N\beta \rceil, \\ 0 & \text{otherwise.} \end{cases}$$

and

$$\tilde{\eta}^w(x, T_\kappa) = \begin{cases} 0 & \text{if } x \leq N - \lceil N\beta \rceil, \\ N[W_{\kappa, \varepsilon} - W_{\kappa-1, \varepsilon} + \gamma_1] & \text{if } x = N - \lceil N\beta \rceil, \\ N(W_{\kappa-1, \varepsilon} - \gamma_1) & \text{if } x = N, \\ 0 & \text{otherwise.} \end{cases}$$

In particular, on the event

$$\mathcal{E} = \{S_{N, k}(T_\kappa) \geq N[W_{\kappa-1, \varepsilon} - \gamma_1]\} \cap \{k - h_{T_\kappa}(N - \lceil N\beta \rceil) \geq N(R_{\kappa, \varepsilon} - \gamma_2)\},$$

the height functions are ordered

$$\tilde{h}_{T_\kappa}^b + \tilde{h}_{T_\kappa}^w \geq h_{T_\kappa},$$

where $\tilde{h}^b, \tilde{h}^w, h_{T_\kappa}$ are the corresponding height functions.

Regarding the dynamics, the white particles evolve only on $\llbracket N - \lceil N\beta \rceil, N \rrbracket$ with a modified rate function $\tilde{g} = (pg(1) + q\bar{g})\mathbb{1}_{\geq 1}$ and a bias $\tilde{p} = \frac{pg(1)}{pg(1) + q\bar{g}}$. The black particles evolve only on $I_{N - \lceil N\beta \rceil}$ according to a ZRP (g, p) . Let us emphasise that the black and white particles coexist only on the site $N - \lceil N\beta \rceil$. For this particular site, the jump rate for $\tilde{\eta}^b$ to the left is $qg(\tilde{\eta}(N - \lceil N\beta \rceil))$. In other words, only black particles can see the white ones. We claim the following:

Lemma 12. *There exists a coupling which ensures almost surely on \mathcal{E} :*

$$\forall \tau \geq t \geq T_\kappa, \quad \tilde{h}_t^b + \tilde{h}_t^w \geq h_t,$$

where

$$\tau = \inf \{s \geq T_\kappa, \quad \tilde{\eta}_s^b(N - \lceil N\beta \rceil) \leq 1\}.$$

The exact form of the generator of $(\tilde{\eta}^b, \tilde{\eta}^w)$ and the proof of this lemma are delayed to the appendix.

From now on, the proof is similar to its counterpart (Lemma 10). We obtain the following inequality:

$$\begin{aligned} \mathbb{P}(S_{N, k}(T_{\kappa+1}) \leq N[W_{\kappa, \varepsilon} - \gamma]) \\ \leq \mathbb{P}(\{\tilde{\eta}_{T_{\kappa+1}}^w(N) \leq N[W_{\kappa, \varepsilon} - \gamma]\} \cap \mathcal{E}) + \mathbb{P}(\mathcal{E}^c) + \mathbb{P}(\{\tau \leq N\varepsilon\} \cap \mathcal{E}). \end{aligned} \quad (35)$$

The hydrodynamic limit of (\wedge_t) ensures that

$$\lim_{N \rightarrow +\infty} \mathbb{P}(k - h_{T_\kappa}(N - \lceil N\beta \rceil) \geq N(R_{\kappa, \varepsilon} - \gamma_2)) = 1.$$

Thus, thanks to the induction assumption $H_{\kappa, 2}$, the event \mathcal{E} occurs with probability converging to one.

If $\tau \leq N\varepsilon$ on \mathcal{E} , there was at least $NR_{\kappa, \varepsilon} - N\gamma_2 - NW_{\kappa, \varepsilon} - 1$ jump to the left from the site $N - \lceil N\beta \rceil$. Thus

$$\mathbb{P}(\{\tau \leq N\varepsilon\} \cap \mathcal{E}) \leq \mathbb{P}(\mathcal{P}(N\varepsilon\bar{g}) \geq NR_{\kappa, \varepsilon} - N\gamma_2 - NW_{\kappa, \varepsilon} - 1).$$

By using the convexity of \mathcal{S}_p^α ,

$$R_{\kappa, \varepsilon} - W_{\kappa, \varepsilon} \geq R_{0, \varepsilon} - W_{0, \varepsilon} \geq 6\varepsilon\bar{g}q.$$

If we fix $\gamma_2 \leq \varepsilon \bar{g} q$, $\mathbb{P}(\{\tau \leq N\varepsilon\} \cap \mathcal{E})$ converges to zero.

Regarding the third term of (35), we remark that $\tilde{\eta}^w$ behaves like a ZRP (\tilde{g}, \tilde{p}) on a segment of size βN and with $NW_{\kappa, \varepsilon}$ particles. By mapping the process into an ASEP (Lemma 3), [19, Proposition 13] implies:

$$\forall t \geq \frac{(\sqrt{\beta} + \sqrt{W_{\kappa, \varepsilon} - W_{\kappa-1, \varepsilon} + \gamma_1})^2}{pg(1) - q\bar{g}}, \quad \lim_{N \rightarrow \infty} \mathbb{P}(\tilde{\eta}^w(Nt + T_{\kappa}, N) \geq N[W_{\kappa, \varepsilon} - \gamma]) = 1.$$

As \mathcal{S}_p^α is convex, we have

$$W_{\kappa, \varepsilon} - W_{\kappa-1, \varepsilon} \leq \varepsilon(\mathcal{S}_p^\alpha)'(T_{eq, p, \alpha}) = \varepsilon(p - q)\Phi\left((-\Phi')^{-1}\left(\frac{-1}{T_{eq, 1, \alpha}}\right)\right)$$

and we assumed

$$pg(1) - q\bar{g} > (p - q)\Phi\left((-\Phi')^{-1}\left(\frac{-1}{T_{eq, 1, \alpha}}\right)\right).$$

We can choose β, γ_1 small enough to verify

$$\varepsilon \geq \frac{(\sqrt{\beta} + \sqrt{W_{\kappa, \varepsilon} - W_{\kappa-1, \varepsilon} + \gamma_1})^2}{pg(1) - q\bar{g}}, \quad (36)$$

which concludes the proof up to Lemma 12. \square

Remark 10. We observe that the flux of ZRP (g, p) when density blows up converges to $(p - q)\bar{g}$. By slowing down the white particles, the maximal flux reachable is the maximal flux of a ZRP (\tilde{g}, \tilde{p}) , that is $pg(1) - q\bar{g}$. So, (7) means that we impose to the highest flux reached at $(\mathcal{S}_p^\alpha)'(T_{eq, p, \alpha})$ to be below the critical level $pg(1) - q\bar{g}$.

APPENDIX A. PROOF OF LEMMA 12

Let us first make explicit the generator of the process $(\tilde{\eta}^w, \tilde{\eta}^b)$:

$$\begin{aligned} \mathcal{A}^c J(\tilde{\eta}^w, \tilde{\eta}^b) &= \sum_{i=1}^{N - \lceil N\beta \rceil - 1} pg(\eta_i^b) [J(\eta^w, (\eta^b)^{i, i+1}) - J(\eta^w, \eta^b)] \\ &+ \sum_{i=N - \lceil N\beta \rceil}^{N-1} pg(1)\mathbb{1}_{\eta_i^w > 1} [J((\eta^w)^{i, i+1}, \eta^b) - J(\eta^w, \eta^b)] \\ &+ \sum_{i=2}^{N - \lceil N\beta \rceil - 1} qg(\eta_i^b) [J(\eta^w, (\eta^b)^{i, i-1}) - J(\eta^w, \eta^b)] \\ &+ qg(\eta_{N - \lceil N\beta \rceil}^w + \eta_{N - \lceil N\beta \rceil}^b)\mathbb{1}_{\eta_{N - \lceil N\beta \rceil}^b \geq 1} [J(\eta^w, (\eta^b)^{N - \lceil N\beta \rceil, N - \lceil N\beta \rceil - 1}) - J(\eta^w, \eta^b)] \\ &+ \sum_{i=N - \lceil N\beta \rceil + 1}^N q\bar{g}\mathbb{1}_{\eta_i^w > 1} [J((\eta^w)^{i, i-1}, \eta^b) - J(\eta^w, \eta^b)]. \end{aligned}$$

Proof of Lemma 12. In order to justify the existence of the coupling, we compare the rate jumps that may inverse the order of the height functions.

- The departure site $i < N - \lceil N\beta \rceil$, only black particle can jump.
 - A jump to the right of a black particle when $\tilde{h}^b(i) = h(i)$. In such case, $\tilde{\eta}_i^b \leq \eta_i$ which implies that $g(\tilde{\eta}_i^b) \leq g(\eta_i)$.
 - A jump to the left of (\wedge_t) -particle when $\tilde{h}^b(i-1) = h(i-1)$. In such case, $\tilde{\eta}_i^b \geq \eta_i$ which implies that $g(\tilde{\eta}_{i+1}^b) \geq g(\eta_{i+1})$.

- When $i = N - \lceil N\beta \rceil$,
 - A jump to the right of a white particle when $\tilde{h}^b(i) + \tilde{h}^w(i) = h(i)$. In such case, $\tilde{\eta}_i^w \leq \tilde{\eta}_i^w + \tilde{\eta}_i^b \leq \eta_i$ which implies that $g(1)\mathbb{1}_{\tilde{\eta}_i^w > 1} \leq g(\eta_i)$.
 - A jump to the left of (\wedge_t) -particle when $\tilde{h}^b(i-1) = h(i-1)$. In such case, $\tilde{\eta}_i^w + \tilde{\eta}_i^b \geq \eta_i$ which implies that $g(\tilde{\eta}_i^w + \tilde{\eta}_i^b) \geq g(\eta_i)$. Moreover, the stopping time ensures the presence of a black particle on this site. Thus, this jump cannot occur.
- The departure site $i > N - \lceil N\beta \rceil$, only black particle can jump.
 - A jump to the right of a black particle when $\tilde{h}^b(i) + \tilde{h}^w(i) = h(i)$. In such case, $\tilde{\eta}_i^w \leq \eta_i$ which implies that $g(1)\mathbb{1}_{\tilde{\eta}_i^w > 1} \leq g(\eta_i)$.
 - A jump to the left of (\wedge_t) -particle when $\tilde{h}^b(i-1) + \tilde{h}_t^w(i-1) = h(i-1)$. In such case, $\tilde{\eta}_i^w \geq \eta_i$ which implies that $\bar{g}\mathbb{1}_{\tilde{\eta}_i^w > 1} \geq g(\eta_{i+1})$.

From this discussion, we conclude the existence of a coupling analogous to \mathcal{C} and we omit its tedious expression. \square

APPENDIX B. A TECHNICAL LEMMA

For the sake of completeness, we provide proof of the concentration of Poisson variables that we used on multiple occasions.

Lemma 13. *Given any $\alpha > 0$ and $C > 0$, let (X_i) a sequence of independent variable distributed according to $\mathcal{P}(\alpha N)$, then*

$$\lim_{N \rightarrow \infty} \mathbb{P} \left(\max_{i \leq CN} X_i \geq 3\alpha N \right) = 0.$$

Proof. Let $\gamma = \ln \left[\frac{\ln N}{N\alpha} + 1 \right]$, Markov's inequality implies

$$\begin{aligned} \mathbb{P} \left(\max_{i \leq CN} X_i \geq 3\alpha N \right) &\leq \frac{\mathbb{E} \left(e^{\gamma \max_{i \leq CN} X_i} \right)}{e^{3\gamma\alpha N}} \\ &\leq CN \frac{\mathbb{E} \left(e^{\gamma X_1} \right)}{e^{3\gamma\alpha N}} \leq CN \frac{e^{\alpha N(e^\gamma - 1)}}{e^{3\gamma\alpha N}} \\ &\leq CN^2 e^{-3\alpha N \ln \left[\frac{\ln N}{N\alpha} + 1 \right]}. \end{aligned}$$

which converges to zero. \square

LIST OF SYMBOLS

N : The size of the segment.

k : The number of particles.

g : The rate function of the ZRP

\mathcal{A}_N : Generator of the ZRP on the segment $\llbracket 1, N \rrbracket$.

$\Omega_{N,k}^0$: The space of configurations of k particles on the segment $\llbracket 1, N \rrbracket$.

$\Omega_{N,k}$: The space of height functions of configurations in $\Omega_{N,k}^0$.

π_N^k : The equilibrium measure of the ZRP on the segment with k particles.

α : An asymptotic density.

Φ : The asymptotic flux function.

Ψ : The convex conjugate of Φ if it is convex and of $-\Phi$ if Φ is concave.

$d_{N,k}$: The worst-case total-variation distance of ZRP to the equilibrium.

$T_{\text{mix}}^{N,k}$: Mixing time of the ZRP on the segment with k particles on the segment $\llbracket 1, N \rrbracket$.

U^α : The Barron–Jensen solution to (HJ).

$T_{eq,p,\alpha}$: The macroscopic equilibrium time of U^α .
 $\wedge_{N,k}$: The configuration where all particles are localized at position 1.
 $\vee_{N,k}$: The configuration where all particles are localized at position N .
 $\ell(\eta)$: The position of the left-most particle in a configuration η .
 $L_{N,k}$: The position of the left-most particle for ZRP started from $\wedge_{N,k}$.
 $S_{N,k}$: The mass accumulated at the position N for ZRP started from $\wedge_{N,k}$.
 \mathcal{L}_p^α : The left-most position where a macroscopic mass is observed starting from $\alpha\delta_0$.
 \mathcal{J}_p^α : The macroscopic mass accumulated at the position 1 starting from $\alpha\delta_0$.
 $\mathcal{P}(\alpha)$: A Poisson distribution of parameter α .
 Cont: The set of continuity point of a given function.
 f^* : The upper semi-continuous envelope of a given function f .

REFERENCES

- [1] D. Aldous and P. Diaconis. Shuffling cards and stopping times. *Amer. Math. Monthly*, 93(5):333–348, 1986.
- [2] I. Armendáriz, S. Grosskinsky, and M. Loulakis. Zero-range condensation at criticality. *Stochastic Process. Appl.*, 123(9):3466–3496, 2013.
- [3] C. Bahadoran. Hydrodynamics and hydrostatics for a class of asymmetric particle systems with open boundaries. *Comm. Math. Phys.*, 310(1):1–24, 2012.
- [4] C. Bahadoran, T. S. Mountford, K. Ravishankar, and E. Saada. Zero-range process in random environment. In *From particle systems to partial differential equations*, volume 352 of *Springer Proc. Math. Stat.*, pages 51–77. Springer, 2021.
- [5] M. Balazs and T. Seppalainen. A convexity property of expectations under exponential weights. *arXiv Preprint*, 2007.
- [6] G. Barles. *Solutions de viscosité des équations de Hamilton-Jacobi*, volume 17 of *Mathématiques & Applications*. Springer-Verlag, Paris, 1994.
- [7] E. N. Barron and R. Jensen. Semicontinuous viscosity solutions for Hamilton-Jacobi equations with convex Hamiltonians. *Comm. Partial Differential Equations*, 15(12):1713–1742, 1990.
- [8] I. Benjamini, N. Berger, C. Hoffman, and E. Mossel. Mixing times of the biased card shuffling and the asymmetric exclusion process. *Trans. Amer. Math. Soc.*, 357(8):3013–3029, 2005.
- [9] M. Bertsch, F. Smarrazzo, A. Terracina, and A. Tesei. A uniqueness criterion for measure-valued solutions of scalar hyperbolic conservation laws. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 30(1):137–168, 2019.
- [10] M. Bertsch, F. Smarrazzo, A. Terracina, and A. Tesei. Radon measure-valued solutions of first order scalar conservation laws. *Adv. Nonlinear Anal.*, 9(1):65–107, 2020.
- [11] M. Bertsch, F. Smarrazzo, A. Terracina, and A. Tesei. Discontinuous solutions of Hamilton-Jacobi equations versus Radon measure-valued solutions of scalar conservation laws: disappearance of singularities. *J. Dynam. Differential Equations*, 35(1):455–491, 2023.
- [12] R. Dandekar and K. Mallick. Macroscopic fluctuations of a driven tracer in the symmetric exclusion process. *J. Phys. A*, 55(43):Paper No. 435001, 21, 2022.
- [13] F. Demengel and D. Serre. Nonvanishing singular parts of measure valued solutions for scalar hyperbolic equations. *Comm. Partial Differential Equations*, 16(2-3):221–254, 1991.
- [14] P. Diaconis and M. Shahshahani. Generating a random permutation with random transpositions. *Z. Wahrsch. Verw. Gebiete*, 57(2):159–179, 1981.
- [15] M. R. Evans and T. Hanney. Nonequilibrium statistical mechanics of the zero-range process and related models. *J. Phys. A*, 38(19):R195–R240, 2005.
- [16] P. Gonçalves, M. Jara, R. Marinho, and O. Menezes. Sharp convergence to equilibrium for the ssep with reservoirs. *arXiv Preprint*, 2021.
- [17] J. Hermon and J. Salez. Cutoff for the mean-field zero-range process with bounded monotone rates. *Ann. Probab.*, 48(2):742–759, 2020.
- [18] C. Kipnis and C. Landim. *Scaling limits of interacting particle systems*, volume 320 of *Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1999.
- [19] C. Labbé and H. Lacoïn. Cutoff phenomenon for the asymmetric simple exclusion process and the biased card shuffling. *Ann. Probab.*, 47(3):1541–1586, 2019.
- [20] C. Labbé and H. Lacoïn. Mixing time and cutoff for the weakly asymmetric simple exclusion process. *Ann. Appl. Probab.*, 30(4):1847–1883, 2020.

- [21] C. Labbé and E. Petit. Hydrodynamic limit and cutoff for the biased adjacent walk on the simplex. *arXiv Preprint*, 2022.
- [22] H. Lacoïn. The cutoff profile for the simple exclusion process on the circle. *Ann. Probab.*, 44(5):3399–3430, 2016.
- [23] H. Lacoïn. Mixing time and cutoff for the adjacent transposition shuffle and the simple exclusion. *Ann. Probab.*, 44(2):1426–1487, 2016.
- [24] H. Lacoïn. The simple exclusion process on the circle has a diffusive cutoff window. *Ann. Inst. Henri Poincaré Probab. Stat.*, 53(3):1402–1437, 2017.
- [25] D. A. Levin and Y. Peres. Mixing of the exclusion process with small bias. *J. Stat. Phys.*, 165(6):1036–1050, 2016.
- [26] D. A. Levin and Y. Peres. *Markov chains and mixing times*. American Mathematical Society, Providence, RI, second edition, 2017. With contributions by E. L. Wilmer, With a chapter on “Coupling from the past” by James G. Propp and David B. Wilson.
- [27] T. M. Liggett. An infinite particle system with zero range interactions. *Ann. Probability*, 1:240–253, 1973.
- [28] P.-L. Lions and P. E. Souganidis. Convergence of MUSCL and filtered schemes for scalar conservation laws and Hamilton-Jacobi equations. *Numer. Math.*, 69(4):441–470, 1995.
- [29] M. Merle and J. Salez. Cutoff for the mean-field zero-range process. *Ann. Probab.*, 47(5):3170–3201, 2019.
- [30] F. Rezakhanlou. Hydrodynamic limit for attractive particle systems on \mathbf{Z}^d . *Comm. Math. Phys.*, 140(3):417–448, 1991.
- [31] F. Rezakhanlou. Microscopic structure of shocks in one conservation laws. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 12(2):119–153, 1995.
- [32] F. Spitzer. Interaction of Markov processes. *Advances in Math.*, 5:246–290, 1970.
- [33] H. Q. Tran. The mean-field zero-range process with unbounded monotone rates: mixing time, cutoff, and Poincaré constant. *Ann. Appl. Probab.*, 33(3):1732–1757, 2023.
- [34] D. B. Wilson. Mixing times of Lozenge tiling and card shuffling Markov chains. *Ann. Appl. Probab.*, 14(1):274–325, 2004.

ONS RAMEH: DMA, ÉCOLE NORMALE SUPÉRIEURE, UNIVERSITÉ PSL, CNRS, 75005 PARIS, FRANCE,
 UNIVERSITÉ PARIS CITÉ AND SORBONNE UNIVERSITÉ, CNRS, LABORATOIRE DE PROBABILITÉS, STATIS-
 TIQUE ET MODÉLISATION, F-75013 PARIS, FRANCE

Email address: orameh@dma.ens.fr