



HAL
open science

C5 Pentacle Structures: A Localization-Delocalization Matrices Approach

Julien Pilmé, Riccardo Spezia

► **To cite this version:**

Julien Pilmé, Riccardo Spezia. C5 Pentacle Structures: A Localization-Delocalization Matrices Approach. *ChemistryOpen*, 2024, 13 (9), pp.e202300277. 10.1002/open.202300277 . hal-04732488

HAL Id: hal-04732488

<https://hal.science/hal-04732488v1>

Submitted on 11 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

C₅ Pentacle Structures: A Localization-Delocalization Matrices Approach

Julien Pilmé*^[a] and Riccardo Spezia*^[a]

This article explores the possible presence of a pentacle valence bond structure in C₅ cyclic molecules. At this end, we have used quantum chemistry tools to elucidate the possible arrangement and the nature of chemical bonds within linear, cyclic, and three-dimensional structures only formed by five carbon atoms. While the linear structure is clearly the most stable one, local minima were obtained for both bi- and three-dimensional structures. Using the localization-delocalization matrices approach, we characterize both the minimum linear structure and

the cyclic ones. Interestingly, the linear structure is a combination of ionic and covalent bonds, albeit the four distances are almost identical, when using Density Functional Theory. For cyclic C₅, the pentacle bonding arrangement emerges as a significant Lewis structure, indicative of an unusual formal configuration characterized by five intersecting C–C bonds. Our calculations show that this pentacle arrangement in cyclic C₅ scheme is also present in the more known cyclo-pentadienyl molecule.

Introduction

The ideas of regular geometries and symmetries had historically a crucial role in imaging and determining chemical structures. From a long time, the concept of valency, the nature of the chemical bond, is often related to structural considerations. One well-known example is given by the Kekule's structure of benzene.^[1] The concept of resonance structures also comes from possible geometrical patterns in which chemical bonds can be disposed fulfilling the valence of atoms constituting the molecules.^[2] Another example of structure discovery motivated by geometrical considerations is the bicyclo[2.2.0]hexa-2,5-diene structure (an isomer of benzene) proposed theoretically by Dewar^[3] and synthesised by van Tamelen and Pappas almost one hundred year after.^[4] This alternative arrangement can be seen as coming from conceptual possible arrangement of bonds which are considered as "linkages between Daltonian atoms".^[2]

We can call this a geometry-driven discovery which we can find in many other examples: from the simple water molecule,^[5] to transition metal complexes, in which group theory, leaded by symmetry, is the common way of describing them, to solid state chemistry (and physics) in which the symmetry is extended to periodicity and spatial groups are used.^[6,7]

In organic chemistry, triangles, squares, rectangles, pentagons, hexagons etc ... made by carbon atoms are ubiquitous

and they are used to simply rationalize bond making and breaking, thanks to the ability of carbon atoms to make multiple bonds. However, in organic chemistry in solution, carbon based structures are composed also by other atoms (at least H atoms), with the relevant exception of fullerenes. Interestingly, their structures were first suggested and observed experimentally only many years after.^[8]


Amongst the possible geometrical figures, one seems missing to us: the pentacle. This is an old geometrical figure composed by five vertices all linked together. If, as usual, each segment is a chemical bond, this will generate a Lewis structure corresponding to a molecule composed by five carbon atoms (and no hydrogen or other atoms). This particular cyclic C₅ molecule can have, in principle, the bonds arranged in a way they form a pentacle motif (see Scheme 1). Here we have considered, in total, three reference Lewis structures for the cyclic form of C₅ which have the same symmetry of a regular pentagon in the bonding pattern. First, the pentacle structure (PE) corresponds to a bonding scheme in which all five electron pairs are fully delocalized inside the cycle. Then, the pentagon structure (PO) corresponds to a bonding carbene-like scheme where all five electron pairs are fully localized on the carbon atoms. Finally, the cyclic structure (CY) corresponds to a benzene-like delocalization scheme where electrons are delocalized around the cycle.

We should remind that the C₅ molecule has also an interest *per se*. In fact, it was studied theoretically and experimentally in

[a] Dr. J. Pilmé, Dr. R. Spezia
Sorbonne Université, Laboratoire de Chimie Théorique, UMR 7616 CNRS, 4
Place Jussieu, 75005 Paris (France)
E-mail: julien.pilme@sorbonne-universite.fr
riccardo.spezia@sorbonne-universite.fr

Supporting information for this article is available on the WWW under
<https://doi.org/10.1002/open.202300277>

© 2024 The Authors. ChemistryOpen published by Wiley-VCH GmbH. This is
an open access article under the terms of the Creative Commons Attribution
License, which permits use, distribution and reproduction in any medium,
provided the original work is properly cited.



Scheme 1. Three Lewis resonance structures of cyclic C₅ of D_{5h} symmetry.

particular due to its astrochemical importance,^[9] finding that the linear structure, belonging to the carbyne family,^[10] is the most stable one.^[11–21] Five atoms can be arranged in different ways, linear, cyclic or with a three-dimensional arrangement.^[22] In the case of C₅, other than the aforementioned linear structure, bi- and three-dimensional geometries were identified, as stable singlet or triplet states.^[12,17] However, no detailed analysis of the chemical bond nature within these structures was conducted in previous studies. In the present contribution, we studied in detail the properties of chemical bonds of C₅ structures, in particular the most stable linear carbyne and the two-dimensional structures which can in principle have a pentacle Lewis structure.

We considered the different possibilities for C₅ to be arranged in space: as chain, as a ring and as a branched structure in order to provide also a general picture on the possible arrangement of C₅, even for non-cyclic structures. Furthermore, the LDM approach when applied to the linear structure can be compared with the more standard QAIM based approach.^[23,24] To this purpose, we used theoretical chemistry and in particular Density Functional Theory (DFT) within the well-parameterized M06-2X functional^[25] to determine the different geometries and to scrutinize the possibility that such particular resonance structure is present in C₅ molecule. Results obtained with DFT could be compared with previous data mainly based on wave-function approaches, thereby offering a degree of reliability in using DFT for larger carbon chains, where highly correlated methods cannot be afforded. Furthermore, we have also quantified the contribution of pentacle resonance structure in the well-known cyclopentadienyl, C₅H₅[−], structure.

Theory

Quantum Chemical Topology

In this paper, we assume that the reader is familiar with the quantum chemical topology (QCT) of scalar fields because numerous presentations of the methodology have already been published in the literature.^[24,26] Briefly, QCT is devoted to answer general questions about the chemical bonding in molecules and solids, and predict or explain chemical reactivity trends. This approach relies on the theory of gradient dynamical systems that enables a partitioning of the molecular space into basins. The most used one is the electron density, giving rise to the quantum theory of atoms-in-molecules (QTAIM).^[27] These basins being only atomic ones, a topological atom can be defined as the union of a nucleus with its associated electron density basin. Another widely used function is the Electron Localization Function (ELF) usually interpreted as a signature of the distribution of electron-pairs in the molecular space.^[28] The ELF topology depicts some non-atomic valence basins in addition to valence and core basins surrounding nuclei with atomic number $Z > 2$. In all the cases, the basins are delimited by zero-flux surfaces and the integration of the electron density over each basin directly provides its corresponding population.

For example, the QAIM atomic charge, q , is calculated by subtracting the electron population of the topological atom. Introduced by Bader and Stephens^[29] and later recovered by Fradera et al.^[30] the delocalization index (δ) is a measure of the electron-sharing between two atoms and can be compared to other bond order indices.

Molecular Similarity

Molecular similarity is a fundamental concept in chemistry. It is crucial to many aspects of chemical reasoning and for example, it stands as a cornerstone assumption in medicinal chemistry.^[31] Conversely, dissimilarity plays a significant role in an expanding array of applications particularly in combinatorial chemistry or for the virtual screening.^[32] Several types of mathematical methods can be used to represent the molecular dissimilarity. Among these various approaches, the electron localization-delocalization real and symmetric matrix (LDM), is a representative graph of a molecule where all atoms (vertices) are interconnected by the QAIM delocalization indices.^[33] The summation of off-diagonal elements of a LDM corresponds to the total delocalized population of the molecule. The trace of a LDM ($N \times N$) is the total localized electron population of the molecule:

$$LDM = \begin{pmatrix} \lambda_A & \delta_{AB}/2 & \dots & \delta_{AN}/2 \\ \delta_{BA}/2 & \lambda_B & \dots & \delta_{BN}/2 \\ \dots & \dots & \dots & \dots \\ \delta_{NA}/2 & \delta_{NB}/2 & \dots & \lambda_N \end{pmatrix} \quad (1)$$

where λ_i are the localization indices and δ_{ij} the delocalization indices (positive when atoms interact). Although several limitations have been identified, the LDM matrix is a powerful tool to measure the similarity/dissimilarity of different molecules themselves or as a predictor in QSAR methods.^[34] There does not exist a unique way to compare matrices. However, the Euclidian-type norm (termed as Frobenius distance) is commonly used to evaluate the dissimilarities between matrices. The dissimilarities between two molecules A and B can be then evaluated using the Frobenius distance $d(A, B)$ applied to the LDM:

$$d(A, B) = \sqrt{\sum_{ij} |a_{ij} - b_{ij}|^2} \quad (2)$$

Where a_{ij} and b_{ij} are the corresponding matrix elements of LDMs of A and B molecules respectively. Smaller the value of the distance, more similar are the two molecules. With this point, the Frobenius distance can be used for measuring the similarity between structures.^[35,36] For convenience, it is also possible to define weights associated to LDM structures. The weights, restricted to [0, 1], can be easily computed from any number N of Frobenius distances $d_1, d_2, \dots, d_{i-1}, d_N$ as follows:

$$\omega_i = \frac{1}{1 + \sum_{j \neq i}^N \frac{D_i}{D_j}} \quad (3)$$

where $D_i = d_i + \varepsilon$, ε being an arbitrary small threshold which prevents the division by zero and has been set to 10^{-6} in this study. When D_i goes to zero, ω_i goes to one. Conversely, if D_i become large, ω_i goes to zero. In this work, the Frobenius distances d_i have been calculated for optimized DFT structures with respect to formal Lewis resonant structures d_j of the same symmetry used as references here.




Figure 1. Optimized C_5 Structures. Geometries have been optimized at the M06-2X/aug-cc-pVTZ level of theory. Distances are given in Å.

Results and Discussion

Structural Parameters

We consider three arrangements of C_5 : linear 1D (or more in general a segment), 2D (a cyclic structure) and 3D (for which we considered different spatial arrangements). The most stable geometry corresponds to a linear 1D structure exhibiting a $D_{\infty h}$ symmetry ($X^1\Sigma_g^+$) in agreement with some previous studies.^[12,13,16,17,21,37] The structure was also detected in the circumstellar shell of IRC + 10216.^[38,39] We have then considered other arrangements for a closed shell singlet configuration.

Even limiting ourselves to singlet states, we found one minimum for 2D cyclic (a planar C_{2v}) and one for 3D structures (a bipyramid trigonal, D_{3h}). All the geometries are shown in Figure 1 while structural and energetic results are gathered in Table 1. More details on geometries, vibrational frequencies and rotational constants for all the minima are detailed in SI, Tables S2–S5. The 2D cyclic minimum (termed as Pentacle-Irregular) corresponds to a distorted pentagon where the bonds are not all the same and with a C_{2v} symmetry and it is 43.8 kcal/mol higher in energy than the linear structure. We also searched for a regular pentagonal structure (termed as Pentagon-Regular) with a D_{5h} symmetry where all bonds and angles are equal. However, we can find only one geometry and it turned out to be a saddle point. Notably, we obtained two imaginary frequencies, one corresponding to an out-of-plane motion consistent with a Jahn-Teller distortion, while the other one leads to the Pentacle-Regular structure.

We should notice that in previous studies, a cyclic minimum structure was obtained as triplet state using B3LYP/6-31G(d) optimization followed by single point CCSD(T)/aug-cc-pVDZ energy calculation^[16] as well as in previous MP2/D95* calculations.^[13] Previous calculations using MP4/6-31G(d) method, have found two single state minima: a cyclic isomer and a bipyrimid trigonal structure, being higher than the linear one by 60 and 95 kcal/mol, respectively.^[12] Similar behaviour are obtained by Roos and co-workers from B3LYP/6-31G(d) opti-

Table 1. Structural and energy parameters of C_5 structures optimized at the M06-2X/aug-cc-pVTZ level of theory. The carbons numbering is given in Figure 2.

| Molecule C_5 | Symmetry | ΔE (kcal/mol) | R_{12} (Å) | R_{23} (Å) | R_{14} (Å) | B (GHz) |
|--------------------|----------------|--------------------------|-----------------------|-----------------------|------------------------|-----------------------|
| Linear | $D_{\infty h}$ | 0.0 ^[a] | 1.278 | 2.556 | 2.558 | 2.574066 |
| | | | 1.3057 ^[d] | 2.6114 ^[d] | 2.6221 ^[d] | 2.5506 ^[e] |
| | | | 1.2819 ^[e] | 2.5638 ^[e] | 2.57149 ^[e] | 2.5704 ^[f] |
| Pentacle-Irregular | C_{2v} | 43.8 ^[a] | 1.602 | 3.161 | 1.446 | |
| Bipyramid trigonal | D_{3h} | 80.6 ^[a] | 1.487 | 2.012 | 1.857 | |
| Pentagon-Regular | D_{5h} | 174.3 ^[b] | 1.333 | 2.158 | 2.158 | |
| Tetrahedron | T_d | 259.5 ^[c] | 1.411 | 2.305 | 1.411 | |
| $C_5H_5^-$ | D_{5h} | | 1.408 | 2.279 | 2.279 | |

[a] Minimum structure. [b] Second-order saddle point. [c] Saddle point with three degenerated imaginary frequencies. [d] CCSD(T)/cc-pVDZ calculations from Ref. [15]. [e] CCSD(T)/cGTOs from Ref. [11]. [f] B3LYP/aug-cc-pVTZ from Ref. [17].

population of 6 electrons in the structures ω_1 and ω_2 , whereas it has a formal population of 5 electrons in ω_3 . Since the QTAIM population $\bar{N}(C_4)$ is 5.5661e, the following equation must be satisfied: $6\omega_1 + 6\omega_2 + 5\omega_3 = 5.5661e$. We can follow the same way for the carbons C_1 and C_2 which leads to the following system of equations:

$$\begin{cases} C_4 : 6\omega_1 + 6\omega_2 + 5\omega_3 = \bar{N}(C_4) = 5.566 \\ C_2 : 6\omega_1 + 7\omega_2 + 6\omega_3 = \bar{N}(C_2) = 6.167 \\ C_1 : 6\omega_1 + 4\omega_2 + 8\omega_3 = \bar{N}(C_1) = 6.531 \end{cases} \quad (4)$$

Where $\bar{N}(C_i)$ is the QTAIM population (electron) of the carbon C_i . The resulting weights are $\omega_1 = 0.40$, $\omega_2 = 0.17$ and $\omega_3 = 0.43$. The strong weight of the covalent structure is in line with the trend observed for other typical linear carbon systems.^[42] However, we can note that the full-ionic weight ($\omega_2 + \omega_3$) of 0.6 is actually predominant. It is possible to compute the weights of w_1 , w_2 and w_3 structures using the corresponding LDM matrices of $D_{\infty h}$ symmetry which can be defined as follows:

$$w_1 = \begin{pmatrix} 4 & 1 & 1 & 0 & 0 \\ 1 & 4 & 0 & 1 & 0 \\ 1 & 0 & 4 & 0 & 1 \\ 0 & 1 & 0 & 4 & 0 \\ 0 & 0 & 1 & 0 & 4 \end{pmatrix} \quad (5)$$

$$w_2 = \begin{pmatrix} 3 & 0.5 & 0.5 & 0 & 0 \\ 0.5 & 5.5 & 0 & 1 & 0 \\ 0.5 & 0 & 5.5 & 0 & 1 \\ 0 & 0.5 & 0 & 5.5 & 0 \\ 0 & 0 & 1 & 0 & 5 \end{pmatrix} \quad (6)$$

$$w_3 = \begin{pmatrix} 7 & 0.5 & 0.5 & 0 & 0 \\ 0.5 & 5 & 0 & 0.5 & 0 \\ 0.5 & 0 & 5 & 0 & 0.5 \\ 0 & 0.5 & 0 & 4.5 & 0 \\ 0 & 0 & 0.5 & 0 & 4.5 \end{pmatrix} \quad (7)$$

The Frobenius distances of the linear structure (see SI) with respect to w_1 , w_2 and w_3 can be calculated together with the corresponding weights given by Eq. 3. The resulting weights are $\omega_1 = 0.42$, $\omega_2 = 0.26$ and $\omega_3 = 0.32$. Again, we can note that the 'full-ionic' weight ($\omega_2 + \omega_3 = 0.58$) is actually predominant and its weight is similar to the 0.6 found by QTAIM-based analysis. This is also in line with a carbyne-like structure where

the terminal carbons are rather electron deficient while the charges of central carbons are rather negatives. We note that the values of delocalization indices (see Table 2) remain surprising in view of the linear structure where all C–C distances are almost identical. As indicated by the computed weights, this is probably due to a subtle resonance picture that highlights a balance between totally symmetrical covalent and ionic symmetric structures.

The Cyclic Structures

As discussed previously, we consider two cyclic C_5 structures, the Pentacle-Irregular, which is a local minimum, and the Pentagon-Regular, which is a saddle point, but the lowest energy structure with the D_{5h} symmetry. Details on vibrational frequencies and rotational constants for the minimum energy structure (Pentacle-Irregular) are reported in SI. This structure was never reported and these values could be useful for further characterization.

We now focus on the two cyclic geometries, which can have potentially a pentacle Lewis structure. As shown in Figure 3, the ELF topology of both Pentacle-Irregular (minima) and Pentagon-Regular (saddle point) yields several valence basins.

Among them, non-bonding basins accounting for the carbon lone-pairs are located on only three carbons for Pentacle-Irregular while all carbons are involved for Pentagon-Regular. For the latter, this aligns perfectly with the formal Lewis structure Pentagon (see Figure 3) where no inner C–C bonds are found. Interestingly, two similar bonding basins accounting for the covalent electron-pairing are identified inside this Pentacle-Irregular which exhibits rather striking resemblances to the pentacle formal structure (see Figure 3).

In order to quantitatively evaluate the similarity between the DFT-optimized structures and the formal Lewis structures (pentacle and pentagon), we propose to use the LDM formalism previously detailed in the theory section. Note that the way used in the previous section to build a resonant system of equations from QTAIM populations cannot be applied to the cyclic forms because all the formal atomic populations in both structures (pentacle and pentagon) are equal to 6 electrons (see Scheme 1). However, we can use the LDM matrices of the




Figure 3. ELF localization domains (isosurface = 0.8) for the C_5 cyclic structures calculated at the M06-2X/aug-cc-pVTZ level of theory. **Color code:** magenta for core basins, red for valence non-bonding basins and green for bonding basins.

formal C_5 Pentacle (PE), Pentagon (PO) and Cyclic (CY) structures of D_{5h} symmetry, which are defined as follows:

$$PE = \begin{pmatrix} 4 & 1/2 & 1/2 & 1/2 & 1/2 \\ 1/2 & 4 & 1/2 & 1/2 & 1/2 \\ 1/2 & 1/2 & 4 & 1/2 & 1/2 \\ 1/2 & 1/2 & 1/2 & 4 & 1/2 \\ 1/2 & 1/2 & 1/2 & 1/2 & 4 \end{pmatrix} \quad (8)$$

$$PO = \begin{pmatrix} 5 & 1/2 & 1/2 & 0 & 0 \\ 1/2 & 5 & 0 & 1/2 & 0 \\ 1/2 & 0 & 5 & 0 & 1/2 \\ 0 & 1/2 & 0 & 5 & 1/2 \\ 0 & 0 & 1/2 & 1/2 & 5 \end{pmatrix} \quad (9)$$

$$CY = \begin{pmatrix} 4 & 1 & 1 & 0 & 0 \\ 1 & 4 & 0 & 1 & 0 \\ 1 & 0 & 4 & 0 & 1 \\ 0 & 1 & 0 & 4 & 1 \\ 0 & 0 & 1 & 1 & 4 \end{pmatrix} \quad (10)$$

We have computed the delocalization indices and corresponding LDM matrices for the different DFT cyclic structures and measure the Frobenius distances with respect to three reference formal matrices of same D_{5h} symmetry, as explained in the Theory section. For example, the following LDM matrix corresponds to the Pentagon-Regular molecule is:

$$\begin{pmatrix} 4.008 & 0.853 & 0.853 & 0.142 & 0.142 \\ 0.853 & 4.008 & 0.142 & 0.853 & 0.142 \\ 0.853 & 0.142 & 4.008 & 0.142 & 0.853 \\ 0.142 & 0.853 & 0.142 & 4.008 & 0.853 \\ 0.142 & 0.142 & 0.853 & 0.853 & 4.008 \end{pmatrix} \quad (11)$$

As the DFT Pentacle-Irregular displays two short C–C and two long C–C distances, we also need to consider an additional average Pentacle-Irregular structure (APEI), which is provided in Figure 4. This latter LDM involves a partial inner delocalized density in agreement with the ELF localization domains provided in Figure 3, where two small inner disynaptic ELF basins are observed along the C_1 – C_4 and C_1 – C_5 bonds.

All the results are reported in Table 3 where we also report the distance for PE, PO and CY as reference as well as what obtained from $C_5H_5^-$.

Results are reported in Table 3 and divided following the symmetry of the different structures: D_{5h} for Pentagon-Regular and $C_5H_5^-$ and C_{2v} for Pentacle-Irregular. Surprisingly, the pentacle Lewis structures make an important contribution in




Figure 4. Average Pentacle-Irregular (PEI) LDM computed as $APEI = \frac{1}{2} (PEI-A + PEI-B)$.

| Molecule C_5 | d_{PE} | d_{PO} | d_{CY} | $d_{APEI}^{[a]}$ | ω_{PE} | ω_{PO} | ω_{CY} | $\omega_{APEI}^{[a]}$ |
|---------------------------------------|----------|----------|----------|------------------|---------------|---------------|---------------|-----------------------|
| D_{2h} Structures | | | | | | | | |
| Pentacle (PE) | 0.0 | 2.738 | 2.236 | | 1.0 | 0.0 | 0.0 | |
| Pentagon (PO) | 2.738 | 0.0 | 2.738 | | 0.0 | 1.0 | 0.0 | |
| Cyclic (CY) | 2.236 | 2.738 | 0.0 | | 0.0 | 0.0 | 1.0 | |
| Pentagon-Regular | 1.588 | 2.524 | 0.650 | | 0.25 | 0.15 | 0.60 | |
| $C_5H_5^-$ | 1.677 | 2.018 | 2.524 | | 0.40 | 0.34 | 0.26 | |
| C_{2v} Structures | | | | | | | | |
| Average Pentacle Irregular (APEI) | 1.224 | 2.236 | 1.870 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 |
| Pentacle-Irregular | 1.441 | 2.017 | 2.049 | 0.868 | 0.25 | 0.17 | 0.17 | 0.41 |

[a] Average Pentacle Irregular (APEI) as shown in Figure 4.

both DFT structures, Pentagon-Regular and Pentacle-Irregular. Notably, structures with *crossing bonds*, i.e. PE and APEI, show an important contribution, with a weight exceeding 60% ($\omega_{PE} + \omega_{APEI}$) for Pentacle-Irregular and also exceeding 40% (ω_{PE}) for the well-known $C_5H_5^-$ molecule. Note that for Pentagon-Regular, it was rather difficult to characterize the contribution of the pentacle Lewis structure from only ELF localization domains (see Figure 3) since none inner C–C bonding basin has been found inside the pentagon. These results illustrate that the rationalization of the electronic structure of cyclic structures needed a thorough analysis of a set of delocalization indices revealing the role of inner delocalized density in the pentacle cycle. Finally, based on the weights obtained from the LDM matrix analysis, the aromatic character of Pentacle Irregular structures appears to be quite comparable to that of the cyclopentadienyl anion, which exhibits maximum aromaticity with five equivalent bonds and a totally delocalized five-membered ring.^[43]

Conclusions and Outlooks

It is remarkable how small and simple systems, such as the C_5 molecule, still offer a rich playground for a better understanding of usual concepts, such as the resonance. In particular, when the five carbon atoms are arranged in a ring, we obtained that a formal structure with bonds passing inside the ring is an important one: this intriguingly corresponds to a pentacle, a particular geometrical motif which was not considered before. This particular bonding pattern is intriguing because the bonds formally *cross* each other. Notably, this pattern is not negligible also in the hydrogenated form, the less exotic $C_5H_5^-$ molecule. While a pentacle is possible only in C_5 systems, it will be interesting to know if other *crossing bonds* structures are possible resonance structures and how this is related (or not) to their reactivity.

Furthermore, we have found that modern DFT functionals with extended basis sets are able to obtain the linear carbyne structure as the most stable minimum, and to locate stable a 2D cyclic structure never considered and/or obtained before, and the well documented 3D bipyramid trigonal structure. Thus, it would be possible to increase the carbon length using DFT and then see how (and if) 2D and 3D geometries become more and more stable and if such crossing bonds are responsible to their stabilization of other structures with growing number of carbon atoms. This question will surely deserve more investigations.

Computational Details

The M06-2X hybrid functional^[25] level with the Gaussian 16 software was used for all calculations.^[44] The standard all-electron aug-cc-pVTZ basis set was used for all atoms.^[45] The C_5 linear ground state structure was fully optimized without symmetry constraints. In a first step, local minima geometries were approached through symmetry constrained calculations. For instance, D_{3h} was used for trigonal bipyramid. We then fully optimized the geometries using

the local minimum-energy structures as initial guesses. Each minimum displays only positive eigenvalues. Optimizing other structures with different symmetry constraints always resulted in higher order saddle points, with two or three imaginary frequencies for second-order and third-order saddle points, respectively. Only the single states for each geometry were considered. The quantum chemical topology analyses and LDM matrices have been performed using the TopChem2 program package.^[46,47]

Acknowledgements

We thank an anonymous Twitter user who brought us to the interest of a possible pentacle bonding structure.

Conflict of Interests

No conflict of interest to declare.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Keywords: Valence Bond Theory · Astrochemical molecules · DFT calculations · QTAIM · ELF

- [1] F. A. Kekulé, *Liebigs Ann. Chem.* **1866**, 137, 129.
- [2] G. N. Lewis, *J. Am. Chem. Soc.* **1916**, 38, 762.
- [3] J. Dewar, *Proc. R. Soc. Edinburgh* **1869**, 6, 82.
- [4] E. E. van Tamelen, S. P. Pappas, *J. Am. Chem. Soc.* **1963**, 85, 3297.
- [5] H. E. Armstrong, *Philos. Trans. R. Soc. London* **1873**, 163, 1.
- [6] A. Hofmann, *Philos. Trans. R. Soc. London* **1862**, 152, 83.
- [7] V. Singh, *J. Chem. Educ.* **2007**, 84, 1919.
- [8] H. Kroto, J. Heath, S. O'Brien, R. Curl, R. Smalley, *Nature* **1985**, 318, 162–163.
- [9] A. Van Ordent, R. Saykally, *Chem. Rev.* **1998**, 98, 2313.
- [10] M. Casari, A. Milani, *MRS Commun.* **2018**, 8, 207.
- [11] P. Botschina, *J. Chem. Phys.* **1987**, 101, 853.
- [12] K. Raghavachari, J. S. Binkley, *J. Chem. Phys.* **1987**, 87, 2191.
- [13] D. Ewing, G. Pfeiffer, *Chem. Phys. Lett.* **1987**, 134, 413.
- [14] N. Moazzen-Ahmadi, S. Flatt, A. McKellar, *Chem. Phys. Lett.* **1991**, 186, 291.
- [15] J. Martin, P. Taylor, *J. Phys. Chem.* **1996**, 100, 6047.
- [16] S. Dua, J. Bowie, *J. Phys. Chem.* **2002**, 106, 1374.
- [17] H. Massó, V. Veryazov, P.-A. Malmqvist, B. O. Roos, M. L. Senent, *J. Chem. Phys.* **2007**, 127, 154318.
- [18] D. W. Arnold, S. E. Bradforth, T. N. Kitsopoulos, D. M. Neumark, *J. Chem. Phys.* **1991**, 95, 8753.
- [19] N. Moazzen-Ahmadi, A. R. W. McKellar, T. Amano, *J. Chem. Phys.* **1989**, 91, 2140.
- [20] N. Moazzen-Ahmadi, A. McKellar, T. Amano, *Chem. Phys. Lett.* **1989**, 157, 1.
- [21] J. Hutter, H. P. Luethi, F. Diederich, *J. Am. Chem. Soc.* **1994**, 116, 750.
- [22] P. Bera, K. Sattelmeyer, M. Saunders, H. Shaefer III, P. Schleyer, *J. Phys. Chem. A* **2006**, 110, 4287.
- [23] C. Gomez Pech, P. A. B. Haase, D. C. Sergentu, A. Borschevsky, J. Pilm, N. Galland, M. Rmi, *J. Comput. Chem.* **2020**, 41, 2055.
- [24] B. Silvi, R. J. Gillespie, C. Gatti, 9.07 - Electron Density Analysis, in J. Reedijk, K. Poeppelmeier (Editors), *Comprehensive Inorganic Chemistry II (Second Edition)*, pages 187–226, Elsevier, Amsterdam, second edition 2013.
- [25] Y. Zhao, D. G. Truhlar, *Theor. Chem. Acc.* **2008**, 120, 215.

- [26] P. L. A. Popelier, Quantum Chemical Topology, in D. M. P. Mingos (Editor), *The Chemical Bond II: 100 Years Old and Getting Stronger*, pages 71–117, Springer International Publishing, Cham 2016.
- [27] R. F. W. Bader, *Chem. Rev.* **1991**, *91*, 893.
- [28] B. Silvi, A. Savin, *Nature* **1994**, *371*, 683–686.
- [29] R. F. W. Bader, M. E. Stephens, *J. Am. Chem. Soc.* **1975**, *97*, 7391.
- [30] X. Fradera, M. A. Austen, R. F. W. Bader, *J. Phys. Chem. A* **1999**, *103*, 304.
- [31] G. Maggiora, M. Vogt, D. Stumpfe, J. Bajorath, *J. Med. Chem.* **2014**, *57*, 3186.
- [32] A. Bender, R. C. Glen, *Org. Biomol. Chem.* **2004**, *2*, 3204.
- [33] C. Matta, I. Sumar, R. Cook, P. Ayers, Localization-Delocalization Matrices and Electron Density-Weighted Adjacency/Connectivity Matrices: A Bridge Between the Quantum Theory of Atoms in Molecules and Chemical Graph Theory, in R. Chauvin, C. Lepetit, B. Silvi, E. Alkhani (Editors), *Applications of Topological Methods in Molecular Chemistry*, Springer International Publishing, Cham, 2016, pages 53–88, .
- [34] R. Cook, *Struct. Chem.* **2017**, *28*, 1525–1535.
- [35] C. F. Matta, *Comput. Theor. Chem.* **2018**, *1124*, 1.
- [36] M. J. Timm, C. F. Matta, L. Massa, L. Huang, *J. Phys. Chem. A* **2014**, *118*, 11304.
- [37] A. Chefai, M. Ben Khalifa, F. Khadri, K. Hammami, *Phys. Chem. Chem. Phys.* **2021**, *23*, 23741.
- [38] P. F. Bernath, K. H. Hinkle, J. J. Keady, *Science* **1989**, *244*, 562.
- [39] G. Galazutdinov, F. Musaev, J. Krelowski, *Mon. Not. R. Astron. Soc.* **2001**, *325*, 1332.
- [40] R. Gillespie, *Coord. Chem. Rev.* **2008**, *252*, 1315.
- [41] B. Silvi, *Phys. Chem. Chem. Phys.* **2004**, *6*, 256.
- [42] F. Pichierri, *J. Phys. Chem. A* **2008**, *112*, 7717, pMID: 18666765.
- [43] J. Poater, M. Duran, M. Solà, B. Silvi, *Chem. Rev.* **2005**, *105*, 3911.
- [44] M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, D. J. Fox, Gaussian 16 Revision C.01 **2016**, gaussian Inc. Wallingford CT.
- [45] T. H. Dunning Jr, *J. Chem. Phys.* **1989**, *90*, 1007.
- [46] D. Kozłowski, J. Pilmé, *J. Comput. Chem.* **2011**, *32*, 3207.
- [47] H. Chevreau, J. Pilmé, *J. Comput. Chem.* **2023**, *44*, 1505.

Manuscript received: November 23, 2023

Revised manuscript received: March 25, 2024

Version of record online: May 16, 2024