



HAL
open science

Gait rehabilitation in patients with lower-limb amputations: a systematic literature review

Lisa Baltus, Kevin Arribart, Alexandra Roren

► To cite this version:

Lisa Baltus, Kevin Arribart, Alexandra Roren. Gait rehabilitation in patients with lower-limb amputations: a systematic literature review. 2024. hal-04737997

HAL Id: hal-04737997

<https://hal.science/hal-04737997v1>

Preprint submitted on 15 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Gait rehabilitation in patients with lower-limb amputations: a systematic literature review

Lisa Baltus*

IFMK Ecole d'ASSAS, Paris, France. Service de Rééducation et de Réadaptation de l'Appareil Locomoteur et des Pathologies du Rachis, Hôpital Cochin, AP-HP, Paris, France.

Kevin Arribart

IFMK Ecole d'ASSAS, Paris, France. Institut Robert Merle d'Aubigné, Service de rééducation et d'appareillage des patients amputés, Valenton, France

Alexandra Roren

Université Paris Cité, INSERM UMR 1153. Service de Rééducation et de Réadaptation de l'Appareil Locomoteur et des Pathologies du Rachis, Hôpital Cochin, AP-HP, Paris, France.

***Corresponding author.**

Email : baltus.lisa@gmail.com

1. Introduction

The higher the level of amputation, the greater the functional consequences (pain, lameness, psychosocial consequences, etc.) and the higher the energy cost of gait. Compared to able bodied individuals, the energy costs 35% more for all amputee on average, 12 to 36% more for transtibial amputee (TTA), 41 to 102% more for a transfemoral amputee (TFA), 120% more for a hemipelvic or disarticulated hip and 100% more for a bilateral below knee amputee. Kinematic gait parameters showed increased pelvic anterior tilt, length of the stance phase on the amputated side and muscular stump contraction. Also, it showed decreased hamstring length-strength relationship and decreased paravertebral, lumbar and stump muscles strength. The objective of this systematic literature review was to determine the most effective rehabilitation protocols on gait parameters in people with lower-limb amputations. Our major hypothesis was that protocols based on standardized exercises significantly improved gait parameters in people with lower-limb amputations compared to non-standardized rehabilitation (usual care).

2. Methods

We only included randomized controlled trials (RCTs) assessing adults, with unilateral or bilateral lower limb amputations (regardless the levels of amputation and the aetiologies) and able to walk with their prosthesis. The intervention was program including standardized rehabilitation methods. The comparator was non-standardized rehabilitation. The outcomes were gait spatio-temporal parameters: speed (m/s), step length (m), step width (m), cadence (step/min) and distance (m). We searched for RCTs from the following electronic databases: MEDLINE, Cochrane Library, PEDro, Science Direct and Google Scholar, in English or French, from June 2022 to March 2023. The RoB2 tool, the PEDRO scale and the AMSTAR 2 tool were used to assess the risk of bias, the overall quality of the RCTs and of the systematic review, respectively.

3. Results and discussion

3.1 Results

The search retrieved 6338 studies. 12 RCT (a total sample of 507 participants), published between 2002 and 2021 were included. The number of patients included in the RCTs varied from 15 and 154. In the intervention group, the standardized rehabilitation program included hip abductor muscle strengthening, proprioceptive neuromuscular facilitation, dual-task training and electrical stimulation therapy. In the comparator group, the participants had usual non-standardized rehabilitation.

Only 3 RCTs reported a higher effectiveness of standardized rehabilitation exercises on gait speed, stride length, cadence, stance and distance compared to non-standardized rehabilitation. [1, 2, 3]. They were all designed in 1-hour sessions twice a week combining muscular work on the lower limbs and trunk, cardiorespiratory work, balance, weight-bearing, and walking trails appear to be effective on spatio-temporal gait parameters.

A RCT including 26 participants, (16 TFA and TTA participants from 18 to 60 years, and 10 able-bodied people) assessed a 4 weeks program with 2 sessions of 60min/week intervention using low-cost basic rehabilitation material: TheraBand, balls... (Almeida et al. 2021) [1]. This RCT showed a clinically meaningful improvement in gait speed, stride length, cadence and stance [Table 1].

A RCT including 15 TFA and TTA 60(+/-12) years assessed a 12 weeks program including a supervised, circuit-style group exercise session twice weekly at the University and personalized exercises at home once a week, progressing to twice-weekly from the mid-point of the intervention (Schafer et al. 2018) [2]. This RCT showed a clinically meaningful improvement in gait speed, length, cadence and stance [Table 1].

A RCT including 16 unilateral TTA 63.25 (+/-11.5) years assessed a 8 weeks, 60min/week program including table of exercises based on AMPPRO tool results (Gailey et al. 2020) [3]. This RCT showed clinically meaningful improvement in distance [Table 1].

The RoB 2 tool varied from moderate [1,3] to high [2] risk of bias. The Pedro score was 4/10 [3], and 7/10 [1,2]. The Amstar-2 tool estimated that this systematic review was of average quality.

Table 1. Improvements in gait, temporal-spatial parameters
credit: Baltus Lisa

Study	Almeida et al. (2021) n=16	Schafer et al. (2018) n=15	Gailey et al. (2020) n=16
Speed	+0.35m/s > MDC=0.11m/s **	+0.25m/s > MDC=0.10m/s**	
Length amputated + sound side	+0.06** and +0.07m**	+0.07m*	
Cadence amputated + sound side	+43 step/min **	+10 step/min* and +12 step/min*	
Stance amputated+ Sound side	57.5% (-7.6)*	56.4% (-3,3)* and 68,6%(-2,7) and 27%(-4,45)*	
Distance			+74m** >MDC=58,2m & MCID=50 m

* means statistically significant

** means clinically relevant

3.1 Discussion

The present systematic literature review showed that 3 standardized exercise programs were more effective on improving gait-speed, stride-length, cadence, stance and gait distance in people with lower limb amputation than non-standardized programs. Nevertheless, in 9/12 RCTs the standardized rehabilitation was not more effective on improvement in gait parameters than non-standardized rehabilitation in people. This result cannot allow us to validate our major hypothesis.

This systematic review has several limits: A single rater performed studies inclusion and data extraction.

The content of the rehabilitation programs and timelines for collection of gait parameters varied between studies which makes it difficult to perform a meta-analysis. The samples of participants in most RCTs remained small.

4. Conclusion

Future studies should explore the long-term effects of multidimensional standardized rehabilitation programs on gait parameters and core outcomes (activity limitation and quality of life). The use of in-prosthesis IMUs and pressure sensors may allow to have real-life data. Providing more detailed rehabilitation protocols and including more homogenous populations may bring valuable information to optimize rehabilitation strategies in people with lower limb amputation.

Funding

No funding was provided for the conduct of the review.

Conflict of Interest Statement

The author declares no potential conflicts of interest.

References

- Berthel M, Ehrler S. (2010). Aspects épidémiologiques de l'amputation de membre inférieur en France. *Kinesither Sci.* 512 : 5-8
- Ettema S, Kal E, Houdijk H. General estimates of the energy cost of walking in people with different levels and causes of lower-limb amputation: a systematic review and meta-analysis. *Prosthet Orthot Int.* 2021 Oct 1;45(5):417-427.
- Almeida LV, Fukuchi CA, Sakanaka TE, Cliquet A.(2021). A low-cost easily implementable physiotherapy intervention clinically improves gait implying better adaptation to lower limb prosthesis: a randomized clinical trial. *Sci Rep.* 27.11:21228.
- Gailey R, Gaunard I, Raya M, Kirk-Sanchez N, Prieto-Sanchez LM, Roach K.(2020). Effectiveness of an Evidence-Based Amputee Rehabilitation Program: A Pilot Randomized Controlled Trial. *Phys Ther.*100(5):773- 87.
- Schafer ZA, Perry JL, Vanicek N.(2018). A personalised exercise programme for individuals with lower limb amputation reduces falls and improves gait biomechanics: A block randomised controlled trial. *Gait Posture.* 63:282- 9

Received date:06/04/2024

Accepted date: 28/06/2024

Published date: XX/XX/2024

Volume: 1

Publication year: 2024