



HAL
open science

Superspace realizations of the Bannai–Ito algebra

Nicolas Crampé, Hendrik de Bie, Plamen Iliev, Luc Vinet

► **To cite this version:**

Nicolas Crampé, Hendrik de Bie, Plamen Iliev, Luc Vinet. Superspace realizations of the Bannai–Ito algebra. *Letters in Mathematical Physics*, 2023, 113 (5), pp.108. <10.1007/s11005-023-01731-6>. <hal-04738441>

HAL Id: hal-04738441

<https://hal.science/hal-04738441v1>

Submitted on 15 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

SUPERSPACE REALIZATIONS OF THE BANNAI–ITO ALGEBRA

NICOLAS CRAMPÉ, HENDRIK DE BIE, PLAMEN ILIEV, AND LUC VINET

ABSTRACT. A model of the Bannai–Ito algebra in a superspace is introduced. It is obtained from the three-fold tensor product of the basic realization of the Lie superalgebra $\mathfrak{osp}(1|2)$ in terms of operators in one continuous and one Grassmanian variable. The basis vectors of the resulting Bannai–Ito algebra module involve Jacobi polynomials.

1. INTRODUCTION

This paper offers a realization of the Bannai–Ito algebra in superspace. Models of algebras with operators acting on spaces of functions are especially relevant in mathematical physics and allow also to enrich the understanding of special functions [1]. This is the basic motivation here. The Bannai–Ito algebra [2] encodes the bispectral properties of the eponymous polynomials which were first introduced in the context of P- and Q- polynomial association schemes [3]. Their characterization as eigenfunctions of shift operators of Dunkl type was given in [4].

The Bannai–Ito polynomials have explicit expressions as combinations of two Racah polynomials [5]. Moreover, the Racah algebra [6] that is abstracted from the relations verified by the difference and recurrence operators corresponding to these last polynomials can be embedded in the Bannai–Ito algebra [7] much like the even Lie algebra part of a superalgebra can be obtained from quadratic expressions in the odd generators.

As a matter of fact, it was also observed that the Racah algebra can be directly embedded in the universal enveloping algebra $\mathcal{U}(\mathfrak{sl}(2))$ of $\mathfrak{sl}(2)$ [8]. Calling upon the Bargmann realization of $\mathfrak{sl}(2)$ in terms of differential operators in one variable [9], this last embedding thus immediately provides a realization of the Racah algebra in terms of differential operators of the hypergeometric type.

Given that the Racah polynomials arise in the $6-j$ recoupling coefficients of the $\mathfrak{sl}(2)$ representations, it is not surprising that a centrally extended version of the Racah algebra arises by taking as generators the intermediate Casimir in a three-fold product of representations [10]. Here, the Bargmann realization produces a model of the Racah algebra in terms of differential operators in three variables. The dimensional reduction of this three-variable realization through separation of variables has been examined in [11] and

shown to yield the one-variable model of the Racah algebra mentioned before. It has been shown in this spirit how the higher rank Racah algebras [12] constructed in this fashion using the n variables associated to an n -fold product of irreducible $\mathfrak{sl}(2)$ representations can be reduced to a model involving differential operators in only $n - 2$ variables [13].

The Bannai-Ito algebra can also be viewed as the commutant of a diagonal embedding of an algebra module in a three-fold tensor product of representations. In this case the underlying algebraic structure is the Lie superalgebra $\mathfrak{osp}(1|2)$ and it is hence understood that the Bannai–Ito polynomials are essentially the Racah coefficients of this superalgebra [14]. Now one extension to $\mathfrak{osp}(1|2)$ of the Bargmann model of $\mathfrak{sl}(2)$ involves thickening the one-dimensional space by the addition of a real Grassmann variable; this leads to using operators acting on functions defined on the resulting superspace [15]. Considering the tensor product of three irreducible (discrete series) representations of $\mathfrak{osp}(1|2)$ [16] in this picture therefore brings three continuous variables and three Grassmannian ones. In view of the corresponding studies of the Racah realizations mentioned before, natural questions we address in this paper are: What is the Bannai–Ito algebra model that the irreducible decomposition of this tensor product representation brings? To how many variables does this reduction yield? What are the special functions involved?

The remainder of the paper is organized as follows. The fundamental superspace model of $\mathfrak{osp}(1|2)$ is recalled in Section 2. The construction of the Bannai–Ito algebra generated by the intermediate Casimir operators of three copies of this $\mathfrak{osp}(1|2)$ realization is described in Section 3. This module decomposes into an even and an odd part each with four components. The dimensional reduction that occurs upon requesting that the representation space be bounded from below is discussed in Section 4. It will be shown that the kernel of the total lowering operator is such that its even and odd part only have two components. The action of the intermediate Casimir operators in each of these 4 parts will be given in Section 5. The construction of the irreducible representations is carried out by considering in turn the even subspace in Section 6 and the odd one in Section 7. In each case, first, the concrete representation basis will be obtained by diagonalizing the total Casimir operator and one of the intermediate Casimir elements and second, the tridiagonal action of the other independent intermediate Casimir element will be computed. The Jacobi polynomials will be seen to appear in the expression of these basis elements. The concluding section will offer summary, remarks and outlook.

2. THE FUNDAMENTAL REALIZATION OF $\mathfrak{osp}(1|2)$ ON SUPERSPACE

Consider the operators

$$\begin{aligned} A_- &= \theta \partial_x + \partial_\theta, \\ A_0 &= 2x \partial_x + \theta \partial_\theta + 2\nu, \\ A_+ &= x \theta \partial_x + x \partial_\theta + 2\nu \theta, \end{aligned}$$

where θ is a Grassmann variable $\theta^2 = 0$ commuting with x . It is easy to check that the operators satisfy the following relations

$$[A_0, A_\pm] = \pm A_\pm \quad \text{and} \quad \{A_+, A_-\} = A_0,$$

where as usual $[a, b] = ab - ba$ denotes the commutator of the operators a and b , and $\{a, b\} = ab + ba$ denotes the anticommutator. Therefore, the operators $\hat{A}_\pm = \sqrt{2}A_\pm$, $\hat{A}_0 = A_0$ define a representation of the superalgebra $\mathfrak{osp}(1|2)$. Moreover

$$A_-^2 = \partial_x.$$

Let P be the operator acting on functions $f(x, \theta)$ by

$$P(f(x, \theta)) = f(x, -\theta).$$

Clearly, P is an involution, i.e. $P^2 = 1$ and

$$[P, A_0] = 0, \quad \{P, A_\pm\} = 0.$$

This operator P is realized as follows

$$P = 1 - 2\theta \partial_\theta.$$

The Casimir operator

$$Q = (A_0 - 2A_+A_- - 1/2)P$$

commutes with A_0 , A_\pm and P . One can also easily check that in our current realization it holds that

$$Q = 2\nu - 1/2.$$

3. THREE COPIES AND THE BANNAI-ITO ALGEBRA

For $j = 1, 2, 3$, let

$$\begin{aligned} A_-^{(j)} &= \theta_j \partial_{x_j} + \partial_{\theta_j}, \\ A_0^{(j)} &= 2x_j \partial_{x_j} + \theta_j \partial_{\theta_j} + 2\nu_j, \\ A_+^{(j)} &= x_j \theta_j \partial_{x_j} + x_j \partial_{\theta_j} + 2\nu_j \theta_j, \end{aligned}$$

where $\{\theta_j\}$ are anticommuting, i.e.

$$\theta_i \theta_j = -\theta_j \theta_i.$$

Note that for $i \neq j$ we have

$$[A_0^{(i)}, A_0^{(j)}] = 0, \quad [A_\pm^{(i)}, A_\pm^{(j)}] = 0, \quad \{A_\pm^{(i)}, A_\pm^{(j)}\} = 0, \quad \{A_\pm^{(i)}, A_\mp^{(j)}\} = 0.$$

In particular, these commutativity relations imply that for every nonempty $S \subset \{1, 2, 3\}$ we obtain another representation of $\mathfrak{osp}(1|2)$ with operators defined by

$$A_0^{(S)} = \sum_{i \in S} A_0^{(i)}, \quad A_{\pm}^{(S)} = \sum_{i \in S} A_{\pm}^{(i)}, \quad P^{(S)} = \prod_{i \in S} P^{(i)},$$

and associated Casimir operator

$$Q^{(S)} = (A_0^{(S)} - 2A_+^{(S)}A_-^{(S)} - 1/2)P^{(S)}.$$

Note that $Q^{(S)}$, usually called intermediate Casimir operator, commutes with $A_0^{(T)}$, $A_{\pm}^{(T)}$ and $P^{(T)}$ for every $T \supset S$ and $Q^{(i)} = 2\nu_i - 1/2$.

The Casimir operators $Q^{(S)}$ provide a realisation of the Bannai–Ito algebra *i.e.* they satisfy the following relations

$$\{Q^{(12)}, Q^{(23)}\} = Q^{(13)} + 2Q^{(1)}Q^{(3)} + 2Q^{(2)}Q^{(123)}, \quad (3.1)$$

$$\{Q^{(12)}, Q^{(13)}\} = Q^{(23)} + 2Q^{(2)}Q^{(3)} + 2Q^{(1)}Q^{(123)}, \quad (3.2)$$

$$\{Q^{(13)}, Q^{(23)}\} = Q^{(12)} + 2Q^{(1)}Q^{(2)} + 2Q^{(3)}Q^{(123)}. \quad (3.3)$$

In this realisation of the Bannai–Ito algebra, the following relation also holds

$$(Q^{(12)})^2 + (Q^{(13)})^2 + (Q^{(23)})^2 + \frac{1}{4} = (Q^{(123)})^2 + (Q^{(1)})^2 + (Q^{(2)})^2 + (Q^{(3)})^2.$$

4. DIMENSIONAL REDUCTION

Below we work with the superspace

$$\mathbb{V} = \mathbb{C}[x_1, x_2, x_3] \langle \theta_1, \theta_2, \theta_3 \rangle.$$

We can decompose \mathbb{V} as the direct sum of the odd subspace \mathbb{V}^o and the even subspace \mathbb{V}^e as follows

$$\mathbb{V} = \mathbb{V}^o \oplus \mathbb{V}^e, \quad (4.1)$$

where

$$\mathbb{V}^o = \mathbb{C}[x_1, x_2, x_3]\theta_1 \oplus \mathbb{C}[x_1, x_2, x_3]\theta_2 \oplus \mathbb{C}[x_1, x_2, x_3]\theta_3 \oplus \mathbb{C}[x_1, x_2, x_3]\theta_1\theta_2\theta_3$$

$$\mathbb{V}^e = \mathbb{C}[x_1, x_2, x_3] \oplus \mathbb{C}[x_1, x_2, x_3]\theta_1\theta_2 \oplus \mathbb{C}[x_1, x_2, x_3]\theta_1\theta_3 \oplus \mathbb{C}[x_1, x_2, x_3]\theta_2\theta_3.$$

These subspaces can be characterized as the spaces consisting of skew-symmetric and symmetric functions with respect to the involution $P^{(123)}$ on \mathbb{V} . Since $P^{(123)}$ anti commutes with $A_-^{(123)}$, we can decompose the $\ker(A_-^{(123)})$ on \mathbb{V} as a direct sum of odd and even functions

$$\ker(A_-^{(123)}) = (\ker(A_-^{(123)}) \cap \mathbb{V}^o) \oplus (\ker(A_-^{(123)}) \cap \mathbb{V}^e).$$

To describe each component, we fix

$$u = x_1 - x_2 \quad \text{and} \quad v = x_2 - x_3,$$

and we consider the linear transformations

$$\mathcal{O}_1, \mathcal{O}_2 : \mathbb{C}[u, v] \rightarrow \mathbb{V}^o$$

$$\mathcal{E}_1, \mathcal{E}_2 : \mathbb{C}[u, v] \rightarrow \mathbb{V}^e$$

defined as follows

$$\begin{aligned}\mathcal{O}_1(h(u, v)) &= h(u, v)(\theta_1 - \theta_2) + h_v(u, v)\theta_1\theta_2\theta_3, \\ \mathcal{O}_2(h(u, v)) &= h(u, v)(\theta_2 - \theta_3) - h_u(u, v)\theta_1\theta_2\theta_3, \\ \mathcal{E}_1(h(u, v)) &= h(u, v)(\theta_1\theta_2 - \theta_1\theta_3 + \theta_2\theta_3), \\ \mathcal{E}_2(h(u, v)) &= h(u, v) + h_u(u, v)\theta_1\theta_2 + h_v(u, v)\theta_2\theta_3,\end{aligned}$$

where $h(u, v) \in \mathbb{C}[u, v]$.

Proposition 4.1. *Let $F \in \mathbb{V}$. Then F solves the equation*

$$A_-^{(123)}F = 0, \quad (4.2)$$

if and only if F can be written as

$$F = F_o + F_e, \quad (4.3)$$

where

$$F_o = \mathcal{O}_1(h_1(u, v)) + \mathcal{O}_2(h_2(u, v)) \quad (4.4)$$

$$F_e = \mathcal{E}_1(g_1(u, v)) + \mathcal{E}_2(g_2(u, v)), \quad (4.5)$$

for some $h_1(u, v), h_2(u, v), g_1(u, v), g_2(u, v) \in \mathbb{C}[u, v]$.

Proof. Since

$$(A_-^{(123)})^2F = (\partial_{x_1} + \partial_{x_2} + \partial_{x_3})F = 0,$$

we see that the x dependence in F is only through the variables u and v , i.e. we can write F as

$$F = \sum_{i_1, i_2, i_3=0}^1 G_{i_1, i_2, i_3}(u, v)\theta_1^{i_1}\theta_2^{i_2}\theta_3^{i_3},$$

where $G_{i_1, i_2, i_3}(u, v) \in \mathbb{C}[u, v]$. Substituting this into (4.2) and equating the coefficients in the different powers of $\theta_1, \theta_2, \theta_3$ shows that F will satisfy (4.2) if and only if the representation in (4.3) holds. \square

5. ACTION OF THE INTERMEDIATE CASIMIR OPERATORS ON $\ker(A_-^{(123)})$

Since for every nonempty $S \subset \{1, 2, 3\}$ the Casimir operator $Q^{(S)}$ commutes with $A_-^{(123)}$, it follows that $Q^{(S)}$ preserves the space of solutions of equation (4.2). Below we compute the action of the intermediate Casimir operators $Q^{(12)}$, $Q^{(13)}$ and $Q^{(23)}$ on the basis of solutions of (4.2) described in Proposition 4.1. For $i \neq j$ we set $\nu_{ij} = \nu_i + \nu_j$.

5.1. Action of $Q^{(12)}$. We have

$$\begin{aligned}Q^{(12)} \circ \mathcal{O}_1 &= -\mathcal{O}_1 \circ (2u\partial_u + 2\nu_{12} + 1/2) - \mathcal{O}_2 \circ (2u\partial_v), \\ Q^{(12)} \circ \mathcal{O}_2 &= \mathcal{O}_1 \circ (2u\partial_u + 4\nu_1) + \mathcal{O}_2 \circ (2u\partial_u + 2\nu_{12} - 1/2), \\ Q^{(12)} \circ \mathcal{E}_1 &= -\mathcal{E}_1 \circ (2u\partial_u + 2\nu_{12} + 1/2) + \mathcal{E}_2 \circ (2u), \\ Q^{(12)} \circ \mathcal{E}_2 &= -\mathcal{E}_1 \circ (2u\partial_u\partial_v + 4\nu_1\partial_v) + \mathcal{E}_2 \circ (2u\partial_u + 2\nu_{12} - 1/2).\end{aligned}$$

5.2. Action of $Q^{(13)}$. We have

$$\begin{aligned} Q^{(13)} \circ \mathcal{O}_1 &= -\mathcal{O}_1 \circ (2\nu_1 - 2\nu_3 - 1/2) - \mathcal{O}_2 \circ (2(u+v)\partial_v + 4\nu_3), \\ Q^{(13)} \circ \mathcal{O}_2 &= -\mathcal{O}_1 \circ (2(u+v)\partial_u + 4\nu_1) - \mathcal{O}_2 \circ (2\nu_1 - 2\nu_3 + 1/2), \\ Q^{(13)} \circ \mathcal{E}_1 &= -\mathcal{E}_1 \circ (2\nu_{13} - 3/2) - \mathcal{E}_2 \circ (2u + 2v), \\ Q^{(13)} \circ \mathcal{E}_2 &= -\mathcal{E}_1 \circ (2(u+v)\partial_u\partial_v + 4\nu_3\partial_u + 4\nu_1\partial_v) + \mathcal{E}_2 \circ (2\nu_{13} - 1/2). \end{aligned}$$

5.3. Action of $Q^{(23)}$. We have

$$\begin{aligned} Q^{(23)} \circ \mathcal{O}_1 &= \mathcal{O}_1 \circ (2v\partial_v + 2\nu_{23} - 1/2) + \mathcal{O}_2 \circ (2v\partial_v + 4\nu_3), \\ Q^{(23)} \circ \mathcal{O}_2 &= -\mathcal{O}_1 \circ (2v\partial_u) - \mathcal{O}_2 \circ (2v\partial_v + 2\nu_{23} + 1/2), \\ Q^{(23)} \circ \mathcal{E}_1 &= -\mathcal{E}_1 \circ (2v\partial_v + 2\nu_{23} + 1/2) + \mathcal{E}_2 \circ (2v), \\ Q^{(23)} \circ \mathcal{E}_2 &= -\mathcal{E}_1 \circ (2v\partial_u\partial_v + 4\nu_3\partial_u) + \mathcal{E}_2 \circ (2v\partial_v + 2\nu_{23} - 1/2). \end{aligned}$$

6. DIAGONALIZATION OF $Q^{(123)}$ AND $Q^{(12)}$ IN THE ODD SUBSPACE

6.1. Spectral equations for $Q^{(123)}$. Consider an odd element

$$f(\mathbf{x}; \theta) = \mathcal{O}_1(h(u, v)) + \mathcal{O}_2(g(u, v)) \in \ker A_-^{(123)}, \quad (6.1)$$

where $\mathbf{x} = (x_1, x_2, x_3)$ and $\theta = (\theta_1, \theta_2, \theta_3)$. Then the spectral equation

$$Q^{(123)} f(\mathbf{x}; \theta) = \mu f(\mathbf{x}; \theta) \quad (6.2)$$

is equivalent to the equations

$$\begin{aligned} uh_u + vh_v &= -(\nu_{123} + \mu/2 + 1/4)h, \\ ug_u + vg_v &= -(\nu_{123} + \mu/2 + 1/4)g, \end{aligned}$$

where $\nu_{123} = \nu_1 + \nu_2 + \nu_3$. The last two equations are satisfied if and only if h and g are homogeneous in u and v of the same degree N , where

$$N = -(\nu_{123} + \mu/2 + 1/4),$$

or equivalently

$$\mu = -(2N + 2\nu_{123} + 1/2).$$

Since $Q^{(123)}$ commutes with each of the operators $Q^{(12)}$, $Q^{(13)}$, $Q^{(23)}$, we can look at their restrictions on the space of solutions of (6.2).

6.2. Spectral equations for $Q^{(12)}$. Consider now the equation

$$Q^{(12)} f(\mathbf{x}; \theta) = \lambda f(\mathbf{x}; \theta), \quad (6.3)$$

where $\lambda \in \mathbb{C}$ and $f(\mathbf{x}; \theta)$ is the element in $\ker A_-^{(123)}$ in (6.1). The coefficient of θ_2 shows that

$$(1 + 2\lambda + 4\nu_1 - 4\nu_2)g = (1 + 2\lambda + 4\nu_{12})h + 4uh_u - 4vh_v. \quad (6.4)$$

It is straightforward to check that if g and h are homogeneous polynomials in u and v of the same degree satisfying (6.3), then $1 + 2\lambda + 4\nu_1 - 4\nu_2$ can be zero only when $\nu_1 = \nu_2 = 0$. Thus for generic ν_1, ν_2 we can assume that

$1 + 2\lambda + 4\nu_1 - 4\nu_2 \neq 0$ and therefore (6.4) determines g uniquely from h as follows

$$g = \frac{(1 + 2\lambda + 4\nu_{12})h + 4uh_u - 4uh_v}{1 + 2\lambda + 4\nu_1 - 4\nu_2}. \quad (6.5)$$

Substituting the last formula into (6.3), we see that (6.3) holds if and only if $h(u, v)$ satisfies the equation

$$(u^2\partial_u^2 - u^2\partial_u\partial_v + (2\nu_{12} + 1)u\partial_u - (2\nu_1 + 1)u\partial_v + (\nu_{12}^2 - (2\lambda + 1)^2/16))h = 0. \quad (6.6)$$

If $h(u, v)$ is homogeneous of degree N we can write it as

$$h(u, v) = u^N\phi(v/u), \quad (6.7)$$

and substituting the last equation in (6.6), we obtain the following equation for $\phi(z)$:

$$\left(z(1+z)\partial_z^2 - ((2\nu_{12} + 2N - 1)z + 2\nu_1 + N)\partial_z + (N + \nu_{12})^2 - (2\lambda + 1)^2/16\right)\phi(z) = 0. \quad (6.8)$$

If the last equation has a solution which is a polynomial of degree $k \leq N$, then the coefficient of z^k yields

$$(N - k + \nu_{12})^2 = \frac{(2\lambda + 1)^2}{16}.$$

This leads to

$$\frac{2\lambda + 1}{4} = \pm(N - k + \nu_{12}), \quad \text{or equivalently} \quad \lambda = \pm 2(N - k + \nu_{12}) - \frac{1}{2} \quad (6.9)$$

and for these values of λ , equation (6.8) reduces to

$$\left(z(1+z)\partial_z^2 - ((2\nu_{12} + 2N - 1)z + 2\nu_1 + N)\partial_z + k(2N + 2\nu_{12} - k)\right)\phi(z) = 0. \quad (6.10)$$

Note that this equation is the hypergeometric differential equation with three regular singular points: 0 , -1 and ∞ . For generic parameters ν_1, ν_2 , up to a constant factor, it has a unique polynomial solution of degree k given in terms of the Jacobi polynomial by

$$\phi(z) = P_k^{(-N-2\nu_1-1, -N-2\nu_2)}(1+2z). \quad (6.11)$$

Let us recall that the Jacobi polynomials are expressed as follows in terms of the hypergeometric functions

$$P_k^{(\alpha, \beta)}(x) = \frac{(\alpha + 1)_k}{k!} {}_2F_1\left(-k, k + \alpha + \beta + 1; \alpha + 1; \frac{1-x}{2}\right). \quad (6.12)$$

Summarizing the above computations we obtain the following theorem describing the common eigenfunctions of the operators $Q^{(123)}$ and $Q^{(12)}$ which can be simultaneously diagonalized on $\ker(A_-^{(123)}) \cap \mathbb{V}^0$.

Theorem 6.1. For $N \in \mathbb{N}_0$ and $k \in \{0, \dots, N\}$, let

$$h_{k,N}(u, v) = u^N P_k^{(-N-2\nu_1-1, -N-2\nu_2)} \left(1 + 2\frac{v}{u}\right), \quad (6.13a)$$

$$g_{k,N}^+(u, v) = \frac{2N + 2\nu_{12} - k}{N + 2\nu_1 - k} u^N P_k^{(-N-2\nu_1, -N-2\nu_2)} \left(1 + 2\frac{v}{u}\right), \quad (6.13b)$$

$$g_{k,N}^-(u, v) = -u^N P_{k-1}^{(-N-2\nu_1, -N-2\nu_2)} \left(1 + 2\frac{v}{u}\right). \quad (6.13c)$$

Then

$$\{f_{k,N}^+(\mathbf{x}; \theta), f_{k,N}^-(\mathbf{x}; \theta) : N \in \mathbb{N}_0, 0 \leq k \leq N\}$$

where

$$f_{k,N}^\pm(\mathbf{x}; \theta) = \mathcal{O}_1(h_{k,N}(u, v)) + \mathcal{O}_2(g_{k,N}^\pm(u, v)),$$

form a basis of $\ker(A_-^{(123)}) \cap \mathbb{V}^o$ and

$$Q^{(123)} f_{k,N}^\pm(\mathbf{x}; \theta) = -(2N + 2\nu_{123} + 1/2) f_{k,N}^\pm(\mathbf{x}; \theta), \quad (6.14a)$$

$$Q^{(12)} f_{k,N}^\pm(\mathbf{x}; \theta) = (\pm 2(N - k + \nu_{12}) - 1/2) f_{k,N}^\pm(\mathbf{x}; \theta). \quad (6.14b)$$

Proof. The result for $h_{k,N}(u, v)$ follows directly from the discussion preceding the theorem. Using this result for $h_{k,N}(u, v)$ and the eigenvalues $\lambda = \pm 2(N - k + \nu_{12}) - 1/2$ in (6.5), one gets the following expressions for $g_{k,N}^\pm(u, v)$:

$$g_{k,N}^+(u, v) = \frac{(N - k + 2\nu_{12})h_{k,N}(u, v) + u(\partial_u h_{k,N}(u, v) - \partial_v h_{k,N}(u, v))}{N - k + 2\nu_1},$$

$$g_{k,N}^-(u, v) = \frac{(N - k)h_{k,N}(u, v) - u(\partial_u h_{k,N}(u, v) - \partial_v h_{k,N}(u, v))}{N - k + 2\nu_2}.$$

Note that

$$\text{if } h = u^N \phi(v/u), \quad \text{then } (u\partial_u - u\partial_v)h = u^N [(N - (t+1)\partial_t)\phi(t)]|_{t=v/u}.$$

Using the above formula and the identity

$$(-k + (t+1)\partial_t) P_k^{(\alpha, \beta)}(1+2t) = (k + \beta) P_{k-1}^{(\alpha+1, \beta)}(1+2t)$$

we obtain the explicit formulas (6.13b)-(6.13c) for $g_{k,N}^\pm(u, v)$. \square

Using the fact that the intermediate Casimir operators satisfy the Bannai-Ito algebra and knowing the spectrum of $Q^{(12)}$ and $Q^{(123)}$, one can deduce that the action of $Q^{(23)}$ is tridiagonal as follows

$$Q^{(23)} f_{k,N}^+(\mathbf{x}; \theta) = \alpha_k^+ f_{k,N}^-(\mathbf{x}; \theta) + \beta_k^+ f_{k,N}^+(\mathbf{x}; \theta) + \gamma_k^+ f_{k+1,N}^-(\mathbf{x}; \theta), \quad (6.16a)$$

$$Q^{(23)} f_{k,N}^-(\mathbf{x}; \theta) = \alpha_{k-1}^- f_{k-1,N}^+(\mathbf{x}; \theta) + \beta_k^- f_{k,N}^-(\mathbf{x}; \theta) + \gamma_k^- f_{k,N}^+(\mathbf{x}; \theta). \quad (6.16b)$$

The coefficients are also constrained by the algebra as follows:

$$\alpha_k^- \gamma_k^+ = \frac{4(k+1)(N-k)(2\nu_{12}+2N-k)(2\nu_{12}+N-k-1)}{(2\nu_{12}+2N-2k-1)^2}, \quad (6.17)$$

$$\alpha_k^+ \gamma_k^- = \frac{(2\nu_1+N-k)(2\nu_2+N-k)(2\nu_3+k)(2\nu_{123}+2N-k)}{(\nu_{12}+N-k)^2}, \quad (6.18)$$

$$\beta_k^\pm = \pm \frac{(\nu_1-\nu_2)(\nu_{12}+2\nu_3+N)}{\nu_{12}+N-k} \mp \frac{(2\nu_{12}-1)(2\nu_{12}+2N+1)}{2(2\nu_{12}+2N-2k \mp 1)}. \quad (6.19)$$

The exact expressions of these coefficients are given in the following corollary.

Corollary 6.2. *The coefficients in relations (6.16) are*

$$\begin{aligned} \alpha_k^+ &= \frac{(2\nu_2+N-k)(2\nu_{123}+2N-k)}{\nu_{12}+N-k}, \\ \alpha_k^- &= \frac{2(2\nu_1+N-k+1)(2\nu_{12}+N-k)}{2\nu_{12}+2N-2k+1}, \\ \gamma_k^+ &= \frac{2(N-k)(k+1)(2\nu_{12}+2N-k)}{(2\nu_1+N-k)(2\nu_{12}+2N-2k-1)}, \\ \gamma_k^- &= \frac{(2\nu_3+k)(2\nu_1+N-k)}{\nu_{12}+N-k}. \end{aligned}$$

Proof. The coefficients of θ_1 in the relations (6.16), computed by using the explicit formulas for $f_{k,N}^\pm(\mathbf{x};\theta)$, provide constraints between the coefficients α^\pm , β^\pm and γ^\pm . Combing these constraints with the expression (6.19) of β_k^\pm , one gets the expressions in the corollary. \square

The values of these coefficients are compatible with the relations (6.17)-(6.18).

7. DIAGONALIZATION OF $Q^{(123)}$ AND $Q^{(12)}$ IN THE EVEN SUBSPACE

7.1. **Spectral equations for $Q^{(123)}$.** Consider an even element

$$f(\mathbf{x};\theta) = \mathcal{E}_1(h(u,v)) + \mathcal{E}_2(g(u,v)) \in \ker A_-^{(123)}. \quad (7.1)$$

Then the spectral equation

$$Q^{(123)} f(\mathbf{x};\theta) = \mu f(\mathbf{x};\theta) \quad (7.2)$$

is equivalent to the equations

$$\begin{aligned} uh_u + vh_v &= (-\nu_{123} + \mu/2 - 3/4)h, \\ ug_u + vg_v &= (-\nu_{123} + \mu/2 + 1/4)g. \end{aligned}$$

The last two equations are satisfied if and only if g is homogeneous of degree N and h is homogeneous of degree $N-1$, where

$$N = -\nu_{123} + \mu/2 + 1/4.$$

or equivalently

$$\mu = 2N + 2\nu_{123} - 1/2.$$

7.2. Spectral equations for $Q^{(12)}$. Consider now the equation

$$Q^{(12)}f(\mathbf{x};\theta) = \lambda f(\mathbf{x};\theta), \quad (7.3)$$

where $\lambda \in \mathbb{C}$ and $f(\mathbf{x};\theta)$ is the element in $\ker A_-^{(123)}$ in (7.1). The free term (i.e. the coefficient of $\theta_1^0\theta_2^0\theta_3^0$) shows that

$$h(u,v) = \frac{(1+2\lambda-4\nu_{12})}{4u}g(u,v) - \partial_u g(u,v). \quad (7.4)$$

Substituting the last formula into (7.3), we see that (7.3) holds if and only if $g(u,v)$ satisfies the equation

$$(u^2\partial_u^2 - u^2\partial_u\partial_v + 2\nu_{12}u\partial_u - 2\nu_1u\partial_v + \nu_{12}(\nu_{12}-1) + 1/4 - (2\lambda-1)^2/16)g = 0. \quad (7.5)$$

If $g(u,v)$ is homogeneous of degree N we can write it as

$$g(u,v) = u^N\phi(v/u), \quad (7.6)$$

and substituting the last formula in (7.5), we obtain the following equation for $\phi(z)$:

$$\begin{aligned} & \left(z(1+z)\partial_z^2 - (2(\nu_{12}+N-1)z + 2\nu_1 + N-1)\partial_z \right. \\ & \left. + (N + \nu_{12} - 1/2)^2 - (2\lambda - 1)^2/16 \right) \phi(z) = 0. \end{aligned} \quad (7.7)$$

If the last equation has a solution $\phi(z)$ which is a polynomial of degree $k \leq N$, then the coefficient of z^k yields

$$(N - k + \nu_{12} - 1/2)^2 = \frac{(2\lambda - 1)^2}{16}.$$

Equivalently

$$\lambda = \pm 2(N - k + \nu_{12} - 1/2) + 1/2 \quad (7.8)$$

and for these values of λ equation (7.7) reduces to

$$\begin{aligned} & \left(z(1+z)\partial_z^2 - (2(\nu_{12}+N-1)z + 2\nu_1 + N-1)\partial_z \right. \\ & \left. + k(2N + 2\nu_{12} - k - 1) \right) \phi(z) = 0. \end{aligned} \quad (7.9)$$

For generic parameters ν_1, ν_2 and up to a constant factor, this equation has a unique polynomial solution of degree k given by

$$\phi(z) = P_k^{(-N-2\nu_1, -N-2\nu_2)}(1+2z), \quad (7.10)$$

where P_k is the Jacobi polynomial (6.12). The next theorem describes the common eigenfunctions of the commuting operators $Q^{(123)}$ and $Q^{(12)}$ which can be simultaneously diagonalized on $\ker(A_-^{(123)}) \cap \mathbb{V}^e$.

Theorem 7.1. For $N \in \mathbb{N}_0$ and $k \in \{0, \dots, N\}$, let

$$g_{k,N}(u, v) = u^N P_k^{(-N-2\nu_1, -N-2\nu_2)} \left(1 + 2\frac{v}{u}\right), \quad (7.11a)$$

$$h_{k,N}^+(u, v) = u^{N-1} (N + 2\nu_1 - k) P_{k-1}^{(-N-2\nu_1, -N-2\nu_2+1)} \left(1 + 2\frac{v}{u}\right), \quad (7.11b)$$

$$h_{k,N}^-(u, v) = (k + 1 - 2N - 2\nu_{12}) u^{N-1} P_k^{(-N-2\nu_1, -N-2\nu_2+1)} \left(1 + 2\frac{v}{u}\right). \quad (7.11c)$$

Then

$$\{f_{k,N}^+(\mathbf{x}; \theta) : N \in \mathbb{N}_0, 0 \leq k \leq N\} \cup \{f_{k,N}^-(\mathbf{x}; \theta) : N \in \mathbb{N}_0, 0 \leq k < N\}$$

where

$$f_{k,N}^\pm(\mathbf{x}; \theta) = \mathcal{E}_1(h_{k,N}^\pm(u, v)) + \mathcal{E}_2(g_{k,N}(u, v)),$$

form a basis of $\ker(A_-^{(123)}) \cap \mathbb{V}^e$ and

$$Q^{(123)} f_{k,N}^\pm(\mathbf{x}; \theta) = (2N + 2\nu_{123} - 1/2) f_{k,N}^\pm(\mathbf{x}; \theta), \quad (7.12a)$$

$$Q^{(12)} f_{k,N}^\pm(\mathbf{x}; \theta) = (\pm 2(N - k + \nu_{12} - 1/2) + 1/2) f_{k,N}^\pm(\mathbf{x}; \theta). \quad (7.12b)$$

Proof. The result for $g_{k,N}(u, v)$ follows directly from the discussion preceding the theorem. Using the formula for $g_{k,N}(u, v)$ and the eigenvalues $\lambda = \pm 2(N - k + \nu_{12} - 1/2) + 1/2$ in (7.4), one gets the following expressions for $h_{k,N}^\pm(u, v)$:

$$h_{k,N}^+(u, v) = \frac{N - k}{u} g_{k,N}(u, v) - \partial_u g_{k,N}(u, v), \quad (7.13)$$

$$h_{k,N}^-(u, v) = -\frac{N - k + 2\nu_{12} - 1}{u} g_{k,N}(u, v) - \partial_u g_{k,N}(u, v). \quad (7.14)$$

Note that

$$\text{if } g = u^N \phi(v/u), \quad \text{then } u \partial_u g = u^N [(N - t \partial_t) \phi(t)]|_{t=v/u}.$$

Using the above formula and the identity

$$(k - t \partial_t) P_k^{(\alpha, \beta)}(1 + 2t) = (\alpha + k) P_{k-1}^{(\alpha, \beta+1)}(1 + 2t)$$

we can write explicit formulas for $h_{k,N}^\pm(u, v)$ in terms of the Jacobi polynomials as stated in the theorem. \square

Similarly to the arguments in the odd space, combining the fact that the intermediate Casimir operators satisfy the Bannai-Ito algebra with the spectrum of $Q^{(12)}$ and $Q^{(123)}$, one can deduce that the action of $Q^{(23)}$ is tridiagonal as follows

$$Q^{(23)} f_{k,N}^+(\mathbf{x}; \theta) = \alpha_k^+ f_{k-1,N}^-(\mathbf{x}; \theta) + \beta_k^+ f_{k,N}^+(\mathbf{x}; \theta) + \gamma_k^+ f_{k,N}^-(\mathbf{x}; \theta), \quad (7.15a)$$

$$Q^{(23)} f_{k,N}^-(\mathbf{x}; \theta) = \alpha_k^- f_{k,N}^+(\mathbf{x}; \theta) + \beta_{k+1}^- f_{k,N}^-(\mathbf{x}; \theta) + \gamma_{k+1}^- f_{k+1,N}^+(\mathbf{x}; \theta). \quad (7.15b)$$

The coefficients are also constrained by the algebra as follows:

$$\alpha_k^+ \gamma_k^+ = \frac{k(2\nu_1 + N - k)(2\nu_2 + N - k)(2\nu_{12} + 2N - k)}{(\nu_{12} + N - k)^2}, \quad (7.16)$$

$$\alpha_k^- \gamma_k^+ = \frac{4(2\nu_3 + k)(N - k)(2\nu_{12} + N - k - 1)(2\nu_{123} + 2N - k - 1)}{(2\nu_{12} + 2N - 2k - 1)^2}, \quad (7.17)$$

$$\beta_k^\pm = \mp \frac{(\nu_1 - \nu_2)(\nu_{12} + N)}{\nu_{12} + N - k} \pm \frac{(2\nu_{12} - 1)(2\nu_{12} + 4\nu_3 + 2N + 1)}{2(2\nu_{12} + 2N - 2k \mp 1)}. \quad (7.18)$$

The exact expressions of these coefficients are given in the following corollary.

Corollary 7.2. *The coefficients in relations (7.15) are*

$$\begin{aligned} \alpha_k^+ &= \frac{(2\nu_1 + N - k)(2\nu_2 + N - k)}{\nu_{12} + N - k}, \\ \alpha_k^- &= \frac{2(2\nu_{12} + N - k - 1)(2\nu_{123} + 2N - k - 1)}{2\nu_{12} + 2N - 2k - 1}, \\ \gamma_k^+ &= \frac{2(2\nu_3 + k)(N - k)}{2\nu_{12} + 2N - 2k - 1}, \\ \gamma_k^- &= \frac{k(2\nu_{12} + 2N - k)}{\nu_{12} + N - k}. \end{aligned}$$

Proof. The coefficients without θ 's in the relations (7.15), computed by using the explicit expressions for $f_{k,N}^\pm(\mathbf{x}; \theta)$, provide constraints between the coefficients α^\pm , β^\pm and γ^\pm . Combining these constraints with formula (7.18) for β_k^\pm , one gets the expressions stated in the corollary. \square

The values of these coefficients are compatible with the relations (7.16)-(7.17).

8. CONCLUSION

Summing up, we have provided a model for the Bannai–Ito algebra on a superspace with \mathbb{C}^2 as body and with soul generated by three anticommuting Grassmann variables. The even and odd basis vectors were found to each have two components and to be realized in terms of Jacobi polynomials. The tridiagonal action of one Bannai–Ito generator in the eigenbasis of the other was explicitly calculated. The formulas thus obtained reflect contiguous relations of the Jacobi polynomials.

It is worth mentioning that the Bannai–Ito algebra is isomorphic to the degenerate double affine Hecke algebra (DAHA) [17] which has thus been modelled here in superspace by the same token.

For the sake of completeness, let us mention that an embedding of the Bannai–Ito algebra into $\mathfrak{osp}(1|2)$ was presented in [18] where an analytic realization of $\mathfrak{osp}(1|2)$ in terms of Dunkl operators was used to connect to some -1 -polynomials. It should be stressed that the Bannai–Ito model that results from combining the realization of $\mathfrak{osp}(1|2)$ used throughout in

this paper with this embedding is clearly not the one that is attained from the dimensional reduction performed here.

Finally, it would definitely be of interest to develop along the lines pursued here realizations on superspaces of the higher rank Bannai–Ito algebras that have been constructed from the intermediate Casimir operators arising in manifold tensor products of $\mathfrak{osp}(1|2)$ [19].

ACKNOWLEDGMENTS

The authors wish to acknowledge the enlightening discussions they had with Wouter van de Vijver in the early stages of the project. NC and PI are grateful to the Centre de Recherches Mathématiques (CRM) for supporting visits during which the research reported here was carried out. NC is funded by the international research project AAPT of the CNRS and the ANR Project AHA ANR-18-CE40-0001. HDB is supported by the EOS-FWO Research Project grant 30889451. PI is supported in part by Simons Foundation Grant #635462. LV is supported in part by a Discovery Grant from NSERC.

REFERENCES

- [1] Willard Miller. *Lie theory and special functions*, volume 43. Academic Press New York, 1968.
- [2] Hendrik De Bie, Vincent X Genest, Satoshi Tsujimoto, Luc Vinet, and Alexei Zhedanov. The Bannai–Ito algebra and some applications. In *Journal of Physics: Conference Series*, volume 597, page 012001. IOP publishing, 2015.
- [3] Eiichi Bannai, Etsuko Bannai, Tatsuro Ito, and Rie Tanaka. Algebraic combinatorics. In *Algebraic Combinatorics*. De Gruyter, 2021.
- [4] Satoshi Tsujimoto, Luc Vinet, and Alexei Zhedanov. Dunkl shift operators and Bannai–Ito polynomials. *Advances in Mathematics*, 229(4):2123–2158, 2012.
- [5] Roelof Koekoek, Peter A Lesky, René F Swarttouw, Roelof Koekoek, Peter A Lesky, and René F Swarttouw. *Hypergeometric orthogonal polynomials*. Springer, 2010.
- [6] Vincent X Genest, Luc Vinet, and Alexei Zhedanov. The Racah algebra and superintegrable models. In *Journal of Physics: Conference Series*, volume 512, page 012011. IOP Publishing, 2014.
- [7] Vincent X Genest, Luc Vinet, and Alexei Zhedanov. Embeddings of the Racah algebra into the Bannai–Ito algebra. *SIGMA. Symmetry, Integrability and Geometry: Methods and Applications*, 11:050, 2015.
- [8] Suogang Gao, Yan Wang, and Bo Hou. The classification of Leonard triples of Racah type. *Linear algebra and its applications*, 439(7):1834–1861, 2013.
- [9] Askold Perelomov. *Generalized coherent states and their applications*. Springer Science & Business Media, 2012.
- [10] Vincent X Genest, Luc Vinet, and Alexei Zhedanov. Superintegrability in two dimensions and the Racah–Wilson algebra. *Letters in Mathematical Physics*, 104:931–952, 2014.
- [11] Vincent X Genest, Luc Vinet, and Alexei Zhedanov. The equitable Racah algebra from three algebras. *Journal of Physics A: Mathematical and Theoretical*, 47(2):025203, 2013.
- [12] Hendrik De Bie, Vincent X Genest, Wouter van de Vijver, and Luc Vinet. A higher rank Racah algebra and the Laplace–Dunkl operator. *Journal of Physics A: Mathematical and Theoretical*, 51(2):025203, 2017.

- [13] Hendrik De Bie, Plamen Iliev, and Luc Vinet. Bargmann and Barut-Girardello models for the Racah algebra. *Journal of Mathematical Physics*, 60(1):011701, 2019.
- [14] Vincent Genest, Luc Vinet, and Alexei Zhedanov. The Bannai-Ito polynomials as Racah coefficients of the $\mathfrak{sl}_{-1}(2)$ algebra. *Proceedings of the American Mathematical Society*, 142(5):1545–1560, 2014.
- [15] Bryce DeWitt. *Supermanifolds*. Cambridge University Press, 1992.
- [16] L Frappat, A Sciarrino, and Paul Sorba. Dictionary on Lie superalgebras. *arXiv preprint hep-th/9607161*, 1996.
- [17] Vincent Genest, Luc Vinet, and Alexei Zhedanov. The non-symmetric Wilson polynomials are the Bannai–Ito polynomials. *Proceedings of the American Mathematical Society*, 144(12):5217–5226, 2016.
- [18] Pascal Baseilhac, Vincent X Genest, Luc Vinet, and Alexei Zhedanov. An embedding of the Bannai–Ito algebra in $\mathcal{U}(\mathfrak{osp}(1, 2))$ and -1 -polynomials. *Letters in Mathematical Physics*, 108:1623–1634, 2018.
- [19] Hendrik De Bie, Vincent X Genest, and Luc Vinet. The \mathbb{Z}_2^3 Dirac–Dunkl operator and a higher rank Bannai–Ito algebra. *Advances in Mathematics*, 303:390–414, 2016.

NC, INSTITUT DENIS-POISSON CNRS/UMR 7013 - UNIVERSITÉ DE TOURS - UNIVERSITÉ D'ORLÉANS, PARC DE GRANDMONT, 37200 TOURS, FRANCE
Email address: `crampe1977@gmail.com`

HDB, CLIFFORD RESEARCH GROUP, DEPARTMENT OF ELECTRONICS AND INFORMATION SYSTEMS, FACULTY OF ENGINEERING AND ARCHITECTURE, GHENT UNIVERSITY, KRIJGSLAAN 281-S8, 9000, GENT (BELGIUM)
Email address: `Hendrik.DeBie@UGent.be`

PI, SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160, USA
Email address: `iliev@math.gatech.edu`

LV, IVADO AND CENTRE DE RECHERCHES MATHÉMATIQUES, UNIVERSITÉ DE MONTRÉAL, P.O. BOX 6128, CENTRE-VILLE STATION, MONTRÉAL, QC H3C 3J7, CANADA
Email address: `vinet@crm.umontreal.ca`