



HAL
open science

Subthalamic high-frequency deep brain stimulation reduces addiction-like alcohol use and the possible negative influence of a peer presence

Lucie Vignal, Cassandre Vielle, Maya Williams, Nicolas Maurice, Mickael Degoulet, Christelle Baunez

► To cite this version:

Lucie Vignal, Cassandre Vielle, Maya Williams, Nicolas Maurice, Mickael Degoulet, et al.. Subthalamic high-frequency deep brain stimulation reduces addiction-like alcohol use and the possible negative influence of a peer presence. *Psychopharmacology*, 2024, 10.1007/s00213-024-06532-w . hal-04741200

HAL Id: hal-04741200

<https://hal.science/hal-04741200v1>

Submitted on 17 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **Title: SUBTHALAMIC HIGH-FREQUENCY DEEP BRAIN STIMULATION REDUCES**
2 **ADDICTION-LIKE ALCOHOL USE AND THE POSSIBLE NEGATIVE INFLUENCE OF A**
3 **PEER PRESENCE**

4
5 **Authors:** Lucie Vignal^{#1}, Cassandre Vielle^{#1}, Maya Williams¹, Nicolas Maurice¹, Mickael Degoulet¹,
6 Christelle Baunez ^{1*}

7 **Affiliations:**

8 ¹ Institut de Neurosciences de la Timone, UMR 7289 CNRS & Aix-Marseille Université ; 13005
9 Marseille, France.

10 * Corresponding author. Email: christelle.baunez@univ-amu.fr

11 #: equal contribution

12 This work was funded by the Institut de Recherche en Santé Publique in collaboration with Alliance
13 Aviesan , and by the Institut de lutte contre le Cancer.

14
15 **Abstract:**

16 *Rationale:* The immediate social context significantly influences alcohol consumption in humans.
17 Recent studies have revealed that peer presence could modulate drugs use in rats. The most efficient
18 condition to reduce cocaine intake is the presence of a stranger peer, naive to drugs. Deep brain
19 stimulation (DBS) of the Subthalamic Nucleus (STN), which was shown to have beneficial effects on
20 addiction to cocaine or alcohol, also modulate the protective influence of peer's presence on cocaine
21 use.

22 *Objectives:* This study aimed to: 1) explore how the presence of an alcohol-naive stranger peer affects
23 recreational and escalated alcohol intake, and 2) assess the involvement of STN on alcohol use and in
24 the modulation induced by the presence of an alcohol-naive stranger peer.

25 *Methods:* Rats with STN DBS and control animals self-administered 10% (v/v) ethanol in presence, or
26 absence, of an alcohol-naive stranger peer, before and after escalation of ethanol intake (observed after
27 intermittent alcohol (20% (v/v) ethanol) access).

28 *Results:* Neither STN DBS nor the presence of an alcohol-naive stranger peer modulated significantly
29 recreational alcohol intake. After the escalation procedure, STN DBS reduced ethanol consumption. The
30 presence of an alcohol-naive stranger peer increased consumption only in low drinkers, which effect
31 was suppressed by STN DBS.

32 *Conclusions:* These results highlight the influence of a peer's presence on escalated alcohol intake, and
33 confirm the role of STN in addiction-like alcohol intake and in the social influence on drug consumption.
34

35 **Keyword:** alcohol, social context, subthalamic nucleus, deep brain stimulation, peer's presence
36

37 **Main text:**

38 **Introduction**

39 Drug consumption is a complex behavior influenced by a multitude of factors, including
40 environmental context and social interactions. Social context has been demonstrated to play a crucial
41 role in modulating drug intake in both human and animal models. Indeed, social factors are known to
42 affect multiples aspects of drug use such as initiation, maintenance, attempts to quit, and relapse,
43 potentially boosting, mediating, or reducing addictive behaviors (Bardo et al. 2013; Pelloux et al. 2019;
44 Venniro et al. 2018). Proximal social factors, such as the presence of peers during drug consumption,
45 have a significant influence by decreasing or increasing drug intake (Pelloux et al. 2019; Strickland and
46 Smith 2014; 2015). The identity of the peer is also a critical element that can modulate drug
47 consumption. For cocaine, the presence of a non-familiar (stranger) peer leads to a stronger reduction in
48 cocaine intake than the presence of a familiar one in both rats and humans (Giorla et al. 2022). The
49 largest reduction was observed in presence of a stranger cocaine-naive peer. A similar beneficial effect

50 of presence of such a peer has been also shown on addictive-like (i.e. escalated) cocaine consumption
51 in rats (Vielle et al. preprint).

52 However, while the presence of peers is protective against cocaine use, its effect might be opposite for
53 alcohol. Indeed, rats chose alcohol over social interaction in a choice procedure (Augier et al. 2022;
54 Marchant et al. 2022), and consumed more alcohol in presence of a non-naive alcohol peer, regardless
55 of their familiarity (Sicre et al., unpublished data). These results suggest that the influence of social
56 presence on drug intake depends on the substance used. Nevertheless, the effect of the presence of an
57 alcohol-naive stranger has not been assessed. Furthermore, the influence of social presence has only
58 been examined in the context of recreational alcohol use. Consequently, the modulation of recreational
59 and escalated ethanol intake by the presence of a stranger peer, naive to alcohol remains to be explored.
60 Over the past few years, the Subthalamic Nucleus (STN), the only glutamatergic structure of the basal
61 ganglia, has gained increasing significance in the reward circuit, in motivational processes and in drug
62 addiction (Baunez and Lardeux 2011). STN inactivation induced by lesions, deep brain stimulation
63 (DBS) or optogenetic manipulations, has shown beneficial effects across various addiction criteria:
64 motivation (Baunez et al. 2005; Lardeux and Baunez 2008; Rouaud et al. 2010; Pelloux and Baunez
65 2017), escalation of drug intake (Pelloux and Baunez 2017; Pelloux et al. 2018), maintenance of
66 escalated drug intake (Vielle et al. preprint) and compulsive drug-seeking behavior (Degoulet et al.
67 2021). Additionally, it has been demonstrated that the STN is involved in mediating the influence of the
68 social context on cocaine and alcohol use. Indeed, inactivation of the STN modulates the modifications
69 caused by the social context (presence of a peer or playback of ultrasonic vocalizations) on cocaine and
70 alcohol intake (Giorla et al. 2022; Montanari et al. 2018; Sicre et al., unpublished data , Vielle et al.
71 preprint; 2021).

72 In this study, we thus examined the influence of the presence of an alcohol-naive male stranger peer on
73 both recreational and escalated alcohol intake in male rats. Furthermore, we investigated the contribution
74 of the STN in these processes by using high-frequency (130 Hz) DBS.

75 **Materials and Methods**

76 **Animals**

77 32 Adult Lister-Hooded rats (~400 g, Charles River Laboratories, Saint-Germain-sur-l'Arbresle,
78 France) were used in these behavioral experiments. Only male rats were used to avoid inter-variability
79 of the hormonal cycle. Animals were paired housed, in Plexiglas cages with unlimited access to food
80 and water. Temperature- and humidity-controlled environments were maintained with an inverted 12 h
81 light/dark cycle (light -onset at 7 pm). All experiments were conducted during the dark cycle (8am-
82 8pm). Animals were handled daily. All animal care and use were conformed to the French regulation
83 (Decree 2013-118) and were approved by local ethic committee and the National French Agriculture
84 Ministry (authorization #3129.01). Experimental group sizes were determined by taking into accounts
85 the 3Rs rules to reduce the number of animals used.

86 **Electrode design for STN DBS**

87 Electrodes were made of Platinum-Iridium wires coated with Teflon (75 μm). Coating was removed
88 over 0.5 mm at the tips and two wires were inserted into a 16 mm stainless steel tubing to form an
89 electrode. Two electrodes, separated by 4.8 mm (twice the STN laterally), were soldered to an electric
90 connector, allowing connection with the stimulator. Electrodes (impedance= $20 \pm 2.25 \text{ k}\Omega$) and
91 connector were subsequently deeply bound, using a custom mold and dental cement. Finally, electrodes
92 were checked with an isolated battery to avoid electrical short circuits.

93 **Surgery**

94 All the animals underwent bilateral electrode implantation into the STN. They were anesthetized with
95 ketamine (Imalgen, Merial, 100 mg/kg, *i.p.*) and medetomidine (Domitor, Janssen, 0.5 mg/kg, *i.p.*). They
96 also received an antibiotic treatment by injection of amoxicillin (Citramox, LA, Pfizer, 100 mg/kg, *s.c.*)
97 and an injection of meloxicam (Metacam, Boehringer Ingelheim, 0.5 mg/kg, *s.c.*) for analgesia. Animals
98 were then placed in the stereotaxic frame (David Kopf apparatus) for the bilateral electrodes'
99 implantation into the STN (coordinates in mm: -3.7 AP, ± 2.4 L from bregma, -8.35 DV from skull

100 surface (Paxinos and Watson 2014), with the incisor bar at -3.3 mm). The electrodes were maintained
101 with a head-cap made of dental cement and screws anchored on the skull. Rats were awakened with an
102 injection of atipamezole (Antisedan, Janssen, 0.15 mg/kg, i.m.), an antagonist of medetomidine. Rats
103 were allowed to recover from the surgery for at least 7 days.

104 **Behavioral apparatus**

105 Behavioral experiment took place in homemade self-administration chambers (60x30x35 cm) divided
106 into two compartments by a metallic grid. The wall of one compartment per cage was equipped with
107 two non-retractable levers and a light above each. A recessed magazine (3.8-cm high and 3.8-cm wide
108 located 5.5 cm above the grid floor) containing a cup liquid receptacle with an 18-gauge pipe was
109 positioned between the two levers and connected to a pump outside the chamber. A syringe filled with
110 ethanol was positioned on the pump and connected with the 18-gauge pipe of the receptacle via a
111 catheter. All the chambers were controlled by a custom-built interface and associated software (built
112 and written by Y. Pelloux).

113 **Behavioral procedure**

114 An overview of the behavioral procedure is illustrated in **Fig. 1**.

115 **1-Experiment 1: recreational ethanol self-administration**

116 At least one week after surgery, rats began ethanol self-administration training. The ethanol delivery
117 was assigned to one of the two levers ('active lever') and counterbalanced between rats. An FR1
118 schedule of reinforcement was used, so each lever press delivered 100 μ L of 10 % ethanol in the cup
119 receptacle and switched on the cue-light above the lever. Each delivery was followed by a 20-s time-
120 out period, during which any press on the active lever was recorded as a perseveration but had no
121 consequence. Presses on the other lever ('inactive lever') had no consequence either. Each daily session
122 lasted 1h.

123 Rats were first trained to self-administer ethanol with sucrose, using a fading sucrose technique in which
124 the sucrose concentration in the 10% ethanol solution gradually decrease over sessions until zero. They
125 were also habituated to be connected to an electric cable (but without stimulation). Once consumption
126 became stable (<25% of variability in the number of ethanol rewards for 5 consecutive days), animals
127 were exposed to the testing period, during which rats were either STN-stimulated with DBS (ON group,
128 n=16) or remained unstimulated (OFF control group, n=16). This testing period consists of 5x1h
129 consecutive daily sessions of ethanol self-administration alone, followed by 5x1h-sessions in presence
130 of an alcohol-naive stranger peer ('With peer' condition), placed on the other side of the grid. To prevent
131 the emergence of a familiarity effect resulting from constantly using the same social stimulus, the peer
132 was changed every day.

133 **2-Escalation of ethanol intake through intermittent access to alcohol (IAA)**

134 After this initial testing period, animals (n=30, two animals were excluded at this point, see results
135 section) underwent an escalation of their ethanol intake, using the intermittent access to alcohol (IAA)
136 protocol for 15 sessions, as previously used in Pelloux and Baunez (2017). During each 12h-session
137 (from 8:00 am to 8:00 pm), animals were isolated in a homecage with access to food and two bottles.
138 One bottle contained water, and the other held a 20% ethanol solution. Both bottles were weighted before
139 and after each session to calculate consumption levels. The weight of each rat was also measured prior
140 to every IAA session. Following the 12h-period of IAA, rats were returned to their homecage with their
141 cagemate, and observed a 36h-period without ethanol access before the next session. The placement of
142 the bottles in the cage was alternated from session to session to avoid side preferences. There was no
143 DBS stimulation applied during this period.

144 **3- Experiment 2: Addiction-like ethanol consumption**

145 36 hours after the last session of IAA, rats were subjected to the same 10% ethanol self-administration
146 procedure than for Experiment 1, for 10x1h other sessions: 5 consecutive sessions alone ('Alone') and
147 5 consecutive sessions in presence of an alcohol-naive stranger peer ('With peer'). Operant chambers
148 were the same as those used in the first procedure. The order of the conditions 'Alone' vs. 'With peer'

149 were counterbalanced between rats. To avoid a possible long-term effect of former STN DBS (i.e.,
150 during recreational alcohol use) on addiction-like alcohol intake, we counterbalanced animals subjected
151 to STN DBS between experiment 1 and 2. Thus, half of the animals subjected to STN DBS OFF during
152 experiment 1 remained OFF for experiment 2, while the other half were subjected to DBS ON for
153 experiment 2, and vice versa for the other group. The DBS was turned ON during the 10 sessions for
154 the STN DBS group (n=15) and remained OFF for the control group (n=15).

155 **Deep Brain Stimulation**

156 Electrodes were connected to homemade electric stimulation cables, themselves connected to a digital
157 stimulator (DS8000, WPI) via a stimulus isolator (DLS100, WPI) and a rotating commutator (Plastic-
158 One). Stimulation parameters were adapted from previous studies (Degoulet et al. 2021; Pelloux et al.
159 2018). Individual stimulation intensity was determined using 130 Hz frequency and 80 μ s pulse width
160 stimulation. Intensity was progressively increased until the appearance of hyperkinetic movements or
161 abnormal behavior. Stimulation intensity (50 to 150 μ A) was set just below the abnormal behavior
162 threshold. In any case, the maximal intensity was set at 150 μ A in order to prevent heating and tissue
163 damage.

164 Before each behavioral session, animals were connected to the stimulation device, DBS was turned ON,
165 and stimulation intensity was progressively increased to reach the predetermined stimulation intensity.
166 Stimulation at 130 Hz was applied continuously during the 1-hour session.

167 **Histology**

168 At the end of the experiments, rats were deeply anesthetized with an injection of pentobarbital (Euthasol
169 Vet, Dechra, 300 mg/kg *i.p.*). Then, brains were extracted and rapidly frozen into liquid isopentane (-
170 40°C) and cut in 40 μ m thick frontal slices using a cryostat. A staining with cresyl violet allowed to
171 perform histological control. Animals with incorrect electrode placement were excluded from the ON
172 group. Representative correct electrodes implantation is illustrated in **Fig. 1**.

173 **Statistical Analyses**

174 Only the number of rewards obtained (i.e., rewarded active lever presses) are shown and expressed as
175 mean number \pm SEM. Using R software, and p-value threshold set at $\alpha=0.05$, data were analyzed by
176 performing two-tailed, one, two or three-way mixed ANOVA, followed by Bonferroni post-hoc tests
177 when applicable. Graphs were performed using GraphPad Prism 8 (version 8.0.2).

178 The consumption of ethanol or water in the IAA protocol was calculated as difference in bottle weight
179 before and after each session. This difference was then divided by the weight of the rat.

180 In Experiment 1 and 2, the outlier's test was conducted using the mean baseline of ethanol intake of all
181 rats in the "alone" condition.

182 Correlation between the amount of ethanol intake during the last 3 sessions of IAA and the number of
183 lever presses/rewards obtained during the first session of operant self-administration after the IAA were assessed
184 by a Pearson's correlation analysis. A Cohen's d was used to confirm that our sample size of low drinkers control
185 animals (n=5) was sufficient to detect reliable changes in social context (power >95% at a level of confidence of
186 $p<0.05$)

187 **Results**

188 In total, 28 rats were included in the analyses of Experiment 1. Four animals were excluded: electrodes'
189 mislocation (n=2, ON group), one rat (OFF group) was identified as an outlier (estimation based on the
190 mean baseline of ethanol intake of all rats), one rat (ON group) did not drink ethanol.

191 To avoid a possible long-term effect of former STN DBS during Experiment 1, we counterbalanced
192 animals subjected to STN DBS between experiment 1 and 2. In Experiment 2 (escalated alcohol
193 consumption), out of 30 rats, 24 were included in the analyses. 5 rats (n=2 ON and 3 OFF) did not
194 escalate their ethanol intake in the IAA protocol, and one rat (ON group) was identified as an outlier
195 (estimation based on the mean baseline of ethanol intake of all rats).

196 **Experiment 1:**

197 **Influence of a Peer's Presence and Contribution of STN on Recreational Alcohol Intake**

198 To study the influence of social context and the role of STN, a three-way ANOVA was used (social
199 context x STN DBS + session) to assess the number of rewards obtained (i.e. active lever presses outside
200 of the timeout period). No main effects were observed for social context ($F(1, 270)=0.015$; $p=0.901$),
201 STN DBS ($F(1, 270)=0.040$; $p=0.842$) and session ($F(4, 270)=0.649$; $p=0.628$). The interactions of STN
202 DBS and social context were also non-significant (STN DBS x social context ($F(1, 270)=0.001$;
203 $p=0.972$)).

204 Indeed, in the alone condition, **control rats** ($n=15$) had a stable intake of ethanol (*mean* 7.88 ± 0.29
205 *rewards*), that remained similar in presence of a peer (*mean* 8.36 ± 0.48 *rewards*). Thus, the presence of
206 an alcohol-naive stranger peer, had no effect on recreational ethanol consumption.

207 In **STN stimulated rats** ($n=13$), the alcohol intake in the alone condition (*mean* 7.60 ± 0.25 *rewards*)
208 was comparable to the control group. STN DBS seems not to influence recreational alcohol use in this
209 experiment. In presence of an alcohol-naive unfamiliar peer, STN DBS did not change significantly the
210 ethanol consumption compared to when STN-stimulated rats were in the alone condition (*mean* $8.09 \pm$
211 0.15 *rewards*), and was similar to the control group's intake (**Fig. 2a**).

212 **Influence of the Alcohol Preference**

213 Because substantial variations in alcohol consumption and preference were observed between rats, they
214 were split into two groups: high drinkers (HD) and low drinkers (LD) depending on their mean ethanol
215 consumption in the alone condition (Spoelder et al. 2015). The effects of the social context and STN
216 DBS were assessed with a three-way ANOVA (social context x STN DBS + sessions) independently
217 for each group according to their preference for ethanol.

218 In **HD rats** ($n=14$, **Fig. 2b**), there were no effect of the STN DBS ($F(1, 130)=0.061$; $p=0.805$), nor of
219 the sessions ($F(4, 130)=0.759$; $p=0.554$) nor of the interaction STN DBS x social context ($F(1,$
220 $130)=0.159$; $p=0.691$), although a trend for the social context effect ($F(1, 130)=3.518$; $p=0.063$).
221 Indeed, HD control rats ($n=8$) had similar ethanol consumption in the alone condition (*mean* $10.95 \pm$
222 0.36 *rewards*) and in a presence of an alcohol-naive stranger peer (*mean* 11.05 ± 0.72 *rewards*). STN
223 DBS did not affect alcohol consumption in HD animals ($n=6$) (*alone: mean* 10.37 ± 0.40 , *with a peer:*
224 *mean* 9.73 ± 0.53 *rewards*), that was equivalent to the control group in the same circumstances.

225 In **LD rats** ($n=14$, **Fig. 2c**), the ANOVA analysis indicated a trend for an STN DBS effect ($F(1,$
226 $130)=3.669$; $p=0.058$) and for the social context ($F(1, 130)=3.415$; $p=0.067$), but no significant effect of
227 the repeated sessions ($F(4, 130)=0.459$; $p=0.766$), nor interaction STN DBS x social context ($F(1,$
228 $130)=0.356$; $p=0.552$).

229 LD control rats ($n=7$) consumed ethanol at the same level in the alone condition (*mean* 4.37 ± 0.23
230 *rewards*) and in presence of a peer (*mean* 5.29 ± 0.39 *rewards*). STN DBS in LD rats ($n=7$) did not
231 change ethanol consumption when compared to that of control animals, regardless of whether they were
232 in the alone condition (*mean* 5.23 ± 0.27 *rewards*) or in a social context (*mean* 6.69 ± 0.19 *rewards*).

233 Overall, our results showed that despite tendencies for slight modulations, neither the presence of an
234 alcohol-naive stranger peer, nor the STN DBS alters significantly recreational alcohol consumption,
235 regardless of alcohol preference (i.e., low vs. high drinkers).

236 **Experiment 2: Influence of peer's presence and STN contribution following escalation of ethanol** 237 **use**

238 *Rats subjected to IAA procedure escalated their ethanol consumption over sessions*

239 Alcohol consumption during IAA was analyzed with a two-way ANOVA with STN DBS group as
240 between factors (control group vs. "to be stimulated group" although there is no stimulation during IAA)

241 and sessions as within factors. During the intermittent access to alcohol, rats showed a progressive
242 increase of their alcohol consumption (**Fig. 3a**) over the sessions ($F(4.34, 95.43)=18.083$; $p<0.0001$),
243 with no difference between the control group and the “to be stimulated” group ($F(1, 22)=1.197$; $p=0.286$)
244 and no interaction between these two factors ($F(4.34, 95.43)=1.749$; $p=0.140$). Post-hoc analyses
245 confirmed the progressive increase of alcohol intake over the sessions, reaching a level of consumption
246 during the last session (*mean* 3.01 ± 0.35 g/kg ethanol) that was significantly higher than that of the first
247 session (*mean* 1.66 ± 0.13 g/kg ethanol, Bonferroni’s test: $p=0.00015$).

248 Two-way ANOVA (STN DBS group x sessions) of water consumption during IAA (**Fig. 3b**) revealed
249 a main effect of session ($F(5.40, 118.80)=7.893$; $p<0.0001$) and no difference between the two future
250 groups ($F(1, 22)=0.170$; $p=0.5$) nor the interaction between session x STN DBS ($F(5.40, 118.80)=1.011$;
251 $p=0.418$). The session effect is due to inter-session variability but no difference between the first (*mean*
252 16.22 ± 0.92 g/kg water) and the last sessions (*mean* 22.39 ± 2.27 g/kg water, Bonferroni’s test, $p=1$)
253 was found.

254 As a result, both control and ‘to be stimulated’ groups exhibited comparable escalation and loss of
255 control only over alcohol intake.

256 *Influence of Peer’s Presence and STN Contribution on Escalated Alcohol Intake*

257 To assess the effect of the peer’s presence and STN DBS after a loss of control over alcohol use, a three-
258 way ANOVA was conducted (STN DBS x social context x sessions). This investigation disclosed
259 significant main effects of STN DBS ($F(1, 215)=39.752$; $p < 0.0001$) but no effect of the social context
260 ($F(1, 215)=0.232$; $p=0.631$), and sessions ($F(4, 215)=0.337$; $p=0.852$). No significant interaction was
261 found (STN DBS x social context ($F(1, 215)=1.230$; $p=0.269$; STN DBS x sessions ($F(4, 215)=1.090$;
262 $p=0.363$; social context x session ($F(4, 215)=0.073$; $p=0.990$) and STN DBS x social context x session
263 ($F(4, 215)=0.250$; $p=0.909$)).

264 In the **control group** ($n=12$, **Fig. 3c black**), after escalation of alcohol intake, the introduction of an
265 alcohol-naive stranger peer, did not modulate ethanol consumption; control rats displayed equivalent
266 ethanol intake whether in the alone condition (*mean* 9.30 ± 1.10 rewards) or in presence of the peer
267 (*mean* 10.58 ± 1.08 rewards, Bonferroni’s test, $p=0.341$).

268 When **STN DBS** ($n=12$, **Fig. 3c**) was applied, post-hoc analysis revealed that in the alone condition, rats
269 consumed a significantly less ethanol (*mean* 6.13 ± 0.55 rewards, Bonferroni’s test, $p=0.004$) than the
270 control group in the same alone condition. This decreased consumption persisted in the presence of a
271 peer, where STN-stimulated rats consumed significantly less (*mean* 5.63 ± 0.62 rewards, Bonferroni’s
272 test, $p=0.0004$) than the control groups in the same condition. However, within the stimulated group, no
273 significant difference was observed between the level of consumption in the two conditions ‘alone’ and
274 ‘with peer’ (Bonferroni’s test, $p=0.567$). Together, these results suggest that STN DBS reduces escalated
275 ethanol intake, irrespective of the social context.

276 To confirm that STN DBS reduced ethanol intake independently of the social context, we reordered the
277 sessions in the proper sequence independently of the social conditions (i.e., the order of passage in the
278 two social conditions were counterbalanced between animals, **Fig. 3d**). We then performed a two-way
279 ANOVA including the controls and STN DBS rats (STN DBS x reordered session) to assess the effects
280 of the STN DBS over time on alcohol intake. Our results revealed a significant effect of the STN DBS
281 ($F(1, 219)=43.299$; $p<0.0001$) and an effect of the reordered session ($F(9, 219)=5.924$; $p<0.0001$ but no
282 effect of the interaction STN DBS x reordered session ($F(9, 219)=0.559$; $p=0.83$)), confirming that the
283 STN DBS decreased alcohol intake whatever the social context.

284 285 *Correlation between the amount of ethanol intake during the last 3 sessions of IAA and the number* 286 *of lever presses/rewards obtained during the first session of operant self-administration* 287

288 In order to make sure that the slight non-significant difference between the “off control” and the “to be
289 stimulated/ ON” groups in alcohol intake during the last 3 sessions of the intermittent access protocol
290 would not affect the number of lever presses/rewards obtained during the first session, we tested a

291 possible correlation between these measures by performing a Pearson's correlation. For both groups,
292 there was no correlation between the number of lever presses and the 13th session of the intermittent
293 access (OFF: $r(10)=-0.0210$; $p=0.95$; ON: $r(10)=0.24$; $p=0.45$), nor the 14th sessions of IAA (OFF:
294 $r(10)=0.27$; $p=0.40$; ON : $r(10)=0.25$; $p=0.43$), nor the 15th sessions of IAA (OFF: $r(10)=0.4$; $p=0.20$;
295 ON: $r(10)=0.14$; $p=0.66$) nor the average of the last 3 IAA sessions (OFF: $r(10)=0.25$; $p=0.44$; ON:
296 $r(10)=0.22$; $p=0.49$). It thus suggests that the difference between the two groups during the operant
297 protocol is due to the STN DBS and not to a baseline difference

298 **Impact of Alcohol preference Following a Loss of Control over Alcohol Intake**

299 An additional possible contributor to inter-individual variation is the individual alcohol preference. This
300 alcohol preference is calculated on the level of alcohol consumption in the alone condition after the
301 escalation procedure. This alcohol preference allows splitting rats into two categories: High Drinkers
302 (HD) and Low Drinkers (LD).

303 **For HD rats** ($n=12$, **Fig. 3e**), a three-way ANOVA was run to explore the influence of social context,
304 STN DBS, and repeated sessions. While no impact of the social context ($F(1, 95)=0.724$, $p=0.397$) and
305 sessions ($F(4, 95)=0.698$; $p=0.595$) emerged, effect of STN DBS ($F(1, 95)=10.003$; $p=0.002$) was
306 significant. No interaction effects were found (STN DBS x social context: $F(1, 95)=0.792$; $p=0.376$),
307 STN DBS x session: $F(4, 95)=0.434$; $p=0.784$, social context x session: $F(4, 95)$; $p=0.248$; $p=0.910$ and
308 STN DBS x social context x session: $F(4, 95)=0.192$; $p=0.942$).

309 HD control animals ($n=7$) had a similar ethanol consumption in the alone condition and in the presence
310 of a peer (*mean* 12.83 ± 1.59 and 12.69 ± 1.78 rewards respectively, Bonferroni's test, $p=0.948$),
311 revealing that the peer presence did not influence escalated alcohol use in HD rats. The STN DBS
312 reduced the consumption in HD STN-stimulated animals ($n=5$) compared to the HD control group in
313 both social conditions: alone (Bonferroni's test, $p=0.037$, *mean* 9.48 ± 0.92 rewards for stimulated
314 animals) and in presence of a peer (Bonferroni's test, $p=0.021$, *mean* 6.72 ± 1.15 rewards for STN-
315 stimulated rats). However, no significant difference was found in alcohol intake between the two social
316 conditions within the HD STN-stimulated group (Bonferroni's test, $p=0.117$), revealing the beneficial
317 effect of the STN DBS on escalated alcohol use in HD rats, independently of the social context.

318 **For LD rats** ($n=12$, **Fig. 3f**), the three-way ANOVA revealed a significant effect of the social context
319 ($F(1, 95)=10.631$; $p=0.002$) and STN DBS ($F(1, 95)=10.108$; $p=0.002$), but no influence of the session
320 ($F(4, 95)=0.443$; $p=0.778$). No significant interaction was found (STN DBS x social context: $F(1,$
321 $95)=2.919$; $p=0.091$, STN DBS x session: $F(4, 95)=0.395$; $p=0.812$, social context x session: $F(4,$
322 $95)=0.317$; $p=0.867$ and STN DBS x social context x session: $F(4, 95)=0.315$; $p=0.868$).

323 LD control animals ($n=5$) consumed more alcohol in presence of a peer (*mean* 7.64 ± 0.34 rewards)
324 than when they were in the alone condition (*mean* 4.40 ± 0.42 rewards, Bonferroni's test, $p=0.0006$,
325 with a power effect above 80% (Cohen's effect size value $d=0.99$)). Although STN DBS ($n=7$) did not
326 affect ethanol intake when rats were in the alone condition (Bonferroni's test compared to LD controls,
327 $p=0.366$, *mean* 3.74 ± 0.45 rewards), it prevented the overconsumption induced by the presence of a
328 peer (*mean* 4.86 ± 0.47 rewards) (lower consumption in LD STN DBS compared with LD controls,
329 Bonferroni's test, $p=0.005$). Consequently, STN DBS effectively suppressed the excessive consumption
330 triggered by the presence of a peer in LD rats.

331 **Discussion**

332 The present study showed that neither the presence of an alcohol-naive stranger peer, nor STN
333 DBS, significantly modulate the recreational use of alcohol. In contrast, after a loss of control over
334 ethanol intake, STN DBS induced a decrease of the alcohol use over time. After escalation of ethanol
335 intake, the presence of an alcohol-naive stranger peer modulated the consumption depending on the
336 initial alcohol preference. HD rats were insensitive to the presence of the peer, and maintained a stable
337 high level of ethanol intake, while LD rats exhibited an overconsumption in presence of the peer. STN
338 DBS decreased ethanol intake in HD rats in both conditions (alone and with peer), while it only reduced
339 the overconsumption induced by the peer's presence in LD rats.

340
341

Presence of a peer during alcohol consumption: Importance of the peer's identity

342 Studies investigating the effect of a peer's presence on alcohol self-administration have yielded
343 mixed results. Several showed that the presence of a social partner drinking alcohol promotes alcohol
344 drinking in humans (Dallas et al. 2014; Larsen et al. 2009). Our recent study in rats further revealed that
345 the presence of a peer, whether familiar or stranger, but who had a regular consumption of alcohol,
346 induced an increase of recreational alcohol consumption (Sicre et al., unpublished data). In the present
347 study, the presence of an alcohol-naive stranger peer during recreational use of alcohol had no significant
348 influence. These results highlight that the peer's history with alcohol plays an important role. If peers
349 consumed alcohol orally recently before being used as the "peer", it is possible that olfactory cues of
350 alcohol consumption could be transmitted to the subject rat through mouth-to-mouth contacts. These
351 ethanol cues might induce a social transmission of ethanol preference, as observed for food (Strupp and
352 Levitsky 1984), and this would result in an overconsumption of alcohol. Such olfactory cues could not
353 be transmitted by a completely alcohol-naive peer, preventing alcohol preference facilitation and
354 therefore no changes in consumption for the subject rat. On the other hand, the absence of increase of
355 alcohol intake during the recreational experiment could be due to a ceiling effect that does not leave
356 room for an increase. However, we have observed an increased effect by the presence of a peer in LD
357 rats after escalation of alcohol consumption and the level reached then by these LD was equivalent to
358 that recorded in the recreational experiment. So, although there was room for an increase in consumption
359 in the recreational experiment, the peer presence had no significant effect. The ceiling effect is thus
360 probably to be ruled out.

361 Once the control over ethanol intake was lost, the influence of an alcohol-naive peer varied
362 depending on alcohol preference. For LD rats, the presence of a peer induced an overconsumption of
363 ethanol while for HD rats, there was no modulation. An appealing hypothesis would be that after an
364 escalation of alcohol intake, LD rats had an enhanced sensitivity to the presence of a peer. Indeed, social
365 interactions are rewarding (Krach 2010), and after undergoing intermittent ethanol exposure during late
366 adolescence, male rats having low doses of ethanol in adulthood displayed an increased social behavior
367 (Varlinskaya et al. 2014). The potentiation between the rewarding effect of low doses of alcohol and
368 social interaction could be magnified, resulting in the overconsumption even if the peer is an alcohol-
369 naive stranger peer. In the case of HD rats after a loss of control, their lack of sensitivity to the peer's
370 presence could be attributed to a preference for alcohol consumption over social interaction. It has been
371 shown that when rats had a high dose of ethanol, they show less social investigation, contact behavior
372 and social avoidance (Varlinskaya and Spear 2002). Moreover, after a protocol of intermittent access to
373 alcohol rats exhibit preference for alcohol over social rewards (Augier et al. 2022; Marchant et al. 2022).
374 In our experiment, however, rats were not offered the choice, the social presence was imposed, and the
375 rats could consume alcohol while interacting with a peer as much as they liked.

376 Nonetheless, from our results, the presence of a stranger peer that never consumed ethanol had
377 not much detrimental effect, which is in contrast to the consequences of presence of a peer that has
378 previously consumed alcohol. These results confirm the significant influence of the immediate social
379 context on drug consumption. This effect is influenced by factors such as the peer's identity (e.g.,
380 familiarity, drug history) and the specific type of drug being used (e.g., psychostimulant vs. depressive
381 drug).

STN as a neurobiological substrate ...

...Of escalated but not recreational alcohol use...

382 During recreational alcohol use (Experiment 1), the present study showed no significant effect
383 of STN DBS on ethanol intake. These results are consistent with a previous work showing that lesion of
384 STN did not affect rats' alcohol intake in a forced consumption test, and in a two-bottle choice protocol
385 (Lardeux and Baunez 2008). Moreover, similar to alcohol, STN manipulation was ineffective at
386 decreasing recreational cocaine use (Baunez et al. 2005; Rouaud et al. 2010; Giorla et al. 2022; Vielle
387 et al. preprint). However, while no effort was required in the present study that used continuous
388 reinforcement, when the demand was increased, as in a progressive ratio task, STN manipulation was
389 efficient at reducing cocaine motivation (Baunez et al. 2005; Rouaud et al. 2010). For alcohol, this effect
390
391

392 has been shown to depend on the loss of control over ethanol intake. If rats did not undergo an IAA
393 protocol, the effect of STN lesion depended on the rat's alcohol preference (Lardeux and Baunez 2008).
394 After an IAA protocol, STN lesion decreased the level of ethanol motivation (Pelloux and Baunez 2017).
395 Thus, like for cocaine, STN manipulation could blunt alcohol addiction-like criteria, without altering
396 recreational drug use.

397 Indeed, applying DBS at high frequency to the STN after escalation of ethanol intake reduced
398 alcohol consumption. There is a possibility that the disparities observed between the control group and
399 the STN stimulated group could be explained by the slight difference, although non-significant, during
400 the last three days of IAA protocol. However, as indicated in the result section, further analyses have
401 shown a lack of correlation between the level of intake during the last 3 sessions of the intermittent
402 access and the number of rewards obtained during the first session, suggesting that the decreased lever
403 pressing for alcohol is due to STN DBS and not to a lower baseline consumption. These effects of STN
404 DBS are in line with previous works showing that lesion of STN prevented escalation of ethanol intake
405 and diminished the rebound intake after abstinence (Pelloux and Baunez 2017). This has been shown
406 for cocaine, heroin and alcohol (Pelloux and Baunez 2017; Pelloux et al. 2018; Vielle et al. preprint;
407 Wade et al. 2016). Interestingly, in contrast to our former results with cocaine that failed to show an
408 effect of STN DBS on the maintained escalated intake (Pelloux et al. 2018), it was here reduced for
409 alcohol. This disparity in the impact of STN DBS on escalated alcohol and cocaine consumption could
410 potentially be attributed to the type of drug available. Cocaine is a psychostimulant, while alcohol has
411 depressant effects. The road of administration could also explain the differences observed between
412 alcohol and cocaine. Alcohol is consumed orally, leading to different pharmacokinetics and longer
413 absorption time compared to intravenously administration that enters directly in the bloodstream
414 (Robinson et al. 2002). This delay deferred the activation of the reward system and pleasurable effects
415 (Robinson et al. 2002). Introducing a delay before obtaining a reward has been shown to decrease
416 motivation (Jarmolowicz and Hudnall 2014). Consequently, it might thus be comparatively easier to
417 reduce motivation for alcohol compared to cocaine with STN DBS. However, one might also consider
418 that having a direct sensory input with alcohol in the mouth, but also its smell, could lead to anticipation
419 of ethanol effects, while intravenous injection might be less efficient at anticipating the physiological
420 and psychoactive effects of cocaine. The differences are therefore difficult to simply explain with the
421 mean of administration. It is interesting to note however that, although the DBS or lesion procedure did
422 not lead to changes in sustained escalated intake of cocaine, a precise targeting of the STN using
423 optogenetics has been shown to be efficient (Vielle et al. preprint).

424 Finally, differences in the experimental protocol would account for such differences. Indeed,
425 cocaine escalation, and maintenance in Pelloux et al. (2018), were performed in the same context
426 (operant cages). In the present experiment, escalation of alcohol intake occurred in different cages than
427 the maintenance, and the access (intermittent *vs.* daily), and concentrations of ethanol were also different
428 between escalation procedure and the operant measure of consumption (20 *vs.* 10%). Thus, in this
429 experiment, such differences between escalation and maintenance could have facilitated STN DBS
430 effect on the maintenance of ethanol use in comparison with the cocaine experiment (Pelloux et al.
431 2018).

432 The level of consumption seems to be an important factor to ensure STN DBS efficiency since
433 STN DBS decreased the consumption of alcohol only in HD rats. The lack of impact of STN DBS in
434 LD rats might be attributed to a floor effect, implying that the LD rats did not consume enough ethanol
435 to exhibit further a reduction triggered by STN DBS. Another hypothesis would be that the
436 neurobiological effect of alcohol consumption does not induce the same changes in LD and HD rats.
437 Indeed, when consuming alcohol, HD rats exhibit an increased power in gamma range and coherence in
438 the medial prefrontal cortex (mPFC) compared to LD rats (Henricks et al. 2019). The mPFC is directly
439 connected to the STN via the hyperdirect pathway (Maurice et al. 1998; Nambu et al. 2002).
440 Additionally, abnormal increase in low frequency oscillations has been observed within the STN during
441 cocaine escalation, and STN DBS was shown to reduce STN oscillatory activity (Pelloux et al. 2018).
442 If such abnormal activity raised in the STN during alcohol escalation of HD rats, as that seen with
443 cocaine, STN high frequency DBS could suppress it and also disturb the pathological oscillatory
444 coherence between STN-mPFC as observed in PD (de Hemptinne et al. 2015; Delaville et al. 2015;
445 Moran et al. 2012). This would lead to a decrease of alcohol intake only for HD rats after the loss of
446 control.

447 Nonetheless, STN DBS appears to offer a potential therapeutic benefit for alcohol-related
448 disorders, as suggested previously for other drugs.

449 ...*And of social influence during alcohol drinking*

450 The influence of the social context was observed here in LD rats having escalated their alcohol
451 consumption. Interestingly, in these animals, STN DBS effectively abolished the overconsumption
452 induced by the presence of the peer. These results align with precedent findings, in which STN DBS has
453 suppressed recreational alcohol overconsumption triggered by different types of peer (Sicre et al.,
454 unpublished data). STN optogenetic inhibition was also found to suppress the protective effect of a peer
455 on cocaine recreational use (Vielle et al. preprint). The STN could thus be part of the neurobiological
456 substrate of the influence of the social context on behavior. Indeed, studies focusing on the effects of
457 ultrasonic vocalizations (USV) playback on cocaine intake showed that STN lesion abolished the
458 beneficial influence of positive USV emitted by a stranger rat (Montanari et al. 2018; Vielle et al. 2021).
459 Additionally, STN lesion suppressed the rewarding effect of positive USVs when rats were pressing a
460 lever to listen to USVs (Vielle et al. 2021). It is important to note however that the combination of social
461 context and STN manipulation does not systematically lead to neutralization or additivity of opposite
462 effects, nor potentiation of similar effects. For example, STN lesions could decrease further the cocaine
463 intake in presence of a peer (Giorla et al 2022), while STN neuronal optogenetic inhibition did not
464 potentiate the decreased cocaine intake after escalation (Vielle et al preprint). All these findings suggest
465 that presence of a peer effects on drug use probably involves STN as a neurobiological substrate. In
466 human studies, STN DBS has been shown to induce deficits in facial and voice emotional decoding
467 (Biseul et al. 2005; Brück, Kreifelts, and Wildgruber 2011; Kalampokini et al. 2020; Péron et al. 2010;
468 2013). A recent study has also shown that STN activity assessed with fMRI, was modulated by the
469 presence of an observer during the performance of a stop-signal reaction time task by cocaine-addicted
470 individuals (Terenzi et al. preprint). All these results converge to suggest the involvement of STN in
471 mediating the impact of social interactions on drug-related behavior. Social interaction has been shown
472 to trigger the release of oxytocin within the brain's reward circuit (Dölen et al. 2013; Hung et al. 2017).
473 Additionally, there are evidences suggesting that the STN possesses oxytocin receptors that can regulate
474 locally dopamine transmission (Baracz and Cornish 2013, 2016). Furthermore the injection of oxytocin
475 into the STN has been demonstrated to reduce methamphetamine-induced reinstatement (Baracz et al.
476 2015). Through oxytocin release induced by social interaction, the STN activity could thus be modulated
477 and potentially affect drug consumption behavior.

478 Additional researches are required to elucidate further the role of the STN in social behaviors
479 and its impact on drug-related effects.

480

481 **References**

- 482 Augier, Gaëlle, Veronika Schwabl, Asmae Lguensat, Mihai Atudorei, Osamudiamen Consoler Iyere, Sandra
483 Eriksson Solander, and Eric Augier. 2022. "Wistar Rats Choose Alcohol over Social Interaction in a
484 Discrete-choice Model." *Neuropsychopharmacology* 48 (7) (December 31): 1098–1107.
485 doi:10.1038/s41386-022-01526-8.
- 486 Baracz, Sarah J., and Jennifer L. Cornish. 2013. "Oxytocin Modulates Dopamine-mediated Reward in the Rat
487 Subthalamic Nucleus." *Hormones and Behavior* 63 (2) (February): 370–375.
488 doi:10.1016/j.yhbeh.2012.12.003.
- 489 Baracz, Sarah J., and Jennifer L. Cornish 2016. "The Neurocircuitry Involved in Oxytocin Modulation of
490 Methamphetamine Addiction." *Frontiers in Neuroendocrinology* 43 (October): 1–18.
491 doi:10.1016/j.yfrne.2016.08.001.
- 492 Baracz, Sarah Jane, Nicholas Adams Everett, and Jennifer Louise Cornish. 2015. "The Involvement of Oxytocin
493 in the Subthalamic Nucleus on Relapse to Methamphetamine-Seeking Behaviour." Edited by Rita
494 Fuchs. *PLOS ONE* 10 (8) (August 18): e0136132. doi:10.1371/journal.pone.0136132.
- 495 Bardo, M. T., J. L. Neisewander, and T. H. Kelly. 2013. "Individual Differences and Social Influences on the
496 Neurobehavioral Pharmacology of Abused Drugs." Edited by Michael A. Nader. *Pharmacological
497 Reviews* 65 (1) (January 23): 255–290. doi:10.1124/pr.111.005124.

498 Baunez, Christelle, Carine Dias, Martine Cador, and Marianne Amalric. 2005. "The Subthalamic Nucleus Exerts
499 Opposite Control on Cocaine and 'Natural' Rewards." *Nature Neuroscience* 8 (4) (March 27): 484–489.
500 doi:10.1038/nn1429.

501 Baunez, Christelle, and Sylvie Lardeux. 2011. "Frontal Cortex-Like Functions of the Subthalamic Nucleus."
502 *Frontiers in Systems Neuroscience* 5. doi:10.3389/fnsys.2011.00083.
503 <http://journal.frontiersin.org/article/10.3389/fnsys.2011.00083/abstract>.

504 Biseul, Isabelle, Paul Sauleau, Claire Haegelen, Pascale Trebon, Dominique Drapier, Sylvie Raoul, Sophie
505 Drapier, et al. 2005. "Fear Recognition Is Impaired by Subthalamic Nucleus Stimulation in Parkinson's
506 Disease." *Neuropsychologia* 43 (7) (January): 1054–1059.
507 doi:10.1016/j.neuropsychologia.2004.10.006.

508 Brück, Carolin, Benjamin Kreifelts, and Dirk Wildgruber. 2011. "Emotional Voices in Context: A
509 Neurobiological Model of Multimodal Affective Information Processing." *Physics of Life Reviews* 8 (4)
510 (December): 383–403. doi:10.1016/j.plev.2011.10.002.

511 Dallas, Rebecca, Matt Field, Andrew Jones, Paul Christiansen, Abi Rose, and Eric Robinson. 2014. "Influenced
512 but Unaware: Social Influence on Alcohol Drinking Among Social Acquaintances." *Alcoholism:
513 Clinical and Experimental Research* 38 (5) (May): 1448–1453. doi:10.1111/acer.12375.

514 De Hemptinne, Coralie, Nicole C Swann, Jill L Ostrem, Elena S Ryapolova-Webb, Marta San Luciano, Nicholas
515 B Galifianakis, and Philip A Starr. 2015. "Therapeutic Deep Brain Stimulation Reduces Cortical Phase-
516 amplitude Coupling in Parkinson's Disease." *Nature Neuroscience* 18 (5) (April 13): 779–786.
517 doi:10.1038/nn.3997.

518 Degoulet, Mickael, Alix Tiran-Cappello, Etienne Combrisson, Christelle Baunez, and Yann Pelloux. 2021.
519 "Subthalamic Low-frequency Oscillations Predict Vulnerability to Cocaine Addiction." *Proceedings of
520 the National Academy of Sciences* 118 (14) (March 30): e2024121118. doi:10.1073/pnas.2024121118.

521 Delaville, Claire, Alex J. McCoy, Colin M. Gerber, Ana V. Cruz, and Judith R. Walters. 2015. "Subthalamic
522 Nucleus Activity in the Awake Hemiparkinsonian Rat: Relationships with Motor and Cognitive
523 Networks." *The Journal of Neuroscience* 35 (17) (April 29): 6918–6930.
524 doi:10.1523/JNEUROSCI.0587-15.2015.

525 Dölen, Gül, Ayehe Darvishzadeh, Kee Wui Huang, and Robert C. Malenka. 2013. "Social Reward Requires
526 Coordinated Activity of Nucleus Accumbens Oxytocin and Serotonin." *Nature* 501 (7466) (September
527 11): 179–184. doi:10.1038/nature12518.

528 Giorla, Elodie, Sandra Nordmann, Cassandre Vielle, Yann Pelloux, Perrine Roux, Camelia Protopopescu,
529 Christine Manrique, et al. 2022. "Peer Presence and Familiarity as Key Factors to Reduce Cocaine
530 Intake in Both Rats and Humans: An Effect Mediated by the Subthalamic Nucleus." *Psychopharmacology* 239 (4) (January 11): 1097–1113. doi:10.1007/s00213-021-06033-0.

531 Henricks, Angela M., Lucas L. Dwiell, Nicholas H. Deveau, Amanda A. Simon, Metztl J. Ruiz-Jaquez, Alan I.
532 Green, and Wilder T. Doucette. 2019. "Corticostriatal Oscillations Predict High Vs. Low Drinkers in a
533 Rat Model of Limited Access Alcohol Consumption." *Frontiers in Systems Neuroscience* 13 (August
534 13). doi:10.3389/fnsys.2019.00035. <https://www.frontiersin.org/article/10.3389/fnsys.2019.00035/full>.

535 Hung, Lin W., Sophie Neuner, Jai S. Polepalli, Kevin T. Beier, Matthew Wright, Jessica J. Walsh, Eastman M.
536 Lewis, et al. 2017. "Gating of Social Reward by Oxytocin in the Ventral Tegmental Area." *Science* 357
537 (6358) (September 29): 1406–1411. doi:10.1126/science.aan4994.

538 Jarmolowicz, David P., and Jennifer L. Hudnall. 2014. "Concurrent Progressive-ratio Schedules: Built-in
539 Controls in the Study of Delayed Reward Efficacy." *Journal of Neuroscience Methods* 233 (August):
540 172–176. doi:10.1016/j.jneumeth.2014.06.024.

541 Kalampokini, S., E. Lyros, P. Lochner, K. Fassbender, and M. M. Unger. 2020. "Effects of Subthalamic Nucleus
542 Deep Brain Stimulation on Facial Emotion Recognition in Parkinson's Disease: A Critical Literature
543 Review." *Behavioural Neurology* 2020 (July 17): 1–18. doi:10.1155/2020/4329297.

544 Krach, Sören. 2010. "The Rewarding Nature of Social Interactions." *Frontiers in Behavioral Neuroscience*.
545 doi:10.3389/fnbeh.2010.00022. <http://journal.frontiersin.org/article/10.3389/fnbeh.2010.00022/abstract>.

546 Lardeux, Sylvie, and Christelle Baunez. 2008. "Alcohol Preference Influences the Subthalamic Nucleus Control
547 on Motivation for Alcohol in Rats." *Neuropsychopharmacology* 33 (3) (February 1): 634–642.
548 doi:10.1038/sj.npp.1301432.

549 Larsen, H., R. C. M. E. Engels, I. Granic, and G. Overbeek. 2009. "An Experimental Study on Imitation of
550 Alcohol Consumption in Same-Sex Dyads." *Alcohol and Alcoholism* 44 (3) (February 23): 250–255.
551 doi:10.1093/alcalc/agg002.

552 Marchant, Nathan J., Allison J. McDonald, Rie Matsuzaki, Yvar van Mourik, Dustin Schettters, and Taco J. De
553 Vries. 2022. "Rats Choose Alcohol over Social Reward in an Operant Choice Procedure." *Neuropsychopharmacology* 48 (4) (September 15): 585–593. doi:10.1038/s41386-022-01447-6.

554 Maurice, Nicolas, Jean-Michel Deniau, Jacques Glowinski, and Anne-Marie Thierry. 1998. "Relationships
555 Between the Prefrontal Cortex and the Basal Ganglia in the Rat: Physiology of the Corticosubthalamic
556
557

558 Circuits.” *The Journal of Neuroscience* 18 (22) (November 15): 9539–9546.
559 doi:10.1523/JNEUROSCI.18-22-09539.1998.

560 Montanari, Christian, Elodie Giorla, Yann Pelloux, and Christelle Baunez. 2018. “Subthalamic Nucleus
561 Mediates the Modulation on Cocaine Self-administration Induced by Ultrasonic Vocalization Playback
562 in Rats.” *Addiction Biology* 25 (1) (December 28). doi:10.1111/adb.12710.
563 <https://onlinelibrary.wiley.com/doi/abs/10.1111/adb.12710>.

564 Moran, A., E. Stein, H. Tischler, and I. Bar-Gad. 2012. “Decoupling Neuronal Oscillations During Subthalamic
565 Nucleus Stimulation in the Parkinsonian Primate.” *Neurobiology of Disease* 45 (1) (January): 583–590.
566 doi:10.1016/j.nbd.2011.09.016.

567 Nambu, Atsushi, Hironobu Tokuno, and Masahiko Takada. 2002. “Functional Significance of the Cortico–
568 subthalamo–pallidal ‘hyperdirect’ Pathway.” *Neuroscience Research* 43 (2) (June): 111–117.
569 doi:10.1016/S0168-0102(02)00027-5.

570 Paxinos, George, and Charles Watson. 2014. “The Rat Brain in Stereotaxic Coordinates.” *Elsevier*.

571 Pelloux, Yann, and Christelle Baunez. 2017. “Targeting the Subthalamic Nucleus in a Preclinical Model of
572 Alcohol Use Disorder.” *Psychopharmacology* 234 (14) (April 11): 2127–2137. doi:10.1007/s00213-
573 017-4618-5.

574 Pelloux, Yann, Mickaël Degoulet, Alix Tiran-Cappello, Candie Cohen, Sylvie Lardeux, Olivier George, George
575 F. Koob, Serge H. Ahmed, and Christelle Baunez. 2018. “Subthalamic Nucleus High Frequency
576 Stimulation Prevents and Reverses Escalated Cocaine Use.” *Molecular Psychiatry* 23 (12) (June 7):
577 2266–2276. doi:10.1038/s41380-018-0080-y.

578 Pelloux, Yann, Elodie Giorla, Christian Montanari, and Christelle Baunez. 2019. “Social Modulation of Drug
579 Use and Drug Addiction.” *Neuropharmacology* 159 (November): 107545.
580 doi:10.1016/j.neuropharm.2019.02.027.

581 Péron, Julie, Sascha Frühholz, Marc Vérin, and Didier Grandjean. 2013. “Subthalamic Nucleus: A Key Structure
582 for Emotional Component Synchronization in Humans.” *Neuroscience & Biobehavioral Reviews* 37 (3)
583 (March): 358–373. doi:10.1016/j.neubiorev.2013.01.001.

584 Péron, Julie, Didier Grandjean, Florence Le Jeune, Paul Sauleau, Claire Haegelen, Dominique Drapier, Tiphaine
585 Rouaud, Sophie Drapier, and Marc Vérin. 2010. “Recognition of Emotional Prosody Is Altered after
586 Subthalamic Nucleus Deep Brain Stimulation in Parkinson’s Disease.” *Neuropsychologia* 48 (4)
587 (March): 1053–1062. doi:10.1016/j.neuropsychologia.2009.12.003.

588 Robinson, Donita L., Lane J. Brunner, and Rueben A. Gonzales. 2002. “Effect of Gender and Estrous Cycle on
589 the Pharmacokinetics of Ethanol in the Rat Brain.” *Alcoholism: Clinical and Experimental Research* 26
590 (2) (February): 165–172. doi:10.1111/j.1530-0277.2002.tb02521.x.

591 Rouaud, T., S. Lardeux, N. Panayotis, D. Paleressompouille, M. Cador, and C. Baunez. 2010. “Reducing the
592 Desire for Cocaine with Subthalamic Nucleus Deep Brain Stimulation.” *Proceedings of the National
593 Academy of Sciences* 107 (3) (January 19): 1196–1200. doi:10.1073/pnas.0908189107.

594 Spoelder, Marcia, Peter Hesselting, Annemarie M. Baars, José G. Lozeman-van ‘t Klooster, Marthe D. Rotte,
595 Louk J. M. J. Vanderschuren, and Heidi M. B. Lesscher. 2015. “Individual Variation in Alcohol Intake
596 Predicts Reinforcement, Motivation, and Compulsive Alcohol Use in Rats.” *Alcoholism: Clinical and
597 Experimental Research* 39 (12) (December): 2427–2437. doi:10.1111/acer.12891.

598 Strickland, Justin C., and Mark A. Smith. 2014. “The Effects of Social Contact on Drug Use: Behavioral
599 Mechanisms Controlling Drug Intake.” *Experimental and Clinical Psychopharmacology* 22 (1): 23–34.
600 doi:10.1037/a0034669.

601 Strickland, Justin C., and Mark A. Smith. 2015. “Animal Models of Social Contact and Drug Self-
602 administration.” *Pharmacology Biochemistry and Behavior* 136 (September): 47–54.
603 doi:10.1016/j.pbb.2015.06.013.

604 Strupp, Barbara J., and David A. Levitsky. 1984. “Social Transmission of Food Preferences in Adult Hooded
605 Rats (*Rattus Norvegicus*).” *Journal of Comparative Psychology* 98 (3) (September): 257–266.
606 doi:10.1037/0735-7036.98.3.257.

607 Terenzi, Damiano, Simon Nicolas, Michael Joe Munyua Gachomba, Jeanne Laure de Peretti, Bruno Nazarian,
608 Julien Sein, Jean-Luc Anton, Didier Grandjean, Christelle Baunez, and Thierry Cheminade. 2023.
609 “Social Context and Drug Cues Modulate Inhibitory Control in Cocaine Addiction: Involvement of the
610 STN Evidenced through Functional Magnetic Resonance Imaging.” *Preprint* (September 7).
611 doi:10.1101/2023.09.06.556336. <https://www.biorxiv.org/content/10.1101/2023.09.06.556336v2>.

612 Varlinskaya, Elena I., and Linda P. Spear. 2002. “Acute Effects of Ethanol on Social Behavior of Adolescent and
613 Adult Rats: Role of Familiarity of the Test Situation.” *Alcoholism: Clinical and Experimental Research*
614 26 (10) (October): 1502–1511. doi:10.1111/j.1530-0277.2002.tb02449.x.

615 Varlinskaya, Elena I., Eric Truxell, and Linda P. Spear. 2014. “Chronic Intermittent Ethanol Exposure During
616 Adolescence: Effects on Social Behavior and Ethanol Sensitivity in Adulthood.” *Alcohol* 48 (5)
617 (August): 433–444. doi:10.1016/j.alcohol.2014.01.012.

- 618 Venniro, Marco, Michelle Zhang, Daniele Caprioli, Jennifer K. Hoots, Sam A. Golden, Conor Heins, Marisela
619 Morales, David H. Epstein, and Yavin Shaham. 2018. “Volitional Social Interaction Prevents Drug
620 Addiction in Rat Models.” *Nature Neuroscience* 21 (11) (October 15): 1520–1529. doi:10.1038/s41593-
621 018-0246-6.
- 622 Vielle, Cassandre, Christian Montanari, Yann Pelloux, and Christelle Baunez. 2021. “Evidence for a Vocal
623 Signature in the Rat and Its Reinforcing Effects: a Key Role for the Subthalamic Nucleus.” *Proceedings
624 of the Royal Society B: Biological Sciences* 288 (1965) (December 15). doi:10.1098/rspb.2021.2260.
625 <https://royalsocietypublishing.org/doi/10.1098/rspb.2021.2260>.
- 626 Vielle, Cassandre, Alix Tiran-Cappello, Lucie Vignal, Nicolas Maurice, Mickael Degoulet, Cécile Brocard,
627 Florence Pelletier, Yann Pelloux, and Christelle Baunez. 2023. “Peer’s Presence Reverses Escalated
628 Cocaine Intake in Rats.” *bioRxiv* (August 2).
629 <https://www.biorxiv.org/content/10.1101/2023.02.07.527550v1.article-info>.
- 630 Wade, Carrie L, Marsida Kallupi, Daniel O Hernandez, Emmanuel Breysse, Giordano de Guglielmo, Elena
631 Crawford, George F Koob, Paul Schweitzer, Christelle Baunez, and Olivier George. 2016. “High-
632 Frequency Stimulation of the Subthalamic Nucleus Blocks Compulsive-Like Re-Escalation of Heroin
633 Taking in Rats.” *Neuropsychopharmacology* 42 (9) (December 5): 1850–1859.
634 doi:10.1038/npp.2016.270.

635
636
637
638 **Acknowledgments:** The authors thank Yann Pelloux for the custom-built interface and associated
639 software, Joel Baurberg for technical assistance with all electronic devices, and the technical
640 staff of the animal facility for ensuring the well-being of our animals. This work was funded by
641 the Institut de Recherche en Santé Public and Alliance Aviesan in the call for proposal on
642 substance use disorders and by the Institut de lutte contre le Cancer.

643

644 **Funding:**

645 Centre National de la Recherche Scientifique (CB, MD, NM)
646 Aix-Marseille Université (CB, CV, LV, MD, NM)
647 French Ministry of Higher Education, Research and Innovation (LV)
648 Institut de Recherche en Santé Public (IRES-19-ADDICTIONS-02) (CB)
649 Institut National de lutte contre le Cancer (INCA_16032) (LV)


650 **Author contributions:**

651 Conceptualization: CB, CV, LV
652 Research performance: CV, LV, MW, NM
653 Data analyses: CV, LV
654 Funding acquisition: CB, LV
655 Writing – original draft: CB, CV, LV
656 Writing – review & editing: CB, CV, LV, MD, MW, NM

657 **Competing interests:** Authors declare that they have no competing interests.

658 **Data and materials availability:** All data are available in the main text, or on request to the authors.

1



659

660 **Fig. 1 Timeline of the behavioral experiment**

661 Top: Behavioral experiment timeline


662 Down: (left) schematic representations of the electrode implanted in the STN,

663 (right) Picture from a representative electrode implanted in the STN (delineated by the red dashed line)

664


2

a




b

High drinkers



c

Low drinkers



665

666

667 **Fig. 2 Neither peer's presence nor STN DBS modulate recreational ethanol intake.**668 Results are presented as mean \pm SEM number of rewards obtained (i.e. number of active lever presses)

669 per 1h-session.

670 (a) Number of rewards obtained for non-STN-stimulated (black dots – OFF - control) and stimulated

671 (red dots – ON) groups during five sessions in the alone condition (7.88 ± 0.29 vs. 7.60 ± 0.25 672 *rewards* respectively) and five sessions with an alcohol-naive stranger peer (8.36 ± 0.48 vs. $8.09 \pm$ 673 0.15 *rewards* respectively)

674 (b) Number of rewards obtained in control (black dots – OFF - control) and STN-stimulated (red dots

675 – ON) High drinkers in the alone condition (10.95 ± 0.40 vs. 10.37 ± 0.40 *rewards* respectively)676 and in a presence of an alcohol-naive stranger peer (11.05 ± 0.72 vs. 9.73 ± 0.53 *rewards*,

677 respectively)

678 (c) Number of rewards obtained in control (black dots – OFF - control) and stimulated (red dots - ON)

679 Low drinkers in the alone condition (4.37 ± 0.23 vs. 5.23 ± 0.27 *rewards*, respectively)680 and in a presence of an alcohol-naive stranger peer (5.29 ± 0.39 vs. 6.69 ± 0.19 *rewards*, respectively)

681

682

683


684

685

686

687

3



689 **Fig. 3 STN DBS reduces the escalated intake of alcohol and suppresses the increased consumption**
690 **induced by the peer's presence.**

691 Results are presented as mean \pm SEM.

- 692 (a) Ethanol intake in control (black dots - OFF) and "to be stimulated" (red dots - ON) groups across
693 the 15 sessions of intermittent access to 20% ethanol. Both groups increased their consumption
694 from session 1 (1.66 ± 0.13 g/kg ethanol) to session 15 (3.01 ± 0.35 g/kg ethanol).
- 695 (b) Water intake in OFF (black dots - OFF) and "to be stimulated" (red dots - ON) groups across the
696 15 sessions of intermittent access to 20% ethanol. Both groups consumed the same amount of water
697 between session 1 (16.22 ± 0.92 g/kg water) and session 15 (22.39 ± 2.27 g/kg water).
- 698 (c) Number of rewards obtained after the loss of control of alcohol intake for non-stimulated (black
699 dots - OFF - control) and STN-stimulated rats (red dots - ON) during five 1h-sessions in the alone
700 condition (9.30 ± 1.10 vs. 6.13 ± 0.55 rewards, respectively) and five 1h-sessions with an alcohol-
701 naive stranger peer (10.58 ± 1.08 vs. 5.63 ± 0.62 rewards, respectively)
- 702 (d) Number of rewards obtained by the non-stimulated group and stimulated rats after reordering the
703 sessions in the proper sequence irrespective of the social context. ANOVA revealed a general
704 session and a group effect.
- 705 (e) Number of rewards obtained in control (black dots - OFF - control) and STN-stimulated (red dots
706 - ON) High Drinkers after a loss of control, in the alone condition (12.83 ± 1.59 vs. 9.48 ± 0.92
707 rewards respectively) and in a presence of an alcohol-naive stranger peer (12.69 ± 1.78 vs. $6.72 \pm$
708 1.15 rewards, respectively)
- 709 (f) Number of rewards obtained in control (black dots - OFF - control) and stimulated (red dots - ON)
710 Low Drinkers after a loss of control, in the alone condition (4.40 ± 0.42 vs. 3.74 ± 0.45 rewards,
711 respectively) and in a presence of an alcohol-naive stranger peer (7.64 ± 0.34 vs. 4.86 ± 0.48
712 rewards, respectively)

713

714 * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$

715 ### $p < 0.001$ between "alone" and "with peer" conditions in the non-stimulated (OFF - control) group

716